

Chapter 17

Expected Values, Covariance and Correlation

Lecture 17 | Devore 9e: 5.2

17.1 Expectation of a function of two variables

Theorem 17.1. For $h : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$\mathbb{E}[h(X, Y)] = \begin{cases} \sum_x \sum_y h(x, y) p(x, y), & \text{discrete,} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x, y) f(x, y) dx dy, & \text{continuous.} \end{cases}$$

Example 17.2 (Seats). Five seats in a row are numbered $1, \dots, 5$. You and one other person sit in two distinct seats chosen at random; X is your seat, Y theirs. So

$$p(x, y) = \begin{cases} \frac{1}{20}, & x, y \in \{1, \dots, 5\}, x \neq y, \\ 0, & \text{else,} \end{cases}$$

since there are $5 \times 4 = 20$ equally likely ordered pairs. Let $h(x, y) = |x - y| - 1$ be the number of seats *between* you. Then

$$\mathbb{E}[h(X, Y)] = \sum_x \sum_y h(x, y) p(x, y) = \frac{1}{20} \sum_{x \neq y} (|x - y| - 1) = \frac{20}{20} = 1.$$

The sum is easiest to organize by the gap $d = |x - y|$: there are $2(5 - d)$ ordered pairs with gap d , each contributing $d - 1$, so

$$\sum_{x \neq y} (|x - y| - 1) = \sum_{d=1}^4 2(5 - d)(d - 1) = 8 \cdot 0 + 6 \cdot 1 + 4 \cdot 2 + 2 \cdot 3 = 20.$$

Example 17.3 (Cost of a can of nuts). Almonds cost \$1.50/lb, cashews \$2.25/lb, peanuts \$0.75/lb. With X, Y the almond and cashew weights and $1 - X - Y$ the peanut weight, the cost of a can is

$$h(X, Y) = 1.5X + 2.25Y + 0.75(1 - X - Y) = 0.75 + 0.75X + 1.5Y.$$

With $f(x, y) = 24xy$ on the triangle,

$$\mathbb{E}[h(X, Y)] = \int_0^1 \int_0^{1-x} (0.75 + 0.75x + 1.5y) 24xy \, dy \, dx = \$1.65.$$

Remark 17.4 (Shortcut). Because h is *linear*, the double integral was avoidable. Using the marginal means $\mathbb{E}[X] = \mathbb{E}[Y] = \frac{2}{5}$ computed later this lecture,

$$\mathbb{E}[h(X, Y)] = 0.75 + 0.75 \mathbb{E}[X] + 1.5 \mathbb{E}[Y] = 0.75 + 0.30 + 0.60 = \$1.65,$$

and note that this needed *no* assumption of independence — expectation is linear regardless. Doing the integral once in full is still worth it, so that you trust the shortcut when you use it later.

17.2 Conditional expectation

Definition 17.5 (Conditional expectation).

$$\mathbb{E}[h(X) \mid Y = y] = \int_{-\infty}^{\infty} h(x) f_{X|Y}(x \mid y) \, dx.$$

Remark 17.6. This is a *function of y* , not a number: for each value of Y you get a different average. Writing $\mathbb{E}[h(X) \mid Y]$ (with a capital Y) therefore denotes a random variable.

Theorem 17.7 (Law of total expectation / tower property).

$$\mathbb{E}[\mathbb{E}[h(X) \mid Y]] = \mathbb{E}[h(X)].$$

Proof. The $f_Y(y)$ cancels, and then the order of integration is swapped:

$$\begin{aligned} \mathbb{E}[\mathbb{E}[h(X) \mid Y]] &= \int \left(\int h(x) \frac{f(x, y)}{f_Y(y)} \, dx \right) f_Y(y) \, dy = \iint h(x) f(x, y) \, dx \, dy \\ &= \int h(x) \left(\int f(x, y) \, dy \right) \, dx = \int h(x) f_X(x) \, dx. \end{aligned} \quad \square$$

This is the expectation version of the law of total probability: average within each case, then average the case averages.

Example 17.8 (Waiting for HH, and for HT). A fair coin is tossed until two consecutive tosses show HH. How many tosses does that take on average? What if we wait for HT instead?

Let N be the number of tosses until HH and $m = \mathbb{E}[N]$. Split on how the game starts, as in the first-step analysis of Lecture 6:

- first toss T (probability $\frac{1}{2}$): one toss is used up and we are back at the start, so $\mathbb{E}[N \mid T] = 1 + m$;
- first tosses HT (probability $\frac{1}{4}$): two tosses are used up and we are back at the start, so $\mathbb{E}[N \mid HT] = 2 + m$;

- first tosses HH (probability $\frac{1}{4}$): done, so $\mathbb{E}[N \mid \text{HH}] = 2$.

Averaging the case averages,

$$m = \frac{1}{2}(1 + m) + \frac{1}{4}(2 + m) + \frac{1}{4} \cdot 2 \implies 4m = 6 + 3m \implies m = 6.$$

For HT the wait is shorter. The first H takes 2 tosses on average (a geometric wait, Lecture 10), and after that the first T completes the pattern, another 2 on average; extra H's in between do no harm. So the expected wait is $2 + 2 = 4$.

Remark 17.9. The two patterns occur equally often. In 10 tosses, for instance, each of the 9 adjacent pairs is HH with probability $\frac{1}{4}$ and HT with probability $\frac{1}{4}$, so by the indicator method of Lecture 9 each pattern is expected $\frac{9}{4}$ times. Yet HH takes longer to show up. The difference is overlap: HH can overlap itself (HHH contains two), so its occurrences come in clumps, and the same long-run frequency spread over clumps leaves longer gaps between them.


17.3 Covariance

Variance measured how far X strays from its own mean. Covariance measures whether X and Y stray *together*.

Definition 17.10 (Covariance).

$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mu_X)(Y - \mu_Y)] = \begin{cases} \sum_x \sum_y (x - \mu_X)(y - \mu_Y)p(x, y), & \text{discrete,} \\ \iint (x - \mu_X)(y - \mu_Y)f(x, y) \, dx \, dy, & \text{continuous.} \end{cases}$$

Remark 17.11 (Intuition). The product $(X - \mu_X)(Y - \mu_Y)$ is positive when both variables are on the same side of their means and negative when they are on opposite sides. Averaging picks out which happens more often.



- Theorem 17.12** (Properties).
1. $\text{Cov}(X, X) = \text{Var}(X)$;
 2. $\text{Cov}(X, Y) = \mathbb{E}[XY] - \mu_X \mu_Y$ (the computational formula);
 3. $\text{Cov}(aX + b, cY + d) = ac \, \text{Cov}(X, Y)$;
 4. $X \perp\!\!\!\perp Y \implies \text{Cov}(X, Y) = 0$. **The converse is false.**

Property (2) gives the recipe: from the joint pdf get $\mathbb{E}[XY]$ by a double integral; from the marginals get μ_X and μ_Y ; subtract.

$$f(x, y) \longrightarrow \mathbb{E}[XY], \quad f(x, y) \longrightarrow f_X, f_Y \longrightarrow \mu_X, \mu_Y.$$

Example 17.13 (Can of nuts). $f(x, y) = 24xy$ on $\{x, y \geq 0, x + y \leq 1\}$.

$$\mathbb{E}[XY] = \int_0^1 \int_0^{1-x} xy \cdot 24xy \, dy \, dx = 8 \int_0^1 x^2(1-x)^3 \, dx = \frac{2}{15},$$

$$f_X(x) = 12x(1-x)^2, \quad f_Y(y) = 12y(1-y)^2 \implies \mu_X = \mu_Y = \int_0^1 12x^2(1-x)^2 \, dx = 12 \cdot \frac{1}{30} = \frac{2}{5}.$$

Therefore

$$\text{Cov}(X, Y) = \frac{2}{15} - \frac{2}{5} \cdot \frac{2}{5} = \frac{10}{75} - \frac{12}{75} = \boxed{-\frac{2}{75}}.$$

Negative, as it must be: the can weighs one pound, so more almonds means fewer cashews.

17.4 Correlation

Covariance has units — change pounds to grams and it changes by a factor of 454². The strength of a relationship should not depend on the choice of units.

Definition 17.14 (Correlation coefficient).

$$\text{Corr}(X, Y) = \rho_{X,Y} := \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Cov}(X, X)}\sqrt{\text{Cov}(Y, Y)}}.$$

Theorem 17.15 (Properties). 1. $\text{Corr}(aX + b, cY + d) = \text{Corr}(X, Y)$ when $a, c > 0$ — units do not matter;

2. $-1 \leq \rho_{X,Y} \leq 1$;

3. $\rho_{X,Y} = \pm 1$ if and only if $Y = aX + b$ exactly, for some $a \neq 0$;

4. $X \perp\!\!\!\perp Y \Rightarrow \rho = 0$ (“uncorrelated”), but not conversely.

Example 17.16 (Property 3 in action — the slope is invisible). Let $Y = 100X + 2$. Then $\text{Cov}(X, Y) = 100 \text{Var}(X)$ by property 3 of covariance, and $\sigma_Y = 100 \sigma_X$, so

$$\rho_{X,Y} = \frac{100 \text{Var}(X)}{\sigma_X \cdot 100 \sigma_X} = \frac{100 \sigma_X^2}{100 \sigma_X^2} = 1.$$

Now let $Y = -0.00001X + 10,000$. The slope is minuscule and the intercept enormous, yet by the same cancellation ($\sigma_Y = 0.00001 \sigma_X$)

$$\rho_{X,Y} = \frac{-0.00001 \text{Var}(X)}{\sigma_X \cdot 0.00001 \sigma_X} = -1.$$

Correlation records only *whether* the relationship is exactly linear and in which direction — never how steep it is. A nearly flat line and a nearly vertical one both give $|\rho| = 1$.

Remark 17.17. The last expression in the definition should look familiar: it is

$$\cos \theta = \frac{\langle u, v \rangle}{\|u\| \|v\|}$$

with covariance as the inner product. Correlation is the cosine of the angle between the centered variables — which is exactly why it lands in $[-1, 1]$ and why ± 1 means perfect alignment.

Example 17.18 (Uncorrelated but dependent). Let (X, Y) take the four values $(-4, 1)$, $(4, -1)$, $(2, 2)$, $(-2, -2)$, each with probability $\frac{1}{4}$. Then $\mu_X = \mu_Y = 0$ and

$$\mathbb{E}[XY] = \frac{1}{4}(-4 + (-4) + 4 + 4) = 0,$$

so $\text{Cov}(X, Y) = 0$ and $\rho = 0$. But X and Y are wildly dependent: knowing $X = -4$ tells you $Y = 1$ exactly.

Remark 17.19 (The one-sentence summary). ρ measures the strength of a **linear** relationship only. A perfect non-linear relationship can have $\rho = 0$. So “uncorrelated” is strictly weaker than “independent”, and $\rho = 0$ is never evidence of independence.

Remark 17.20 (Looking ahead). Writing $Y = aX + b + \varepsilon$ with ε a noise term and asking for the a, b that fit best is **linear regression**. The answer turns out to be $a = \rho \sigma_Y / \sigma_X$, which is the sense in which ρ “is” the slope.

17.4.1 How correlations constrain each other

The cosine picture has a practical consequence: correlations cannot be chosen freely. To use it, standardize: $\tilde{X} = (X - \mu_X) / \sigma_X$, and similarly $\tilde{Y}$ and $\tilde{Z}$. Then $\mathbb{E}[\tilde{X}^2] = 1$ and $\mathbb{E}[\tilde{X}\tilde{Y}] = \rho_{X,Y}$.

Example 17.21 (Three variables). Can three random variables have correlation -0.9 between every pair?

No. Expanding a square and using linearity,

$$0 \leq \mathbb{E}[(\tilde{X} + \tilde{Y} + \tilde{Z})^2] = 3 + 2(\rho_{X,Y} + \rho_{Y,Z} + \rho_{X,Z}),$$

so the three correlations add up to at least $-\frac{3}{2}$, and a common value must be at least $-\frac{1}{2}$. That bound is attained: take $\tilde{X}, \tilde{Y}$ with correlation $-\frac{1}{2}$ and set $\tilde{Z} = -(\tilde{X} + \tilde{Y})$. (Check that $\mathbb{E}[\tilde{Z}^2] = 1$ and $\mathbb{E}[\tilde{X}\tilde{Z}] = \mathbb{E}[\tilde{Y}\tilde{Z}] = -\frac{1}{2}$.)

Example 17.22 (Correlation is not transitive, but almost). If $\rho_{X,Y} = 0.9$ and $\rho_{Y,Z} = 0.9$, how small can $\rho_{X,Z}$ be?

In the cosine picture, X and Y meet at an angle θ with $\cos \theta = 0.9$, and so do Y and Z . The angle between X and Z is at most 2θ , so

$$\rho_{X,Z} \geq \cos 2\theta = 2 \cos^2 \theta - 1 = 2(0.81) - 1 = 0.62.$$

The same bound without geometry: write $\tilde{X} = 0.9\tilde{Y} + W$ and $\tilde{Z} = 0.9\tilde{Y} + V$. The leftovers W, V are uncorrelated with $\tilde{Y}$ and have $\mathbb{E}[W^2] = \mathbb{E}[V^2] = 1 - 0.81 = 0.19$, so $\rho_{X,Z} = \mathbb{E}[\tilde{X}\tilde{Z}] = 0.81 + \mathbb{E}[WV]$, and $\mathbb{E}[WV] \geq -0.19$ because $|\text{Corr}(W, V)| \leq 1$. With 0.5 in place of 0.9 the bound is $2(0.25) - 1 = -0.5$: two moderate positive correlations say nothing about the sign of the third.