

High-order fluctuations of temperature in hot QCD matter

Jinhui Chen,^{1,2,*} Wei-jie Fu,^{3,†} Shi Yin,^{4,‡} and Chunjian Zhang^{1,2,§}

¹Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), and
Institute of Modern Physics, Fudan University, Shanghai 200433, P.R. China

²Shanghai Research Center for Theoretical Nuclear Physics, NSFC and Fudan University, Shanghai 200438, P.R. China

³School of Physics, Dalian University of Technology, Dalian, 116024, P.R. China

⁴Institut für Theoretische Physik, Justus-Liebig-Universität Gießen, 35392 Gießen, Germany

A new thermodynamic state function is introduced to describe the thermodynamics relevant for the mean transverse momentum fluctuations of charged particles in heavy-ion collisions, which allows us to compute the temperature fluctuations of different orders in hot quantum chromodynamics (QCD) matter for the first time. Consequently, it is found that the temperature fluctuations are suppressed remarkably as the system transitions from the hadron resonance gas (HRG) to the quark-gluon plasma (QGP) with increasing temperature or baryon chemical potential, alongside a negative skewness. This is attributed to the general fact that the heat capacity of QCD matter increases significantly in QGP in comparison to that in HRG. These predictions provide a unique signature to discover the thermodynamical temperature fluctuations in upcoming heavy-ion collision experiments, which also paves a novel way to study QCD thermodynamics and QCD phase diagram through measurements of the mean transverse momentum fluctuations of charged particles.

Introduction. The temperatures and densities reached in relativistic heavy-ion collisions are expected to be similar to those thought to have prevailed in the very early universe, prior to the formation of protons and neutrons. The observation and study of QCD matter, the quark-gluon plasma (QGP) under these conditions has been one of the main science goals of high energy nuclear physics facilities [1–7]. Many questions have been addressed and developed in heavy-ion research programs [6]. The occurrence of a phase transition from the QGP to a hadron resonance gas (HRG) or the existence of a critical end point in the phase diagram of QCD matter is an exciting scientific endeavor [8–12], which may potentially be revealed by measurements of thermodynamic fluctuations [13–15], such as net-baryon or net-proton number fluctuations [16–24] and temperature fluctuations [25, 26]. Event-by-event (EbE) fluctuations in charged particle momentum distributions serve as probes of thermalization and the statistical nature of particle production in such collisions [27–31].

Recent advances in heavy-ion collision experiments now enable the isolation of the thermal fluctuations from confounding effects, such as the initial state geometry fluctuations [32–36], flow contributions, and other non-thermal sources, allowing the direct measurements of thermodynamical properties of hot QCD matter [37], e.g., the speed of sound [38–41]. High-order thermodynamical quantities, e.g., the temperature fluctuations, could be used to probe the QCD thermodynamics and phase transitions, since they are more sensitive to the critical fluctuations, i.e. the singular part of the thermodynamic potential during the phase transition in com-

parison to low-order thermodynamical quantities, such as the temperature itself [42–45]. This is similar to the case of ordinary net-proton number fluctuations. In principle, temperature fluctuations can be extracted from EbE mean transverse momentum fluctuations of final-state charged particles [25], isolated from other effects, e.g., the hydrodynamic effect [46]. To that end, one has to study the thermodynamical properties of temperature fluctuations from the theoretical side, in particular the smoking-gun signature of temperature fluctuations, in measurements of EbE mean transverse momentum fluctuations that have been extensively done across collision energies and systems in various heavy-ion facilities [35, 47–52].

To systematically investigate temperature fluctuations in hot QCD matter, we develop a theoretical framework which is general and applicable to temperature fluctuations of arbitrary order. As a specific application, this approach is applied to the QCD thermodynamics described by a 2+1 flavor low energy effective field theory (LEFT) [53], where quantum and thermal fluctuations are encoded self-consistently through the functional renormalization group (fRG). The fRG has proven to be a powerful nonperturbative theoretical method, and is well suited for studies of properties of the hot QCD matter including the QCD phase diagram, critical end point, and real-time dynamics, etc., [10, 54–59].

We first introduce a new thermodynamic state function to characterize the thermodynamics related to the mean transverse momentum fluctuations of charged particles, from which we derive analytic expressions for the temperature fluctuations to arbitrary order. Numerical results are obtained by applying this framework to a 2+1 flavor LEFT within the fRG approach. Our approach demonstrates that temperature fluctuations would be suppressed remarkably as the matter evolved from HRG to QGP with the increase in temperature or the baryon chemical potential.

A new thermodynamic state function. We begin with

* chenjinhui@fudan.edu.cn

† wjfu@dlut.edu.cn

‡ Shi.Yin@theo.physik.uni-giessen.de

§ chunjianzhang@fudan.edu.cn




FIG. 1. Variance of temperature fluctuations as a function of the temperature with several different values of baryon chemical potential.

a total derivative of the thermodynamic potential Ω

$$d\Omega = -SdT - pdV - N_B d\mu_B, \quad (1)$$

with the entropy S , temperature T , pressure p , volume V , baryon number N_B , and the baryon chemical potential μ_B . Although we explicitly show μ_B as a representative of the conserved charge, Eq. (1) is readily generalized to include additional chemical potentials when other conserved charges are presented. The thermodynamic potential Ω is a state function of T , V and μ_B . By implementing a Legendre transformation upon Ω w.r.t. the conjugate pair S and T , we introduce a new state function as

$$W = \Omega + TS. \quad (2)$$

One immediately recognizes that there is another relation for the state function W , that is,

$$W = U - \mu_B N_B, \quad (3)$$

resulting from general thermodynamical relations, where U denotes the energy. Inserting Eq. (2) into Eq. (1), one arrives at

$$dW = TdS - pdV - N_B d\mu_B, \quad (4)$$

indicating that W is a state function of S , V and μ_B .

In experimental measurements of mean transverse momentum fluctuations, finite acceptance cuts in the rapidity (y) and transverse momentum (p_T) range are applied, which signifies that the system volume in Eq. (4) is approximately constant. The chemical potential can also be regarded as a constant, which is determined by, e.g., the collision energy, centrality, etc. While this approximation holds for high-energy collisions, we note that μ_B may vary in low-energy regions, e.g., fixed target collisions at RHIC, due to global baryon number conservation effects [23, 60, 61]. For the present study, we neglect

these corrections and maintain the constant approximation.

Recent measurements of mean transverse momentum fluctuations at RHIC and the LHC were made at a fixed multiplicity of charged particles N_{ch} , as shown in [35, 52, 62, 63]. Since N_{ch} scales directly with the entropy of the system ($N_{\text{ch}} \sim S$). Consequently, the state function W in Eq. (4) becomes the appropriate thermodynamic potential to describe these experimental observables, since its natural variables correspond directly to the constrained quantities in the measures.

Temperature fluctuations derivations. Having established the relevance of the state function W in Eq. (4) for heavy-ion collisions, we now derive the temperature fluctuations, or equivalently, the mean transverse momentum fluctuations of charged particles, computed from the derivative of W w.r.t. S for different orders.

For a fixed volume V , we define the intensive quantities: the thermodynamic potential density $w = W/V$ and the entropy density $s = S/V$, one arrives at

$$w = -p + Ts, \quad (5)$$

where $\Omega = -pV$ is used and the entropy density can be obtained from $s = \frac{\partial p}{\partial T}$. The first-order derivative of w w.r.t. s produces the temperature

$$\frac{\partial w}{\partial s} = T. \quad (6)$$

Then, the n -th order fluctuation of temperature is obtained from the n -th order derivatives of w w.r.t. s , to wit,

$$\langle (\Delta T)^n \rangle = T^{4n-4} \frac{\partial^n w}{\partial s^n}, \quad (7)$$

with $\Delta T = T - \langle T \rangle$ and $n \geq 2$ ($n \in \mathbb{Z}$), where $\langle \dots \rangle$ denotes the ensemble average. It is convenient to adopt a dimensionless temperature fluctuation

$$c_n = \frac{\langle (\Delta T)^n \rangle}{T^n}. \quad (8)$$

Cumulants c_n can be expressed in terms of the temperature derivatives of pressure through fundamental thermodynamic relations. The first three nontrivial orders corresponding to the variance, skewness, and kurtosis of temperature fluctuations are given by,

$$\begin{aligned} c_2 &= T^2 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-1} \\ c_3 &= -T^5 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-3} \frac{\partial^3 p}{\partial T^3} \\ c_4 &= T^8 \left[3 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-5} \left(\frac{\partial^3 p}{\partial T^3} \right)^2 - \left(\frac{\partial^2 p}{\partial T^2} \right)^{-4} \frac{\partial^4 p}{\partial T^4} \right]. \end{aligned} \quad (9)$$




FIG. 2. Skewness (left panel) and kurtosis (right panel) of temperature fluctuations, i.e., c_3 and c_4 in Eq. (8), as functions of the temperature with several different values of baryon chemical potential.

This systematic approach can be extended to higher-order cumulants, e.g., the fifth and sixth hyper-order ones, which are presented in the supplement.

In relativistic heavy-ion collisions, the temperature fluctuations are measured through EbE mean transverse momentum fluctuations of charged particles, leveraging the approximate linear dependence between the mean transverse momentum and the system temperature, i.e., $\langle p_T \rangle = aT$ [37–39, 46]. As explicitly derived for a free relativistic gas in thermal equilibrium with Boltzmann statistics in the supplement, the coefficient is found to be $a = 2.356$. In realistic cases such as in experiments, the coefficient might deviate from this value in the ideal case. Therefore, it is useful to construct dimensionless ratios of different orders, thereby eliminating the dependence on the coefficient a . The numerical results for those ratios of temperature fluctuation cumulants are presented in the supplement.

Numerical results. We investigate QCD thermodynamics employing a 2+1 flavor LEFT within the fRG approach. As demonstrated in Ref. [53], this approach yields an equation of state (EoS) and baryon number fluctuations consistent with lattice QCD calculations. The setup of our LEFT is also recapitulated in the supplemental materials of this Letter.

To proceed, we systematically calculate the temperature derivatives of pressure:

$$\chi_n = T^{n-4} \frac{\partial^n p}{\partial T^n}, \quad (10)$$

which is dimensionless by normalization with appropriate powers of T . From the state function Ω in Eq. (1), we identify the first and second order derivatives, χ_1 and χ_2 , are just related to the entropy and heat capacity, respectively. Higher-order χ_n ($n \geq 2$) can be interpreted as entropy fluctuations of different orders. The numerical results of χ_n from the first to sixth orders calculated in the 2+1 flavor LEFT-fRG framework are presented in

the supplement. We found that the entropy fluctuations increase and oscillate near the chiral crossover, and the strength and amplitude of the oscillation increase with the order of fluctuations or the value of the baryon chemical potential.

The temperature fluctuations in Eq. (8) can be reformulated in terms of χ_n , defined in Eq. (10). For the lowest-order cumulants, we obtain

$$c_2 = \frac{1}{\chi_2}, \quad c_3 = -\frac{\chi_3}{\chi_2^3}, \quad c_4 = 3\frac{\chi_3^2}{\chi_2^5} - \frac{\chi_4}{\chi_2^4}. \quad (11)$$

The variance of temperature fluctuations, c_2 , is inversely proportional to the variance of entropy fluctuations, i.e., the heat capacity χ_2 , as demonstrated in Fig. 1. We observe that c_2 decreases with increasing temperature, reflecting the opposite trend of χ_2 as shown in the supplement. This behavior indicates a significant suppression of temperature fluctuations in QGP phase compared to those in HRG phase. The suppression is more remarkable for high-order temperature fluctuations, as is evident in Fig. 2 (note the logarithmic y -axis). A direct consequence of the suppression of temperature fluctuations at high temperature is that the distribution of temperature is wider in the region of lower temperature, which implies a negative skewness, as confirmed in the left panel of Fig. 2. While the kurtosis remains positive in most cases, its sign may reverse near the chiral crossover as it is sharpened continuously with the increase in the baryon chemical potential. The numerical results of the hyper-order c_5 and c_6 cumulants are presented in the supplement.

We further compute the temperature fluctuations on the chemical freeze-out curves, which allow us to investigate their dependence on collision energy. The relevant results are shown in Fig. 3. In order to study the errors arising from the determination of the freeze-out curves, we utilize three representative freeze-out curves, whose details are introduced in [22] and also used in [23]. The




FIG. 3. Variance (panel a), skewness (panel b), and kurtosis (panel c) of temperature fluctuations, i.e., c_2 , c_3 and c_4 , as functions of the collision energy, calculated on the three representative freeze-out curves [22, 23, 64].

freeze-out curve of Andronic *et al.* is obtained by parameterizing the freeze-out data from [64]. The other two freeze-out curves (STAR Fit I and II) are based on the freeze-out data from the STAR experiment [65], where all the freeze-out data from STAR are used for the former, and for the latter, some flawed data of collision energy are dropped from general considerations, see [22] for more details.

From Fig. 3 one observes that with the decrease of collision energy from high to low, all the temperature fluctuations investigated here, c_2 , c_3 and c_4 , exhibit mild dependence on collision energy firstly. When the collision energy is reduced to below, approximately 14.5 GeV, the magnitude of c_2 and c_3 increases significantly, while skewness c_3 remaining negative. The kurtosis c_4 in the regime of low collision energy is highly sensitive to the choice of freeze-out curves, resulting in uncertainties too large to draw a definitive conclusion.

Conclusions. We have studied temperature fluctuations in hot QCD matter through a newly introduced thermodynamic state function that directly connects to the mean transverse momentum fluctuations measured in heavy-ion collisions. Our approach yields analytic expressions for arbitrary-order temperature fluctuations, revealing their fundamental relationship with entropy, heat capacity, and high-order entropy fluctuations. Implementing this general formalism in a 2+1 flavor LEFT within the fRG, we obtain numerical results that quantify the temperature fluctuations across different thermodynamic regimes for the first time.

As the system transitions from the HRG to QGP phase with the increasing temperature or baryon chemical potential, the heat capacity of QCD matter increases substantially. This implies that a tiny change of the temperature would cost a huge amount of energy in the regime of high temperature. Therefore, the temperature tends not to change in comparison to the case in the regime of low temperature. In other words, the temperature fluctuations would be suppressed remarkably as the matter evolves from the HRG phase to the QGP phase with the

increase in temperature or baryon chemical potential, as demonstrated in our calculations. The fact that temperature fluctuations at high temperature are smaller than those at low temperature leads to another direct consequence, i.e., a negative skewness of temperature fluctuations. Such a signature emerges because the increasingly narrow fluctuation distribution at high temperature creates an asymmetric probability density weighted toward lower temperatures.

Note that these findings are general and model independent. They arise from the fact that the heat capacity of the QCD matter increases significantly from the HRG phase to the QGP phase. In the meantime, they provide a unique signature for discovering the thermodynamical temperature fluctuations in upcoming heavy-ion collision experiments at e.g., RHIC-BES, FAIR-CBM, NICA, and HIAF, which also paves a novel way to study QCD thermodynamics and QCD phase diagram through measurements of the mean transverse momentum fluctuations of charged particles.

Acknowledgements. We thank Fei Gao, Xuguang Huang, Jan M. Pawłowski for discussions and comments. W.J. Fu and S. Yin also would like to thank the members of the fQCD collaboration [66], especially Rui Wen, for collaborations on related projects. J.H. Chen is supported by the National Key Research and Development Program of China under Contract No. 2022YFA1604900, by the National Natural Science Foundation of China under Contract No. 12025501. W.J. Fu is supported by the National Natural Science Foundation of China under Contract Nos. 12447102, 12175030. S. Yin is supported by the Alexander v. Humboldt Foundation. S. Yin acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the CRC-TR 211 “Strong-interaction matter under extreme conditions” – project number 315477589 – TRR 211. C. Zhang is supported by the National Key Research and Development Program of China under Contract No. 2024YFA1612600, National Natural Science Foundation of China under Contract No. 12147101, Shanghai Pujiang Talents Program under Contract No. 24PJ A009.

-
- [1] E. V. Shuryak, Quantum Chromodynamics and the Theory of Superdense Matter, *Phys. Rept.* **61**, 71 (1980).
 - [2] J. W. Harris and B. Muller, The Search for the quark - gluon plasma, *Ann. Rev. Nucl. Part. Sci.* **46**, 71 (1996), [arXiv:hep-ph/9602235](#).
 - [3] J. Adams *et al.* (STAR), Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration's critical assessment of the evidence from RHIC collisions, *Nucl. Phys. A* **757**, 102 (2005), [arXiv:nucl-ex/0501009](#).
 - [4] K. Adcox *et al.* (PHENIX), Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX collaboration, *Nucl. Phys. A* **757**, 184 (2005), [arXiv:nucl-ex/0410003](#).
 - [5] C. W. Fabjan *et al.* (ALICE), ALICE: Physics Performance Report, *J. Phys. G* **32**, 1295 (2006).
 - [6] W. Busza, K. Rajagopal, and W. van der Schee, Heavy Ion Collisions: The Big Picture, and the Big Questions, *Ann. Rev. Nucl. Part. Sci.* **68**, 339 (2018), [arXiv:1802.04801 \[hep-ph\]](#).
 - [7] M. I. Abdulhamid *et al.* (STAR), Observation of the electromagnetic field effect via charge-dependent directed flow in heavy-ion collisions at the Relativistic Heavy Ion Collider, *Phys. Rev. X* **14**, 011028 (2024), [arXiv:2304.03430 \[nucl-ex\]](#).
 - [8] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, Signatures of the tricritical point in QCD, *Phys. Rev. Lett.* **81**, 4816 (1998), [arXiv:hep-ph/9806219](#).
 - [9] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, Event-by-event fluctuations in heavy ion collisions and the QCD critical point, *Phys. Rev. D* **60**, 114028 (1999), [arXiv:hep-ph/9903292](#).
 - [10] W.-j. Fu, J. M. Pawłowski, and F. Rennecke, QCD phase structure at finite temperature and density, *Phys. Rev. D* **101**, 054032 (2020), [arXiv:1909.02991 \[hep-ph\]](#).
 - [11] F. Gao and J. M. Pawłowski, Chiral phase structure and critical end point in QCD, *Phys. Lett. B* **820**, 136584 (2021), [arXiv:2010.13705 \[hep-ph\]](#).
 - [12] P. J. Gunkel and C. S. Fischer, Locating the critical end-point of QCD: Mesonic backcoupling effects, *Phys. Rev. D* **104**, 054022 (2021), [arXiv:2106.08356 \[hep-ph\]](#).
 - [13] M. M. Aggarwal *et al.* (STAR), An Experimental Exploration of the QCD Phase Diagram: The Search for the Critical Point and the Onset of De-confinement, (2010), [arXiv:1007.2613 \[nucl-ex\]](#).
 - [14] A. Bzdak, S. Esumi, V. Koch, J. Liao, M. Stephanov, and N. Xu, Mapping the Phases of Quantum Chromodynamics with Beam Energy Scan, *Phys. Rept.* **853**, 1 (2020), [arXiv:1906.00936 \[nucl-th\]](#).
 - [15] J. Chen *et al.*, Properties of the QCD matter: review of selected results from the relativistic heavy ion collider beam energy scan (RHIC BES) program, *Nucl. Sci. Tech.* **35**, 214 (2024), [arXiv:2407.02935 \[nucl-ex\]](#).
 - [16] J. Adam *et al.* (STAR), Nonmonotonic Energy Dependence of Net-Proton Number Fluctuations, *Phys. Rev. Lett.* **126**, 092301 (2021), [arXiv:2001.02852 \[nucl-ex\]](#).
 - [17] M. S. Abdallah *et al.* (STAR), Measurements of Proton High Order Cumulants in $\sqrt{s_{NN}} = 3$ GeV Au+Au Collisions and Implications for the QCD Critical Point, *Phys. Rev. Lett.* **128**, 202303 (2022), [arXiv:2112.00240 \[nucl-ex\]](#).
 - [18] B. Aboona *et al.* (STAR), Beam Energy Dependence of Fifth and Sixth-Order Net-proton Number Fluctuations in Au+Au Collisions at RHIC, *Phys. Rev. Lett.* **130**, 082301 (2023), [arXiv:2207.09837 \[nucl-ex\]](#).
 - [19] M. Abdallah *et al.* (STAR), Higher-order cumulants and correlation functions of proton multiplicity distributions in sNN=3 GeV Au+Au collisions at the RHIC STAR experiment, *Phys. Rev. C* **107**, 024908 (2023), [arXiv:2209.11940 \[nucl-ex\]](#).
 - [20] B. E. Aboona *et al.* (STAR), Precision Measurement of Net-Proton-Number Fluctuations in Au+Au Collisions at RHIC, *Phys. Rev. Lett.* **135**, 142301 (2025), [arXiv:2504.00817 \[nucl-ex\]](#).
 - [21] W.-j. Fu, J. M. Pawłowski, F. Rennecke, and B.-J. Schaefer, Baryon number fluctuations at finite temperature and density, *Phys. Rev. D* **94**, 116020 (2016), [arXiv:1608.04302 \[hep-ph\]](#).
 - [22] W.-j. Fu, X. Luo, J. M. Pawłowski, F. Rennecke, R. Wen, and S. Yin, Hyper-order baryon number fluctuations at finite temperature and density, *Phys. Rev. D* **104**, 094047 (2021), [arXiv:2101.06035 \[hep-ph\]](#).
 - [23] W.-j. Fu, X. Luo, J. M. Pawłowski, F. Rennecke, and S. Yin, Ripples of the QCD critical point, *Phys. Rev. D* **111**, L031502 (2025), [arXiv:2308.15508 \[hep-ph\]](#).
 - [24] Y. Lu, F. Gao, Y.-x. Liu, and J. M. Pawłowski, Finite density signatures of confining and chiral dynamics in QCD thermodynamics and fluctuations of conserved charges, (2025), [arXiv:2504.05099 \[hep-ph\]](#).
 - [25] S. Gavin, Traces of thermalization from transverse momentum fluctuations in nuclear collisions, *Phys. Rev. Lett.* **92**, 162301 (2004), [arXiv:nucl-th/0308067](#).
 - [26] R.-X. Cao, S. Zhang, and Y.-G. Ma, Specific heat and its high-order moments in relativistic heavy-ion collisions from a multiphase transport model, *Phys. Rev. C* **106**, 014910 (2022), [arXiv:2112.10299 \[nucl-th\]](#).
 - [27] H. Heiselberg, Event-by-event physics in relativistic heavy ion collisions, *Phys. Rept.* **351**, 161 (2001), [arXiv:nucl-th/0003046](#).
 - [28] S. Jeon and V. Koch, Charged particle ratio fluctuation as a signal for QGP, *Phys. Rev. Lett.* **85**, 2076 (2000), [arXiv:hep-ph/0003168](#).
 - [29] S. A. Voloshin, V. Koch, and H. G. Ritter, Event-by-event fluctuations in collective quantities, *Phys. Rev. C* **60**, 024901 (1999), [arXiv:nucl-th/9903060](#).
 - [30] M. Asakawa, U. W. Heinz, and B. Muller, Fluctuation probes of quark deconfinement, *Phys. Rev. Lett.* **85**, 2072 (2000), [arXiv:hep-ph/0003169](#).
 - [31] L. Stodolsky, Temperature fluctuations in multiparticle production, *Phys. Rev. Lett.* **75**, 1044 (1995).
 - [32] F. G. Gardim, F. Grassi, M. Luzum, and J.-Y. Ollitrault, Mapping the hydrodynamic response to the initial geometry in heavy-ion collisions, *Phys. Rev. C* **85**, 024908 (2012), [arXiv:1111.6538 \[nucl-th\]](#).
 - [33] B. Schenke, P. Tribedy, and R. Venugopalan, Initial-state geometry and fluctuations in Au + Au, Cu + Au, and U + U collisions at energies available at the BNL Relativistic Heavy Ion Collider, *Phys. Rev. C* **89**, 064908 (2014), [arXiv:1403.2232 \[nucl-th\]](#).
 - [34] M. I. Abdulhamid *et al.* (STAR), Imaging shapes of atomic nuclei in high-energy nuclear collisions, *Nature* **635**, 67 (2024), [arXiv:2401.06625 \[nucl-ex\]](#).

- [35] G. Aad *et al.* (ATLAS), Disentangling Sources of Momentum Fluctuations in Xe+Xe and Pb+Pb Collisions with the ATLAS Detector, *Phys. Rev. Lett.* **133**, 252301 (2024), [arXiv:2407.06413 \[nucl-ex\]](#).
- [36] L. Zhang, J. Chen, and C. Zhang, Energy dependence of transverse momentum fluctuations in Au+Au collisions from a multiphase transport model, *Phys. Rev. C* **111**, 024911 (2025), [arXiv:2501.08209 \[nucl-th\]](#).
- [37] F. G. Gardim, G. Giacalone, M. Luzum, and J.-Y. Ollitrault, Thermodynamics of hot strong-interaction matter from ultrarelativistic nuclear collisions, *Nature Phys.* **16**, 615 (2020), [arXiv:1908.09728 \[nucl-th\]](#).
- [38] F. G. Gardim, G. Giacalone, and J.-Y. Ollitrault, The mean transverse momentum of ultracentral heavy-ion collisions: A new probe of hydrodynamics, *Phys. Lett. B* **809**, 135749 (2020), [arXiv:1909.11609 \[nucl-th\]](#).
- [39] F. G. Gardim, A. V. Giannini, and J.-Y. Ollitrault, Accessing the speed of sound in relativistic ultracentral nucleus-nucleus collisions using the mean transverse momentum, *Phys. Lett. B* **856**, 138937 (2024), [arXiv:2403.06052 \[nucl-th\]](#).
- [40] A. Hayrapetyan *et al.* (CMS), Extracting the speed of sound in quark-gluon plasma with ultrarelativistic lead-lead collisions at the LHC, *Rept. Prog. Phys.* **87**, 077801 (2024), [arXiv:2401.06896 \[nucl-ex\]](#).
- [41] Y.-S. Mu, J.-A. Sun, L. Yan, and X.-G. Huang, Extracting the Speed of Sound in Heavy-Ion Collisions: A Study of Quantum-Initiated Fluctuations and Thermalization, *Phys. Rev. Lett.* **135**, 162301 (2025), [arXiv:2501.02777 \[nucl-th\]](#).
- [42] R. Arnaldi *et al.* (NA60), Evidence for the production of thermal-like muon pairs with masses above 1-GeV/c² in 158-A-GeV Indium-Indium Collisions, *Eur. Phys. J. C* **59**, 607 (2009), [arXiv:0810.3204 \[nucl-ex\]](#).
- [43] J. Adamczewski-Musch *et al.* (HADES), Probing dense baryon-rich matter with virtual photons, *Nature Phys.* **15**, 1040 (2019).
- [44] J. Churchill, L. Du, C. Gale, G. Jackson, and S. Jeon, Virtual Photons Shed Light on the Early Temperature of Dense QCD Matter, *Phys. Rev. Lett.* **132**, 172301 (2024), [arXiv:2311.06951 \[nucl-th\]](#).
- [45] B. E. Aboona *et al.* (STAR), Temperature measurement of Quark-Gluon plasma at different stages, *Nature Commun.* **16**, 9098 (2025), [arXiv:2402.01998 \[nucl-ex\]](#).
- [46] G. Giacalone, F. G. Gardim, J. Noronha-Hostler, and J.-Y. Ollitrault, Skewness of mean transverse momentum fluctuations in heavy-ion collisions, *Phys. Rev. C* **103**, 024910 (2021), [arXiv:2004.09799 \[nucl-th\]](#).
- [47] H. Appelshäuser *et al.* (NA49), Event-by-event fluctuations of average transverse momentum in central Pb + Pb collisions at 158-GeV per nucleon, *Phys. Lett. B* **459**, 679 (1999), [arXiv:hep-ex/9904014](#).
- [48] D. Adamova *et al.* (CERES), Event by event fluctuations of the mean transverse momentum in 40, 80 and 158 A GeV / c Pb - Au collisions, *Nucl. Phys. A* **727**, 97 (2003), [arXiv:nucl-ex/0305002](#).
- [49] S. S. Adler *et al.* (PHENIX), Measurement of nonrandom event by event fluctuations of average transverse momentum in s(NN)^{1/2} = 200-GeV Au+Au and p+p collisions, *Phys. Rev. Lett.* **93**, 092301 (2004), [arXiv:nucl-ex/0310005](#).
- [50] J. Adams *et al.* (STAR), Incident energy dependence of pt correlations at RHIC, *Phys. Rev. C* **72**, 044902 (2005), [arXiv:nucl-ex/0504031](#).
- [51] T. Anticic *et al.* (NA49), Energy dependence of transverse momentum fluctuations in Pb+Pb collisions at the CERN Super Proton Synchrotron (SPS) at 20A to 158A GeV, *Phys. Rev. C* **79**, 044904 (2009), [arXiv:0810.5580 \[nucl-ex\]](#).
- [52] S. Acharya *et al.* (ALICE), Skewness and kurtosis of mean transverse momentum fluctuations at the LHC energies, *Phys. Lett. B* **850**, 138541 (2024), [arXiv:2308.16217 \[nucl-ex\]](#).
- [53] R. Wen, C. Huang, and W.-J. Fu, Baryon number fluctuations in the 2+1 flavor low energy effective model, *Phys. Rev. D* **99**, 094019 (2019), [arXiv:1809.04233 \[hep-ph\]](#).
- [54] J. Braun, W.-j. Fu, J. M. Pawłowski, F. Rennecke, D. Rosenblüh, and S. Yin, Chiral susceptibility in (2+1)-flavor QCD, *Phys. Rev. D* **102**, 056010 (2020), [arXiv:2003.13112 \[hep-ph\]](#).
- [55] J. Braun *et al.*, Soft modes in hot QCD matter, (2023), [arXiv:2310.19853 \[hep-ph\]](#).
- [56] Y.-y. Tan, Y.-r. Chen, W.-j. Fu, and W.-J. Li, Universality of pseudo-Goldstone damping near critical points, *Nature Commun.* **16**, 2916 (2025), [arXiv:2403.03503 \[hep-th\]](#).
- [57] W.-j. Fu, J. M. Pawłowski, R. D. Pisarski, F. Rennecke, R. Wen, and S. Yin, The QCD moat regime and its real-time properties, (2024), [arXiv:2412.15949 \[hep-ph\]](#).
- [58] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. M. Pawłowski, M. Tissier, and N. Wschebor, The nonperturbative functional renormalization group and its applications, *Phys. Rept.* **910**, 1 (2021), [arXiv:2006.04853 \[cond-mat.stat-mech\]](#).
- [59] W.-j. Fu, QCD at finite temperature and density within the fRG approach: an overview, *Commun. Theor. Phys.* **74**, 097304 (2022), [arXiv:2205.00468 \[hep-ph\]](#).
- [60] P. Braun-Munzinger, B. Friman, K. Redlich, A. Rustamov, and J. Stachel, Relativistic nuclear collisions: Establishing a non-critical baseline for fluctuation measurements, *Nucl. Phys. A* **1008**, 122141 (2021), [arXiv:2007.02463 \[nucl-th\]](#).
- [61] V. Vovchenko, V. Koch, and C. Shen, Proton number cumulants and correlation functions in Au-Au collisions at sNN=7.7–200 GeV from hydrodynamics, *Phys. Rev. C* **105**, 014904 (2022), [arXiv:2107.00163 \[hep-ph\]](#).
- [62] R. Manikandhan (STAR Collaboration), Dynamical transverse momentum fluctuations at high baryon density measured by the STAR Experiment, *Quark Matter* **2025**.
- [63] Y. Gao (STAR Collaboration), Collision energy dependence of mean transverse momentum fluctuations in Au+Au collisions at STAR, *Quark Matter* **2025**.
- [64] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, Decoding the phase structure of QCD via particle production at high energy, *Nature* **561**, 321 (2018), [arXiv:1710.09425 \[nucl-th\]](#).
- [65] L. Adamczyk *et al.* (STAR), Bulk Properties of the Medium Produced in Relativistic Heavy-Ion Collisions from the Beam Energy Scan Program, *Phys. Rev. C* **96**, 044904 (2017), [arXiv:1701.07065 \[nucl-ex\]](#).
- [66] fQCD collaboration, <https://fqcd-collaboration.github.io>.
- [67] P. M. Lo, B. Friman, O. Kaczmarek, K. Redlich, and C. Sasaki, Polyakov loop fluctuations in SU(3) lattice gauge theory and an effective gluon potential, *Phys. Rev. D* **88**, 074502 (2013), [arXiv:1307.5958 \[hep-lat\]](#).

Supplemental Materials

The supplemental materials provide some details of the 2+1 flavor low energy effective field theory within the functional renormalization group approach, [Section S.1](#), preliminary results of temperature fluctuations calculated in QCD within fRG, [Section S.2](#), hyper-order fluctuations of temperature, [Section S.3](#), ratios of temperature fluctuation cumulants, [Section S.4](#), relation between the temperature T and the mean transverse momentum $\langle p_T \rangle$, [Section S.5](#).

S.1. 2+1 flavor low energy effective field theory within the fRG

In this appendix, we recapitulate the setup of the 2+1 flavor low energy effective field theory (LEFT) within the functional renormalization group used in this work. More details about the 2+1 LEFT can be found in [\[53\]](#). The effective action reads

$$\Gamma_k[\Phi] = \int_x \left\{ \bar{q} [\gamma_\mu \partial_\mu - \gamma_0 (\mu + i g A_0)] q + h_k \bar{q} \Sigma_5 q + \text{tr}(\partial_\mu \Sigma \cdot \partial_\mu \Sigma^\dagger) + V_{\text{matt},k}(\phi) + V_{\text{glue}}(L, \bar{L}) \right\}, \quad (12)$$

with the shorthand notation $\int_x = \int_0^\beta dx_0 \int d^3x$ and $\beta = 1/T$, where T stands for the temperature. The subscript k in Γ_k indicates that an infrared (IR) cutoff is applied to the effective action, such that quantum and thermal fluctuations of momenta $p \lesssim k$ are suppressed. The full effective action is resolved as $k \rightarrow 0$, and thus k plays a role as the renormalization group (RG) scale. The field $\Phi = (q, \bar{q}, \phi)$ includes the three-flavor quark field $q = (q_u, q_d, q_s)^\top$ and the scalar and pseudoscalar meson fields $\phi = (\sigma, \pi)$. The mesons are in the adjoint representation of the $U(N_f = 3)$ group, which reads

$$\Sigma = T^a (\sigma^a + i \pi^a), \quad a = 0, 1, \dots, 8, \quad (13)$$

with

$$T^0 = \frac{1}{\sqrt{2N_f}} \mathbb{1}_{N_f \times N_f}, \quad (14)$$

and

$$T^a = \frac{\lambda^a}{2} \quad (a = 1, \dots, 8), \quad (15)$$

where λ^a are the Gell-Mann matrices. The quark and meson fields interact with each other through the Yukawa coupling h_k with

$$\Sigma_5 = T^a (\sigma^a + i \gamma_5 \pi^a). \quad (16)$$

The quark chemical potential μ in [\(12\)](#) is related to the baryon chemical potential μ_B with $\mu = \mu_B/3$, where other chemical potentials, e.g., the chemical potentials for the electric charge and strangeness, are assumed to be vanishing.

The mesonic potential in [\(12\)](#), i.e., the matter sector of the effective potential, reads

$$V_{\text{matt},k}(\phi) = \tilde{V}_k(\rho_1, \rho_2) - c_A \xi - c_l \sigma_l - c_s \sigma_s, \quad (17)$$

with

$$\rho_1 = \text{tr}(\Sigma \cdot \Sigma^\dagger), \quad (18)$$

$$\rho_2 = \text{tr} \left(\Sigma \cdot \Sigma^\dagger - \frac{1}{3} \rho_1 \mathbb{1}_{3 \times 3} \right)^2, \quad (19)$$

where ρ_1 and ρ_2 are invariant under the transformations $SU_V(3) \times SU_A(3) \times U_V(1) \times U_A(1)$ in the flavor space. Here σ_l and σ_s indicate the scalar mesons of light and strange quarks, respectively. The relevant strength constants c_l and c_s result in explicit breaking of the chiral symmetry, as well as the breaking of the flavor symmetry from the three-flavor case to that of 2+1 flavors. The $U_A(1)$ symmetry is broken by the Kobayashi-Maskawa-'t Hooft determinant, viz.,

$$\xi = \det(\Sigma) + \det(\Sigma^\dagger), \quad (20)$$




FIG. 4. Yukawa coupling, normalized to unity for $k = 0$, as a function of the RG scale k , computed from the first-principles QCD within the FRG in the vacuum [10].




FIG. 5. Constituent light quark mass (left panel) and its derivative to the temperature (right panel) as functions of the temperature with several values of baryon chemical potential.

arising from quantum fluctuations, whose strength is controlled by the constant c_A .

The glue dynamics is encoded in the glue potential V_{glue} in (12), also known as the Polyakov loop potential. The Polyakov loop is related to the temporal gluon background field A_0 , that reads

$$L(\mathbf{x}) = \frac{1}{N_c} \langle \text{Tr } \mathcal{P}(\mathbf{x}) \rangle, \quad \bar{L}(\mathbf{x}) = \frac{1}{N_c} \langle \text{Tr } \mathcal{P}^\dagger(\mathbf{x}) \rangle, \quad (21)$$

with

$$\mathcal{P}(\mathbf{x}) = \mathcal{P} \exp \left(ig \int_0^\beta d\tau A_0(\mathbf{x}, \tau) \right), \quad (22)$$

where $\mathcal{P}$ on the right side denotes the path ordering, and g is the strong coupling constant.

In this work, we employ the Haar glue potential [53, 67],

$$V_{\text{glue}}(L, \bar{L}) = T^4 \bar{V}_{\text{glue-Haar}}, \quad (23)$$

with

$$\bar{V}_{\text{glue-Haar}} = -\frac{\bar{a}(T)}{2} \bar{L}L + \bar{b}(T) \ln M_H(L, \bar{L}) + \frac{\bar{c}(T)}{2} (L^3 + \bar{L}^3) + \bar{d}(T)(\bar{L}L)^2, \quad (24)$$




FIG. 6. Pressure normalized by T^4 as a function of the temperature at baryon chemical potential $\mu_B = 0, 100, 200, 300, 400$ and 500 MeV. These results are obtained in the 2+1 flavor LEFT.




FIG. 7. Dimensionless entropy (left panel) and heat capacity (right panel) normalized by appropriate powers of T , i.e., χ_1 and χ_2 , as functions of the temperature with several values of baryon chemical potential.

where the Haar measure reads

$$M_H(L, \bar{L}) = 1 - 6\bar{L}L + 4(L^3 + \bar{L}^3) - 3(\bar{L}L)^2. \quad (25)$$

The temperature dependence of the coefficients $\bar{a}$, $\bar{c}$, $\bar{d}$ in (24) is parameterized as

$$x(T) = \frac{x_1 + x_2/(t+1) + x_3/(t+1)^2}{1 + x_4/(t+1) + x_5/(t+1)^2}, \quad x \in (\bar{a}, \bar{c}, \bar{d}), \quad (26)$$

and that of $\bar{b}$ as

$$\bar{b}(T) = \bar{b}_1(t+1)^{-\bar{b}_4} \left(1 - e^{\bar{b}_2/(t+1)^{\bar{b}_3}}\right). \quad (27)$$

Here t in (26) and (27) is the reduced temperature, that is

$$t = \alpha(T - T_c^{\text{glue}})/T_c^{\text{glue}}, \quad (28)$$

where the parameters $T_c^{\text{glue}} = 250$ MeV and $\alpha = 0.6$ are used throughout this work. The values of other parameters in (26) and (27) can be found in [53, 67].




FIG. 8. Skewness (left panel) and kurtosis (right panel) of the entropy fluctuations, i.e., χ_3 and χ_4 , as functions of the temperature with several values of baryon chemical potential.




FIG. 9. Fifth (left panel) and sixth (right panel) order fluctuations of the entropy, i.e., χ_5 and χ_6 , as functions of the temperature with several values of baryon chemical potential.

Moreover, using the same method in [23], we employ the dependence of the Yukawa coupling on the RG scale k calculated from the first-principles QCD [10], as an input for the LEFT. Then the Yukawa coupling in the LEFT now reads

$$h_k = h_0 \frac{h_k^{\text{QCD}}}{h_{k=0}^{\text{QCD}}}, \quad (29)$$

where h_k^{QCD} is computed by using the fRG approach to the first-principles QCD in the vacuum [10], as shown in Fig. 4. Here the parameter in the LEFT $h_0 = 12$ is determined by fitting the constituent light u and d quark mass $m_l = 311$ MeV. Furthermore, we use the same values of parameters in the matter sector in (17) as those in [53].

In the left panel of Fig. 5, we show the constituent masses for the u , d light quarks calculated in the 2+1 flavor LEFT, depicted as functions of the temperature at several values of μ_B . Their respective derivatives with respect to the temperature are shown in the right panel of Fig. 5, from which one can determine the pseudo-critical temperature for the chiral crossover through the location of the peak. Figure 6 displays the temperature dependence of pressure calculated in our 2+1 flavor LEFT-fRG framework for baryon chemical potential μ_B ranging from $\mu_B = 0$ to 500 MeV, from which one can compute the temperature derivatives of pressure. The relevant results, from the first to sixth order derivatives, are presented in Figs. 7 to 9, which stand for the entropy and its fluctuations of different orders.




FIG. 10. Comparison between the LEFT and QCD computations for the second- and third-order fluctuations of temperature, i.e., c_2 (left panel) and c_3 (right panel), as functions of the temperature with several values of baryon chemical potential. The LEFT and QCD results are denoted by the solid and dashed lines, respectively.




FIG. 11. Hyper-order fluctuations of temperature, c_5 (left panel) and c_6 (right panel), as functions of the temperature with several different values of baryon chemical potential. The inset on the right shows the plot by using the linear y -axis, where the zero-crossing for c_6 at high baryon chemical potential is clear.

S.2. Preliminary results of temperature fluctuations calculated in QCD within fRG

In addition, we also compare the temperature fluctuations obtained in the LEFT with the preliminary results obtained in QCD within the fRG, whose setup is detailed in [10]. The relevant results are presented in Fig. 10. One can see that although there are some differences, the variance and skewness of temperature fluctuations calculated from the LEFT and QCD are consistent with each other qualitatively. Note that the main conclusions obtained from the calculations in LEFT are corroborated by the QCD results, e.g., the skewness of temperature fluctuations is negative and the variance decreases significantly with the increase of temperature.




FIG. 12. Ratios between high-order temperature fluctuations and the variance, $R_{32} = c_3/c_2^2$, $R_{42} = c_4/c_2^3$, $R_{52} = c_5/c_2^4$, $R_{62} = c_6/c_2^5$, as functions of the temperature with several different values of baryon chemical potential.

S.3. Hyper-order fluctuations of temperature

The fifth- and sixth-order fluctuations of temperature are given by

$$\begin{aligned}
 c_5 &= T^{11} \left[-15 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-7} \left(\frac{\partial^3 p}{\partial T^3} \right)^3 - \left(\frac{\partial^2 p}{\partial T^2} \right)^{-5} \frac{\partial^5 p}{\partial T^5} + 10 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-6} \frac{\partial^3 p}{\partial T^3} \frac{\partial^4 p}{\partial T^4} \right] \\
 c_6 &= T^{14} \left[105 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-9} \left(\frac{\partial^3 p}{\partial T^3} \right)^4 - 105 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-8} \left(\frac{\partial^3 p}{\partial T^3} \right)^2 \frac{\partial^4 p}{\partial T^4} + 10 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-7} \left(\frac{\partial^4 p}{\partial T^4} \right)^2 \right. \\
 &\quad \left. + 15 \left(\frac{\partial^2 p}{\partial T^2} \right)^{-7} \frac{\partial^3 p}{\partial T^3} \frac{\partial^5 p}{\partial T^5} - \left(\frac{\partial^2 p}{\partial T^2} \right)^{-6} \frac{\partial^6 p}{\partial T^6} \right].
 \end{aligned} \tag{30}$$

The relevant numerical results are shown in Fig. 11.

S.4. Ratios of temperature fluctuation cumulants

In relativistic heavy-ion collisions, the event-averaged mean transverse momentum $\langle p_T \rangle$ of charged particles exhibits an approximate linear dependence on the system temperature, $\langle p_T \rangle = aT$. The parameter a represents the proportionality coefficient. In order to eliminate the influence from this coefficient that is not determined quite well, we instead analyze dimensionless ratios of temperature fluctuation cumulants:

$$R_{32} = \frac{c_3}{c_2^2}, \quad R_{42} = \frac{c_4}{c_2^3}, \quad R_{52} = \frac{c_5}{c_2^4}, \quad R_{62} = \frac{c_6}{c_2^5}. \tag{31}$$

where the powers of the variance in the denominators are chosen to balance the powers of T . The relevant ratios, presented in Fig. 12, reveal that the cumulant ratios develop an increasingly rich nonmonotonic structure while exhibiting systematically reduced amplitudes with the increase of μ_B , reflecting competing effects where enhanced critical fluctuations near the sharpened phase boundary emerge concurrently with the overall suppression of the magnitude of temperature fluctuation.

S.5. Relation between the temperature T and the mean transverse momentum $\langle p_T \rangle$

Considering a free relativistic gas in thermal equilibrium at temperature T with Boltzmann statistics, where the mass of particle, e.g., the pion $m_\pi \sim 150$ MeV, is negligible in comparison to the typical value of the transverse momentum $p_T \sim 1$ GeV, the particle number reads

$$N = (2s + 1)V \int \frac{d^3p}{(2\pi)^3} e^{-p/T}, \quad (32)$$

where s denotes the spin of the particle and V the volume of system. One readily finds for the mean transverse momentum of particles

$$\langle p_T \rangle = \frac{1}{N} (2s + 1)V \int \frac{d^3p}{(2\pi)^3} p_T e^{-p/T}, \quad (33)$$

where the momentum and the transverse momentum is related through

$$p = \sqrt{p_T^2 + p_z^2}, \quad (34)$$

with the longitudinal momentum p_z . Combining Eqs. (32) and (33) we have

$$\langle p_T \rangle = \frac{\int \frac{d^3p}{(2\pi)^3} p_T e^{-p/T}}{\int \frac{d^3p}{(2\pi)^3} e^{-p/T}}. \quad (35)$$

From the dimensional analysis in Eqs. (35), one immediately arrives at

$$\langle p_T \rangle = aT, \quad (36)$$

that is, $\langle p_T \rangle$ is linearly proportional to the temperature, with the proportionality coefficient a . Note that this is different from the non-relativistic case, where it is easy to find that $\langle p_T \rangle \sim T^{1/2}$. In order to determine the coefficient a in Eq. (36), one has to compute the expression in Eq. (35). The integral in the numerator in Eq. (35) cannot be computed analytically, but one can expand the momentum in the exponent in powers of p_z/p_T , i.e.,

$$\begin{aligned} p &= p_T \left(1 + \frac{p_z^2}{p_T^2} \right)^{1/2} \\ &= p_T + \frac{1}{2} \frac{p_z^2}{p_T} + \dots \end{aligned} \quad (37)$$

Expanding Eq. (37) to the second order, and substituting into Eq. (35), one finds

$$\langle p_T \rangle \approx \frac{15}{16\sqrt{2}} \pi T \approx 2.08 T. \quad (38)$$

Note that this value is close to the exact value

$$a = 2.356, \quad (39)$$

obtained by directly computing Eq. (35) numerically.