
A

PLAT

Bulk logFC = -1.216; q = 8.3×10^{-6}

Malignant logFC = -6.314; q = 0.039





B


S100A4

Bulk logFC = -0.306; q = 0.042


Malignant logFC = -5.056; q = 0.039





C
ABCB4
ADGRL4
ANXA1
ANXA2
+ APOD
ART4
+ ATF3
BAG3
BNC2
CAV1
CCL21
CD63
CDK5R1
CDKN2A
CETP
CHI3L1
+ COCH
CR2
CRISPLD2
CXCL8
CYP7B1
CYR61
DCN
DNASE1L3
F2R
F5
FCER2
FCMR
FMO1
FOXF1
FRMD4B
GP1BA
GPR18
HTR2B
IFI27
IFITM2
IFITM3
B
CAFs
CD4T
CD8T
DCs
Mac
NK
TEC
PlasmaB
Author-defined population
D
IGFBP3
IGHV5-78
+ ITGB1
KLF6
LARP6
MCTP1
MMP1
MMP13
MMP3
MMP7
MYL9
NTS
P2RX5
+ PDCD4-AS1
+ PDLIM4
+ PLA2G16
PLAG1
PLPP3
PLSCR1
+ PSD4
+ ROR2
S100A4
S100A8
S100A9
SELE
SERPINA3
SLC16A10
SPARCL1
SPINK2
STOM
STON1
TAC1
TCL1A
TNFRSF12A
TNFSF4
TPM1
TXK
B
CAFs
CD4T
CD8T
DCs
Mac
NK
TEC
PlasmaB
Author-defined population



E
PLAT
S100A4
APOD
ATF3
COCH
ITGB1
PDCD4-AS1
PDLIM4
PLA2G16
PSD4
ROR2
-5
0
5
Five-patient malignant log2-CPM Δ
1 5 6 7 8
F
2.5
0.0
-2.5
-5.0
Malignant log2 fold change
-2
-1
0
1
Bulk log2 fold change
Other evaluable
Supported concordant
G
-3
-2
-1
0
1
2
Cell-threshold subset mean Δ
30 40 50
H
-2.5
0.0
Leave-one-patient-out mean Δ
1 5 6 7 8

I Additional ESTIMATE outputs






J Integrated candidate classification



K Nonmalignant evaluability



A: malignant-compatible; B: context-compatible; C: unresolved; D: malignant non-evaluable.

Blue: supported; white: evaluable, not supported; hatched: not evaluable.

L **Full nonmalignant support coverage, including ineligible Mast cells**

