



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Economics Letters 82 (2004) 341–347

**economics  
letters**

[www.elsevier.com/locate/econbase](http://www.elsevier.com/locate/econbase)

# The information matrix test with bootstrap-based covariance matrix estimation

Geert Dhaene\*, Dirk Hoorelbeke

*Department of Economics, Katholieke Universiteit Leuven, Naamsestraat 69, 3000 Leuven, Belgium*

Received 24 July 2003; accepted 2 September 2003

---

## Abstract

We propose an information matrix (IM) test in which the covariance matrix of the vector of indicators is estimated using the parametric bootstrap. Monte Carlo results and theoretical arguments show that its small sample performance is comparable with that of the efficient score form.

© 2003 Elsevier B.V. All rights reserved.

*Keywords:* Information matrix test; Parametric bootstrap

*JEL classification:* C12; C15

---

## 1. Introduction

While the information matrix (IM) test, introduced by [White \(1982\)](#), is well known as a general test for misspecification of a parametric likelihood function, its use in applied econometric research is still limited. A major drawback of the test is that the asymptotic  $\chi^2$  distribution is a very poor approximation to the finite sample distribution of the test statistic, as evidenced by the Monte Carlo experiments reported in [Taylor \(1987\)](#), [Orme \(1990\)](#), [Chesher and Spady \(1991\)](#), [Davidson and MacKinnon \(1992, 1998\)](#), and [Horowitz \(1994\)](#). Several approaches have been proposed to overcome this problem. [Chesher and Spady \(1991\)](#) derive, for specific models, critical values for the IM test statistic that are based on a higher order Edgeworth expansion. [Davidson and MacKinnon \(1992\)](#) propose a variant of the IM test based on double length artificial regressions. [Horowitz \(1994\)](#) advocates the use of bootstrap-based critical values. Despite these efforts, computing the correct critical value of an IM test statistic for an arbitrary model is still not particularly easy.

---

\* Corresponding author. Tel.: +32-16-32-67-98; fax: +32-16-32-67-96.

E-mail address: [geert.dhaene@econ.kuleuven.ac.be](mailto:geert.dhaene@econ.kuleuven.ac.be) (G. Dhaene).

All existing versions of the IM test rely on some estimate of the asymptotic covariance matrix of the vector of indicators. In Section 2, we propose a new form of the IM test, making use of the parametric bootstrap to estimate the finite sample covariance matrix. Although this approach is based on simulations, the computational demands are very modest and there are no analytical requirements at all. The proposed method eliminates the approximation errors that result in other IM tests from the use of asymptotic covariance matrix formulae and from approximating expectations by sample averages. Section 3 reports Monte Carlo results for the linear model and the probit model, showing that the new test performs well compared to existing IM tests. Section 4 concludes.

## 2. Estimating the covariance matrix of the IM test

Consider a parametric model with log-density  $F(y; \theta)$ , where  $\theta$  is a  $p \times 1$  vector of parameters. Let  $F_i = [\partial F / \partial \theta_i]_{\theta = \theta_0}$  and  $F_{ij} = [\partial^2 F / \partial \theta_i \partial \theta_j]_{\theta = \theta_0}$ , where  $\theta_0$  maximizes  $E[F(y; \theta)]$  with respect to  $\theta$  and  $E[\cdot]$  denotes expectation. The null hypothesis underlying the class of IM tests is

$$H_0 : E[F_i F_j + F_{ij}] = 0 \quad (1 \leq i, j \leq p). \quad (1)$$

Given a sample of observations  $y_1, \dots, y_n$ , define the indicators

$$\hat{D}_{ij} = n^{-1/2} \sum_{t=1}^n (\hat{F}_i \hat{F}_j + \hat{F}_{ij}), \quad (2)$$

where a hat indicates evaluation at  $y_t$  and  $\hat{\theta}$ , the MLE of  $\theta_0$ . Most existing IM tests are based on an asymptotically  $\chi_q^2$  distributed statistic of the form

$$\omega = \hat{D}' \hat{V}^{-1} \hat{D}, \quad (3)$$

where  $\hat{D}$  is a  $q \times 1$  vector of appropriately selected indicators  $\hat{D}_{ij}$  and  $\hat{V}$  is a consistent estimate of its covariance matrix under  $H_0$  (alternatively, under the stronger assumption that  $F(y; \theta)$  is the correct log-density).

Orme (1990) reviews many alternative choices of  $\hat{V}$ , including those leading up to White's (1982) form, the Chesher (1983) and Lancaster (1984) form, and the efficient score form of the IM test. All these choices of  $\hat{V}$  are based on equivalent analytical formulae for the asymptotic covariance matrix of  $\hat{D}$ , the differences arising essentially from replacing expectations with sample averages in different parts of those formulae. Available Monte Carlo evidence, in settings where  $F(y; \theta)$  is the correct log-density, shows that the ensuing IM test statistics have finite sample distributions that are poorly approximated by the  $\chi_q^2$  distribution. Four sources of possible error may be involved in the approximation:

- (i) the finite sample distribution of  $\hat{D}$  may be non-normal;
- (ii) the finite sample covariance matrix of  $\hat{D}$ , say  $V_n$ , may differ from its asymptotic covariance matrix,  $V_\infty$ ;
- (iii)  $\hat{\theta}$  is used in place of  $\theta_0$  in formulae for  $V_\infty$ ;
- (iv) sample averages replace expectations in parts of formulae for  $V_\infty$ .

In most circumstances, the error sources (i)–(iii) effectively apply to the IM tests discussed so far. The efficient score form is the only one not vulnerable to (iv).

Rather than relying on an asymptotic covariance matrix formula, one may choose  $\hat{V}$  to estimate the *exact* finite sample covariance matrix of  $\hat{D}$ , denoted  $V_n(\theta_0)$ , since it typically depends on  $\theta_0$ . Although it is simple enough to write  $V_n(\theta_0)$  as an integral, working out the integral analytically is bound to be impossible in all but the simplest models. A simple and feasible alternative is to estimate  $V_n(\theta_0)$  by the parametric bootstrap, which involves the following steps, after computing the MLE  $\hat{\theta}$ :

1. for  $b = 1, \dots, B$ : generate an i.i.d. sample  $y_{1b}, \dots, y_{nb}$  from the density  $\exp F(\cdot; \theta)$ , and compute the corresponding MLE  $\hat{\theta}_b$  and the vector of selected indicators  $\hat{D}_b$ ;
2. compute  $\hat{V}_B = (B - 1)^{-1} \sum_{b=1}^B (\hat{D}_b - \bar{D})(\hat{D}_b - \bar{D})'$ , where  $\bar{D} = B^{-1} \sum_{b=1}^B \hat{D}_b$ .

It is obvious that  $E[\hat{V}_B | \hat{\theta}] = V_n(\hat{\theta})$  and that for fixed  $n$ ,  $\hat{V}_B \xrightarrow{\text{a.s.}} V_n(\hat{\theta})$  as  $B \rightarrow \infty$ . Thus, through the choice of the number of bootstrap replications  $B$ ,  $\hat{V}_B$  approximates  $V_n(\hat{\theta})$  to any desired accuracy. Taking  $\hat{V} = \hat{V}_B$  in Eq. (3) yields the IM test statistic

$$\omega_B = \hat{D}' \hat{V}_B^{-1} \hat{D}. \quad (4)$$

Under the assumption that  $F(y; \theta)$  is the correct log-density,  $\omega_B$  has the following limit behavior. As  $n \rightarrow \infty$  and  $B \rightarrow \infty$ ,  $\omega_B \xrightarrow{d} \chi_q^2$ , where  $\xrightarrow{d}$  denotes convergence in distribution. For fixed  $B \geq q + 1$  and  $n \rightarrow \infty$ ,

$$\omega_B \xrightarrow{d} T_{q, B-1}^2 \quad (5)$$

(Hotelling's  $T^2$ ), since  $\hat{D} \xrightarrow{d} N(0, V_\infty(\theta_0))$ ,  $(B - 1)\hat{V}_B \xrightarrow{d} W(V_\infty(\theta_0), B - 1)$  (central Wishart), and  $\hat{D}$  and  $\hat{V}_B$  are asymptotically independent. Note that Eq. (5) may also be stated as  $((B - q)/(B - 1)q)\omega_B \xrightarrow{d} F_{q, B-q}$ . Using the IM test statistic  $\omega_B$  and critical values from the  $T_{q, B-1}^2$  distribution, (ii) is eliminated as a source of approximation error. This IM test is closest in spirit to the efficient score form of the IM test as it replaces  $V_\infty$  with  $V_n$  in the latter. With finite  $B$ ,  $V_n$  is estimated with some noise, but the  $T^2$  critical values correct for this. Since the test based on  $\omega_B$  has less sources of error, we expect it in general to exhibit smaller errors in rejection probability (ERP)<sup>1</sup> than the IM tests based on the  $\chi_q^2$  approximation to  $\omega$  already discussed. Furthermore, for large enough  $B$ , we expect the power of the IM test based on  $\omega_B$  to be no less than the power of the efficient score test, which, given the available evidence, appears to be the most powerful of existing IM tests (Davidson and MacKinnon, 1998). The efficient score test, however, requires calculating certain expectations analytically and hence is only available for models where this has proven feasible. In other models, Orme (1995) proposes to compute the expectations by means of simulation. The difference between his approach and ours lies in the fact that he uses simulations to estimate  $V_\infty$ , whereas we use them to estimate the exact infinite sample covariance matrix,  $V_n$ .

<sup>1</sup> The ERP of a test is the actual minus the nominal (i.e. chosen) probability of rejecting the null hypothesis when it is true. The ERP often depends on the parameter that indexes the distributions constituting the null hypothesis.

We have two final remarks. First, the only computational requirement to obtain  $\omega_B$  is that observations can be generated from the density  $\exp F$  and that the vector of indicators can be computed. The latter can often be extracted without effort from econometric software packages, either as the difference between two information matrix estimates, or as the difference between the inverses of two estimates of the covariance matrix of the MLE. Thus, no analytical work is required before the test can be applied. Second, although Monte Carlo results show that the ERP of the newly proposed test is moderate, it may be advisable in situations with few observations to use bootstrap-based critical values, as suggested by Horowitz (1994) in the context of the IM test. Although this requires a nested bootstrap—the inner bootstrap serves to calculate  $\hat{V}_B$ —this is nowadays quite feasible: 50 inner and 99 outer bootstrap replications will often suffice.

### 3. ERP and power: Monte Carlo evidence

Here we report comparative Monte Carlo results on the finite sample properties of the statistic  $\omega_B$ , White's (1982) IM test statistic  $\omega_W$ , Chesher (1983) and Lancaster's (1984) IM test statistic  $\omega_{CL}$ , Orme's (1990)  $\omega_3$ , here  $\omega_O$ , and the efficient score IM test statistic  $\omega_{EFF}$ . Without ambiguity, we refer to  $\omega_B$ ,  $\omega_W$ , etc. as IM test statistics and IM tests, with the understanding that  $\omega_B$  is used with  $T_{q,B-1}^2$  critical values, and the other statistics with  $\chi_q^2$  critical values. We study the ERP under the null of correct specification as well as the power against a heteroskedastic alternative, both in the linear model and in the probit model. Throughout, the IM tests are based on the maximum number of linearly independent indicators. Published Monte Carlo results show that  $\omega_W$  and  $\omega_{CL}$ , and to a lesser extent  $\omega_O$  and  $\omega_{EFF}$ , suffer from substantial ERPs in these models and that, after bootstrap-correcting the critical values,  $\omega_{EFF}$  is the most powerful. Our results show that  $\omega_B$  offers an improvement on  $\omega_{EFF}$  in terms of smaller ERP, and that it is very close to  $\omega_{EFF}$  in terms of power. These findings support the intuitive arguments advanced in Section 2.

#### 3.1. The normal linear regression model

The conditional density in this model is  $\phi((y_t - x_t' \beta)/\sigma)$ , with  $\phi$  the standard normal density,  $x_t$  a  $k \times 1$  vector of regressors, and parameters  $\beta$  ( $k \times 1$ ) and  $\sigma > 0$ . Hall (1987) shows that the IM test is a combined test against heteroskedasticity, conditional skewness, and non-normal kurtosis.

We use the following Monte Carlo design. The regressor matrix  $X$ , which is kept fixed across Monte Carlo replications, consists of a vector of ones and independent drawings from  $N(0, 1)$  elsewhere. We note that all IM test statistics are invariant under transformations  $X \rightarrow XA$  with  $A$  non-singular. Therefore, the results extend to any case where the  $k - 1$  non-constant regressors are generated from a non-singular  $(k - 1)$ -variate normal distribution. Moreover, since all statistics considered are pivotal under the null, the results concerning ERP are independent of  $\beta$  and  $\sigma$ . We implement a full factorial design with  $k = 2, 3, 4, 5$  and  $n = 50, 100, 250, 500, 1000$ . Throughout,  $B = 50$ . All results are based on 10,000 Monte Carlo replications.

The ERP is displayed using  $p$ -value plots (Davidson and MacKinnon, 1998). A  $p$ -value plot gives the (estimated) actual rejection probability (RP) of a test as a function of the nominal RP. On the  $45^\circ$  line actual and nominal RP agree, so one would hope to see a  $p$ -value plot close to the  $45^\circ$  line.




Fig. 1. Linear model:  $p$ -value plots and power curves for  $n=100$ ,  $k=2$  (left panels) and  $n=100$ ,  $k=4$  (right panels).

The  $p$ -value plots for  $n=100$  and  $k=2, 4$  are given in Fig. 1, panels (a) and (b). The ERP is largest for  $\omega_W$  and  $\omega_{CL}$ . This is in fact true for all design points.<sup>2</sup> The performance of  $\omega_B$  is in general comparable with that of  $\omega_{EFF}$ , although overall  $\omega_B$  has the smallest ERP of all tests considered. The behavior of  $\omega_O$  is better than that of  $\omega_W$  and  $\omega_{CL}$ , although its convergence to the  $45^\circ$  line as  $n$  grows is remarkably slow. The IM test is a good example of how bad a first-order asymptotic approximation can work: even for  $n=1000$  the ERP of  $\omega_W$ ,  $\omega_{CL}$ ,  $\omega_O$  and even  $\omega_{EFF}$  is too large to use these tests in practice. The  $T_{q,B-1}^2$  approximation to the distribution of  $\omega_B$ , on the other hand, works fine for larger sample sizes. Since the statistics are pivotal, all the tests considered can be turned into exact tests using bootstrap-based critical values.

We investigated the power of the IM tests against a heteroskedastic alternative with density  $\phi((y_t - x_t' \beta) / |x_t' \beta|^{1/2})$ . In order to correct for ERP, we plot power as a function of actual RP under

<sup>2</sup> Detailed results are available from the authors.

the null (Davidson and MacKinnon, 1998). Fig. 1, panels (c) and (d) give the power curves for  $n=100$  and  $k=2, 4$ . From the results at these and other design points, the following patterns emerge. The tests  $\omega_W$  and  $\omega_{CL}$  have similar power, and are in most cases dominated by  $\omega_O$ . The tests  $\omega_{EFF}$  and  $\omega_B$  have very similar power, with  $\omega_{EFF}$  being slightly better. These tests always outperform the others. We note that taking  $B$  larger would increase the power of  $\omega_B$ , but a relatively small  $B$  already yields a powerful test.

### 3.2. The probit model

In this model  $y_i$  is binary with conditional mean  $\Pr[y_i=1]=\Phi(x_i'\beta)$ , where  $\Phi$  is the standard normal distribution function. Orme (1988) gives the efficient score form of the IM test. In the Monte Carlo experiment, we set  $\beta=(0.5, 1, \dots, 1)'$  and  $\sigma=1$ , and choose  $X$ ,  $n$ ,  $k$ ,  $B$ , and the number of Monte Carlo




Fig. 2. Probit model:  $p$ -value plots and power curves for  $n=100$ ,  $k=2$  (left panels) and  $n=100$ ,  $k=4$  (right panels).

replications as in the linear model. None of the IM statistics is pivotal, so the results are specific to the choice of  $\beta$ . A further consequence is that the bootstrap-based critical values are not exact anymore, although their use ensures that the ERP vanishes quickly as the sample size gets larger (Horowitz, 1994).

In panels (a) and (b) of Fig. 2 the  $p$ -value plots for  $n=100$  and  $k=2, 4$  are given. More or less the same patterns are observed as in the linear model:  $\omega_W$  and  $\omega_{CL}$  severely over-reject;  $\omega_O$ ,  $\omega_{EFF}$ , and  $\omega_B$  have much smaller ERP, although for small  $n$  and large  $k$  it may still be substantial; and most of the time  $\omega_B$  has the smallest ERP.

We studied the power of the IM tests against the heteroskedastic probit  $\Pr[y_t=1] = \Phi(x_t'\beta/|x_t'\beta|^{1/2})$ . Following Horowitz (1994), we plot power as a function of actual RP under the pseudo-true null. Panels (c) and (d) of Fig. 2 display the power curves for  $n=100$  and  $k=2, 4$ . The powers of  $\omega_B$  and  $\omega_{EFF}$  are in all design points extremely close to each other, and well above the power of  $\omega_W$ ,  $\omega_{CL}$  and  $\omega_O$ .

#### 4. Conclusion

We have introduced an IM test which uses the parametric bootstrap to estimate the covariance matrix of the indicator vector. The test is easy to compute using standard econometric software and requires no analytical derivations. Its analytical simplicity comes at a small cost, namely it requires a limited number of simulations. When one wants to use bootstrap-based critical values, a nested bootstrap becomes necessary. In the models analyzed here, its performance was found to be comparable to that of the efficient score form of the IM test.

#### Acknowledgements

Financial support from the Flemish Fund for Scientific Research (Grant G.0366.01) is gratefully acknowledged.

#### References

- Chesher, A., 1983. The information matrix test: simplified calculation via a score test interpretation. *Economics Letters* 13, 45–48.
- Chesher, A., Spady, R., 1991. Asymptotic expansions of the information matrix test statistic. *Econometrica* 59, 787–815.
- Davidson, R., MacKinnon, J.G., 1992. A new form of the information matrix test. *Econometrica* 60, 145–157.
- Davidson, R., MacKinnon, J.G., 1998. Graphical methods for investigating the size and power of hypothesis tests. *The Manchester School* 66, 1–26.
- Hall, A., 1987. The information matrix test for the linear model. *The Review of Economic Studies* 54, 257–263.
- Horowitz, J.L., 1994. Bootstrap-based critical values for the information matrix test. *Journal of Econometrics* 61, 395–411.
- Lancaster, T., 1984. The covariance matrix of the information matrix test. *Econometrica* 52, 1051–1053.
- Orme, C., 1988. The calculation of the information matrix test for binary data models. *The Manchester School* 56, 370–376.
- Orme, C., 1990. The small-sample performance of the information-matrix test. *Journal of Econometrics* 46, 309–331.
- Orme, C., 1995. Simulated conditional moment tests. *Economics Letters* 49, 239–245.
- Taylor, L.W., 1987. The size bias of the information matrix test. *Economics Letters* 24, 63–67.
- White, H., 1982. Maximum likelihood estimation of misspecified models. *Econometrica* 50, 1–26.