

Tegravance Film-Coated Tablets

Module 3 Quality Documentation — Formulation F-832/B

Regional Information

No excipients of human or animal origin are used in the drug product with the exception of lactose monohydrate, for which a TSE/BSE compliance declaration is provided in Module 3.2.A.2. Certificates of suitability are maintained where applicable.

Annual product quality reviews evaluate batch trends, deviations, complaints, and stability data to confirm the continued state of control. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation. Statistical evaluation of batch release data demonstrates process capability well within specification limits. No changes to the approved analytical procedures are proposed in this submission. Environmental monitoring of the manufacturing areas is performed per the site monitoring program, with alert and action limits established from historical data. Risk assessments were performed in accordance with ICH Q9 principles, and residual risks were judged acceptable.

Cleaning verification swab results were below the health-based exposure limits derived for the product. Annual product quality reviews evaluate batch trends, deviations, complaints, and stability data to confirm the continued state of control. Risk assessments were performed in accordance with ICH Q9 principles, and residual risks were judged acceptable. Training records for personnel involved in manufacturing and testing are maintained under the site quality system. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation.

Process Validation Summary

Process performance qualification was executed on three consecutive commercial-scale batches. All predefined acceptance criteria for critical process parameters and critical quality attributes were met, supporting a conclusion of a validated state of control.

Supplier qualification includes periodic audits, review of certificates of analysis, and trending of incoming material test results. The finished product is stored and distributed under controlled ambient conditions consistent with the approved label storage statement. Analytical results are trended electronically, and out-of-trend results trigger a documented laboratory investigation. The control strategy links critical quality attributes to critical process parameters identified during development. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation. Hold-time studies established acceptable storage durations for intermediate blends and cores under defined conditions.

Section 3.2.P.4 — Excipients

For the commercial formulation F-832/B, Explotab (Sodium starch glycolate, disintegrant) is used.

Attribute	Acceptance approach
Identity	Compendial monograph
Assay / purity	Compendial monograph
Functionality-related characteristics	Supplier certificate plus in-house verification*

* Control approach per USP-NF, FDA Guidance for Industry: Immediate Release Solid Oral Dosage Forms, ICH Q8(R2).

Selection Justification

The formulation development rationale for Explotab rests on reduced sticking and picking observed during tableting trials.

Current Commercial Specification

Component	Function	Grade per current specification
Sodium starch glycolate	disintegrant	Primojel
Reference		SPEC-EX-5198 Rev 09

Process Validation Summary

Process performance qualification was executed on three consecutive commercial-scale batches. All predefined acceptance criteria for critical process parameters and critical quality attributes were met, supporting a conclusion of a validated state of control.

The control strategy links critical quality attributes to critical process parameters identified during development. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation. Cleaning verification swab results were below the health-based exposure limits derived for the product. Environmental monitoring of the manufacturing areas is performed per the site monitoring program, with alert and action limits established from historical data.

Control of Drug Product

The drug product specification includes tests for description, identification, assay, uniformity of dosage units, dissolution, related substances, water content, and microbial limits. Analytical procedures are compendial where available; non-compendial procedures were validated in accordance with ICH Q2(R1).

Hold-time studies established acceptable storage durations for intermediate blends and cores under defined conditions. Training records for personnel involved in manufacturing and testing are maintained under the site quality system. Risk assessments were performed in accordance with ICH Q9 principles, and residual risks were judged acceptable. Comparability of pilot- and commercial-scale batches was demonstrated with respect to dissolution and impurity profiles. Cleaning verification swab results were below the health-based exposure limits derived for the product. Annual product quality reviews evaluate batch trends, deviations, complaints, and stability data to confirm the continued state of control.

Stability Summary

Primary stability studies were conducted under ICH long-term (25°C/60% RH) and accelerated (40°C/75% RH) conditions on three registration batches. All quality attributes remained within acceptance criteria through the tested intervals. Photostability testing per ICH Q1B confirmed that the product is not photolabile in the commercial pack.

The finished product is stored and distributed under controlled ambient conditions consistent with the approved label storage statement. Deviations observed during the campaign were investigated and closed with no impact to product quality. Statistical evaluation of batch release data demonstrates process capability well within specification limits. Cleaning verification swab results were below the health-based exposure limits derived for the product. No changes to the approved analytical procedures are proposed in this submission.

Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation. Statistical evaluation of batch release data demonstrates process capability well within specification limits. No changes to the approved analytical procedures are proposed in this submission. Annual product quality reviews evaluate batch

trends, deviations, complaints, and stability data to confirm the continued state of control. Deviations observed during the campaign were investigated and closed with no impact to product quality.

Batch Formula

The representative commercial batch formula is provided for a nominal batch size of 400,000 tablets. Quantities of all components are expressed per tablet and per batch. Purified water is used as granulation fluid and is removed during processing; it does not appear in the final composition.

Comparability of pilot- and commercial-scale batches was demonstrated with respect to dissolution and impurity profiles. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation. Environmental monitoring of the manufacturing areas is performed per the site monitoring program, with alert and action limits established from historical data. Cleaning verification swab results were below the health-based exposure limits derived for the product. No changes to the approved analytical procedures are proposed in this submission. Deviations observed during the campaign were investigated and closed with no impact to product quality.

Reference Standards

Primary reference standards for the drug substance and specified impurities are qualified against compendial standards where available. Working standards are qualified against the primary standard under an approved protocol and are stored under controlled conditions.

The control strategy links critical quality attributes to critical process parameters identified during development. Comparability of pilot- and commercial-scale batches was demonstrated with respect to dissolution and impurity profiles. Supplier qualification includes periodic audits, review of certificates of analysis, and trending of incoming material test results. Deviations observed during the campaign were investigated and closed with no impact to product quality. Change control procedures ensure that any modification to materials, methods, or equipment is assessed for regulatory impact prior to implementation.