

Ebrantix — Post-Approval Stability Protocol

Protocol STAB-P-2021-86

Sample Inventory and Pull Windows

Sample inventory per condition and timepoint is defined in the pull schedule appendix. Scheduled pulls are performed within ± 7 days of the nominal date; pull records are maintained in the site stability management system.

Study Storage Conditions

The primary (zone IVb condition) storage condition for this protocol is $30^{\circ}\text{C} \pm 2^{\circ}\text{C}$ / $75\% \text{ RH} \pm 5\% \text{ RH}$. Accelerated studies at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}$ / $75\% \text{ RH} \pm 5\% \text{ RH}$ were completed through six months during development and are not part of this protocol.

Timepoints

Interval (months)	0	1	3	6	9	12	18
Long-term testing	X	X	X	X	X	X	X

Acceptance Criteria

The stability acceptance criterion applied at each interval is — Dissolution: $Q = 80\%$ in 30 minutes. All other registered specification tests apply unchanged.

Protocol Deviations

No deviations from this protocol have been recorded to date.

Stability Commitments

A post-approval commitment is made to enrol one commercial batch of Ebrantix per year into the ongoing stability program and to report confirmed out-of-specification results within 30 days.

Bracketing and Matrixing

In accordance with ICH Q1D, bracketing is applied: the 20 mg and 75 mg strengths are tested, and the intermediate 50 mg strength is covered because all strengths are direct scale-equivalents of a common blend in

the same container closure system.

Sample Inventory and Pull Windows

Sample inventory per condition and timepoint is defined in the pull schedule appendix. Scheduled pulls are performed within ± 7 days of the nominal date; pull records are maintained in the site stability management system.

Data Evaluation and Reporting

Stability data are evaluated per ICH Q1E. Regression analysis is applied to quantitative attributes exhibiting change over time; poolability across batches is assessed at the 0.25 significance level. Annual reports summarize all data generated in the reporting interval.

Batch Selection

The protocol covers three primary batches manufactured at commercial scale using the approved process. Batch genealogy, manufacturing dates, and packaging lots are recorded in the batch appendix. All batches were released against the approved specification prior to set-down.

Container Closure and Orientation

Samples are stored in the commercial container closure system. Bottles are stored upright; a subset of inverted samples is included at the initial and final timepoints to assess closure-liner interaction. Induction seal integrity is verified at each pull.