

MULTI-MODAL AI MEDICAL TRIAGE SUMMARY

Clinical Decision Support Report | Hospital Triage | UN SDG 3

Patient Name: Anonymous / Unregistered	Assessment Date: 2026-10-07 10:42:28
Age / Sex: 72 yrs / Male	Verified Mobile: N/A
Resting BP: 185 mmHg	Blood Glucose: 240 mg/dL

OVERALL URGENCY: HIGH

Calibrated Score: 100%

Clinical Synthesis: URGENT CLINICAL ALERT: Patient triaged as HIGH urgency (100%) due to acute indicators in Symptom Triage (NLP Red-Flag / Acute), Vitals Anomaly (Severe Physiological Instability), Risk Screening (High Chronic Liability). Vitals presentation reveals **CRITICAL SAFETY OVERRIDE: Critical Hypoxia:** SpO2 dropped to 86.0% (< 90.0%). Immediate medical attention required while symptoms indicate **CRITICAL OVERRIDE: Emergency Red-Flag detected:** 'crushing chest pain'. Seek immediate emergency care. Immediate physician evaluation or emergency care transport is strongly indicated.

Multi-Modal Component Breakdown

Modality	Model Tier	Score	Clinical Findings
Risk Screening	HIGH	98%	Baseline chronic risk is HIGH (Score: 98.1%). Primary risk driver is Type 2 Diabetes (Heart Risk: 35.6%, Diabetes Risk: 98.1%). Plasma Glucose (240.0), Age (72.0), Body Mass Index (BMI) (36.5) were the biggest factors elevating risk. Conversely, Pregnancies (0.0), Serum Insulin (85.0) provided protective effects.
Vitals Anomaly	HIGH	100%	CRITICAL SAFETY OVERRIDE: Critical Hypoxia: SpO2 dropped to 86.0% (< 90.0%). Immediate medical attention required.
Symptom Triage	HIGH	100%	CRITICAL OVERRIDE: Emergency Red-Flag detected: 'crushing chest pain'. Seek immediate emergency care.

Actionable Care Recommendations

CRITICAL ACTION: Consult a doctor or proceed to the nearest emergency room immediately.

Emergency Dispatcher: 112 (National Emergency) / 108 (Ambulance)

While waiting for medical responders:

- Keep patient resting in a comfortable position (semi-upright if breathing difficulty, lying flat if faint).
- Loosen tight collar, belt, or restrictive clothing around the neck and chest.
- Do NOT administer solid foods or excessive water if consciousness is reduced or choking risk exists.
- Ensure front entrance is unlocked and clearly visible for incoming emergency responders.
- Keep patient's medication list, recent vitals, and identity card ready for paramedics.

DISCLAIMER: This multi-modal triage tool is an AI-powered clinical decision-support prototype created to assist healthcare access and prioritization in underserved and rural areas. It does NOT constitute a definitive medical diagnosis or replacement for licensed clinical consultation. In cases of acute pain or emergency, seek immediate professional care.