


# SYSTEMATIC IDENTIFICATION OF TOR DOWNSTREAM EFFECTORS

## USING RANDOM-WALKS ON THE YEAST INTERACTOME

Shahin Mohammadi<sup>1</sup>, Shankar Subramaniam<sup>2</sup>, Ananth Grama<sup>1</sup>

<sup>1</sup>Computer Science Department, Purdue University, <sup>2</sup>Department of Bioengineering, University of California, San Diego

Reducing glucose concentration in yeast growth medium from 2% to 0.05%, known as severe dietary restriction (DR), had been shown recently to increase both RLS and CLS in a Sir2-independent manner[1-2]. The target of Rapamycin (TOR) signaling pathway plays a key role in fine-tuning cellular response to nutrient-availability by coordinately orchestrating various aspects of cellular machinery, including cell growth, translation initiation, ribosome biogenesis, and autophagy, among others[3]. Additionally, inhibition of the TOR signaling pathway by Rapamycin treatment exhibits similar phenotype and the lifespan can not be further extended by subjecting yeast to severe DR. These changes, as a whole, shift the energy allocation from growth and reproduction towards cell maintenance/repair. Regardless of the difference in the DR protocol, this mechanism is conserved in distant species, ranging from yeast to humans[4].



### REFERENCES


- [1] M. Kaeberlein, et al., *Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients*, Science (New York, N.Y.), 310(5751):1193-6, Nov. 2005
- [2] N. A. Bishop and L. Guarente, *Genetic links between diet and lifespan: shared mechanisms from yeast to humans*, Nature reviews, Genetics, 8(11):835-44, Nov. 2007
- [3] Y. Wei and X. F. S. Zheng, *Nutritional control of cell growth via TOR signaling in budding yeast*, Methods in molecular biology (Clifton, N.J.), 759:307-19, Jan. 2011
- [4] L. Fontana, L. Partridge, and V. D. Longo, *Extending healthy life span from yeast to humans*, Science (New York, N.Y.), 328(5976):321-6, Apr. 2010
- [5] N. L. Samara and C. Wolberger, *A new chapter in the transcription SAGA*, Current opinion in structural biology, 21(6):767-74, Dec. 2011

### ACKNOWLEDGEMENTS

This work supported by the Center for Science of Information(CSoI), an NSF Science and Technology Center, under grant agreement CCF-0939370, and by NSF grants DBI 0835677 and 0641037.



### Network integration



### Tracing information flow

Random walk with restart (preferential PageRank)

#### ➤ Constructing transition matrix:

- ✓ Normalizing out-going edges:

$$\tilde{P}_{ji} = \begin{cases} 0 & \text{if } A_{ik} = 0; \forall 1 \leq k \leq n, \\ \frac{A_{ij}}{\sum_{k=1}^n A_{ik}} & \text{O.W.} \end{cases}$$

- ✓ Adding self-loop to dangling nodes:

$$P = \tilde{P} + \text{diag}(\mathbf{1}^T - \mathbf{1}^T \tilde{P})$$

#### ➤ Computing stationary distribution:


- ✓ Iterative method:

$$\pi_v(\alpha) \equiv \alpha P \pi_v(\alpha) + (1 - \alpha) e_v$$

- ✓ Direct method:

$$\pi_v(\alpha) \equiv \frac{(1 - \alpha)(I - \alpha P)^{-1} e_v}{Q}$$

### Distribution of Scores



#### ➤ Interpretation:


Expansion using Neumann series:

$$\pi_v(\alpha) = (1 - \alpha) \sum_{i=0}^{\infty} (\alpha P)^i e_v$$


Thus,  $\pi_v$  is a function of:

1. Distance to source node
2. Multiplicity of paths


### Experimental Validation



- Dataset: Transcriptional changes in response to Rapamycin treatment
- Tail of hypergeometric distribution as a function of cut-off
- Exact p-value is computed using dynamic programming



### Enrichment Map of GO Slim terms



- ✓ Each node represents a GO term
- ✓ Edges represent the overlap between genesets of GO terms.
- ✓ Color intensity of nodes encodes the mHG p-value enrichment
- ✓ Node size represents the number of enriched genes

### Enrichment Analysis

1. We sort all nodes in the interactome ( $N$  nodes) based on their random walk distance to TORC1
2. For each functional term, we encode all annotated nodes ( $n$  true-positives) using binary vector  $\lambda$
3. For each binary vector  $\lambda$ , we compute the enrichment score as:

$$mHG(\lambda) = \min_{1 \leq l \leq N} \text{HGT}(b_l(\lambda); N, n, l)$$

where  $b_l(\lambda) = \sum_{i=1}^l \lambda_i$

### Transcription initiation

