

Numerical Linear Algebra Workshop

Duo Cao

Department of Mathematics, Purdue University

cao157@purdue.edu


March 26, 2016

Overview

- 1 Background and PDE formulation
 - Diffusion Equation
 - Miscellaneous Equations
- 2 Numerical methods for solving PDEs
 - Finite Difference Method
 - Finite Element Method
 - Miscellaneous Numerical Methods
- 3 Numerical Method for solving PDEs
- 4 Numerical Linear Algebra


Diffusion Equations

Diffusive Transport: Dye in water, pollution, heat, perfume...




Diffusion Equations Cont'd

- Consider temperature in a long thin tube of constant cross section.
- The tube is perfectly insulated laterally. Heat only flow along the tube.
- Its ends maintain at zero temperature.



Diffusion Equations Cont'd




Suppose

- thermal conductivity in the wire is K .
- Cross sectional area is A .
- Material density ρ .
- Heat capacity is σ .
- Temperature at point x at time t is $u(x, t)$.

Then the heat flow into bar across face at x : $-KA \frac{\partial u}{\partial x} \Big|_x$.

At the face $x + \delta x$: $-KA \frac{\partial u}{\partial x} \Big|_{x+\delta x}$

Diffusion Equations Cont'd



- The net flow out is: $KA \frac{\partial^2 u}{\partial x^2} \delta x$
- $Q = \sigma m \Delta T$
- So, the conservation of heat gives: $KA \frac{\partial^2 u}{\partial x^2} \delta x = \sigma \rho A \frac{\partial u}{\partial t} \delta x$

$$\boxed{\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}}$$

Boundary Conditions

- Homogeneous Dirichlet Boundary Condition: $u(0, t) = u(L, t) = 0$
- Homogeneous Neumann Boundary Condition: $\frac{\partial u}{\partial x}(0, t) = \frac{\partial u}{\partial x}(L, t) = 0$

Miscellaneous Equations

- Navier-Stokes Equation: $\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla p + \gamma \nabla^2 \mathbf{u} + \frac{1}{\rho} \mathbf{F}$
- Fisher's Equation: $\frac{\partial u}{\partial t} = u(1 - u) + \frac{\partial^2 u}{\partial x^2}$
- Nonlinear Schrodinger Equation: $i \partial_t \phi = -\frac{1}{2} \partial_x^2 \phi + \kappa |\phi|^2 \phi$
- Black-Scholes Equation: $\frac{\partial V}{\partial t} + \frac{\sigma^2 S^2}{2} \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0$
- etc

Finite Difference Method

We consider the two-point boundary value problem:

$$Au : \quad = -au'' + bu' + cu = f \text{ in } \Omega = (0, 1), \quad (1)$$

$$u(0) = u_0, \quad u(1) = u_1 \quad (2)$$

where the coefficients $a = a(x)$, $b = b(x)$, and $c = c(x)$ are smooth functions satisfying $a(x) > 0$ and $c(x) \geq 0$ in $\overline{\Omega}$. And f, u_0, u_1 are given.

To find numerical solution of (2) we introduce $M + 1$ grid points $0 = x_0 < x_1 < \dots < x_M = 1$ by setting $x_j = jh, j = 0, \dots, M$, where $h = 1/M$. We denote the approximation of $u(x_j)$ by U_j and use the following finite difference approximation for derivatives.

$$\partial U_j = \frac{U_{j+1} - U_j}{h}, (\text{forward difference})$$

$$\partial \overline{U_j} = \frac{U_j - U_{j-1}}{h}, (\text{backward difference})$$

$$\hat{\partial} U_j = \frac{U_{j+1} - U_{j-1}}{2h}, (\text{central difference})$$

$$\partial \partial U_j = \frac{U_{j+1} - 2U_j + U_{j-1}}{h^2}$$

Setting also $a_j = a(x_j)$, $b_j = b(x_j)$, $c_j = c(x_j)$, $f_j = f(x_j)$, we now define a finite difference approximation of (2) by

$$A_h U_j := -a_j \partial \bar{\partial} U_j + b_j \hat{\partial} U_j + c_j U_j = f_j, \text{ for } j = 1, \dots, M-1, \quad (3)$$

$$U_0 = u_0, \quad U_M = u_1. \quad (4)$$

Then, after simplification, the equation at the interior point x_j may be written as

$$(2a_j + h^2 c_j) U_j - (a_j - \frac{1}{2} h b_j) U_{j+1} - (a_j + \frac{1}{2} h b_j) U_{j-1} = h^2 f_j \quad (5)$$

for all j .

Put (5) into a matrix form:

$$AU = g$$

We finally comes to LINEAR ALGEBRA!! OH YEAH!!

In our system $AU = g$:

$U = (U_1, \dots, U_{M-1})^T$ and the first and last components of the vector $g = (g_1, \dots, g_{M-1})^T$ contain contributions from the boundary values u_0, u_1 as well as f_1 and f_{M-1} , respectively. The $(M-1) \times (M-1)$ matrix A is tridiagonal and diagonally dominant for h sufficiently small.

Finite Element Method for BVP

We consider the special case $b = 0$ of the two-point boundary value problem of (2),

$$Au := -(au')' + cu = f \text{ in } \Omega := (0, 1), \text{ with } u(0) = u(1) = 0,$$

where $a = a(x)$, $c = c(x)$ are smooth functions with $a(x) \geq q_0 > 0$, $c(x) \geq 0$ in $\overline{\Omega}$ and $f \in L_2 = L_2(\Omega)$.

Recall the variational formulation of this problem is to find $u \in H_0^1$ such that

$$a(u, \phi) = (f, \phi), \quad \forall \phi \in H_0^1,$$

where

$$a(v, w) = \int_{\Omega} (av'w' + cvw)dx \text{ and } (f, v) = \int_{\Omega} fvdv,$$

and that this problem has a unique solution $u \in H^2$.

For the purpose of finding an approximate solution of (15) we introduce a partition of Ω ,

$$0 = x_0 < x_1 < \dots < x_M = 1,$$

and set

$$h_j = x_j - x_{j-1}, K_j = [x_{j-1}, x_j], \text{ for } j = 1, \dots, M, \text{ and } h = \max_j h_j.$$

The discrete solution will be sought in the finite-dimensional space of functions

$$S_h = \{v \in C = C(\overline{\Omega}) : v \text{ linear on each } K_j, v(0) = v(1) = 0\}.$$

The set $\{\Phi_i\}_{i=1}^{M-1} \subset S_h$ is defined by

$$\Phi_i(x_j) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

and any $v \in S_h$ may be written as

$$v(x) = \sum_{i=1}^{M-1} v_i \Phi_i(x), \text{ with } v_i = v(x_i).$$

Now we pose the finite-dimensional problem to find $u_h \in S_h$ such that

$$a(u_h, \chi) = (f, \chi), \quad \forall \chi \in S_h. \quad (6)$$

In terms of the basis $\{\Phi_i\}_{i=1}^{M-1}$ we write $u_h(x) = \sum_{j=1}^{M-1} U_j \Phi_j(x)$ and insert this into (6) to find that this equation is equivalent to

$$\sum_{j=1}^{M-1} U_j a(\Phi_j, \Phi_i) = (f, \Phi_i), \quad \text{for } i = 1, \dots, M-1.$$

This linear system of equations could be expressed in matrix form as

$$AU = b$$

Finished!!!

In our system $U = (U_j)$, $A = (a_{ij})$ is the stiffness matrix with elements $a_{ij} = a(\Phi_j, \Phi_i)$, and $b = (b_i)$ with elements $b_i = (f, \Phi_i)$. The matrix A is symmetric and positive definite, because for $V = (V_i)$ and $v(x) = \sum_{i=1}^{M-1} V_i \Phi_i(x)$ we have

$$V^T A V = \sum_{j=1}^{M-1} V_j a_{ij} V_j = a\left(\sum_{j=1}^{M-1} V_j \Phi_j, \sum_{i=1}^{M-1} V_i \Phi_i\right) = a(v, v) \geq a_0 \|v'\|^2,$$

and hence $V^T A V = 0$ implies $v' = 0$, so that v is 0. Matrix A is tridiagonal since $a_{ij} = 0$ when x_i and x_j are not neighbors, i.e., when $|i - j| \geq 2$.

Miscellaneous Numerical Methods

- Finite Volume Method
- Spectral Method
- Meshfree Methods
- Multigrid
- etc.

$$AU = b$$

Blocks of Highlighted Text

Block 1

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Integer lectus nisl, ultricies in feugiat rutrum, porttitor sit amet augue. Aliquam ut tortor mauris. Sed volutpat ante purus, quis accumsan dolor.

Block 2

Pellentesque sed tellus purus. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos. Vestibulum quis magna at risus dictum tempor eu vitae velit.

Block 3

Suspendisse tincidunt sagittis gravida. Curabitur condimentum, enim sed venenatis rutrum, ipsum neque consectetur orci, sed blandit justo nisi ac lacus.

Heading

- 1 Statement
- 2 Explanation
- 3 Example

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Integer lectus nisl, ultricies in feugiat rutrum, porttitor sit amet augue. Aliquam ut tortor mauris. Sed volutpat ante purus, quis accumsan dolor.

Theorem

Theorem (Mass–energy equivalence)

$$E = mc^2$$

Example (Theorem Slide Code)

```
\begin{frame}  
\frametitle{Theorem}  
\begin{theorem}[Mass--energy equivalence]  
$E = mc^2$  
\end{theorem}  
\end{frame}
```

Figure

Uncomment the code on this slide to include your own image from the same directory as the template .TeX file.

An example of the `\cite` command to cite within the presentation:

This statement requires citation [Erwin Kreyszig, 2014].



Erwin Kreyszig, Advanced Engineering Mathematics, 8th Edn, Sections 11.4b

The End