

Towards a taxonomy of atlases and of morphisms between them*

Shmuel (Seymour J.) Metz

September 1, 2026

Abstract

Manifolds and fiber bundles, while superficially different, have strong parallels; in particular, they are both defined in terms of equivalence classes of atlases or in terms of maximal atlases, with the atlases treated as mere adjuncts. This paper presents a unified view of atlases for manifolds and fiber bundles as mathematical entities in their own right. It defines some convenient notation, defines categories of, e.g., atlases, and constructs functors among them.

{LCS}

[Metz, 2018] introduced the ideas presented here, but many of the details are not needed there. This paper fleshes out the concepts in more detail than would be relevant there.

Part I

Introduction

{part:intro}

Historically, manifolds were formalized in terms of atlases based on Euclidean spaces and fiber bundles were formalized in terms of atlases based on product spaces, using either equivalence classes of atlases or maximal atlases. The concept of pseudogroups allowed unifying manifolds and manifolds with boundary. The definitions of fiber bundles and manifolds have strong parallels, and can be unified in a similar fashion. The standard presentations treat these atlases as secondary adjuncts.

{LCS}

[Metz, 2018] introduced the concept of an m-atlas and of morphisms among them only as preliminaries to unifying manifolds and fiber bundles as special cases of local coordinate spaces. This paper treats m-atlases as objects of interest in their own right, and includes concepts not relevant to local coordinate spaces.

This paper defines an approach using categories and commutative diagrams that is designed for easy exposition at the possible expense of abstractness and generality. In particular, I have chosen to simplify proofs by stating stricter assumptions than necessary and have assumed the Axiom of Choice (AOC).

*I wish to gratefully thank Walter Hoffman (z"l), Milton Parnes, Dr. Stanley H. Levy (z"l), the Mathematics department of Wayne State University, the Mathematics department of the State University of New York at Buffalo and others who guided my education.

Although this paper incidentally defines partial equivalents to manifolds and fiber bundles using model spaces and model atlases, it does not explicitly reflect the role of the group in fiber bundles.

This paper defines functors among categories of atlases, categories of model spaces, categories of manifolds and categories of fiber bundles; it constructs more machinery than is customary in order to facilitate the presentation of those categories and functors.

Remark 0.1. The definitions of morphisms given here are nonstandard, but there are also definitions of classical morphisms that are much closer to the standard ones.

Parts II to V present nomenclature and give basic results.

Part VI defines m-atlases, m-atlas morphisms and categories of them; lemma 11.23 proves that the defined categories are indeed categories.

part VII (Equivalence of manifolds) on page 114 and part VIII (Equivalence of fiber bundles) on page 137 show the equivalence of manifolds and fiber bundles with m-atlases by explicitly exhibiting functors to and from the m-atlases.

Remark 0.2. The unconventional definitions of manifold and fiber bundle are intended to make their relationship to local coordinate spaces more natural. See [Metz, 2018] for details.

Most of the lemmata, theorems and corollaries in this paper should be substantially identical to results that are familiar to the reader. What is novel is the perspective and the material directly related to local coordinate spaces. The presentation assumes only a basic knowledge of Category Theory, such as may be found in the first chapter of [Mac Lane, 1998] or [Adamek, Herrlich, Strecker, 1990].

Many of the proofs are obvious and may be omitted from other versions of this paper.

1 New concepts and notation

This paper introduces a significant number of new concepts and some modifications of the definitions for some conventional concepts. It also introduces some notation of lesser importance. The following are the most important.

1. **Nearly commutative diagram (NCD)**, NCD at a point, locally NCD and special cases with related nomenclature. These are cases where a diagram can be modified to make it commutative.
2. **Model space** and related concepts. A **Model space**¹ is a topological space with a category of permissible open sets and mappings satisfying specified conditions. It is similar to a pseudo-group, with some important differences.
3. **Model topology** and **M-paracompactness**
4. An m-atlas is a generalization of atlases in manifolds.

¹The phrase has been used before, but with a different meaning.

5. An **m-atlas near morphism** (**m-atlas morphism**) is a generalization of a C^k map between differentiable manifolds. It differs in using both a function between the manifolds and a function between the coordinate spaces.
6. **Linear space and related concepts** A linear space is a subspace of a Banach space or of a Fréchet space that is suitable for defining coordinate patches in a manifold.
7. **Trivial C^k linear model space** and related concepts. A trivial C^k linear model space is a maximal model space with morphisms C^k maps between open subsets of a linear space.
8. **G - ρ bundle atlases²** and related concepts

Part II

Conventions

{part:conv}

An arrow with an Equal-Tilde ($A \xrightarrow[\phi]{\cong} B$) represents an isomorphism. One with a hook ($A \hookrightarrow_i B$) represents an inclusion map. One with a tail ($A \rightarrowtail_i B$) represents a monomorphism. One with a double head ($A \twoheadrightarrow_\pi B$) represents a surjection.

When a superscript ends in -1 , e.g., f^{i-1} , it is to be taken as function inverse rather than subtraction of 1.

All diagrams shown are commutative; none are exact.

A definition of a base term and several related terms may specify modifiers in parenthese in the base definition and then give the definitions for each modifier; usually the modifiers will be restrictions and the base definition will not be repeated. E.g., “ **f** is a (semi-strict, strict) prestructure morphism of P^1 to P^2 iff” is followed by the definition of prestructure morphism, then by the restrictions for strict and semi-strict prestructure morphisms.

When a definition defines a base propositional function and variant propositional functions with a qualifier given as a superscript or underset, then the form $\text{base}^{(qualifiers)}$ will refer to either $\text{base}^{qualifier}$ or base and the form $\text{base}_{(qualifiers)}$ will

refer to either $\text{base}_{qualifier}$ or base, e.g., $\text{isAtl}_{Ar}^{(classic)}$ refers to either isAtl_{Ar} , $\text{isAtl}_{Ar}^{classic}$, $\text{isAtl}_{Ar}^{semi-strict}$, $\text{isAtl}_{Ar}^{strict}$, $\text{isAtl}_{Ar}^{classic,strict}$ or $\text{isAtl}_{Ar}^{strict}$.

The form $\text{base}^{qualifier(qualifiers)}$ will refer to either $\text{base}^{qualifier}$ or to the variant with one or more of the additional qualifiers in the superscript, e.g., $\text{isAtl}_{Ar}^{classic(near)}$ refers to either $\text{isAtl}_{Ar}^{classic}$ or $\text{isAtl}_{Ar}^{classic,near}$.

strict strict

²Similar to coordinate bundles

Alternatively, a definition may specify a numbered list of alternatives, and subsequently specify additional numbered lists with items corresponding to those in the first list, e.g., $\mathbf{f}$ is also a

1. full
2. semi-maximal
3. maximal
4. full semi-maximal
5. full maximal

E^1 - E^2 C^k morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as

1. abbreviation for full
2. abbreviation for semi-maximal
3. abbreviation for maximal
4. abbreviation for full semi-maximal
5. abbreviation for full maximal

iff

1. definition for full
2. definition for semi-maximal
3. definition for maximal
4. definition for full semi-maximal
5. definition for full maximal

A Corollary, Lemma or Theorem that applies to related terms defined with the above convention will specify the modifiers in parentheses to indicate that it applies to the base term and to the variant terms, e.g., “a (semi-strict, strict) E^i - E^{i+1} m-atlas morphism” If it applies only to more restrictive terms then it will specify the first relevant modifier followed by the remaining relevant modifiers in parentheses, e.g.,

“If $\mathcal{S}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}'^i$ and $\mathbf{f}^i$ is a semi-strict (strict) prestructure morphism”.

Blackboard bold upper case will denote specific sets, e.g., the Naturals.

Bold lower case italic letters will refer to sets, sequences and tuples of functions, e.g., $\mathbf{f} \stackrel{\text{def}}{=} (f_1, f_2)$.

Bold lower case Latin letters will refer to sequence valued functions of sequences and tuple valued functions of tuples, e.g., **range** yields the sequence of ranges of a sequence of functions.

Bold upper case italic letters will refer to sequences or tuples, e.g., $\mathbf{A} = (x, y, z)$, to sets of them, to sets of topological spaces or to sets of open sets.

Bold upper case script letters will refer to sequences of categories, e.g.,

$\mathcal{A} \stackrel{\text{def}}{=} (\mathcal{A}_\alpha, \alpha \in A)$.

Fraktur will refer to topologies and to topology-valued functions, e.g., $\mathfrak{Top}$.

Functions have a range, domain and relation, not just a relation. Unless otherwise stated, they are assumed or proven to be continuous.

Groups are assumed to be topological groups. The ambiguous notation x^{-1} will be used when it is obvious from context what the group operation $\star$ and the group identity $\mathbf{1}_G$ are.

Lower case Greek letters other than π, ρ, σ, ϕ and ψ will refer to ordinals, possibly transfinite, and to formal labels. A letter with a Greek superscript and a letter with a Latin or numeric superscript always refer to distinct variables.

Lower case π will refer to a projection operator

Lower case ρ will refer to a continuous effective group action, i.e., a continuous representation of a group in a homeomorphism group.

Lower case σ will refer to a sequence of ordinals, referred to as a signature.

Lower case ϕ will refer to a coordinate function.

Lower case italic and Latin letters will refer to

1. elements of a set or sequence

2. formal labels

When the letters u and v , with or without a superscript, refer to an element of a set then they may also be used as formal labels for sets associated with that element.

3. functions

4. morphisms of a category

5. natural numbers

6. objects of a category

7. ordinal numbers

Upper case Greek letters other than Σ may refer to

1. ordinal used as the limit of a sequence of consecutive ordinals, e.g., $x_\alpha, \alpha \leq A$

2. ordinal used as the order type of a sequence of consecutive ordinals, e.g., $x_\alpha, \alpha < A$

Upper case Σ will refer to a sequence of signatures

Upper case Latin letters will refer to

1. Natural numbers

2. Topological spaces

3. Open sets
4. Formal labels for elements of a sequence or tuple of functions, e.g., f_E might be $f_0: E_1 \longrightarrow E_2$.

Upper case Script Latin letters will refer to categories and functors.
Upright Latin letters will be used for long names. In particular,

1. $\mathbf{Ar}$, $\mathbf{Hom}$, $\mathbf{Mor}$ and $\mathbf{Ob}$ are the conventional set-valued functions on small categories:

$\mathbf{Ar}(\mathcal{C})$ All morphisms of $\mathcal{C}$

$\mathbf{Hom}_{\mathcal{C}}(a, b)$ All morphisms of $\mathcal{C}$ from a to b

$\mathbf{Mor}(\mathcal{C})$ All morphisms of $\mathcal{C}$. Synonymous with $\mathbf{Ar}(\mathcal{C})$.

$\mathbf{Ob}(\mathcal{C})$ All objects of $\mathcal{C}$

They will never be applied to large categories in this paper.

2. **Set** and **Top** are the conventional categories of sets and topological spaces.
3. The letters E , G , X and Y will be used for formal labels.

The term C^k includes C^∞ (smooth) and C^ω (analytic).

This paper uses the term morphism in preference to arrow, but uses the conventional $\mathbf{Ar}$.

This paper uses the terms empty, null and void interchangeably.

The term sequence without an explicit reference to $\mathbb{N}$ will refer to a general ordinal sequence, possibly transfinite.

Sequence numbering, unlike tuple numbering, starts at 0 and the exposition assumes a von Neumann definition of ordinals, so that $\alpha \in \beta \equiv \alpha < \beta$.

Except where explicitly stated otherwise, all categories mentioned are small categories with underlying sets, but the morphisms will often not be set functions between the objects and there will not always be a forgetful functor to **Set** or **Top**. By abuse of language no distinction will consistently be made between a category $\mathcal{A}$ of topological spaces and the concrete category $(\mathcal{A}, \mathcal{U})$ over **Top**. Similarly, no distinction will consistently be made among the object $U \in \mathbf{Ob}(\mathcal{A})$, the topological space $\mathcal{U}(U)$ and the underlying set.

The notation G^V will refer only to the set of continuous functions from V to G , never to the set of all functions from V to G .

When defining a category, the ordered pair $(\mathbf{O}, \mathbf{M})$ refers to the smallest concrete category over **Set** or **Top** whose objects are the elements in $\mathbf{O}$, whose morphisms include all of the elements of $\mathbf{M}$ and whose morphisms from $o^1 \in \mathbf{O}$ to $o^2 \in \mathbf{O}$ are functions $f: o^1 \longrightarrow o^2$ and whose composition is function composition.

When defining a category, the ordered triple $(\mathbf{O}, \mathbf{M}, C)$ refers to the small category whose objects are in $\mathbf{O}$, whose morphisms are in $\mathbf{M}$, whose $\mathbf{Hom}$ is

$$\mathbf{Hom}_{(\mathbf{O}, \mathbf{M}, C)}(o_1 \in \mathbf{O}, o_2 \in \mathbf{O}) \stackrel{\text{def}}{=} \{(f, o_1, o_2) \in \mathbf{M}\} \quad (1.1)$$

and whose composition is C .

By abuse of language I may write “ $\mathcal{S}$ ” for $\text{Ob}(\mathcal{S})$, “ $A \in \mathcal{A}$ ” for $A \in \text{Ob}(\mathcal{A})$, “ $A \subset \mathcal{A}$ ” for $A \subset \text{Ob}(\mathcal{A})$, “ $A \in \mathcal{A} \subset B \in \mathcal{B}$ ” for “the underlying set of A is contained in the underlying set of B and the inclusion $i: x \in A \hookrightarrow x \in B$ is a morphism” and “ $f: A \longrightarrow B$ ” for $f \in \text{Hom}_{\mathcal{C}}(A, B)$, where $\mathcal{C}$ is understood by context.

By abuse of language I shall use the same nomenclature for sequences and tuples, and will use parentheses around a single expression both for grouping and for a tuple with a single element; the intent should be clear from context.

By abuse of language I will omit parentheses around the operands of Functors when they can be assumed by context.

By abuse of language I shall use the $\times$ and $\times$ symbols for both Cartesian products of sets and Cartesian products of functions on those sets.

By abuse of language, and assuming AOC, I shall refer to some sets as ordinal sequences, e.g., “ $(C_\alpha, \alpha \in A)$ ” for “ $\{C_\alpha \mid \alpha \in A\}$ ”, in cases where the order is irrelevant.

Where a proof requires proving intermediate propositions, a bold leadin to a paragraph will indicate such a proposition, e.g.

Intermediate proposition 1: Intermediate proof 1.

Intermediate proposition 2: Intermediate proof 2.

By abuse of language, I may omit universal quantifiers in cases where the intent is clear.

In some cases I define a notion similar to a conventional notion and also need to refer to the conventional notion. In those cases I prefix a letter or phrase to the term, e.g., m-paracompact versus paracompact.

Part III

General notions

{part: notions}

This part of the paper describes nomenclature used throughout the paper. In some cases this reflects new nomenclature or notions, in others it simply makes a choice from among the various conventions in the literature.

{def: catprop}

Definition 1.1 (Operations on categories). If $\mathcal{C}$ is a category then

1. $x \in^{\text{Ob}} \mathcal{C} \equiv x \in \text{Ob}(\mathcal{C})$
2. $x \in^{\text{Ar}} \mathcal{C} \equiv x \in \text{Ar}(\mathcal{C})$
3. If $\mathcal{C}$ is a concrete category over **Set** or **Top** and $o \in^{\text{Ob}} \mathcal{C}$, then $\mathbf{Set}_{\mathcal{C}}(o)$ is the underlying set of o . By abuse of language we will write $\mathbf{Set}(o)$ when the category is understood from context.

4. If $\mathcal{C}$ is a concrete category over **Top** and $o \in \mathcal{C}$, then $\text{Top}_{\mathcal{C}}(o)$ is the underlying topological space of o . By abuse of language we will write $\text{Top}(o)$ when the category is understood from context.

If $\mathcal{S}$ and $\mathcal{T}$ are categories then $\mathcal{S} \stackrel{\text{cat}}{\subseteq} \mathcal{T}$ iff $\mathcal{S}$ is a subcategory of $\mathcal{T}$ and $\mathcal{S} \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}$ iff $\mathcal{S}$ is a full subcategory of $\mathcal{T}$.

If $\mathcal{S}$ and $\mathcal{T}$ are categories then the category union of $\mathcal{S}$ and $\mathcal{T}$, abbreviated $\mathcal{S} \cup^{\text{cat}} \mathcal{T}$, is the category whose objects are in $\mathcal{S}$ or in $\mathcal{T}$ and whose morphisms are in $\mathcal{S}$ or in $\mathcal{T}$. {def: Id}

Definition 1.2 (Identity). $\text{Id}_{\mathcal{S}}$ is the identity function on the space $\mathcal{S}$,

Id_o is the identity morphism for the object o ³,

Id_V^U , for $U \subseteq V$, is the inclusion map⁴. $\text{Id}_{U,V}$ is synonymous with Id_V^U .

$\text{Id}_{\mathcal{C}}$ is the identity functor on the category $\mathcal{C}$.

$\mathbf{Id}_{\mathcal{S}^i}$, $i = 1, 2$, is the sequence of identity functions for the elements of the sequence $\mathcal{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_{\alpha}^i, \alpha < A)$. Let $\mathcal{S}^1 \stackrel{\circ}{\subseteq} \mathcal{S}^2$. Then $\mathbf{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}$ is the sequence of inclusion maps $(\text{Id}_{\mathcal{S}_{\alpha}^2}^{\mathcal{S}_{\alpha}^1}, \alpha < A)$ for the elements of the sequences $\mathcal{S}^i$. $\mathbf{Id}_{\mathcal{S}^1, \mathcal{S}^2}$ is synonymous with $\mathbf{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}$. A similar definition applies for tuples, except in a context where they are treated as objects of a category.

$\mathbf{Id}_{(A \dots Z)}$ is the tuple of identity maps $(\text{Id}_A, \dots, \text{Id}_Z)$. $\mathbf{Id}_{(A^1, \dots, Z^1)}^{(A^2, \dots, Z^2)}$ is the tuple of inclusion maps $(\text{Id}_{A^2}^{A^1}, \dots, \text{Id}_{Z^2}^{Z^1})$.

Definition 1.3 (Images). $f[U] \stackrel{\text{def}}{=} \{f(x) | x \in U\}$ is the image of U under f and $f^{-1}[V] \stackrel{\text{def}}{=} \{x | f(x) \in V\}$ is the inverse image of V under f .

Remark 1.4. This notation, adopted from [Kelley, 1955], avoids the ambiguity in the traditional $f(U)$ and $f^{-1}(V)$. {GenTop}

Definition 1.5 (Projections). π_{α} is the projection function that maps a sequence into element α of the sequence. π_i is also the projection function that maps a tuple into element i of the tuple. {projections}

Definition 1.6 (Topological category). A topological category is a small subcategory of **Top** or its concrete category over **Set**. {def: topcat}

$\mathcal{T}$ is a full topological category iff it is a topological category and whenever $U^i, V^i \in \mathcal{T}, i = 1, 2, V^i \stackrel{\text{subsp}}{\subseteq} U^i, f: U^1 \longrightarrow U^2 \in \mathcal{T}$ and $f[V^1] \subseteq V^2$ then $f: V^1 \longrightarrow V^2 \in \mathcal{T}$. {lem: topInc}

Lemma 1.7 (Inclusions in topological categories are morphisms). Let $\mathcal{T}$ be a full topological category, $\mathcal{S}^i \in \mathcal{T}, i = 1, 2$, and $\mathcal{S}^1 \stackrel{\text{subsp}}{\subseteq} \mathcal{S}^2$. Then $\text{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}$ is a morphism of $\mathcal{T}$.

³The object is often expressed as a tuple, e.g., $\text{Id}_{(A,B)}$ is the identity morphism for the object (A, B)

⁴ U need not have the subspace topology.

Proof. 1. $U^i \stackrel{\text{def}}{=} S^2 \in \mathcal{T}^{\text{Ob}}, i = 1, 2$, by hypothesis.

2. $V^1 \stackrel{\text{def}}{=} S^1 \in \mathcal{T}$ and $V^2 \stackrel{\text{def}}{=} S^2 \in \mathcal{T}$ by hypothesis.

3. $V^i \stackrel{\text{subsp}}{\subseteq} U^i$,

4. $f \stackrel{\text{def}}{=} \text{Id}_{S^2}: U^1 \longrightarrow U^2 \in \mathcal{T}^{\text{Ar}}$.

5. $f[V^1] = V^1 \stackrel{\text{subsp}}{\subseteq} V^2$.

Thus $\text{Id}_{S^2}^{S^1} = f: V^1 \longrightarrow V^2 \in \mathcal{T}$ by definition 1.6. $\square$

$\{\text{def:topcat}\}$
 $\{\text{def:topLocal}\}$

Definition 1.8 (Local morphisms). Let $\mathcal{S}^i, i = 1, 2$, be a full topological category and $S^i \in \mathcal{S}^i$. A continuous function $f: S^1 \longrightarrow S^2$ is a local $\mathcal{S}^1$ - $\mathcal{S}^2$ morphism of S^1 to S^2 iff $\mathcal{S}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}^2$ and for every $u \in S^1$ there is an open neighborhood U_u for u and an open neighborhood V_u for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathcal{S}^2$.

Remark 1.9. The condition $f: U_u \longrightarrow V_u \in \mathcal{S}^2$ ensures that $U_u \in \mathcal{S}^1$ and $V_u \in \mathcal{S}^2$

Let $\mathcal{T}$ be a full topological category and $S^i \in \mathcal{T}, i = 1, 2$. A continuous function $f: S^1 \longrightarrow S^2$ is a local $\mathcal{T}$ morphism of S^1 to S^2 iff it is a local $\mathcal{T}$ - $\mathcal{T}$ morphism of S^1 to S^2 .

$\{\text{lem:topLocal}\}$

Lemma 1.10 (Local morphisms). Let $\mathcal{T}^i, i \in [1, 3]$, be a full topological category, $S^i \in \mathcal{T}^i$ and $\mathcal{T}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^{i+1}$.

If $f^i: S^i \longrightarrow S^{i+1} \in \mathcal{T}^{i+1}$ then f^i is a local $\mathcal{T}^i$ - $\mathcal{T}^{i+1}$ morphism of S^i to S^{i+1} .

Proof. Let $u \in S^i$ and $v \stackrel{\text{def}}{=} f^i(u) \in S^{i+1}$. S^i is an open for u , S^{i+1} is an open neighborhood of v and $f^i: S^i \longrightarrow S^{i+1} \in \mathcal{T}^{i+1}$ by hypothesis. $\square$

If each $f^i: S^i \longrightarrow S^{i+1}$, is a local $\mathcal{T}^i$ - $\mathcal{T}^{i+1}$ morphism of S^i to S^{i+1} then $f^2 \circ f^1: S^1 \longrightarrow S^3$ is a local $\mathcal{T}^1$ - $\mathcal{T}^3$ morphism of S^1 to S^3 .

Proof. Since $\mathcal{T}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^2$ and $\mathcal{T}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^3$, $\mathcal{T}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^3$. Let $u \in S^1$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist an open neighborhood U_u for u , open neighborhoods V_u, V'_v for v and an open neighborhood W_v of w such that $f^1[U_u] \subseteq V_u$, $f^1: U_u \longrightarrow V_u$ is a morphism of $\mathcal{T}^2$, $f^2[V'_v] \subseteq W_v$ and $f^2: V'_v \longrightarrow W_v$ is a morphism of $\mathcal{T}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_u \cap V'_v \neq \emptyset$, $\hat{V}_u$ is an open neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}_u]$ is an open neighborhood of u . $f^1: \hat{U}_u \longrightarrow \hat{V}_u$ and $f^2: \hat{V}_u \longrightarrow W_v$ are morphisms of $\mathcal{T}^3$ by definition 1.6 (Topological category) on page 8 and thus $f^2 \circ f^1: \hat{U}_u \longrightarrow W_v$ is a morphism of $\mathcal{T}^3$. $\square$

$\{\text{def:topcat}\}$

{cor:topLocal}

Corollary 1.11 (Local morphisms). Let $\mathcal{T}^i, i = 1, 2$, be a full topological category, $\mathcal{T}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^{i+1}, S^i \in \mathcal{T}^i$ and $S^1 \subseteq S^2$. Then $\text{Id}_{S^2}^{S^1}$ is a local $\mathcal{T}^1$ - $\mathcal{T}^2$ morphism of S^1 to S^2 and Id_{S^i} is a local $\mathcal{T}$ morphism of S^i to S^i .

Proof. $S^1 \stackrel{\text{Ob}}{\in} \mathcal{T}^2$ because $S^1 \stackrel{\text{Ob}}{\in} \mathcal{T}^1$ and $\mathcal{T}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{T}^2, S^2 \stackrel{\text{Ob}}{\in} \mathcal{T}^2$ by hypothesis and $S^1 \subseteq S^2$ by hypothesis, so $\text{Id}_{S^2}^{S^1} \in \mathcal{T}^2$ by lemma 1.7.

{lem:topInc}

$$\text{Id}_{S^i} \stackrel{\text{def}}{=} \text{Id}_{S^i}^{S^i}.$$

□

Definition 1.12 (Sequence functions). Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\alpha, \alpha < A)$ be a sequence of functions. Then

$$\mathbf{domain}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{domain}(s_\alpha), \alpha < A) \quad (1.2)$$

$$\mathbf{range}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{range}(s_\alpha), \alpha < A) \quad (1.3)$$

Let $\mathbf{T} \stackrel{\text{def}}{=} (t_\alpha, \alpha < A)$ be a sequence of functions with $\mathbf{range}(\mathbf{S}) = \mathbf{domain}(\mathbf{T})$. Then their composition is the sequence $\mathbf{T} \stackrel{\circ}{\circ} \mathbf{S} \stackrel{\text{def}}{=} (t_\alpha \circ s_\alpha, \alpha < A)$,

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\gamma, \gamma \leq \Gamma)$, then these functions extract information about the sequence:

$$\text{head}(\mathbf{S}, \Omega) \stackrel{\text{def}}{=} (s_\gamma, \gamma < \Omega) \quad (1.4)$$

$$\text{head}(\mathbf{S}) \stackrel{\text{def}}{=} \text{head}(\mathbf{S}, \Gamma) \quad (1.5)$$

$$\text{length0}(\mathbf{S}) \stackrel{\text{def}}{=} \Gamma \quad (1.6)$$

$$\text{tail}(\mathbf{S}) \stackrel{\text{def}}{=} S_\Gamma \quad (1.7)$$

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\gamma, \gamma < \Gamma)$, then

$$\text{length}(\mathbf{S}) \stackrel{\text{def}}{=} \Gamma \quad (1.8)$$

Remark 1.13. If $\text{length0}(\mathbf{S})$ is defined then $\text{length}(\mathbf{S}) = \text{length0}(\mathbf{S}) + 1$. $\text{length0}(\mathbf{S})$ is the ordinal type of $\text{head}(\mathbf{S})$, not the ordinal type of $\mathbf{S}$.

Let $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$ and $\mathcal{T} \stackrel{\text{def}}{=} (\mathcal{T}_\alpha, \alpha < A)$ be sequences of categories. Then $\mathcal{S}$ is a subcategory sequence of $\mathcal{T}$, abbreviated $\mathcal{S} \stackrel{\text{cat}}{\subseteq} \mathcal{T}$, iff every category in $\mathcal{S}$ is a subcategory of the corresponding category in $\mathcal{T}$, i.e., $\left(\bigvee_{\alpha < A}\right) \mathcal{S}_\alpha \stackrel{\text{cat}}{\subseteq} \mathcal{T}_\alpha$, and $\mathcal{S}$ is a full subcategory sequence of $\mathcal{T}$, abbreviated $\mathcal{S} \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}$, iff every category in $\mathcal{S}$ is a full subcategory of the corresponding category in $\mathcal{T}$, i.e., $\left(\bigvee_{\alpha < A}\right) \mathcal{S}_\alpha \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}_\alpha$.

The category sequence union of $\mathcal{S}$ and $\mathcal{T}$, abbreviated $\mathcal{S} \stackrel{\text{cat}}{\cup} \mathcal{T}$, is the sequence of category unions of corresponding categories in $\mathcal{S}$ and $\mathcal{T}$, i.e., $(\mathcal{S}_\alpha \stackrel{\text{cat}}{\cup} \mathcal{T}_\alpha)$.

{lem:seqfunc}

Lemma 1.14 (Sequence functions). Let $\mathbf{f}^i \stackrel{\text{def}}{=} (f_\alpha^i, \alpha < A)$, $i \in [1, 3]$, be sequences of functions with $\mathbf{domain}(\mathbf{f}^2) = \mathbf{range}(\mathbf{f}^1)$ and $\mathbf{domain}(\mathbf{f}^3) = \mathbf{range}(\mathbf{f}^2)$. Then $(\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 = \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1)$.

Proof.

$$\begin{aligned} (\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 &= ((f_\alpha^3 \circ f_\alpha^2) \circ f_\alpha^1, \alpha < A) \\ &= (f_\alpha^3 \circ (f_\alpha^2 \circ f_\alpha^1), \alpha < A) \\ &= \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1) \end{aligned}$$

□

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha, \alpha < A)$ be a sequence of functions, $\mathbf{D} = \mathbf{domain}(\mathbf{f})$ and $\mathbf{R} = \mathbf{range}(\mathbf{f})$. Then $\mathbf{Id}_\mathbf{R}$ is a left $\circ$ identity for $\mathbf{f}$ and $\mathbf{Id}_\mathbf{D}$ is a right $\circ$ identity for $\mathbf{f}$.

Proof.

$$\begin{aligned} \mathbf{Id}_\mathbf{R} \circ \mathbf{f} &= (\text{Id}_{\mathbf{range}(f_\alpha)} \circ f_\alpha, \alpha < A) \\ &= (f_\alpha, \alpha < A) \\ &= \mathbf{f} \end{aligned}$$

$$\begin{aligned} \mathbf{f} \circ \mathbf{Id}_\mathbf{D} &= (f_\alpha \circ \text{Id}_{\mathbf{domain}(f_\alpha)}, \alpha < A) \\ &= (f_\alpha, \alpha < A) \\ &= \mathbf{f} \end{aligned}$$

□

Definition 1.15 (Tuple functions). Let $\mathbf{S} \stackrel{\text{def}}{=} (s_n, n \in [1, N])$ be a tuple of functions. Then

$$\mathbf{domain}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{domain}(s_n), n \in [1, N]) \quad (1.9)$$

$$\mathbf{range}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{range}(s_n), n \in [1, N]) \quad (1.10)$$

Let $\mathbf{T} \stackrel{\text{def}}{=} (t_n, n \in [1, N])$ be a tuple of functions with $\mathbf{range}(\mathbf{S}) = \mathbf{domain}(\mathbf{T})$. Then their composition is the tuple $\mathbf{T} \circ \mathbf{S} \stackrel{\text{def}}{=} (t_n \circ s_n, n \in [1, N])$

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_m, m \in [1, M])$ and $\mathbf{T} \stackrel{\text{def}}{=} (t_n, n \in [1, N])$ be tuples. Then the following are tuple functions

$$\text{head}(\mathbf{S}, I) \stackrel{\text{def}}{=} (s_m, m \in [1, I]) \quad (1.11)$$

$$\text{head}(\mathbf{S}) \stackrel{\text{def}}{=} \text{head}(\mathbf{S}, M - 1) \quad (1.12)$$

$$\text{tail}(T, I) \stackrel{\text{def}}{=} (t_n, n \in [I, N]) \quad (1.13)$$

$$\text{tail}(T) \stackrel{\text{def}}{=} t_N \quad (1.14)$$

$$\text{join}(S, T) \stackrel{\text{def}}{=} (s_1, \dots, s_M, t_1, \dots, t_N) \quad (1.15)$$

{lem:tupfunc}

Lemma 1.16 (Tuple functions). Let $\mathbf{f} \stackrel{\text{def}}{=} (f_n, n \in [1, N])$ be a tuple of functions, $\mathbf{D} \stackrel{\text{def}}{=} \mathbf{domain}(\mathbf{f})$ and $\mathbf{R} \stackrel{\text{def}}{=} \mathbf{range}(\mathbf{f})$. Then $\mathbf{Id}_{\mathbf{R}}$ is a left $\circ$ identity for $\mathbf{f}$ and $\mathbf{Id}_{\mathbf{D}}$ is a right $\circ$ identity for $\mathbf{f}$.

Proof.

$$\begin{aligned} \mathbf{Id}_{\mathbf{R}} \circ \mathbf{f} &= (\text{Id}_{\text{range}(f_n)} \circ f_n, n \in [1, N]) \\ &= (f_n, n \in [1, N]) \\ &= \mathbf{f} \end{aligned}$$

$$\begin{aligned} \mathbf{f} \circ \mathbf{Id}_{\mathbf{D}} &= (f_n \circ \text{Id}_{\text{domain}(f_n)}, n \in [1, N]) \\ &= (f_n, n \in [1, N]) \\ &= \mathbf{f} \end{aligned}$$

□

{lem:tupcomp}

Lemma 1.17 (Tuple composition for unlabeled morphisms). Let

$\mathbf{f}^i \stackrel{\text{def}}{=} (f_n^i, n \in [1, N])$, $i \in [1, 3]$, be tuples of functions with $\mathbf{domain}(\mathbf{f}^2) = \mathbf{range}(\mathbf{f}^1)$ and $\mathbf{domain}(\mathbf{f}^3) = \mathbf{range}(\mathbf{f}^2)$. Then $(\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 = \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1)$.

Proof.

$$\begin{aligned} (\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 &= ((f_n^3 \circ f_n^2) \circ f_n^1, n \in [1, N]) \\ &= (f_n^3 \circ (f_n^2 \circ f_n^1), n \in [1, N]) \\ &= \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1) \end{aligned}$$

□

{def:labcomp}

Definition 1.18 (Tuple composition for labeled morphisms). Let $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o_1^i, o_2^i)$, $i = 1, 2$, be tuples such that each $\mathbf{f}^i$ is a sequence of functions or each $\mathbf{f}^i$ is a tuple of functions, $\mathbf{range}(\mathbf{f}^1) = \mathbf{domain}(\mathbf{f}^2)$ and $o_2^1 = o_1^2$. Then

$$\mathbf{M}^2 \circ^A \mathbf{M}^1 \stackrel{\text{def}}{=} (\mathbf{f}^2 \circ \mathbf{f}^1, o_1^1, o_2^2) \quad (1.16)$$

{lem:labcomp}

Lemma 1.19 (Tuple composition for labeled morphisms). Let $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o_1^i, o_2^i)$, $i \in [1, 3]$, be a tuple such that $\mathbf{f}^i$ is a sequence or tuple of functions, $\mathbf{range}(\mathbf{f}^i) = \mathbf{domain}(\mathbf{f}^{i+1})$ and $o_2^i = o_1^{i+1}$, $i = 1, 2$. Then

$$\mathbf{M}^3 \circ^A (\mathbf{M}^2 \circ^A \mathbf{M}^1) = (\mathbf{M}^3 \circ^A \mathbf{M}^2) \circ^A \mathbf{M}^1 \quad (1.17)$$

Proof. From definition 1.18 (Tuple composition for labeled morphisms), lemma 1.14 (Sequence functions) on page 11 and lemma 1.16 (Tuple functions), we have

$$\begin{aligned} \mathbf{M}^3 \circ^A (\mathbf{M}^2 \circ^A \mathbf{M}^1) &= \mathbf{M}^3 \circ^A (\mathbf{f}^2 \circ^0 \mathbf{f}^1, o_1^1, o_2^2) \\ &= (\mathbf{f}^3 \circ^0 \mathbf{f}^2 \circ^0 \mathbf{f}^1, o_1^1, o_2^3) \\ &= (\mathbf{f}^3 \circ^0 \mathbf{f}^2, o_1^2, o_2^3) \circ^A \mathbf{M}^1 \\ &= (\mathbf{M}^3 \circ^A \mathbf{M}^2) \circ^A \mathbf{M}^1 \end{aligned}$$

□

Let $\mathbf{D}^i \stackrel{\text{def}}{=} \mathbf{domain}(\mathbf{f}^i)$ and $\mathbf{R}^i \stackrel{\text{def}}{=} \mathbf{range}(\mathbf{f}^i)$. Then

$$(\mathbf{Id}_{\mathbf{R}^i}, o_2^i, o_2^i) \circ^A \mathbf{M}^i = \mathbf{M}^i \quad (1.18)$$

$$\mathbf{M}^i \circ^A (\mathbf{Id}_{\mathbf{D}^i}, o_1^i, o_1^i) = \mathbf{M}^i \quad (1.19)$$

Proof.

$$\begin{aligned} (\mathbf{Id}_{\mathbf{R}^i}, o_2^i, o_2^i) \circ^A \mathbf{M}^i &= (\mathbf{Id}_{\mathbf{R}^i} \circ^0 \mathbf{f}^i, o_1^i, o_2^i) \\ &= (\mathbf{f}^i, o_1^i, o_2^i) \\ &= \mathbf{M}^i \end{aligned}$$

$$\begin{aligned} \mathbf{M}^i \circ^A (\mathbf{Id}_{\mathbf{D}^i}, o_1^i, o_1^i) &= (\mathbf{f}^i \circ^0 \mathbf{Id}_{\mathbf{D}^i}, o_1^i, o_2^i) \\ &= (\mathbf{f}^i, o_1^i, o_2^i) \\ &= \mathbf{M}^i \end{aligned}$$

□

{def:Cart}

Definition 1.20 (Cartesian product of sequence). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence and $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha < A)$ be a sequence of functions, then $\times \mathbf{S}^i \stackrel{\text{def}}{=} \times_{\alpha < A} S_\alpha^i$ is the generalized Cartesian product of the sequence $\mathbf{S}^i$ and $\times \mathbf{f}: \mathbf{S}^1 \longrightarrow \mathbf{S}^2 \stackrel{\text{def}}{=} \times_{\alpha < A} f_\alpha$ is the generalized Cartesian product of the function sequence $\mathbf{f}$.

Definition 1.21 (Topology functions). Let S and T be a topological spaces and Y a subset of S . Then

1. $S \overset{\text{subsp}}{\subseteq} T$ iff S is a subspace of T .
2. $S \overset{\text{open-subsp}}{\subseteq} T$ iff S is an open subspace of T .
3. $\mathbf{Set}(S)$ is the underlying set of S .
4. $\mathbf{Top}(S)$ is the topology of S .
5. $\mathbf{Top}(Y, S) \stackrel{\text{def}}{=} \{U \cap Y \mid U \in \mathbf{Top}(S)\}$ is the relative topology of Y .
6. $\mathbf{Top}(Y, S) \stackrel{\text{def}}{=} (Y, \mathbf{Top}(Y, S))$ is Y with the relative topology.
7. $S \overset{\text{def}}{\underset{\text{op}}{=}} \{(U, \mathbf{Top}(U, S)) \mid U \in \mathbf{Top}(S) \setminus \{\emptyset\}\}$ is the set of all non-null open subspaces of S .

Let $\mathbf{S}$ be a set of topological spaces. Then $\mathbf{S} \overset{\text{def}}{\underset{\text{op}}{=}} \bigcup_{S \in \mathbf{S}} S$ is the set of open subspaces in $\mathbf{S}$.

Let S and T' be spaces, $T \subseteq T'$ be a subspace and $f: S \longrightarrow T$ a function. Then $f: S \longrightarrow T' \stackrel{\text{def}}{=} \text{Id}_{T'}^T \circ f$ is f considered as a function from S to T' .

Let S' and T' be spaces, $S \subseteq S'$, $T \subseteq T'$ be subspaces and $f': S' \longrightarrow T'$ a function such that $f'[S] \subseteq T$. Then $f': S \longrightarrow T$, also written $f' \upharpoonright_{S, T}$, is $f' \upharpoonright_S$ considered as a function from S to T .

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha \leq A)$, $i = 1, 2$, be a sequence of spaces, $\mathbf{S}^1 \overset{0}{\subseteq} \mathbf{S}^2$ and $f^2: \text{head}(\mathbf{S}^2) \longrightarrow \text{tail}(\mathbf{S}^2)$ a function. $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)} \stackrel{\text{def}}{=} f^2 \upharpoonright_{\times \text{head}(\mathbf{S}^1)}$. If $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)} [\times \text{head}(\mathbf{S}^1)] \subseteq \text{tail}(\mathbf{S}^1)$ then $f^2: \text{head}(\mathbf{S}^1) \longrightarrow \text{tail}(\mathbf{S}^1)$, also written $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1), \text{tail}(\mathbf{S}^1)}$, is $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)}$ considered as a function from $\times \text{head}(\mathbf{S}^1)$ to $\text{tail}(\mathbf{S}^1)$. {def:seqin}

Definition 1.22 (Sequence inclusion). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ and $\mathbf{T} \stackrel{\text{def}}{=} (T_\alpha, \alpha < A)$ be sequences. $\mathbf{S} \overset{0}{\in} \mathbf{T}$ iff $\left(\bigvee_{\alpha < A} S_\alpha \in T_\alpha\right)$ or $\left(\bigvee_{\alpha < A} S_\alpha \overset{\text{Ob}}{\in} T_\alpha\right)$. {lem:seqin}

Lemma 1.23 (Sequence inclusion). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ be a sequence, $\mathcal{T}^i \stackrel{\text{def}}{=} (\mathcal{T}_\alpha^i, \alpha < A)$, $i = 1, 2$, a sequence of categories, $\mathcal{T}^1 \overset{\text{cat}}{\subseteq} \mathcal{T}^2$ and $\mathbf{S} \overset{0}{\in} \mathcal{T}^1$. Then $\mathbf{S} \overset{0}{\in} \mathcal{T}^2$.

Proof. If $\left(\bigvee_{\alpha < A} S_\alpha \overset{\text{Ob}}{\in} \mathcal{T}_\alpha^1\right)$ then $\left(\bigvee_{\alpha < A} S_\alpha \overset{\text{Ob}}{\in} \mathcal{T}_\alpha^2\right)$. □




Figure 1: Uncompleted nearly commutative diagram

{fig:NCDb}

Part IV

Nearly commutative diagrams

The notion of commutative diagrams is very useful in, e.g., Algebraic Topology. Often one encounters commutative diagrams in which two outgoing terminal nodes can be connected by a bridging function such that the resulting diagram is still commutative. This paper uses the term nearly commutative to describe a restricted class of such diagrams.

{part:ncd}

Let $\mathcal{C}$ be a full topological category and D a tree with two branches, whose nodes are topological spaces U_i and V_j and whose links are continuous functions $f_i: U_i \rightarrow U_{i+1}$ and $f'_j: V_j \rightarrow V_{j+1}$ between the sets:

$$D \stackrel{\text{def}}{=} \{ U_0 = V_0 \xrightarrow{f_0} U_1, \dots, U_{m-1} \xrightarrow{f_{m-1}} U_m, \quad (1.20) \quad \{\text{eq:NCD}\} \\ U_0 = V_0 \xrightarrow{f'_0} V_1, \dots, V_{n-1} \xrightarrow{f'_{n-1}} V_n \}$$

with $U_0 = V_0, U_m \in \mathcal{C}$ and $V_n \in \mathcal{C}$, as shown in fig. 1.

{def:NCD}

Definition 1.24 (Nearly commutative diagrams in category $\mathcal{C}$). D is left (right) nearly commutative in category $\mathcal{C}$ iff the two final nodes are in $\mathcal{C}$ and there is a morphism

$\hat{f}: U_m \xrightarrow{\text{Ar}} V_n \in \mathcal{C}$ ($\hat{f}': V_n \xrightarrow{\text{Ar}} U_m \in \mathcal{C}$) making the graph a commutative diagram, as shown in fig. 2. D is nearly commutative in category $\mathcal{C}$ iff it is either left nearly commutative or right nearly commutative.

{fig:NCDa}

D is strongly nearly commutative in category $\mathcal{C}$ iff the morphism is an isomorphism of $\mathcal{C}$,




Figure 2: Completed nearly commutative diagram

{fig:NCDa}

Definition 1.25 (Nearly commutative diagrams in category $\mathcal{C}$ at a point). Let D be as above and x be an element of the initial node. D is (left,right,strongly) nearly commutative in $\mathcal{C}$ at x iff there are subobjects of the nodes such that the tree D' formed by replacing the nodes is (left,right,strongly) nearly commutative in $\mathcal{C}$, and x is in the new initial node, as shown in fig. 3 (Local nearly commutative diagram) on page 17 below:

{def:NCDp}

$$x \in U'_0 = V'_0 \quad (1.21) \quad \{\text{eq:NCD}\}$$

$$U'_i \subseteq U_i \quad (1.22)$$

$$V'_j \subseteq V_j \quad (1.23)$$

$$f_i \upharpoonright_{U'_i}: U'_i \longrightarrow U'_{i+1} \quad (1.24)$$

$$f'_j \upharpoonright_{V'_j}: V'_j \longrightarrow V'_{j+1} \quad (1.25)$$

$$\begin{aligned} D' &\stackrel{\text{def}}{=} \{U'_0 = V'_0 \xrightarrow{f_0} U'_1, \dots, U'_{m-1} \xrightarrow{f_{m-1}} U'_m, \\ &\quad U'_0 = V'_0 \xrightarrow{f'_0} V'_1, \dots, V'_{n-1} \xrightarrow{f'_{n-1}} V'_n\} \end{aligned} \quad (1.26) \quad \{\text{eq:NCDp}\}$$

{def:NCD1}

Definition 1.26 (Locally nearly commutative diagrams in category $\mathcal{C}$). Let D be as above. D is (left,right,strongly) locally nearly commutative in $\mathcal{C}$ iff it is (left,right,strongly) nearly commutative in $\mathcal{C}$ at x for every x in the initial node.

Remark 1.27. It will often be clear from context what the relevant categories are. This paper may use the term “nearly commutative” without explicitly identifying the categories in which the nodes are found.




Figure 3: Local nearly commutative diagram

{fig:NCD1}

Lemma 1.28 (Locally nearly commutative diagrams in category $\mathcal{C}$). Let $\mathcal{C}$ and D be as above. If $U_0 = \emptyset$ then D is (left, right, strongly) locally nearly commutative in $\mathcal{C}$.

{lem:NCD}

Proof. D is vacuously locally nearly commutative at every $x \in U_0$ since there is no such x . $\square$

Let D be locally nearly commutative in $\mathcal{C}$ and let $\hat{U}_0 = \hat{V}_0 \subseteq U_0$ and $\hat{D}$ be D with $U_0 = V_0$ replaced by $\hat{U}_0 = \hat{V}_0$. Then $\hat{D}$ is locally nearly commutative in $\mathcal{C}$.

Proof. If $x \in \hat{U}_0$ then $x \in U_0$ and hence D is locally nearly commutative in $\mathcal{C}$ at x . Replacing U'_0 with $U'_0 \cap \hat{U}_0$ in the definition shows that $\hat{D}$ is locally nearly commutative in $\mathcal{C}$ at x . $\square$

{lem:NCDsubcat}

Lemma 1.29 (Nearly commutative diagrams in category $\mathcal{C}'$). Let $\mathcal{C} \subseteq^{\text{cat}} \mathcal{C}'$ and D be as in diagram (1.20) above.

If D is (left, right, strongly) nearly commutative in $\mathcal{C}$ then D is (left, right, strongly) commutative in $\mathcal{C}'$.

Proof. If $U_m \in^{\text{Ob}} \mathcal{C}$ then $U_m \in^{\text{Ob}} \mathcal{C}'$. If $V_n \in^{\text{Ob}} \mathcal{C}$ then $V_n \in^{\text{Ob}} \mathcal{C}'$. If $\hat{f}: U_m \xrightarrow{\sim} V_n \in^{\text{Ar}} \mathcal{C}$ then $\hat{f} \in^{\text{Ar}} \mathcal{C}'$ $\square$

Corollary 1.30 (Locally nearly commutative diagrams in category $\mathcal{C}'$ at a point). Let $\mathcal{C} \stackrel{\text{cat}}{\subseteq} \mathcal{C}'$ and D be as in diagram (1.20) above.

If D is locally nearly commutative in $\mathcal{C}$ at x then D is locally nearly commutative in $\mathcal{C}'$ at x .

Proof. Let U'_i, V'_j and diagram (1.26) be as in definition 1.25 (Nearly commutative diagrams in category $\mathcal{C}$ at a point) on page 16, as shown in diagram fig. 3 (Local nearly commutative diagram). D' is nearly commutative in category $\mathcal{C}$, and hence is nearly commutative in category $\mathcal{C}$. $\square$

Corollary 1.31 (Locally nearly commutative diagrams in category $\mathcal{C}'$). Let $\mathcal{C} \stackrel{\text{cat}}{\subseteq} \mathcal{C}'$ and D be as in diagram (1.20) above.

If D is locally nearly commutative in $\mathcal{C}$ then D is locally nearly commutative in $\mathcal{C}'$.

Proof. Let $x \in U_0 = V_0$. D is locally nearly commutative in $\mathcal{C}$ at x , hence locally nearly commutative in $\mathcal{C}'$ at x . $\square$

Part V

Model spaces and allied notions

Let S be a topological space. We need to formalize the notions of an open cover by sets that are “well behaved” in some sense, e.g., convex, sufficiently small, and of “well behaved” functions among those sets, e.g., preserving fibers, smooth. We do this by associating a category of acceptable sets and functions.

Remark 1.32. Using pseudo-groups, as in [Kobayashi, 1996, p. 1], would not allow restricting model neighborhoods to, e.g., convex sets.

2 Model spaces

Definition 2.1 (Model spaces). Let S be a topological space and $\mathcal{S}$ a small category whose objects are open subsets of S and whose morphisms are continuous functions. $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is a model space for S iff

1. $\text{Ob}(\mathcal{S})$ is an open cover for S . Note that it need not be a basis for S .
2. $\text{Ob}(\mathcal{S})$ is closed under finite intersections.
3. The morphisms of $\mathcal{S}$ are continuous functions in S .
4. If $f: A \rightarrow B$ is a morphism, $A' \in \mathcal{S} \subseteq A \stackrel{\text{Ob}}{\in} \mathcal{S}, B' \in \mathcal{S} \subseteq B \stackrel{\text{Ob}}{\in} \mathcal{S}$ and $f[A'] \subseteq B'$ then $f|_{A'}: A' \rightarrow B'$ is a morphism.

5. If $A' \in \mathcal{S}^{\text{Ob}} \subseteq A \in \mathcal{S}$ then the inclusion map $\text{Id}_A^{A'} : A' \hookrightarrow A$ is a morphism. {mod:inclusion}
Remark 2.2. This is actually a consequence of item 4, but it is convenient to give it here. {mod:restriction}

6. Restricted sheaf condition: informally, consistent morphisms can be glued together. Whenever {mod:sheaf}

- (a) U_α and $V_\alpha, \alpha < A$, are objects of $\mathcal{S}$.
- (b) $f_\alpha : U_\alpha \longrightarrow V_\alpha$ are morphisms of $\mathcal{S}$.
- (c) $U = \bigcup_{\alpha < A} U_\alpha$ is an object of $\mathcal{S}$.
- (d) $V = \bigcup_{\alpha < A} V_\alpha$ is an object of $\mathcal{S}$.
- (e) $f : U \longrightarrow V$ is a continuous function and for every $\alpha < A$, f agrees with f_α on U_α

then f is a morphism of $\mathcal{S}$.

By abuse of language we write “ $U \in \mathcal{S}$ ” for “ $U \in \text{Cat}\mathcal{S}^{\text{Ob}}$ ” and “ $f \in \mathcal{S}$ ” for “ $f \in \text{Cat}\mathcal{S}^{\text{Ar}}$ ”.

Remark 2.3. A model space is similar to a pseudo-group, but has some important differences:

- 1. Open subsets of objects of $\mathcal{S}$ need not be objects of $\mathcal{S}$.
- 2. The morphisms of $\mathcal{S}$ need not be homeomorphisms or even 1-1, thus they need not have inverses.
- 3. The morphisms of $\mathcal{S}$ need not satisfy the sheaf condition, but only the restricted sheaf condition in item 6. {mod:sheaf}

$\mathcal{S}$ is fine grained iff every open subset of an object of $\mathcal{S}$ is an object of $\mathcal{S}$. Note that it is not sufficient that $\text{Ob}(\mathcal{S})$ be a basis for $\mathcal{S}$.

$\text{Top}(\mathcal{S}) \stackrel{\text{def}}{=} S = \pi_1(\mathcal{S})$ is the topological space of $\mathcal{S}$.

Let $\mathcal{S}^i, i = 1, 2$, be a model space. $\text{Cat}(\mathcal{S}^i) \stackrel{\text{def}}{=} \mathcal{S}^i = \pi_2(\mathcal{S}^i)$ is the category of model space $\mathcal{S}^i$. By abuse of language we write $\mathcal{S}^1 \stackrel{\text{subsp}}{\subseteq} \mathcal{S}^2$ ($\mathcal{S}^1 \stackrel{\text{open-subsp}}{\subseteq} \mathcal{S}^2$) for

$\text{Top}(\mathcal{S}^1) \stackrel{\text{subsp}}{\subseteq} \text{Top}(\mathcal{S}^2)$ ($\text{Top}(\mathcal{S}^1) \stackrel{\text{open-subsp}}{\subseteq} \text{Top}(\mathcal{S}^2)$).

Let $U \in \mathcal{S}$. Then $\text{Top}(U, \mathcal{S}) \stackrel{\text{def}}{=} \text{Top}(U, S)$ is U with the relative topology.

$\mathcal{Top}(\mathcal{S}) \stackrel{\text{def}}{=} \left(\left\{ \text{Top}(U, S) \mid U \in \mathcal{S}^{\text{Ob}} \right\}, \text{Ar}(\mathcal{S}) \right)$ is the topological category of $\mathcal{S}$.

By abuse of language we write $U \subseteq \mathcal{S}$ for $U \in \mathcal{S}$, $\mathcal{Top}(\mathcal{S})$ for $\mathcal{Top}(\text{Top}(\mathcal{S}))$, f is a morphism of $\mathcal{S}$ for f is a morphism of $\mathcal{S}$ and f is an isomorphism of $\mathcal{S}$ for f is an isomorphism of $\mathcal{S}$.

Let $\mathcal{C}$ be a set of model spaces. $\mathcal{C}$ is fine grained iff every $(C, \mathcal{C}) \in \mathcal{C}$ is fine grained.

Lemma 2.4 (The topological category of a model space is a full topological category). {lem:modtopcat}

Let $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then $\mathcal{Top}(\mathbf{M})$ is a full topological category.

Proof. $\mathcal{Top}(\mathbf{M})$ is a small subcategory of $\mathbf{Top}$ by construction. $\mathcal{Top}(\mathbf{M})$ is a full topological category by item 4 of definition 2.1. □ {mod:restriction}

Definition 2.5 (Model neighborhoods). Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then the objects of $\mathcal{S}$ are model neighborhoods of $\mathbf{S}$. If $u \in U \stackrel{\text{Ob}}{\in} \mathcal{S}$ then U is a model neighborhood of $\mathbf{S}$ for u . If $U_i \stackrel{\text{Ob}}{\in} \mathcal{S}$, $i = 1, 2$, and $U_1 \subseteq U_2$ then U_1 is a model subneighborhood of U_2 ; if $u^1 \in U_1$ then U_1 is also a model subneighborhood for u^1 .

If $\emptyset \stackrel{\text{Ob}}{\in} \mathcal{S}$ then $\emptyset$ is a degenerate model neighborhood of $\mathbf{S}$; any nonvoid model neighborhood of $\mathbf{S}$ is a nondegenerate model neighborhood of $\mathbf{S}$. {def:modsub}

Definition 2.6 (Model subspaces). $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is a model subspace of $\mathbf{T} \stackrel{\text{def}}{=} (T, \mathcal{T})$, abbreviated $\mathbf{S} \stackrel{\text{mod}}{\subseteq} \mathbf{T}$, iff

1. $\mathbf{S}$ and $\mathbf{T}$ are model spaces

2. S is a subspace of T {modsub:sub}

3. $\mathcal{S} \stackrel{\text{cat}}{\subseteq} \mathcal{T}$ {modsub:subcat}

4. every intersection of S with an object of $\mathcal{T}$ is an object of $\mathcal{S}$ {modsub:rest}

$$\left(\bigvee_{U \stackrel{\text{Ob}}{\in} \mathcal{T}} \right) U \cap S \stackrel{\text{Ob}}{\in} \mathcal{S} \quad (2.1)$$

{modsub:full}

5. Every object of $\mathcal{T}$ contained in S is an object of $\mathcal{S}$

$$\left(\bigvee_{U \stackrel{\text{Ob}}{\in} \mathcal{T} : U \subseteq S} \right) U \stackrel{\text{Ob}}{\in} \mathcal{S} \quad (2.2)$$

Remark 2.7. This is actually a consequence of item 4, but it is convenient to state it here. {modsub:rest}

$\mathbf{S}$ is also a full model subspace of $\mathbf{T}$, abbreviated $\mathbf{S} \stackrel{\text{full-mod}}{\subseteq} \mathbf{T}$, iff $S = T$.

$\mathbf{S}$ is also a strict model subspace of $\mathbf{T}$, abbreviated $\mathbf{S} \stackrel{\text{strict-mod}}{\subseteq} \mathbf{T}$, iff S is a model neighborhood of $\mathbf{T}$.

Remark 2.8. S is also a model neighborhood of $\mathbf{S}$ by item 5. {modsub:full}

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space and T a model neighborhood of $\mathbf{S}$. Then $\text{Mod}(T, \mathbf{S})$, the relative model space of T , is $(T, \mathcal{T})$, where $\mathcal{T}$ is the full subcategory of $\mathcal{S}$ containing all model subneighborhoods of T .

By abuse of language we write T for $\text{Mod}(T, \mathbf{S})$ when $\mathbf{S}$ is understood by context.

{lem:modsub}

Lemma 2.9 (Model subspaces). 1. $\overset{\text{mod}}{\subseteq}$, $\overset{\text{full-mod}}{\subseteq}$ and $\overset{\text{strict-mod}}{\subseteq}$ are transitive.

Proof. Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$ $i = 1, 2, 3$, be a model space. If $\mathbf{S}^i \overset{\text{mod}}{\subseteq} \mathbf{S}^{i+1}$, $i = 1, 2$, then

- (a) $\mathbf{S}^1$ and $\mathbf{S}^3$ are model spaces by hypothesis.
- (b) Since S^i is a subspace of S^{i+1} , $i = 1, 2$, then S^1 is a subspace of S^3 .
- (c) Each $\mathcal{S}^i \overset{\text{cat}}{\subseteq} \mathcal{S}^{i+1}$ by [item 3 of definition 2.6 \(Model subspaces\)](#) above. {def:modsub}
Then $\mathcal{S}^1 \overset{\text{cat}}{\subseteq} \mathcal{S}^3$.
- (d) Each intersection of S^i with an object of $\mathcal{S}^{i+1}$ is an object of $\mathcal{S}^i$ by [item 4 of definition 2.6](#). Then every intersection of S^1 with an object of $\mathcal{S}^3$ is an object of $\mathcal{S}^1$. {def:modsub}
- (e) Let $U \overset{\text{Ob}}{\in} \mathbf{S}^3$ and $U \subseteq S^1$. $U \cap S^1 = U \overset{\text{Ob}}{\in} \mathbf{S}^1$.

If each $\mathbf{S}^i \overset{\text{full-mod}}{\subseteq} \mathbf{S}^{i+1}$, then each $S^i = S^{i+1}$ by [definition 2.6](#); then $S^1 = S^3$ {def:modsub}
and thus $\mathbf{S}^1 \overset{\text{full-mod}}{\subseteq} \mathbf{S}^3$.

If each $\mathbf{S}^i \overset{\text{strict-mod}}{\subseteq} \mathbf{S}^{i+1}$, then each S^i is a model neighborhood of $\mathbf{S}^{i+1}$ by [definition 2.6](#), and S^1 is a model neighborhood of $\mathbf{S}^3$ by [item 5 of definition 2.6](#). {def:modsub}
{def:modsub} $\square$

- 2. Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space and $T \subseteq S$. Then $\text{Mod}(T, \mathbf{S}) \overset{\text{mod}}{\subseteq} \mathbf{S}$

Proof. Let $\mathbf{T} = ((T, \mathfrak{T}), \mathcal{T}) \stackrel{\text{def}}{=} \text{Mod}(T, \mathbf{S})$.

- (a) $\mathbf{S}$ and $\mathbf{T}$ are model spaces.
- (b) T is a subspace of S .
- (c) $\mathcal{T} \overset{\text{cat}}{\subseteq} \mathcal{S}$ by construction.
- (d) Let $U \overset{\text{Ob}}{\in} \mathbf{S}$. $U \cap T \overset{\text{Ob}}{\in} \mathbf{S}$. Since $U \cap T \subseteq T$ is a model neighborhood of $\mathbf{S}$, it is a model neighborhood of $\text{Mod}(T, \mathbf{S})$ by construction.
- (e) Let $U \overset{\text{Ob}}{\in} \mathbf{S}$ be contained in T . Then U is a model neighborhood of $\text{Mod}(T, \mathbf{S})$ by construction.

$\square$

- 3. Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, and $\mathbf{S}^1 \overset{\text{mod}}{\subseteq} \mathbf{S}^2$. Then S^1 is an open subspace of S^2 .

Proof. The model neighborhoods of S^1 are model neighborhoods of S^2 , hence open in S^2 . Their union, which is open in S^2 , is S^1 by item 1 of definition 2.6 (Model subspaces) on page 20. $\square$ {mod:cover}

4. Let $S^i, i = 1, 2$, be a topological space and $S^1 \stackrel{\text{open-subsp}}{\subseteq} S^2$. Then $\text{Id}_{S^2}^{S^1}$ is open and continuous.

Proof. Let U be open in S^1 . Since $S^1 \stackrel{\text{open-subsp}}{\subseteq} S^2$, there is a V open in S^2 such that $U = S^1 \cap V$. Since V and S^1 are open in S^2 , so is $U = S^1 \cap V$.

Let V be open in S^2 . Since $S^1 \stackrel{\text{subsp}}{\subseteq} S^2$, $\text{Id}_{S^2}^{S^1}[V] = V \cap S^1$ is open in S^1 . $\square$

{cor:modsub}

Corollary 2.10 (Model subspaces). Let $S^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i), i = 1, 2, S'^2 \stackrel{\text{def}}{=} (S'^2, \mathcal{S}'^2)$ be model spaces, and $S^1 \stackrel{\text{mod}}{\subseteq} S^2 \stackrel{\text{mod}}{\subseteq} S'^2$.

1. $S^1 \stackrel{\text{open-subsp}}{\subseteq} S'^2$.

Proof. $S^1 \stackrel{\text{open-subsp}}{\subseteq} S^2 \stackrel{\text{open-subsp}}{\subseteq} S'^2$. $\square$

2. Let $f: S^1 \rightarrow S^2$ be continuous. Then $f: S^1 \rightarrow S'^2$ is continuous.

Proof. $S^1 \stackrel{\text{open-subsp}}{\subseteq} S'^2$, Let $V \subseteq S'^2$ be an arbitrary open set. Since $S^2 \stackrel{\text{mod}}{\subseteq} S'^2$ by hypothesis, $V \cap S^2$ is open in S^2 . Since $f: S^1 \rightarrow S^2$ is continuous by hypothesis, $f^{-1}[V] = f^{-1}[V \cap S^2]$ is open in S^1 . $\square$

3 M-nearly commutative diagrams

{sec:M-NCD}

Let $M \stackrel{\text{def}}{=} (M, \mathcal{M})$ be a model space and D a tree with two branches, whose nodes are topological spaces U_i and V_j and whose links are continuous functions $f_i: U_i \rightarrow U_{i+1}$ and $f'_j: V_j \rightarrow V_{j+1}$ between the spaces:

$$D \stackrel{\text{def}}{=} \{U_0 = V_0 \xrightarrow{f_0} U_1, \dots, U_{m-1} \xrightarrow{f_{m-1}} U_m, \\ U_0 = V_0 \xrightarrow{f'_0} V_1, \dots, V_{n-1} \xrightarrow{f'_{n-1}} V_n\}$$

with $U_0 = V_0, U_m \in \mathcal{M}$ and $V_n \in \mathcal{M}$, as shown in fig. 1 (Uncompleted nearly commutative diagram) on page 15. {fig:NCDb}

{def:M-NCD}

Definition 3.1 (M-nearly commutative diagrams). D is (left, right, strongly) M-nearly commutative in model space M iff D is (left, right, strongly) nearly commutative in category $\mathcal{M}$.

Definition 3.2 (M-nearly commutative diagrams at a point). Let $\mathbf{M}$ and D be as above and x be an element of the initial node. D is (left,right,strongly) M-nearly commutative in $\mathbf{M}$ at x iff D is (left,right,strongly) nearly commutative in category $\mathcal{M}$ at x . {def:M-NCDp}

Definition 3.3 (M-locally nearly commutative diagrams). Let $\mathcal{C}$ and D be as above. D is (left,right,strongly) M-locally nearly commutative in $\mathbf{M}$ iff D is (left,right,strongly) nearly commutative in category $\mathcal{M}$ at every point. {def:M-NCD1}

4 Trivial model spaces

Informally, a trivial model space of a specific type is one that does not restrict the potential objects and morphisms of its type. {sec:triv}

Definition 4.1 (Trivial model spaces). Let S be a topological space and $\mathcal{S}$ the category of all continuous functions between open sets of S . {def:trivmod}

$\mathcal{S}_{\text{triv}}^{\text{def}}(S, \mathcal{S})$ is the trivial model space of S and $\mathcal{S}_{\text{triv}}$ is a trivial model space.

Let $\mathbf{S}$ be a set of topological spaces.

$\mathcal{S}_{\text{triv}}^{\text{def}} = \left\{ \mathcal{S}_{\text{triv}}' \mid S' \in \mathbf{S} \right\}$ is the set of all trivial model spaces in $\mathbf{S}$.

$\mathcal{S}_{\text{triv}}$, the category of all continuous functions between elements of $\mathcal{S}_{\text{triv}}$, is the trivial model category of $\mathbf{S}$.

$\mathcal{S}_{\text{op-Triv}}$, the category of open trivial model spaces in $\mathbf{S}$, is the category whose

objects are $\left\{ U_{\text{triv}} \mid U \in \mathcal{S}_{\text{op}} \right\}$, the trivial model spaces of non-null open sets of spaces in $\mathbf{S}$, and whose morphisms are all the continuous functions among them. {lem:trivmod}

Lemma 4.2 (Trivial model spaces). Let $S^i, i = 1, 2$, be a topological space.

1. $\mathcal{S}_{\text{triv}}^i$ is a model space.

Proof. Let $\mathcal{S}^i \stackrel{\text{def}}{=} \text{Cat} \left(\mathcal{S}_{\text{triv}}^i \right), f: A \longrightarrow B$ be a morphism of $\mathcal{S}^i, A' \in \mathcal{S}^i \subseteq A$ and $B' \in \mathcal{S}^i \subseteq B$. Then the definitions of continuity and of $\mathcal{S}_{\text{triv}}^i$ imply each of the following.

- (a) $\text{Ob}(\mathcal{S}^i) = \mathfrak{Top}(S^i)$ and thus is an open cover of S^i .
- (b) $\text{Ob}(\mathcal{S}^i) = \mathfrak{Top}(S^i)$ and thus is closed under finite intersections.
- (c) All morphisms in $\text{Ar}(\mathcal{S}^i)$ are continuous.
- (d) Since $f: A \longrightarrow B$ is a morphism of $\mathcal{S}^i, f$ is continuous.
- (e) Since $A' \in \mathcal{S}^i$ then A' is open.

- (f) Since $B' \in \mathcal{S}^i$ then B' is open.
- (g) Since $f: A \rightarrow B$ is continuous then $f|_{A'}: A' \rightarrow B$ is continuous.
- (h) Since $f[A'] \subseteq B'$ then $f|_{A', B'}: A' \rightarrow B'$ is well defined and continuous, hence a morphism.
- (i) If $A' \in \mathcal{S}^i \subseteq A \in \mathcal{S}^i$ then
 - i. A and A' are open.
 - ii. The inclusion map $\text{Id}_A^{A'}$ is continuous.
 - iii. The inclusion map $\text{Id}_A^{A'}$ is a morphism of $\mathcal{S}^i$ by the definition of $\mathcal{S}_{\text{triv}}^i$.
 - iv. $\mathcal{S}_{\text{triv}}^i$ trivially satisfies the restricted sheaf condition because every continuous function between open sets of S^i is a morphism of $\mathcal{S}^i$.

□

2. Let S^1 be an open subspace of S^2 . Then $\mathcal{S}_{\text{triv}}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathcal{S}_{\text{triv}}^2$.

Proof. Let $\mathcal{S}^i \stackrel{\text{def}}{=} \text{Cat}(\mathcal{S}_{\text{triv}}^i)$, $i = 1, 2$. By definition 4.1 (Trivial model spaces) on page 23, $\mathcal{S}^i$ is the category of all continuous functions between open sets of S^i ,

{def:trivmod}

$\mathcal{S}_{\text{triv}}^1$ and $\mathcal{S}_{\text{triv}}^2$ satisfy the six conditions of definition 2.6 (Model subspaces) on page 20:

{def:modsub}

$\mathcal{S}_{\text{triv}}^1$ and $\mathcal{S}_{\text{triv}}^2$ are model spaces: Proved above.

$\mathcal{S}_{\text{triv}}^1$ is a subspace of $\mathcal{S}_{\text{triv}}^2$:

$\text{Top}(\mathcal{S}_{\text{triv}}^i) = S^i$ by definition 4.1 (Trivial model spaces) on page 23. S^1 is an open subspace of S^2 by hypothesis.

{def:trivmod}

$\mathcal{S}_{\text{triv}}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}_{\text{triv}}^2$:

If $U \in \mathcal{S}_{\text{triv}}^1$ then U is open in S^1 . Since S^1 is an open subspace of S^2 , U is open in S^2 and thus $U \in \mathcal{S}_{\text{triv}}^2$.

If $f: U \rightarrow V \in \mathcal{S}_{\text{triv}}^1$ then f is a continuous function between open sets of S^1 . Since S^1 is an open subspace of S^2 , f is a continuous function between open sets of S^2 and thus a morphism of $\mathcal{S}_{\text{triv}}^2$.

If $f: U \rightarrow V \in \mathcal{S}_{\text{triv}}^2$ then f is a continuous function between open sets of S^2 . If $U \in \mathcal{S}_{\text{triv}}^1$ and $V \in \mathcal{S}_{\text{triv}}^1$, then U and V are open sets of S^1 . Since S^1 is an open subspace of S^2 , f is a continuous function between open sets of S^1 and thus a morphism of $\mathcal{S}_{\text{triv}}^1$.

Every intersection of S^1 with an object of $\mathcal{S}^2$ is an object of $\mathcal{S}^1$:

Let $U \in \mathcal{S}^2$. U and S^1 are open in S^2 . $U/\text{cup} \text{Set}(S^1)$ is open in S^2 and contained in S^1 . Since S^1 is an open subspace of S^2 , $U/\text{cup} \text{Set}(S^1)$ is open in S^1 and thus an object of $\mathcal{S}^1$.

Every object of $\mathcal{S}^2$ contained in S^1 is an object of $\mathcal{S}^1$:

Let $U \in \mathcal{S}^2$. U is open in S^2 . If U is contained in S^1 then it is open in S^1 and thus an object of $\mathcal{S}^1$.

Since S^1 is an open subspace of S^2 , $\text{Set}(S^1)$ is a model neighborhood of S^2_{triv} by definition 4.1 above □ {def:trivmod}

$$S^1_{\text{triv}} \overset{\text{mod}}{\subseteq} S^2_{\text{triv}}.$$

Proof. $S^1 \overset{\text{open-subsp}}{\subseteq} S^2$ by lemma 2.9 (Model subspaces) on page 21. □ {lem:modsub}

3. Let $f: S^1 \longrightarrow S^2$ be continuous. Then f is a model function from S^1_{triv} to S^2_{triv} .

Proof. Let U^i be a model neighborhood of S^i_{triv} , $i = 1, 2$.

(a) Since U^2 is a model neighborhood of S^2_{triv} , U^2 is open, $f^{-1}[U^2]$ is open and thus a model neighborhood of S^1_{triv} .

(b) $f[U^1] \subseteq S^2$. Since S^2 is open, it is a model neighborhood of S^2_{triv} . □

5 Minimal model spaces

Even if a set of open sets fails one of items 1 and 2 or a set of functions among them fails one of items 3 to 6 in the definition of a model space, there is a minimal model space containing them. {sec:min}
{mod:cover,mod:closed}
{mod:continuous}

Definition 5.1 (Minimal model spaces). Let S be a topological space, $\mathcal{O}$ a set of open sets in S , $\mathcal{f}$ a set of continuous functions between elements of $\mathcal{O}$ and $\mathcal{S}$ the smallest concrete category over **Top** having all sets in $\mathcal{O}$ as objects, having all functions in $\mathcal{f}$ as morphisms and satisfying items 3 to 6 of definition 2.1 (Model spaces) on page 18. {def:minmod}
{mod:continuous}

Then

$$\text{Mod}_{\min}(S, \mathcal{O}, \mathcal{f}) \stackrel{\text{def}}{=} \left(\text{Top} \left(\bigcup \mathcal{O}, S \right), \mathcal{S} \right) \quad (5.1)$$

is the minimal model space of S with neighborhoods $\mathcal{O}$ and neighborhood mappings $\mathcal{f}$.

Remark 5.2. The trivial model space $\underset{\text{triv}}{S}$ is a special case.

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space, $\mathbf{O}$ a set of model neighborhoods of $\mathbf{S}$ and $\mathbf{f}$ a set of morphisms between elements of $\mathbf{O}$.

Then

$$\text{Mod}(\mathbf{S}, \mathbf{O}, \mathbf{f}) \stackrel{\text{def}}{=} \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f}) \quad (5.2)$$

is the minimal model space of $\mathbf{S}$ with neighborhoods $\mathbf{O}$ and neighborhood mappings $\mathbf{f}$.

Remark 5.3. $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is not, in general, a model subspace of $\mathbf{S}$, since modsub:rest, modsub:full items 4 and 5 of definition 2.6 (Model subspaces) on page 20 may fail. {def:modsub}

While every model neighborhood of $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is a model neighborhood of $\underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$, the converse need not be true. {lem:minmod}

Lemma 5.4 (Minimal model spaces are model spaces). Let S be a topological space, $\mathbf{O}$ a set of open sets in S and $\mathbf{f}$ a set of continuous functions between elements of $\mathbf{O}$. Then $(C, \mathcal{C}) \stackrel{\text{def}}{=} \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ is a model space.

Proof.

1. Finite intersections of open sets are open and $C = \bigcup_{U \in \mathbf{O}} U$ by construction
2. $\text{Ob}(\mathcal{C})$ is closed under finite intersections by construction
3. Compositions of continuous functions, inclusion maps and restrictions of continuous functions are continuous
4. Restrictions of morphisms are morphisms by construction
5. Inclusion maps are morphisms by construction
6. The restricted sheaf condition holds by construction

□

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space, $\mathbf{O}$ a set of model neighborhoods of $\mathbf{S}$ and $\mathbf{f}$ a set of morphisms between elements of $\mathbf{O}$. Then $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is a model space.

Proof. $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f}) = \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ and $\underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ is a model space. □

6 M-paracompact model spaces

Paracompactness is an important property for topological spaces because of partitions of unity. There is an analogous property for model spaces.

{sec:m-para}

Definition 6.1 (Model topology). Let $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then the model topology $\mathfrak{M}^*$ for M is the topology generated by $\text{Ob}(\mathcal{S})$.

{def:ModTop}

Remark 6.2. $\mathfrak{M}^*$ is not guaranteed to be T0 even if S is T4. However, $\mathfrak{M}^*$ may be normal or regular even if S is not.

{def:M-para}

Definition 6.3 (m-paracompactness). A model space $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is m-paracompact iff $\mathfrak{M}^*$ is regular and every cover of S by model neighborhoods has a locally finite refinement by model neighborhoods.

Remark 6.4. This is a stronger condition than merely requiring $\mathfrak{M}^*$ to be paracompact.

Theorem 6.5 (m-paracompactness and paracompactness). If $\mathbf{M} = (S, \mathcal{S})$ is m-paracompact and the model neighborhoods form a basis for S then S is paracompact.

Proof. Let R be an open cover of S . Since the model neighborhoods form a basis, every set in R is a union of model neighborhoods and thus there is a refinement R_1 by model neighborhoods. Since by hypothesis $\mathbf{M}$ is m-paracompact, R_1 has a locally finite refinement R_2 by model neighborhoods. Since model neighborhoods are open sets, R_2 is a locally finite refinement of R in the conventional sense. $\square$

7 Model functions and model categories

It is convenient to have a notion of mappings between model spaces that are well behaved in some sense, e.g., fiber preserving; using that notion it is then possible to group model spaces into categories.

{sec:modcat}

7.1 Model functions

Definition 7.1 (Model functions). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: S^1 \rightarrow S^2$ be a continuous function. f is a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff the inverse images of model neighborhoods are model neighborhoods.

{def:modfunc}

$$\left(\bigvee_{V \in \mathcal{S}^2}^{\text{Ob}} \right) f^{-1}[V] \in \mathcal{S}^1 \quad (7.1)$$

f is also a constrained model function of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff $f[S^1]$ is contained in a model neighborhood of $\mathbf{S}^2$.

$$\left(\bigvee_{V \in \mathcal{S}^2}^{\text{Ob}} \right) f[S^1] \subseteq V \quad (7.2)$$

Remark 7.2. When $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is constrained, S^1 is a model neighborhood of $\mathbf{S}^1$.

By abuse of language we write $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ both for f considered as a model function and for f considered as a continuous function.

Let $\mathbf{S}^1 \stackrel{\text{def}}{=} (S^1, \mathcal{S}^1)$ and $\mathbf{S}'^2 \stackrel{\text{def}}{=} (S'^2, \mathcal{S}'^2)$ be model spaces, $\mathbf{S}^2 \stackrel{\text{def}}{=} (S^2, \mathcal{S}^2) \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^2$ be a model subspace and $f: S^1 \longrightarrow S^2$ be a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$. Then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}'^2 \stackrel{\text{def}}{=} \text{Id}_{\mathbf{S}'^2} \circ f$ is f considered as a model function from $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Let $\mathbf{S}'^i \stackrel{\text{def}}{=} (S'^i, \mathcal{S}'^i)$, $i = 1, 2$, be a model space, $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^i$ be a model subspace and f' be a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$ such that $f'[S^1] \subseteq S^2$. Then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$, also written $f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$, is $f' \upharpoonright_{\mathbf{S}^1}$ considered as a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$. {lem:modfunc}

Lemma 7.3 (Model functions). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$ and $\mathbf{S}'^i \stackrel{\text{def}}{=} (S'^i, \mathcal{S}'^i)$, $i = 1, 2$, be model spaces, $\mathbf{S}^i \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^i$ and $f': S'^1 \longrightarrow S'^2$ be a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$.

$f \stackrel{\text{def}}{=} f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$ is a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. f' is a model function by hypothesis. f' is continuous by definition 7.1 above, so $f \stackrel{\text{def}}{=} f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$ is continuous. {def:modfunc}

Let U be a model neighborhood of $\mathbf{S}^2$. $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^2$ by hypothesis, so U is a model neighborhood of $\mathbf{S}'^2$. f' is a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$ by hypothesis, so $f'^{-1}[U] \in \mathbf{S}'^1$ by definition 7.1. Then $f^{-1}[U] = f'^{-1}[U] \cap S^1 \in \mathbf{S}^1$ by definition 7.1. {def:modfunc}

Lemma 7.4 (Model functions and trivial model spaces). Let S^i , $i = 1, 2$, be a topological space and $f: S^1 \longrightarrow S^2$ be a continuous function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ is a model function. {lem:modfunctriv}

Proof. Let U^i be a model neighborhood of $\underset{\text{triv}}{S^i}$, $i = 1, 2$.

$f^{-1}[U^2]$ is open, hence a model neighborhood of $\underset{\text{triv}}{S^1}$.

S^2 is a model neighborhood of $\underset{\text{triv}}{S^2}$ and $f[U^1] \subseteq U^2$. □

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ be a model function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ is a model function.

Proof. $f: S^1 \longrightarrow S^2$ is continuous. □

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ be a model function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ is a model function iff $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a constrained model function.

Proof. let U^2 be a model neighborhood of $\mathbf{S}^2$. $f^{-1}[U^2]$ is open, hence a model neighborhood of $\mathbf{S}^1$.

$\mathbf{S}^1$ is a model neighborhood of $\mathbf{S}^1$. If $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a model function then $f[\mathbf{S}^1]$ is contained in a model neighborhood of $\mathbf{S}^2$ and thus f is constrained.

If f is constrained, let U^1 be a model neighborhood of $\mathbf{S}^1$. Then $f[U^1] \subseteq f[\mathbf{S}^1]$ is contained in a model neighborhood of $\mathbf{S}^2$. $\square$

{lem:modcomp}

Lemma 7.5 (Composition of model functions). Let $\mathbf{S}_i \stackrel{\text{def}}{=} (S_i, \mathcal{S}_i)$, $i \in [1, 3]$, be a model space and $f_i: \mathbf{S}_i \rightarrow \mathbf{S}_{i+1}$, $i = 1, 2$, be a model function.

$f_2 \circ f_1$ is a model function.

Proof. Let $U_i \in \mathcal{S}_i$, $i = 1, 2, 3$.

Since f_2 is a model function, $f_2^{-1}[U_3] \in \mathcal{S}_2$. Since f_1 is a model function, $(f_2 \circ f_1)^{-1}[U_3] = f_1^{-1}[f_2^{-1}[U_3]] \in \mathcal{S}_1$. Since f_1 is a model function, $f_1[U_1]$ is contained in a model neighborhood V_2 . Since f_2 is a model function, $f_2[V_2]$ is contained in a model neighborhood V_3 . Then $(f_2 \circ f_1)[U_1] \subseteq V_3$. $\square$

If each f_i is constrained then $f_2 \circ f_1$ is constrained.

Proof. $f_1[\mathbf{S}^1] \subseteq \mathbf{S}^2$ and $f_2[\mathbf{S}^2] \subseteq \mathbf{S}^3$, hence $f_2 \circ f_1[\mathbf{S}^1] \subseteq \mathbf{S}^3$. $\square$

{def:modelhomeo}

Definition 7.6 (Model homeomorphisms). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ be a model function. f is a model homeomorphism iff it is also invertible and its inverse is a model function.

7.2 Model categories

{def:modcat}

Definition 7.7 (Model categories). A category $\mathcal{M}$ is a model category iff

1. the objects of $\mathcal{M}$ are model spaces
2. the morphisms of $\mathcal{M}$ are model functions.
3. composition is functional composition.
4. If $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S}) \in \mathcal{M}$ and $S^i \in \mathcal{S}$, $i = 1, 2$, then $\mathbf{S}^i \stackrel{\text{def}}{=} \text{Mod}(S^i, \mathbf{S}) \in \mathcal{M}$. If further $S^1 \subseteq S^2$ then the inclusion map $\text{Id}_{S^2}^{\mathbf{S}^1}: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{S}$.
5. If $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}$, $U^i \in \mathcal{S}^i$, $i = 1, 2$, $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{M}$ and $f[U^1] \subseteq U^2$ then $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}$.
6. If $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, is a subspace of $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S}) \in \mathcal{M}$ and $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{S}$ then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{M}$.

{modcat:inc}

{modcat:rest}

{modcat:morph}

A model category $\mathcal{M}$ satisfies the restricted sheaf condition iff whenever

1. U and V are objects of $\mathcal{S}$.
2. $U_\alpha \overset{\text{mod}}{\subseteq} U$ and $V_\alpha \overset{\text{mod}}{\subseteq} V$, $\alpha < A$, are objects of $\mathcal{M}$.
3. $\mathbf{Set}(U) = \bigcup_{\alpha < A} \mathbf{Set}(U_\alpha)$
4. $\mathbf{Set}(V) = \bigcup_{\alpha < A} \mathbf{Set}(V_\alpha)$
5. $f_\alpha: U_\alpha \longrightarrow V_\alpha$ are morphisms of $\mathcal{M}$.
6. $f: U \longrightarrow V$ is a model function and for every $\alpha < A$, f agrees with f_α on U_α

then f is a morphism of $\mathcal{M}$.

Let $\mathcal{M}^i$, $i = 1, 2$, be a model category. $\mathcal{M}^1$ is a strict subcategory of $\mathcal{M}^2$, abbreviated $\mathcal{M}^1 \overset{\text{strict-cat}}{\subseteq} \mathcal{M}^2$, iff $\mathcal{M}^1 \overset{\text{fullt-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2$ satisfies the restricted sheaf condition.

{lem:modcat}

Lemma 7.8 (Model categories). Let $\mathcal{M}$ be a model category, $\mathbf{S}^i \overset{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}$.

$i = 1, 2$, $U^i \in \mathcal{S}^i$, $\mathbf{U}^i \overset{\text{def}}{=} \text{Mod}(U^i, \mathbf{S}^i)$ and $f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \in \mathcal{M}$.

$f: \mathbf{U}^1 \longrightarrow \mathbf{S}^2 \overset{\text{Ar}}{\in} \mathcal{M}$.

Proof. Since $\text{Id}_{\mathbf{S}^2} \overset{\text{Ar}}{\in} \mathcal{M}$ by item 4 of definition 7.7 above and $f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \overset{\text{Ar}}{\in} \mathcal{M}$ by hypothesis, $f: \mathbf{U}^1 \longrightarrow \mathbf{S}^2 = \text{Id}_{\mathbf{S}^2} \circ f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \overset{\text{Ar}}{\in} \mathcal{M}$ $\square$

{modcat:inc}

{def:modtrivcat}

Definition 7.9 (Trivial model categories). Let $\mathbf{S} \overset{\text{def}}{=} (S, \mathcal{S})$ be a model space.

$\mathbf{S}_{\text{submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \mathbf{M} \overset{\text{mod}}{\subseteq} \mathbf{S} \right\}$ is the set of all subspaces of $\mathbf{S}$.

$\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}$, the category of all model functions between elements of $\mathbf{S}_{\text{submod-triv}}$, is the trivial subspace model category of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \mathbf{M} \overset{\text{full-mod}}{\subseteq} \mathbf{S} \right\}$ is the set of all full subspaces of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}$, the category of all full model functions between elements of $\mathbf{S}_{\text{full-submod-triv}}$, is the trivial full subspace model category of $\mathbf{S}$.

Let $\mathbf{S}$ be a set of model spaces.

$\mathbf{S}_{\text{submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \left(\exists_{\mathbf{M}' \in \mathbf{S}} \right) \mathbf{M} \overset{\text{mod}}{\subseteq} \mathbf{M}' \right\}$ is the set of all model subspaces of model spaces in $\mathbf{S}$.

$\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}$, the category of all model functions between elements of $\mathbf{S}_{\text{submod-triv}}$, is the trivial subspace model category of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-triv}} \stackrel{\text{def}}{=} \left\{ \mathbf{M} \mid \left(\exists_{\mathbf{M}' \in \mathbf{S}} \mathbf{M} \stackrel{\text{full-mod}}{\subseteq} \mathbf{M}' \right) \right\}$ is the set of all full subspaces in $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}$, the category of all full model functions between elements of $\mathbf{S}_{\text{full-submod-triv}}$, is the trivial full subspace model category of $\mathbf{S}$.

{lem:modtrivcat}

Lemma 7.10 (Trivial model categories). Let $\mathbf{S} \stackrel{\text{def}}{=} (\mathbf{S}, \mathcal{S})$ be a model space.

$$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}} \stackrel{\text{full-cat}}{\subseteq} \mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}.$$

Proof. $\left(\bigvee_{\mathbf{M} \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}} \mathbf{M} \right)^{\text{mod}} \subseteq \mathbf{S}$, so $\text{Ob}(\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}) \subseteq \text{Ob}(\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}})$.

It remains to show that for a pair of objects in both categories, the Hom sets are the same. Let $\mathbf{M}^i \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}$, $i = 1, 2$.

1. Let $f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}^{\text{Ar}}$. Then f is a model function and thus

$$f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{submod-triv}}^{\text{Ar}}.$$

2. Let $f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{submod-triv}}^{\text{Ar}}$. Then f is a model function and thus

$$f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}^{\text{Ar}}.$$

□

7.3 (Local) m-morphisms

{def:modlocalMmorph}

Definition 7.11 (Local m-morphisms). Let $\mathcal{M}^i$ be a model category, $i = 1, 2$, and $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. A function $f: U^1 \longrightarrow U^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 iff $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and for every $u \in U^1$ there is a model neighborhood $U_u \subseteq U^1$ for u and a model neighborhood $V_u \subseteq U^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$ be open. A function $f: U^1 \longrightarrow U^2$ is a local $\mathbf{S}^1$ m-morphism of U^1 to U^2 iff it is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

⁵This definition does not require that $\mathcal{S}^i$ be a model neighborhood of $\mathcal{S}^i$.

Let $U^i \in \mathcal{S}^i$, $i = 1, 2$. A function $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of U^1 to U^2 iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and for every $u \in U^1$ there is a model neighborhood $U_u \subseteq U^1$ for u and a model neighborhood $V_u \subseteq U^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: \text{Mod}(U_u, \mathbf{S}^1) \rightarrow \text{Mod}(V_u, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$, be open. A model function $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 iff it is a local $\mathbf{S}^i\text{-}\mathbf{S}^i\text{-}\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is a local $\mathcal{M}^i$ m-morphism iff it is a local $\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

The phrases “locally a(n) ... m-morphism” and “a local ... m-morphism” are equivalent.

{def:modMmorph}

Definition 7.12 (M-morphisms). Let $\mathcal{M}^i$ be a model category, $i = 1, 2$, and

$\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 iff $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and f is a morphism of $\mathbf{S}^2$.

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is an m-morphism of $\mathbf{S}^i$ iff it is an m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be model neighborhoods. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of U^1 to U^2 iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$, $i = 1, 2$, be model neighborhoods. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 iff it is an $\mathbf{S}^i\text{-}\mathbf{S}^i\text{-}\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an $\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and f is a morphism of $\mathcal{M}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is an $\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ iff it is an $\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

{lem:modMmorph}

Lemma 7.13 ((Local) m-morphisms). Let $\mathcal{M}^i, \mathcal{M}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^i$, $i = 1, 2$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$ and $\mathbf{S}^i \stackrel{\text{mod}}{\subseteq} \mathbf{S}^i$.

{M-morph:openmorphislocal}

1. Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 .

Proof.

- (a) Since f is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 , $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and f is a morphism of $\mathbf{S}^2$,

(b) U^1 and U^2 are model neighborhoods of $\mathbf{S}^2$. U^1 is a model neighborhood of $\mathbf{S}^1$ by item 5 of ^{def:modsub}definition 2.6 (Model subspaces) on page 20.

{modsub:full}

(c) Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f(u^1) \in U^2$, Then U^1 is a model neighborhood for u^1 and U^2 is a model neighborhood for u^2 . $f[U^1] \subseteq U^2$ by hypothesis. $f: U^1 \rightarrow U^2$ is a morphism of $\mathbf{S}^2$ as shown above.

□

{M-morph:openlocalismorph}

2. Let $U^i \subseteq \mathbf{S}^i$ be a model neighborhood of $\mathbf{S}^i$, $i = 1, 2$. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ local m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .

Proof. $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ by hypothesis. U^1 is a model neighborhood of $\mathbf{S}^1$ by hypothesis, thus a model neighborhood of $\mathbf{S}^2$

For each $u \in U^1$, let U_u be a model subneighborhood of U^1 for u and V_u be a model subneighborhood of U^2 for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \xrightarrow{\text{mod:sheaf}} V_u$ is a morphism of $\mathbf{S}^2$. Then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathbf{S}^2$ by item 6 of definition 2.1 (Model spaces) on page 18.

□

{def:model}

{M-morph:opencatmorphisloc}

3. Let $U^i \subseteq \mathbf{S}^i$ be a model neighborhood, $i = 1, 2$. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 .

Proof.

(a) Each U^i is a model neighborhood of $\mathbf{S}^i$ by hypothesis.

(b) Since $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 ,

$\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$ by definition 7.12 (M-morphisms) on page 31.

{def:modMmorph}

(c) Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f(u^1) \in U^2$.

i. Each U^i is a model neighborhood of $\mathbf{S}^i$ for u^i ,

ii. Each $U^i \subseteq U^i$.

iii. $f[U^1] \subseteq U^2$.

□

{M-morph:opencatlocalismorph}

4. Let $\mathcal{M}^2$ satisfy the restricted sheaf condition and each $U^i \subseteq \mathbf{S}^i$ be a model neighborhood, $i = 1, 2$. Every $f: U^1 \rightarrow U^2$ that is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ local m-morphism of U^1 to U^2 is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 .

Proof.

(a) Since each U^i is a model neighborhood of $\mathbf{S}^i$ by hypothesis, each

$\text{Mod}(U^i, \mathbf{S}^i) \in \mathcal{M}^i$ by item 4 of definition 7.7

{modcat:inc}

(b) $f: \text{Mod}(U^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a model function by definition 7.11

{def:modlocalMmorph}

(c) Each $\mathbf{S}^i \in \mathcal{M}^i$ by hypothesis

(d) $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$ by hypothesis

(e) $\mathcal{M}^2$ satisfies the restricted sheaf condition by hypothesis

(f) For every $u^1 \in U^1$ there is a model neighborhood $U_{u^1}^1$ for u^1 and a model neighborhood $U_{u^1}^2$ for $u^2 \stackrel{\text{def}}{=} f(u^1)$ such that $f[U_{u^1}^1] \subseteq U_{u^1}^2$ and $f: \text{Mod}(U_{u^1}^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U_{u^1}^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^i$.

i. $\text{Mod}(U_{u^1}^1, \mathbf{S}^1) \subseteq^{\text{mod}} \mathbf{S}^1$.

ii. Each $f: \text{Mod}(U_{u^1}^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U_{u^1}^2, \mathbf{S}^2)$ agrees with f on $U_{u^1}^1$.

iii. Since $\bigcup_{u^1 \in U^1} U_{u^1}^1 = U^1$ and $\bigcup_{u^1 \in U^1} U_{u^1}^2 = U^2$, then $f \in \mathcal{M}^2$ by definition 7.7 (Model categories) on page 29.

{def:modcat}

□

{M-morph:localislocaltosup}

5. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. $f: \mathbf{S}^1 \longrightarrow \mathbf{S}'^2$ is continuous by corollary 2.10 (Model subspaces) on page 22.

{cor:modsub}

Let $u \in \mathbf{S}^1$, U_u be a model neighborhood of $\mathbf{S}^1$ for u and V_u be a model neighborhood of $\mathbf{S}^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}^2$.

Since $\mathbf{S}^2 \subseteq^{\text{mod}} \mathbf{S}'^2$ by hypothesis, $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}'^2$.

□

{M-morph:morphismorphptosup}

6. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$, then f is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2$ by definition 7.12 (M-morphisms) on page 31 and $\mathbf{S}^2 \subseteq^{\text{mod}} \mathbf{S}'^2$ by hypothesis, so $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}'^2$.

{def:modMmorph}

f is a morphism of $\mathbf{S}^2$ by definition 7.12 (M-morphisms) on page 31, hence a morphism of $\mathbf{S}'^2$.

{def:modMmorph}

□

{M-morph:catlocalislocaltosup}

7. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$, then f is a local $\mathcal{M}'^1$ - $\mathcal{M}'^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. Let $u \in S^1$, U_u be a model neighborhood of S^1 for u and V_u be a model neighborhood of S^2 for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \rightarrow V_u$ is a morphism of S^2 .

Since $S^2 \stackrel{\text{full-cat}}{\subseteq} S'^2$ by hypothesis, $f: U_u \rightarrow V_u$ is a morphism of S'^2 . $\square$

{M-morph: catmorphismphtos}

8. If $f: S^1 \rightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of S^1 to S^2 , then f is an $\mathcal{M}'^1$ - $\mathcal{M}'^2$ m-morphism of S^1 to S'^2 .

Proof. $S^1 \stackrel{\text{full-cat}}{\subseteq} S^2$ by definition 7.12 (M-morphisms) on page 31 and $S^2 \stackrel{\text{full-cat}}{\subseteq} S'^2$ by hypothesis, so $S^1 \stackrel{\text{full-cat}}{\subseteq} S'^2$.

{def: modMmorph}

f is a morphism of S^1 by definition 7.12, hence a morphism of S'^2 . $\square$

{def: modMmorph}
{M-morph: contisstrict}

9. If $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ and $f: S^1 \rightarrow S^2$ is continuous, then f is an m-morphism of S^1_{triv} to S^2_{triv} .

Proof.

f is continuous:

$f: S^1 \rightarrow S^2$ is continuous. $\text{Top}\left(\begin{smallmatrix} S^i \\ \text{triv} \end{smallmatrix}\right) = \text{Top}(S^i) = S^i, i = 1, 2.$

f maps model neighborhoods into model neighborhoods:

S^2 is open in S^2 , hence a model neighborhood of S^2_{triv} . Let U be a model neighborhood of S^1_{triv} . Then $f[U] \subseteq S^2$.

Inverse images of model neighborhoods are model neighborhoods:

Let V be a model neighborhood of S^2_{triv} . V is open in S^2 and f is continuous, so $f^{-1}[V]$ is open in S^1 and hence a model neighborhood of S^1_{triv} .

$\square$

{M-morph: Idislocal}

10. If $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ then $\text{Id}_{S^2}^{S^1}$ is a local m-morphism of S^1 to S^2 . If each S^i is a model neighborhood of S^i then $\text{Id}_{S^2}^{S^1}$ is a morphism of S^2 .

Proof. Let $u \in S^1$ and U_u a model neighborhood of S^1 for u . Since $S^1 \stackrel{\text{mod}}{\subseteq} S^2$, U_u is also a model neighborhood of S^2 for u , and hence $\text{Id}_{U_u}^{\text{Ar}} \in S^2$.

If each S^i is a model neighborhood of S^i , then since $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ by hypothesis, S^1 is a model neighborhood of S^2 . The result follows by item 5 of definition 2.1 (Model spaces) on page 18. $\square$

{def: model}
{mod: inclusion}

{M-morph:Idcatislocal}

11. If $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$ and $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2$ then $\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}$ is a morphism of $\mathcal{M}^2$.

Proof. $\mathbf{S}^1 \in^{\text{Ob}} \mathcal{M}^2$ because $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$. $\text{Id}_{\mathbf{S}^2}^{\text{Ar}} \in \mathcal{M}^2$. Then $\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1} \in \mathcal{M}^2$ by item 4 of definition 7.7 (Model categories) on page 29. $\square$

{def:modcat}
{M-morph:openmorphistrivmor}

12. Let $U^i \subseteq \mathbf{S}^i, i = 1, 2$, be open. If $f: U^1 \longrightarrow U^2$ is a (local) $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 then $f: U^1 \longrightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .
 triv triv

Proof. Since $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2, \mathbf{S}^1 \subseteq^{\text{subsp}} \mathbf{S}^2$. Since $f: U^1 \longrightarrow U^2$ is a model function, it is continuous and hence a morphism of $\mathbf{S}^2$. $\square$

{cor:modMmorph}

Corollary 7.14 ((Local) m-morphisms). Let $\mathcal{M}^i, \mathcal{M}^i \subseteq^{\text{full-cat}} \mathcal{M}^i, i = 1, 2$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in^{\text{Ob}} \mathcal{M}^i, \mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in^{\text{Ob}} \mathcal{M}^i$ and $\mathbf{S}^i \subseteq^{\text{mod}} \mathbf{S}^i$.

{M-morph:morphislocal}

1. If $f: \mathbf{S}^1$ to $\mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1$ to $\mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since $f: \mathbf{S}^1$ to $\mathbf{S}^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1$ to $\mathbf{S}^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{M-morph:sublocalismorph}

2. If U^1 and U^2 are model neighborhoods of $\mathbf{S}^i$ then every function $f: U^1 \longrightarrow U^2$ that is a local $\mathbf{S}^i$ m-morphism of U^1 to U^2 is a $\mathbf{S}^i$ m-morphism of U^1 to U^2 .

Proof. Since $f: U^1 \longrightarrow U^2$ is a local $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^1 to U^2 then it is an $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^1 to U^2 . $\square$

{M-morph:localismorph}

3. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and each $\mathbf{S}^i$ is a model neighborhood of $\mathbf{S}^i$, then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then it is an $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{M-morph:catmorphislocal}

4. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since f is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

5. If $\mathcal{M}^i$ satisfies the restricted sheaf condition and U^j is a model neighborhood of $\mathcal{S}^i$, $j = 1, 2$, then every function $f: U^1 \rightarrow U^2$ that is a local $\mathcal{S}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 is a $\mathcal{S}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 .

{M-morph: subcatlocalismorph}

Proof.

- (a) $f: \text{Mod}(U^1, \mathcal{S}^i) \rightarrow \text{Mod}(U^2, \mathcal{S}^i)$ is a local $\mathcal{S}^i$ - $\mathcal{S}^i$ - $\mathcal{M}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 by definition 7.11 (Local m-morphisms) on page 31
- (b) $f: \text{Mod}(U^1, \mathcal{S}^i) \xrightarrow{\text{lem:modMmorph}} \text{Mod}(U^2, \mathcal{S}^i)$ is a $\mathcal{S}^i$ - $\mathcal{S}^i$ - $\mathcal{M}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 by item 4 of lemma 7.13.
- (c) The result follows from definition 7.11.

{def:modlocalMmorph}

{M-morph: opencatlocalismorph}

{def:modlocalMmorph}

□

{M-morph: catlocalismorph}

6. Let each S^i be a model neighborhood of $\mathcal{S}^i$, Then every $f: S^1 \rightarrow S^2$ that is a $\mathcal{M}^1$ - $\mathcal{M}^2$ local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ is a $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$.

1. If $f: S^1 \rightarrow S^2$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $\mathcal{S}^2 \stackrel{\text{mod}}{\subseteq} \mathcal{S}'^2$ then f is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}'^2$.

Proof. $f: S^1 \rightarrow S^2$ is continuous by corollary 2.10 (Model subspaces) on page 22.

{cor:modsub}

□

2. If $f: S^1 \rightarrow S^2$ is a(n) (local) m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ then f is an m-morphism of S^1 to S^2 .

Proof. f is continuous by definition 7.11 (Local m-morphisms) on page 31.

$\text{Top}_{\text{triv}}(S^i) = \text{Top}(\mathcal{S}^i) = S^i, i = 1, 2.$

{def:modlocalMmorph}

□

3. If S^i is a model neighborhood of $\mathcal{S}^i$ then Id_{S^i} is a morphism of $\mathcal{S}^i$.

Proof. $\mathcal{S}^i \stackrel{\text{mod}}{\subseteq} \mathcal{S}^i.$

□

4. If $f: S^1 \rightarrow S^2$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $f^2: S^2 \rightarrow S^3$ is an m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then $f^2 \circ f$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $f^2: S^2 \rightarrow S^3$ is an m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then f^2 is a local m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$.

□

5. If $f: S^1 \rightarrow S^2$ is an m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $f^2: S^2 \rightarrow S^3$ is a local m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then $f^2 \circ f$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $f: S^1 \longrightarrow S^2$ is an $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^2$. $\square$

6. If $f: S^1 \longrightarrow S^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and $f^2: S^2 \longrightarrow S^3$ is an $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then $f^2 \circ f$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $f^2: S^2 \longrightarrow S^3$ is an $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then f^2 is a local $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$. $\square$

7. If $f: S^1 \longrightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and $f^2: S^2 \longrightarrow S^3$ is a local $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then $f^2 \circ f$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $f: S^1 \longrightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^2$. $\square$

{M-morph:morphistrivmorph}

8. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a (local) $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is an $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Each S^i is open, so $f: S^1 \longrightarrow S^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{lem:modlocalMmorphcomp}

Lemma 7.15 (Composition of local $\mathbf{m}$ -morphisms). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}^i$ and $U^i \subseteq S^i$ be open.

1. If $f^i: U^i \longrightarrow U^{i+1}$, $i = 1, 2$, is a local $\mathbf{S}^i$ - $\mathbf{S}^{i+1}$ $\mathbf{m}$ -morphism of U^i to U^{i+1} then $f^2 \circ f^1$ a local $\mathbf{S}^1$ - $\mathbf{S}^3$ $\mathbf{m}$ -morphism of U^1 to U^{i+1} .

Proof. Since $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$ then $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$, Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathbf{S}^i$ and $\mathbf{S}^{i+1}$ such that that $f^i[U_{u^i}] \subseteq V_{u^i}$ and $f^i: \text{Mod}(U_{u^i}, \mathbf{S}^i) \longrightarrow \text{Mod}(V_{u^i}, \mathbf{S}^{i+1})$ is a morphism of $\mathbf{S}^{i+1}$.

Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^2 \circ f^1)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \longrightarrow V_{u^3}$ is a morphism of $\mathbf{S}^3$. $\square$

2. If $f^i: S^i \longrightarrow S^{i+1}$, $i = 1, 2$, is a local $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ $\mathbf{m}$ -morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1$ a local $\mathcal{M}^1$ - $\mathcal{M}^{i+2}$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Let $u^1 \in \mathcal{S}^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathcal{S}^i$ and $\mathcal{S}^{i+1}$ such that $f^i: U_{u^i} \rightarrow V_{u^i}$ is a morphism of $\mathcal{M}^{i+1}$. Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^3 \circ f^2)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \rightarrow V_{u^3}$ is a morphism of $\mathcal{M}^3$ by [item 5 of definition 7.7 \(Model categories\)](#) on page 29. $\square$ {def:modcat}

If each $f^i: \mathcal{S}^i \rightarrow \mathcal{S}^{i+1}$, $i = 1, 2$, is a local m-morphism of $\mathcal{S}^i$ to $\mathcal{S}^{i+1}$, then $f^2 \circ f^1: \mathcal{S}^1 \rightarrow \mathcal{S}^3$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^3$, $\mathcal{M}^1 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^3$. Let $u \in \mathcal{S}^i$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist a model neighborhood U_u for u , model neighborhoods V_u, V'_u for v and a model neighborhood W_v of w such that $f_1[U_u] \subseteq V'_u$, $f_2[V_v] \subseteq W_v$, $f_1: U_u \rightarrow V'_u$ is a morphism of $\mathcal{S}^2$ and $f_2: V_v \rightarrow W_v$ is a morphism of $\mathcal{S}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_v \cap V'_u \neq \emptyset$, $\hat{V}_u$ is a model neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}_u]$ is a model neighborhood for u . $f_1: \hat{U}_u \rightarrow \hat{V}_u$ and $f_2: \hat{V}_u \rightarrow W_v$ are morphisms of $\mathcal{S}^3$ by [item 4 of definition 2.1 \(Model spaces\)](#) on page 18 and thus $f_2 \circ f_1: \hat{U}_u \rightarrow W_v$ is a morphism of $\mathcal{S}^3$. $\square$ {def:model}

If each $f^i: \mathcal{S}^i \rightarrow \mathcal{S}^{i+1}$ is a local $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathcal{S}^i$ to $\mathcal{S}^{i+1}$ then $f^2 \circ f^1: \mathcal{S}^1 \rightarrow \mathcal{S}^3$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$, $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Let $u \in \mathcal{S}^1$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist a model neighborhood U_u for u , model neighborhoods V_v, V'_u for v and a model neighborhood W_v of w such that $f^1[U_u] \subseteq V'_u$, $f^1: U_u \rightarrow V'_u$ is a morphism of $\mathcal{M}^2$, $f^2[V_v] \subseteq W_v$ and $f^2: V_v \rightarrow W_v$ is a morphism of $\mathcal{M}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_v \cap V'_u \neq \emptyset$, $\hat{V}_u$ is a model neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}_u]$ is a model neighborhood for u . $f^1: \hat{U}_u \rightarrow \hat{V}_u$ and $f^2: \hat{V}_u \rightarrow W_v$ are morphisms of $\mathcal{M}^3$ by [item 4 of definition 7.7 \(Model categories\)](#) on page 29 and thus $f^2 \circ f^1: \hat{U}_u \rightarrow W_v$ is a morphism of $\mathcal{M}^3$. $\square$ {def:modcat}

Corollary 7.16 ([Composition of local m-morphisms]). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories, $\mathcal{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i)^{\text{Ob}} \in \mathcal{M}^i$.

1. Let $U^j \subseteq \mathcal{S}^i$, $j = 1, 2, 3$, be open. If $f^j: U^j \rightarrow U^{j+1}$, $j = 1, 2$, is a local $\mathcal{S}^i$ m-morphism of U^j to U^{j+1} then $f^2 \circ f^1$ is a local $\mathcal{S}^i$ m-morphism of U^1 to U^3 .

Proof. Since each $f^j: U^j \rightarrow U^{j+1}$ is a local $\mathcal{S}^i$ - $\mathcal{S}^i$ m-morphism of U^j to U^{j+1} then $f^2 \circ f^1$ is a local $\mathcal{S}^i$ - $\mathcal{S}^i$ m-morphism of U^1 to U^3 . $\square$

2. If $f^i: \mathcal{S}^i \rightarrow \mathcal{S}^{i+1}$, $i = 1, 2$, is a local m-morphism of $\mathcal{S}^i$ to $\mathcal{S}^{i+1}$ then $f^2 \circ f^1$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$ then $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$. Let $u^1 \in \mathbf{S}^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathbf{S}^i$ and $\mathbf{S}^{i+1}$ such that $f^i: U_{u^i} \rightarrow V_{u^i}$ is a morphism of $\mathbf{S}^{i+1}$. Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^2 \circ f^1)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \rightarrow V_{u^3}$ is a morphism of $\mathbf{S}^3$. $\square$

Lemma 7.17 (Composition of m-morphisms). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories and $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i)^{\text{Ob}} \in \mathcal{M}^i$.

{lem:modMmorphcomp}

1. If $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is an m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since each f^i is a morphism of $\mathbf{S}^{i+1}$ by definition 7.12 (M-morphisms) on page 31 and $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$, then each f^i is a morphism of $\mathbf{S}^3$ by definition 7.12 (M-morphisms) on page 31, and thus $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is a morphism of $\mathbf{S}^3$. $\square$

{def:modMmorph}

{def:modMmorph}

2. If $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is an $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1$ is an $\mathcal{M}^1$ - $\mathcal{M}^{i+2}$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Since each $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$ is a morphism of $\mathcal{S}^3$, then so is $f^2 \circ f^1$. $\square$

3. If each $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$ is an $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is an $\mathcal{M}^1$ - $\mathcal{M}^3$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ and f^i is a morphism of $\mathcal{S}^{i+1}$ by definition 7.12 (M-morphisms) on page 31. Since $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$, f^1 is a morphism of $\mathcal{S}^3$. and thus $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is a morphism of $\mathcal{S}^3$. $\square$

{def:modMmorph}

8 Spaces and proper functions

{sec:spaces}

Several of the following definitions involve spaces of a restrictive character and specific types of mappings among them. Except where otherwise qualified, the word *space* will have this restricted meaning.

{def:space}

Definition 8.1 (Spaces). A space is a topological space, a model space or either with an additional associated structure.

{def:proper}

Definition 8.2 (Proper functions). Let S^1, S^2 be spaces and $f: S^1 \rightarrow S^2$ a continuous function; f need not preserve any associated algebraic structure⁶. f is a proper⁷ function iff

1. S^1 and S^2 are both Truthspace or both **Truthspace**, and $f(\text{True}) = \text{True}$.
2. S^1, S^2 are topological spaces other than Truthspace.
3. S^1 is a topological space other than Truthspace, S^2 is a model space other than **Truthspace** and the images of open sets are contained in model neighborhoods.
4. S^1 is a model space other than **Truthspace**, S^2 is a topological space other than Truthspace and the inverse images of open sets are model neighborhoods.
5. S^1, S^2 are both model spaces other than **Truthspace** and f is a model function.

{def:singcat}

Definition 8.3 (Singleton categories and sequences of singleton categories). Let S be a topological space. Then the singleton category of S , abbreviated $\mathcal{S}_{\text{Sing}}$, is the category whose sole object is S and whose sole morphism is the identity morphism of S to itself.

Let $\mathcal{S}$ be a model space. Then the singleton category of $\mathcal{S}$, abbreviated $\mathcal{S}_{\text{Sing}}$, is the category whose sole object is $\mathcal{S}$ and whose sole morphism is the identity morphism of $\mathcal{S}$ to itself.

Let $\mathcal{S} \stackrel{\text{def}}{=} (S^\alpha, \alpha < A)$ be a sequence of spaces. Then the singleton category sequence of $\mathcal{S}$, abbreviated $\mathcal{S}_{\text{Sing}}$, is $(\mathcal{S}^\alpha, \alpha < A)$.

Let $\mathcal{S}$ be a set of spaces. Then the singleton category of $\mathcal{S}$, abbreviated $\mathcal{S}_{\text{Sing}}$, is

$$\bigcup_{S \in \mathcal{S}} \mathcal{S}_{\text{Sing}}.$$

Remark 8.4. By abuse of language the notation $\mathcal{S}_{\text{Sing}}$ will be used to name sequences of categories constructed with this and similar functions.

⁶However, in practice commutation relations will often enforce the preservation of algebraic structures.

⁷In the spirit of [Kelley, 1955, footnote, p. 112]

{GenTop}

This nomenclature is an excellent example of the time-honored custom of referring to a problem we cannot handle as abnormal, irregular, improper, degenerate, inadmissible, and otherwise undesirable.

Part VI

M-charts and m-atlases

{part:M-charts}

The literature defines fiber bundles and manifolds using the language of charts, atlases and transition functions; it has multiple equivalent definitions. Some authors start with topological spaces and define atlases over them. Some start with abstract sets and define atlases over them, deriving the topology from the atlas. Some start with an indexed set of open patches in a coordinate space and transition functions among them satisfying a cocycle (compatibility) constraint, and then derive the total space as a quotient space of the disjoint union of the patches. Some authors use maximal atlases while others use equivalence classes of atlases.

This paper uses the first approach, explicitly making the relevant spaces topological spaces, but modifies the definitions of charts in order to make them fit more naturally into the context of local coordinate spaces. It adds a prefix, e.g., C^k , M , in order to avoid confusion with the conventional definitions. An m -atlas based on a topological space is closer to the conventional definitions of an atlas for a manifold while an m -atlas based on a model space is suitable for defining both manifolds and fiber bundles. Although simple manifolds and fiber bundles could both be defined directly in terms of maximal m -atlases, this paper has a different perspective, and uses the m -atlases as part of the more general Local Coordinate Space (LCS), presented in [Metz, 2018].

{LCS}

Sections 9 to 10 define m -charts and m -atlases. Maximal m -atlases are essentially the same as manifolds.

Section 11 defines morphisms between m -atlases, referred to as m -atlas morphisms, in a fashion tailored to use in defining local coordinate spaces. It constructs categories of M -atlases and functors, and proves some basic results. It also defines a related concept of m -atlas near morphisms, although the restriction needed to use them as morphisms in a category of m -atlases requires them to be morphisms.

Section 13 defines morphisms between m -atlases in a fashion similar to that conventionally used for C^k maps between differentiable manifolds, in order to provide context. It constructs categories of M -atlases and functors, and proves some basic results.

9 M-charts

sec:M-charts {def:M-chart}

Definition 9.1 (M-charts). Let $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $C \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces. An m -chart (U, V, ϕ) of E in the coordinate space C consists of

1. A nonvoid model neighborhood $U \in \mathcal{E}$, known as a coordinate patch
2. A model neighborhood $V \in \mathcal{C}$
3. A model homeomorphism $\phi: U \xrightarrow[\cong]{\sim} V$, known as a coordinate function

Remark 9.2. I consider it clearer to explicate the range, rather than the conventional usage of specifying only the domain and function or the minimalist usage of specifying only the function.

For every $u \in U$, (U, V, ϕ) is an m-chart of E in the coordinate space $\mathcal{C}$ at u and covers u . E is the total model space or total space for the chart and C is the coordinate model space or coordinate space for the chart.

Let E be a topological space and $C \stackrel{\text{def}}{=} (C, \mathcal{C})$ be a model space. (U, V, ϕ) is an m-chart of E in the coordinate space $\mathcal{C}$ iff it is an m-chart of $\underset{\text{triv}}{E}$ in the coordinate space $\mathcal{C}$.

For every $x \in U$, (U, V, ϕ) is an m-chart of E in the coordinate space $\mathcal{C}$ at x . E is the total space for the chart and C is the coordinate model space or coordinate space for the chart.

Lemma 9.3 (M-charts).

{lem:M-chart}
{M-chart:openC}

1. Let $E \stackrel{\text{def}}{=} (E, \mathcal{C})$ and $C \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces, and (U, V, ϕ) be an m-chart of E in the coordinate space C such that every open subset of V is a model neighborhood of C . Then (U, V, ϕ) is an m-chart of E in the coordinate space C .

Proof. U , V and ϕ satisfy the conditions of the definition:

U is a model neighborhood of $\underset{\text{triv}}{E}$: U is a model neighborhood of E , hence open.

V is a model neighborhood of C : By hypothesis.

ϕ is a model homeomorphism from $\text{Mod}\left(U, \underset{\text{triv}}{E}\right)$ to $\text{Mod}(V, C)$:

- (a) Since ϕ is a model homeomorphism from $\text{Mod}(U, E)$ to $\text{Mod}(V, C)$, it is a homeomorphism.
- (b) If V' is a model subneighborhood of V , then $\phi^{-1}[V']$ is open and hence a model neighborhood of $\underset{\text{triv}}{E}$.
- (c) If U' is a model neighborhood of $\underset{\text{triv}}{E}$ then it is open, hence $\phi[U']$ is an open subset of V and thus by hypothesis a model neighborhood of C .

□

{M-chart:openE}

2. Let (U, V, ϕ) be an m-chart of E in the coordinate space C . (U, V, ϕ) is an m-chart of E in the coordinate space C iff every open subset of U is a model neighborhood of E .

Proof. If (U, V, ϕ) is an m-chart of E in the coordinate space C and also an m-chart of E in the coordinate space C then let $U' \subseteq U$ be open. Since (U, V, ϕ) is an m-chart of E in the coordinate space C , ϕ^{-1} is a model function from

$\text{Mod}(V, \mathbf{C})$ to $\underset{\text{triv}}{E}$ and $\phi[U']$ is a model neighborhood of $\mathbf{C}$. Since (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$, ϕ is a model function from $\text{Mod}(U, E)$ to $\mathbf{C}$ and thus $\phi^{-1}[\phi[U']]$ is a model neighborhood of E .

If (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$ and every open $U' \subseteq U$ is a model neighborhood of E then

- (a) Since $U \subseteq U$ is open, U is a model neighborhood of E by hypothesis.
- (b) V is a model neighborhood of $\mathbf{C}$ by hypothesis.
- (c) If $V' \subseteq V$ is a model neighborhood of $\mathbf{C}$ then $\phi^{-1}[V']$ is open and hence a model neighborhood of E by hypothesis.

ϕ^{-1} is a model function from $\text{Mod}\left(U, \underset{\text{triv}}{E}\right)$ to $\text{Mod}(V, \mathbf{C})$. if $U' \subseteq U$ is a model neighborhood of E then U' is open and hence a model neighborhood of $\underset{\text{triv}}{E}$. Then $\phi[U']$ is a model neighborhood of $\mathbf{C}$.

□

{cor:M-chart}

Corollary 9.4 (M-charts). Let $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces, $\mathbf{C}$ be fine grained and (U, V, ϕ) be an m-chart of E in the coordinate space $\mathbf{C}$. Then (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. Since V is open and $\mathbf{C}$ is fine grained, every open subset of V is a model neighborhood of $\mathbf{C}$. The result follows by item 1 of lemma 9.3 (M-charts) on page 42. □

{M-chart:openC}

{def:M-subchart}

Definition 9.5 (Subcharts). Let (U, V, ϕ) be an m-chart of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and $U' \subseteq U$ a nonvoid model neighborhood of E . Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

Let (U, V, ϕ) be an m-chart of E in the coordinate space $(C, \mathcal{C})$ and $U' \subseteq U$ an open set of E . Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

(U', V', ϕ') is a subchart of (U, V, ϕ) at x iff (U', V', ϕ') is a subchart of (U, V, ϕ) and $x \in U'$.

{lem:M-subchart}

Lemma 9.6 (Subcharts). Let $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces.

Let (U, V, ϕ) be an m-chart of E in the coordinate space $\mathbf{C}$ and (U', V', ϕ') a subchart of (U, V, ϕ) . Then (U', V', ϕ') is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. U' , V' and ϕ' satisfy the conditions of the definition of an m-chart:

1. U' is a model neighborhood of E by the definition of subchart.
2. Since U' is a model neighborhood and ϕ is a model homeomorphism, $V' = \phi[U']$ is a model neighborhood.

3. Since ϕ and ϕ^{-1} are model functions, so are $\phi' = \phi|_{U',V'}$ and $\phi'^{-1} = \phi^{-1}|_{V',U'}$. Thus ϕ' is a model homeomorphism.

□

Let (U, V, ϕ) be an m-chart of E in the coordinate space $\mathbf{C}$ and (U', V', ϕ') a subchart of (U, V, ϕ) . Then (U', V', ϕ') is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. U' , V' and ϕ' satisfy the conditions of the definition of an m-chart:

1. U' is open by the definition of subchart.
2. Since ϕ is a homeomorphism and U' is open, $V' = \phi[U']$ is open.
3. Since ϕ is a homeomorphism that maps open sets into model neighborhoods, $\phi|_{U',V'}$ is a homeomorphism that maps open sets into model neighborhoods.

□

{def:M-compat}

Definition 9.7 (M-compatibility). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces and (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) in the coordinate space $\mathbf{C}$ iff either

1. U_1 and U_2 are disjoint
2. The transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$.

(U_1, V_1, ϕ_1) is also properly m-compatible with (U_2, V_2, ϕ_2) in the space $\mathbf{E}$ and the coordinate space $\mathbf{C}$, and properly m-compatible in the space $\mathbf{E}$ and the coordinate space $\mathbf{C}$ with (U_2, V_2, ϕ_2) , iff for every model neighborhoods $U'_i \subseteq U_i$

1. For every morphism $f: U'_1 \longrightarrow U'_2$ of $\mathbf{E}$, the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \longrightarrow \phi_2[U'_2]$ is a morphism of $\mathbf{C}$.
2. For every morphism $g: U'_2 \longrightarrow U'_1$ of $\mathbf{E}$, the composition $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \longrightarrow \phi_1[U'_1]$ is a morphism of $\mathbf{C}$.

Remark 9.8. This is a stronger condition than requiring that for every $f: U_1 \longrightarrow U_2$ and $g: U_2 \longrightarrow U_1$ the compositions be morphisms of $\mathbf{C}$.

Let E be a topological space, $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be a model space and (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of E in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) iff either

1. U_1 and U_2 are disjoint
2. The transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$

(U_1, V_1, ϕ_1) is also properly m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$, and properly m-compatible in the space E and the coordinate space $\mathbf{C}$ with (U_2, V_2, ϕ_2) , iff for every open $U'_i \subseteq U_i$

1. For every continuous $f: U'_1 \rightarrow U'_2$, the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \rightarrow \phi_2[U'_2]$ is a morphism of $\mathbf{C}$.
2. For every continuous $g: U'_2 \rightarrow U'_1$, the composition $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \rightarrow \phi_1[U'_1]$ is a morphism of $\mathbf{C}$.

In both cases, the space and coordinate space may be omitted when they are clear from context.

{lem:M-compatible}

Lemma 9.9 (M-compatibility).

1. Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be a chart of E in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ iff (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space $\underset{\text{triv}}{E}$ and the coordinate space $\mathbf{C}$.

Proof. $U_1 \cap U_2$ and the transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \rightarrow \phi_2[U_1 \cap U_2]$ is the same for the case of E and the case of $\underset{\text{triv}}{E}$.

Any $U'_i \subseteq U_i$ is a model neighborhood of $\underset{\text{triv}}{E}$ iff it is open.

Any $f: U'_1 \rightarrow U'_2$ is a morphism of $\underset{\text{triv}}{E}$ iff it is continuous.

Any $g: U'_2 \rightarrow U'_1$ is a morphism of $\underset{\text{triv}}{E}$ iff it is continuous.

The compositions $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \rightarrow \phi_2[U'_2]$ and $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \rightarrow \phi_1[U'_1]$ are the same for the case of E and the case of $\underset{\text{triv}}{E}$. $\square$

{cor:M-compatible}

Corollary 9.10. Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be a chart of E in the coordinate space $\mathbf{C}$ and let (U'_i, V'_i, ϕ'_i) be a subchart. If (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. By lemma 9.9 above, (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space E and the coordinate space $\mathbf{C}$ iff (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space $\underset{\text{triv}}{E}$ and the coordinate space $\mathbf{C}$.

{lem:M-compatible}

The result follows from lemma 9.9.

$\square$ {lem:M-compatible}
{lem:M-compat sym}

Lemma 9.11 (Symmetry of m-compatibility). Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(\mathbf{C}, \mathcal{C})$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ iff (U_2, V_2, ϕ_2) is (properly) m-compatible with (U_1, V_1, ϕ_1) .

Proof. It suffices to prove the implication in only one direction.

1. $U_1 \cap U_2 = U_2 \cap U_1$.
2. Since the transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$, so is $t^{-1} = \phi_1 \circ \phi_2^{-1}: \phi_2[U_1 \cap U_2] \longrightarrow \phi_1[U_1 \cap U_2]$.
3. The constraint for proper maximality is intrinsically symmetrical.

□

Let $(U_i, V_i, \phi_i), i = 1, 2$, be an m-chart of E in the coordinate space $(C, \mathcal{C})$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space C iff (U_2, V_2, ϕ_2) is (properly) m-compatible with (U_1, V_1, ϕ_1) .

Proof. The same proof applies as above. □

{lem:M-compatsub}

Lemma 9.12 (M-compatibility of subcharts). Let $(U_i, V_i, \phi_i), i = 1, 2$, be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$, (U_1, V_1, ϕ_1) be (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space C and (U'_i, V'_i, ϕ'_i) be a subchart. Then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. Since subcharts are charts, U'_i and V'_i are model neighborhoods. If $U_1 \cap U_2 = \emptyset$ then $U'_1 \cap U'_2 = \emptyset$ and thus (U'_1, V'_1, ϕ'_1) is m-compatible with (U'_2, V'_2, ϕ'_2) . Otherwise, the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$ and hence $t_2^1: \phi_1[U'_1 \cap U'_2] \xrightarrow{\cong} \phi_2[U'_1 \cap U'_2]$ is an isomorphism of $\mathcal{C}$.

If (U_1, V_1, ϕ_1) is properly m-compatible with (U_2, V_2, ϕ_2) , $U''_i \subseteq U'_i, i = 1, 2$, is a model neighborhood and $f: U''_1 \longrightarrow U''_2$ is a morphism of E , then each $U''_i \subseteq U_i$ and thus by definition 9.7 the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U''_1] \longrightarrow \phi_2[U''_2]$ is a morphism of C . □

{def:M-compat}

Let $(U_i, V_i, \phi_i), i = 1, 2$, be an m-chart of E in the coordinate space $(C, \mathcal{C})$, (U'_i, V'_i, ϕ'_i) be a subchart and (U_1, V_1, ϕ_1) be (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space C . Then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. By lemma 9.9 (M-compatibility) on page 45, (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space C . Then by lemma 9.12 above, (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space E and the coordinate space C . □

{lem:M-compat}

{lem:M-compatsub}

The result follows from lemma 9.9. □

{lem:M-compat}
{cor:M-compatsub}

Corollary 9.13 (M-compatibility of subcharts). Let (U, V, ϕ) be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and (U', V', ϕ') a subchart. Then (U', V', ϕ') is properly m-compatible with (U, V, ϕ) in the space E and the coordinate space C .

Proof. (U, V, ϕ) is m-compatible with itself and is a subchart of itself, □

Let (U, V, ϕ) be an m-chart of E in the coordinate space $(C, \mathcal{C})$ and (U', V', ϕ') a subchart. Then (U', V', ϕ') is m-compatible with (U, V, ϕ) in the space E and the coordinate space $\mathbf{C}$.

Proof. (U, V, ϕ) is m-compatible with itself and is a subchart of itself, □

Definition 9.14 (Covering by m-charts). Let $\mathbf{A}$ be a set of charts of the topological space E in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. $\mathbf{A}$ covers E iff $\pi_1[\mathbf{A}]$ covers E .

Let $\mathbf{A}$ be a set of charts of the model space $\mathbf{E} = (E, \mathcal{E})$ in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. $\mathbf{A}$ covers $\mathbf{E}$ iff $\pi_1[\mathbf{A}]$ covers E .

10 M-atlases

{sec:M-atlas}

A set of charts can be atlases for different coordinate model spaces even if it is for the same total model space. In order to aggregate atlases into categories, there must be a way to distinguish them. Including the two⁸ spaces in the definitions of the categories serves the purpose.

{def:M-atlas}

Definition 10.1 (M-atlases). Let $\mathbf{A}$ be a set of mutually (properly) m-compatible m-charts of $\mathbf{E} = (E, \mathcal{E})$ in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. Then

1. $\mathbf{A}$ is an (a proper) m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C})$, iff $\mathbf{A}$ covers E .
2. $\mathbf{A}$ is a full (proper) atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C})$, iff $\mathbf{A}$ covers E and $\pi_2[\mathbf{A}]$ covers C .
3. The triple $(\mathbf{A}, \mathbf{E}, \mathbf{C})$ refers to $\mathbf{A}$ considered as an m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$.
4. $\mathbf{E}$ is the total model space or total space for the atlas
5. $\mathbf{C}$ is the coordinate model space or coordinate space for the atlas.

Let E be a topological space, $\mathbf{C} = (C, \mathcal{C})$ a model space and $\mathbf{A}$ be a set of mutually m-compatible m-charts of E in the coordinate space $\mathbf{C}$. Then

6. $\mathbf{A}$ is an m-atlas of E in the coordinate space $\mathbf{C}$, abbreviated as $\text{isAtl}_{\text{Ob}}(\mathbf{A}, E, \mathbf{C})$, iff $\mathbf{A}$ is an m-atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\mathbf{C}$.
7. $\mathbf{A}$ is a full atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{\text{full}}(\mathbf{A}, E, \mathbf{C})$, iff $\mathbf{A}$ is a full atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\mathbf{C}$.
8. The triple $(\mathbf{A}, E, \mathbf{C})$ refers to $\mathbf{A}$ considered as an m-atlas of E in the coordinate space $\mathbf{C}$.

⁸The spaces are redundant, but convenient.

9. E is the total space for the atlas.

10. C is the coordinate model space or coordinate space for the atlas.

Let A be an (a proper) m-atlas of E in the coordinate space C . A is fine grained iff whenever $(U, V, \phi) \in A$, every subchart of (U, V, ϕ) is also an atlas of A .

Remark 10.2. A can be a fine graind (proper) atlas of E in the coordinate space C even if C is not fine grained.

Let A be an (a proper) m-atlas of E in the coordinate space C . A is fine grained iff whenever $(U, V, \phi) \in A$, every subchart of (U, V, ϕ) is also an atlas of A .

By abuse of language we write $U \in A$ for $U \in \pi_1[A]$.

{lem:M-atlas}

Lemma 10.3 (M-atlases). Let A be an m-atlas of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $C \stackrel{\text{def}}{=} (C, \mathcal{C})$. If C is fine grained then A is an m-atlas of E in the coordinate space C .

Proof. Let $(U, V, \phi) \in A$. Then (U, V, ϕ) is an m-chart of E in the coordinate space C by corollary 9.4 (M-charts) on page 43. $\square$

{cor:M-chart}
{def:M-CompatATL}

Definition 10.4 (M-compatibility with m-atlases). An m-chart (U, V, ϕ) of $(E, \mathcal{E})$ is (properly) m-compatible with an m-atlas A in the space E and the coordinate space C iff it is (properly) m-compatible in the space E and the coordinate space C with every chart in A .

An m-chart (U, V, ϕ) of E is (properly) m-compatible with an m-atlas A in the space E and the coordinate space C iff it is (properly) m-compatible in the space E and the coordinate space C with every chart in A .

{lem:M-CompatATL}

Lemma 10.5 (M-compatibility with m-atlases). Let A be an m-atlas of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $C \stackrel{\text{def}}{=} (C, \mathcal{C})$. If C is fine grained then A is an m-atlas of E in the coordinate space C .

Proof. $\square$

{lem:M-CompatsubATL}

Lemma 10.6 (M-compatibility of subcharts with atlases). Let A be an (a proper) m-atlas of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and (U_1, V_1, ϕ_1) an m-chart in A . Then any subchart of (U_1, V_1, ϕ_1) is properly m-compatible with A in the space E and the coordinate space C .

Proof. Let (U'^1, V'^1, ϕ'^1) be a subchart of (U^1, V^1, ϕ^1) and (U_2, V_2, ϕ_2) be another chart in A . Then (U_1, V_1, ϕ_1) is properly compatible with (U_2, V_2, ϕ_2) by definition 10.1 (M-atlases) on page 47, (U_2, V_2, ϕ_2) is properly m-compatible with itself and (U_2, V_2, ϕ_2) is a subchart of itself. (U'^1, V'^1, ϕ'^1) is properly m-compatible with (U_2, V_2, ϕ_2) by lemma 9.12 (M-compatibility of subcharts) on page 46. Since (U_2, V_2, ϕ_2) is arbitrary, (U'^1, V'^1, ϕ'^1) is properly m-compatible with A by definition 10.4 (M-compatibility with m-atlases) on page 48. $\square$

{def:M-atlas}

{lem:M-compatsub}
{def:M-CompatATL}

Let A be an m-atlas of E in the coordinate space $(C, \mathcal{C})$ and (U_1, V_1, ϕ_1) an m-chart in A . Then any subchart of (U_1, V_1, ϕ_1) is properly m-compatible with A in the space E and the coordinate space C .

Proof. The same proof applies in as above. $\square$

{lem:M-ATLextensions}

Lemma 10.7 (Extensions of m-atlases). Let $E = (E, \mathcal{E})$ and $C = (C, \mathcal{C})$ be model spaces, A an m-atlas of E in the coordinate space C , and (U_i, V_i, ϕ_i) , $i = 1, 2$, an m-chart of E in the coordinate space C (properly) m-compatible with A in the space E and the coordinate space C . Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) .

Proof. If $U_1 \cap U_2 = \emptyset$ then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) . Otherwise, ϕ_1 is a model homeomorphism, $U_1 \cap U_2$ is a model neighborhood of E , $\phi_1[U_1 \cap U_2]$ is a model neighborhood of C , $\phi_2[U_1 \cap U_2]$ is a model neighborhood of

C and $\phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \xrightarrow{\cong} \phi_2[U_1 \cap U_2]$ is a homeomorphism. It remains to show that $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a model homeomorphism. Let $(U_\alpha, V_\alpha, \phi_\alpha)$, $\alpha < A$,

be charts in A such that $U_1 \cap U_2 \subseteq \bigcup_{\alpha < A} U_\alpha$ and $(\bigvee_{\alpha < A}) U_1 \cap U_2 \cap U_\alpha \neq \emptyset$.

Since the charts are m-compatible with $(U_\alpha, V_\alpha, \phi_\alpha)$, the compositions

$\phi_2 \circ \phi_\alpha^{-1}: \phi_\alpha[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha]$ and $\phi_\alpha \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_\alpha[U_1 \cap U_2 \cap U_\alpha]$ are isomorphisms of C and thus

the composition $\phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha] =$

$\phi_2 \circ \phi_\alpha^{-1} \circ \phi_\alpha \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha]$ is an isomorphisms

of C . Then by [item 6](#), the restricted sheaf property, of [definition 2.1 \(Model spaces\)](#) {def:model}

on page 18, $\phi_2 \circ \phi_1^{-1}$ is an isomorphism of C .

If each (U_i, V_i, ϕ_i) is properly m-compatible with A then let $f: U'_1 \rightarrow U'_2$ and $g: U'_2 \rightarrow U'_1$ be morphisms of E and $U'_i \subseteq U_i$, $i = 1, 2$. Let u be an arbitrary point in U_1 and $(U_u, V_u, \phi_u) \in A$ be an arbitray chart at u . Since (U_1, V_1, ϕ_1) is properly m-compatible with A , it is properly m-compatible with $(U_u, V_u, \phi_u) \in A$, thus the compositions $\phi_2 \circ f \circ \phi_u^{-1}: \phi_u^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2]$ and $\phi_u \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_u^{-1}[U_u]$ are morphisms of C , their composition $\phi_2 \circ f \circ \phi_u^{-1} \circ \phi_u \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2] =$

$\phi_2 \circ f \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2]$ is a morphism of C and by [item 6](#), the restricted sheaf property, of [definition 2.1 \(Model spaces\)](#) on page 18,

{def:model}

$\phi_2 \circ f \circ \phi_1^{-1}: \phi_1^{-1}[U'_1] \rightarrow \phi_2^{-1}[U'_2]$ is a morphism of C .

The same proof applies in the opposite direction. $\square$

Let E be a topological space, $C = (C, \mathcal{C})$ a model space, A an (a proper) m-atlas of E in the coordinate space C and (U_i, V_i, ϕ_i) , $i = 1, 2$, a chart (properly) m-compatible with A in the space E and the coordinate space C . Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) .

Proof. The same proof applies as above. $\square$

{def:M-ATLmax}

Definition 10.8 (Maximal (semi-maximal) m-atlases). Let E and C be model spaces and A an m-atlas of E in the coordinate space C . A is a (proper) maximal m-atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, C)$, iff A cannot be extended by adding an additional (properly) m-compatible chart. A is a (proper) semi-

maximal m-atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob S-max}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C})$.

iff whenever

1. $(U, V, \phi) \in \mathbf{A}$
2. $U' \subseteq U, V' \subseteq V$ and $V'' \in \mathcal{C}$ are model neighborhoods
3. $\phi[U'] = V'$
4. $\phi': V' \xrightarrow{\cong} V''$ is an isomorphism of $\mathcal{C}$

then $(U', V'', \phi' \circ \phi|_{U'}) \in \mathbf{A}$.

$$\text{isAtl}_{\text{Ob max-full}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob full}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob max}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \quad (10.1)$$

$$\text{isAtl}_{\text{Ob S-max-full}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob full}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob S-max}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \quad (10.2)$$

Let E be a topological space, $\mathbf{C} = (C, \mathcal{C})$ be a model space and $\mathbf{A}$ be an m-atlas of E in the coordinate space $\mathbf{C}$. $\mathbf{A}$ is a (properly) maximal m-atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob max}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C})$, iff $\mathbf{A}$ is an m-atlas that cannot be extended by adding an additional (properly) m-compatible chart. $\mathbf{A}$ is a (properly) semi-maximal m-atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob S-max}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C})$, iff whenever

1. $(U, V, \phi) \in \mathbf{A}$
2. $U' \subseteq U, V' \subseteq V$ and $V'' \in \mathcal{C}$ are model neighborhoods
3. $\phi[U'] = V'$
4. $\phi': V' \xrightarrow{\cong} V''$ is an isomorphism of $\mathcal{C}$

then $(U', V'', \phi' \circ \phi|_{U'}) \in \mathbf{A}$.

Remark 10.9. There is no sheaf condition⁹; the union of a set of coordinate patches in the maximal atlas whose coordinate functions match on the intersections need not be a coordinate patch in the atlas.

⁹However, note item 6 (restricted sheaf condition) of [definition 2.1 \(Model spaces\)](#) on page 18.

`{mod:sheaf}`

$$\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \quad (10.3)$$

$$\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, \mathbf{C}) \quad (10.4)$$

{lem:M-ATLmax}

Lemma 10.10 (Maximal m-atlases are semi-maximal m-atlases). Let $E = (E, \mathcal{E})$ and $\mathbf{C} = (C, \mathcal{C})$ be model spaces and $\mathbf{A}$ a maximal m-atlas of E in the coordinate space $\mathbf{C}$. Then $\mathbf{A}$ is a semi-maximal m-atlas of E in the coordinate space $\mathbf{C}$.

Proof. Let $(U, V, \phi) \in \mathbf{A}$, $U' \subseteq U$, $V' \subseteq V$ and $V'' \stackrel{\text{Ob}}{\in} \mathcal{C}$ be model neighborhoods, $\phi[U'] = V'$ and $\phi': V' \xrightarrow{\cong} V''$ be an isomorphism of $\mathcal{C}$. (U', V', ϕ) is a subchart of (U, V, ϕ) and by ?? (??) on ?? is m-compatible with the charts of $\mathbf{A}$. Since ϕ' is a model homeomorphism, $(U', V'', \phi' \circ \phi)$ is m-compatible with the charts of $\mathbf{A}$. Since $\mathbf{A}$ is maximal, $(U', V'', \phi' \circ \phi)$ is a chart of $\mathbf{A}$. $\square$

{lem:M-Compat}

Let E be a topological space, $\mathbf{C} = (C, \mathcal{C})$ a model space and $\mathbf{A}$ a maximal m-atlas of E in the coordinate space $\mathbf{C}$. Then $\mathbf{A}$ is a semi-maximal m-atlas of E in the coordinate space $\mathbf{C}$.

Proof. Let $(U, V, \phi) \in \mathbf{A}$, $U' \subseteq U$ open, $V' \subseteq V$ and $V'' \stackrel{\text{Ob}}{\in} \mathcal{C}$ be model neighborhoods, $\phi[U'] = V'$ and $\phi': V' \xrightarrow{\cong} V''$ be an isomorphism of $\mathcal{C}$. (U', V', ϕ) is a subchart of (U, V, ϕ) and by ?? (??) on ?? is m-compatible with the charts of $\mathbf{A}$. Since ϕ' is a model homeomorphism, $(U', V'', \phi' \circ \phi)$ is m-compatible with the charts of $\mathbf{A}$. Since $\mathbf{A}$ is maximal, $(U', V'', \phi' \circ \phi)$ is a chart of $\mathbf{A}$. $\square$

{lem:M-Compat}

{the:M-ATLmax}

Theorem 10.11 (Existence and uniqueness of maximal m-atlases). Let $\mathbf{A}$ be an (a proper) m-atlas of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. Then there exists a unique (proper) maximal m-atlas $\text{Atlas}[(\text{proper})](\mathbf{A}, E, \mathbf{C})$ of E in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all m-atlases of E in the coordinate space $\mathbf{C}$ containing $\mathbf{A}$ and (properly) m-compatible in the coordinate space $\mathbf{C}$ with all of the m-charts in $\mathbf{A}$. Let $\mathbf{P}_{\max}$ be a maximal chain of $\mathbf{A}$. Then $\text{Atlas}[(\text{proper})](\mathbf{A}, E, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\max} \mathbf{P}_{\max}$ is a (proper) maximal m-atlas of E in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$. Uniqueness follows from lemma 10.7 (Extensions of m-atlases) on page 49. $\square$

{lem:M-ATLextensions}
{cor:M-ATLmax}

Corollary 10.12 (Existence and uniqueness of maximal m-atlases). Let $\mathbf{A}$ be an (a proper) m-atlas of E in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. Then there exists a unique (proper) maximal m-atlas $\text{Atlas}[(\text{proper})](\mathbf{A}, E, \mathbf{C})$ of E in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$.

Proof. $\text{Atlas}[(proper)](A, E, C) \stackrel{\text{def}}{=} \text{Atlas}[(proper)](A, E, C)_{\text{triv}}$

□

{def:M-ATLsets}

Definition 10.13 (Sets of m-atlases). Let E and C be model spaces.

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C) \stackrel{\text{def}}{=} \left\{ (A, E, C) \mid \begin{array}{l} \text{isAtl}_{\text{Ob}}^{(proper)}(A, E, C) \\ \text{(full, S-max, max, S-max-full, max-full)} \end{array} \right\} \quad (10.5)$$

Let E and C be sets of model spaces.

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C) \stackrel{\text{def}}{=} \left\{ (A, (E, \mathcal{E}), (C, \mathcal{C})) \mid \begin{array}{l} (E, \mathcal{E}) \in E \wedge (C, \mathcal{C}) \in C \wedge \text{isAtl}_{\text{Ob}}^{(proper)}(A, (E, \mathcal{E}), (C, \mathcal{C})) \\ \text{(full, S-max, max, S-max-full, max-full)} \end{array} \right\} \quad (10.6)$$

Let E be a set of topological spaces and C a set of model spaces. Then

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C) \stackrel{\text{def}}{=} \left\{ (A, E, (C, \mathcal{C})) \mid \begin{array}{l} E \in E \wedge (C, \mathcal{C}) \in C \wedge \text{isAtl}_{\text{Ob}}^{(proper)}(A, E, (C, \mathcal{C})) \\ \text{(full, S-max, max, S-max-full, max-full)} \end{array} \right\} \quad (10.7)$$

{lem:M-ATLsets}

Lemma 10.14 (Sets of m-atlases). Let E and C be sets of model spaces.

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{(full, S-max, max, S-max-full)}} \quad (10.8)$$

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{full (S-max, max, S-max-full, max-full)}} \subseteq \mathcal{Att}_{\text{Ob}}^{(proper)}(E, C) \quad (10.9)$$

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{S-max-full (max-full)}} \subseteq \mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{S-max (max)}} \quad (10.10)$$

Proof. The result follows from $\stackrel{\text{def:M-atlas}}{\text{definition 10.1 (M-atlases)}}$ on page 47, definition 10.8 (Maximal (semi-maximal) m-atlases) on page 50 and definition 10.13 above. □

{def:M-ATLmax}
{def:M-ATLsets}

$$\mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{S-max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(proper)}(E, C)_{\text{full}} \quad (10.11)$$

Proof. The result follows from definition 10.8 (Maximal (semi-maximal) m-atlases) on page 50 and definition 10.13 above. □

{def:M-ATLmax}
{def:M-ATLsets}

{cor:M-ATLsets}

Corollary 10.15 (Sets of m-atlases). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{(\text{full}, \text{S-max}, \text{max}, \text{S-max-full})} \quad (10.12)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{full (S-max, max, S-max-full, max-full)}} \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (10.13)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{S-max-full (max-full)}} \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{S-max (max)}} \quad (10.14)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{S-max-full}} \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C})_{\text{full}} \quad (10.15)$$

Proof. The result follows from definition 10.1 (M-atlases) on page 47, definition 10.8 (Maximal (semi-maximal) m-atlases) on page 50 and definition 10.13 above. $\square$

{def:M-ATLmax}
{def:M-ATLsets}

11 M-atlas (near) morphisms and functors

{sec:M-ATLmorph}

This section introduces a taxonomy for morphisms between m-atlases, defines categories of m-atlases, defines functors among them and proves some basic results.

11.1 M-atlas (near) morphisms

{sub:M-ATLmorph}

This subsection introduces the notions of m-atlas morphisms, m-atlas near morphisms, strict m-atlas (near) morphisms and semi-strict m-atlas (near) morphisms. Only (strict, semi-strict) m-atlas morphisms are needed for the exposition of local coordinate spaces, as m-atlas near morphisms between maximal atlases are proven to be m-atlas morphisms.

11.1.1 Definitions of m-atlas (near) morphisms

{def:M-ATLmorph}

Definition 11.1 (M-atlas morphisms for model spaces). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i, \mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ a pair of model functions.

$\mathbf{f}$ is an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$ and an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{E}^1, \mathcal{E}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, \mathcal{E}^1, \mathcal{C}^1, \mathbf{A}^2, \mathcal{E}^2, \mathcal{C}^2, f_0, f_1)$, iff for any charts $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$ and any $u^1 \in I$, there exists a sub-chart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , a model neighborhood $U'^2 \subseteq U^2$

of $u^2 \stackrel{\text{def}}{=} f_0(u^1)$ and a chart $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow[\cong]{\approx} \hat{V}'^2) \in \mathbf{A}^2$ at u^2 such that $f_0[U'^1] \subseteq U'^2$ and diagram (11.1) below is commutative, as shown in fig. 4 (Completed m-atlas morphism) on page 54. {fig:AtlM1}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} \{u^1\} \xrightarrow{i} U'^1 \xrightarrow[\cong]{\phi^1} V'^1 \xrightarrow{f_1} \hat{V}'^2, \\ U'^1 \xrightarrow{f_0} U'^2 \xrightarrow[\cong]{\phi'^2} \hat{V}'^2 \end{array} \right. \quad (11.1) \quad \{\text{eq:AtlM1}\}$$




Figure 4: Completed m-atlas morphism

{fig:AtlM1}

The triple $(\mathbf{f}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2))$ will refer to $\mathbf{f}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is also a constrained $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$ and a constrained $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, \mathcal{C}^1, \mathcal{C}^1, \mathbf{A}^2, \mathcal{C}^2, \mathcal{C}^2, f_0, f_1)$, iff f_0 and f_1 are constrained, i.e., $f_0[\mathbf{E}^1]$ is contained in a model neighborhood of $\mathbf{E}^2$ and $f_1[\mathbf{C}^1]$ is contained in a model neighborhood of $\mathbf{C}^2$.

Remark 11.2. If $\mathbf{f}$ is constrained then $\text{Top}(\mathbf{E}^1)$ is a model neighborhood of $\mathbf{E}^1$ and $\text{Top}(\mathbf{C}^1)$ is a model neighborhood of $\mathbf{C}^1$.

{def:M-ATLmorphotop}

Definition 11.3 (M-atlas morphisms for topological spaces). Let $\mathcal{C}^i, i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space and $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$.

A pair of functions $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ is an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas mor-

phism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^i$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^i$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff it is an $\begin{smallmatrix} E^1 & - & E^2 \\ \text{triv} & & \text{triv} \end{smallmatrix}$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$.

The triple $(\mathbf{f}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2))$ will refer to $\mathbf{f}$ considered as an E^1 - E^2 m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.


$\mathbf{f}$ is also a constrained E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and a constrained E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff f_1 is constrained, i.e., $f_1[\mathbf{C}^1]$ is contained in a model neighborhood of $\mathbf{C}^2$.

{def:M-ATLmorphnear}

Definition 11.4 (M-atlas near morphisms for model spaces). Let $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $\mathbf{E}^i \in \mathcal{C}^i$, $\mathbf{C}^i \in \mathcal{C}^i$, $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ a pair of model functions.

$\mathbf{f}$ is an E^1 - E^2 m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{near}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$ and an $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated

as $\text{isAtl}_{\text{Ar}}^{\text{near}}(\mathbf{A}^1, \mathcal{C}^1, \mathcal{C}^1, \mathbf{A}^2, \mathcal{C}^2, \mathcal{C}^2, f_0, f_1)$ iff for any charts $(U^i, V^i, \phi^i: U^i \xrightarrow{\cong} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, diagram (11.2) below is strongly M-locally nearly commutative in $\mathbf{C}^2$, i.e., for any $u^1 \in I$ there are model neighborhoods $U'^1 \subseteq I, V'^1 \subseteq V^1, U'^2 \subseteq U^2, V'^2 \subseteq V^2, \hat{V}'^2 \subseteq \mathbf{C}^2$ and an isomorphism

$\hat{f}: V'^2 \xrightarrow{\cong} \hat{V}'^2$ ¹⁰ such that eqs. (11.3) to (11.8) below hold, as shown in  fig. 5 {eq:xin}

to 6.

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} f_0: I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \longrightarrow U^2, \phi^2: U^2 \xrightarrow{\cong} V^2, \\ \phi^1: I \hookrightarrow V^1, f_1: V^1 \longrightarrow \mathbf{C}^2 \end{array} \right. \quad (11.2) \quad \{\text{eq:AtlN1}\}$$

$$u^1 \in U'^1 \quad (11.3) \quad \{\text{eq:xin}\}$$

$$f_0[U'^1] \subseteq U'^2 \quad (11.4) \quad \{\text{eq:fup1in}\}$$

¹⁰This reverses the direction of the arrow $\hat{f}: U_m \xrightarrow{\cong} V_n$ from definition 1.24 (Nearly commutative diagrams in category $\mathcal{C}$) on page 15. This is permissible since $\hat{f}$ is an isomorphism {def:NCD}

$$\phi^1[U'^1] = V'^1 \quad (11.5) \quad \{\text{eq:fup1in}\}$$

$$f_1[V'^1] \subseteq \hat{V}'^2 \quad (11.6) \quad \{\text{eq:fvp1in}\}$$

$$\phi^2[U'^2] = V'^2 \quad (11.7) \quad \{\text{eq:fvp1in}\}$$

$$\hat{f} \circ \phi^2 \circ f_0 = f_1 \circ \phi^1 \quad (11.8) \quad \{\text{eq:fphigeqgphi}\}$$

Remark 11.5. $(U'^i, V'^i, \phi^i \upharpoonright_{U'^i, V'^i})$ need not be a chart of $\mathbf{A}^i$.

$$\begin{array}{ccc} I = U^1 \cap f_0^{-1}[U^2] & \xrightarrow{f_0} & U^2 \\ \downarrow \phi^1 & & \downarrow \cong \phi^2 \\ V^1 & \xrightarrow{f_1} & C^2 \end{array}$$

Figure 5: Uncompleted m-atlas near morphism

{fig:AtlN1}

$$\begin{array}{ccccccc} & & u^1 & & & & \\ & & \downarrow i & & & & \\ I = U^1 \cap f_0^{-1}[U^2] & \xleftarrow{i} & U'^1 & \xrightarrow{f_0} & U'^2 & \xrightarrow{i} & U^2 \\ \downarrow \phi^1 & & \downarrow \cong \phi^1 & & \downarrow \cong \phi^2 & & \downarrow \cong \phi^2 \\ V^1 & \xleftarrow{i} & V'^1 & \xrightarrow{f_1} & \hat{V}'^2 & \xrightarrow[\cong]{\hat{f}} & V'^2 \xrightarrow{i} V^2 \end{array}$$

Figure 6: Completed m-atlas near morphism

{fig:AtlN2}

The triple $(\mathbf{f}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2))$ will refer to $\mathbf{f}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Definition 11.6 (M-atlas near morphisms for topological spaces). Let $\mathcal{C}^i, i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the

{def:M-ATLmorphptopnear}

coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be a pair of continuous functions.

$\mathbf{f}$ is an E^1 - E^2 m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{near}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and a E^1 - E^2 m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{near}}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff it is an E^1 - E^2 m-atlas near morphism triv triv of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model spaces $\mathbf{C}^1, \mathbf{C}^2$.

The triple $(\mathbf{f}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2))$ will refer to $\mathbf{f}$ considered as an E^1 - E^2 m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

{def:M-ATLident}

Definition 11.7 (M-atlas inclusion and identity morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{C}^i$
3. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
4. $\mathbf{C}^i \in \mathcal{C}^i$
5. $\mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$
6. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$
7. $\mathbf{A}^1 \subseteq \mathbf{A}^2$

The E^1 - E^2 m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\text{Id}_{(E^2, \mathbf{C}^2)}^{(E^1, \mathbf{C}^1)}$ considered as an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The m-atlas inclusion morphism of $(\mathbf{A}^1, E^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, E^2, \mathbf{C}^2)$ is $\text{Id}_{(A^2, E^2, C^2)}^{(A^1, E^1, C^1)} \stackrel{\text{def}}{=} (\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2))$, $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ considered as an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The E^i m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^1, \mathbf{C}^2$ is $\text{Id}_{(E^i, C^i)}$ considered as an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The m-atlas identity morphism of $(\mathbf{A}^i, E^i, \mathbf{C}^i)$ is $\text{Id}_{(A^i, E^i, C^i)} \stackrel{\text{def}}{=} (\text{Id}_{(E^i, C^i)}, (\mathbf{A}^i, E^i, \mathbf{C}^i), (\mathbf{A}^i, E^i, \mathbf{C}^i))$, $\text{Id}_{(E^i, C^i)}$ considered as an E^i m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

Let

1. E^i be a topological space, $i = 1, 2$
2. $E^1 \stackrel{\text{subsp}}{\subseteq} E^2$
3. $\mathcal{C}^i$ be a model category

4. $\mathcal{C}^i \in \mathcal{C}^i$
5. $\mathcal{C}^1 \stackrel{\text{mod}}{\subseteq} \mathcal{C}^2$
6. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathcal{C}^i$
7. $\mathbf{A}^1 \subseteq \mathbf{A}^2$

The E^1 - E^2 m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$ is $\mathbf{Id}_{(\mathcal{C}^2, \mathcal{C}^1)}^{(E^1, \mathcal{C}^1)}$.

The m-atlas inclusion morphism of $(\mathbf{A}^1, E^1, \mathcal{C}^1)$ to $(\mathbf{A}^2, E^2, \mathcal{C}^2)$ is $\mathbf{Id}_{(\mathbf{A}^2, E^2, \mathcal{C}^2)}^{(\mathbf{A}^1, E^1, \mathcal{C}^1)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(E^2, \mathcal{C}^2)}^{(E^1, \mathcal{C}^1)}, (\mathbf{A}^1, E^1, \mathcal{C}^1), (\mathbf{A}^2, E^2, \mathcal{C}^2) \right)$, $\mathbf{Id}_{(E^2, \mathcal{C}^2)}^{(E^1, \mathcal{C}^1)}$ considered as an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$.

The E^i m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathcal{C}^i$ is $\mathbf{Id}_{(\mathcal{C}^i, \mathcal{C}^i)}^{(E^i, \mathcal{C}^i)}$.

The m-atlas identity morphism of $(\mathbf{A}^i, E^i, \mathcal{C}^i)$ is

$\mathbf{Id}_{(\mathbf{A}^i, E^i, \mathcal{C}^i)} \stackrel{\text{def}}{=} (\mathbf{Id}_{(E^i, \mathcal{C}^i)}^{(E^i, \mathcal{C}^i)}, (\mathbf{A}^i, E^i, \mathcal{C}^i), (\mathbf{A}^i, E^i, \mathcal{C}^i))$, $\mathbf{Id}_{(E^i, \mathcal{C}^i)}^{(E^i, \mathcal{C}^i)}$ considered as an E^i m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathcal{C}^i$.

{def:M-ATLmorphequiv}

Definition 11.8 (Equivalence of m-atlas (near) morphisms). Let E^i , $\mathcal{C}^i$, $i = 1, 2$, be model spaces, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathcal{C}^i$, and

$\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathcal{C}^1 \rightarrow \mathcal{C}^2)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_0: E^1 \rightarrow E^2, g_1: \mathcal{C}^1 \rightarrow \mathcal{C}^2)$ be m-atlas (near) morphisms of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$. Then $\mathbf{f}$ is equivalent to $\mathbf{g}$ as an E^1 - E^2 m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$, abbreviated $\mathbf{f}: (\mathbf{A}^1, E^1, \mathcal{C}^1) \rightarrow (\mathbf{A}^2, E^2, \mathcal{C}^2)$ is equivalent to $\mathbf{g}: (\mathbf{A}^1, E^1, \mathcal{C}^1) \rightarrow (\mathbf{A}^2, E^2, \mathcal{C}^2)$ and $(\mathbf{f}, (\mathbf{A}^1, E^1, \mathcal{C}^1), (\mathbf{A}^2, E^2, \mathcal{C}^2))$ is equivalent to $(\mathbf{g}, (\mathbf{A}^1, E^1, \mathcal{C}^1), (\mathbf{A}^2, E^2, \mathcal{C}^2))$ iff $f_0 = g_0$.

By abuse of language we write $\mathbf{f}$ is equivalent to $\mathbf{g}$ when the spaces and atlases are understood by context.

Let E^i , $i = 1, 2$, be a topological space, $\mathcal{C}^i$ be a model space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathcal{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathcal{C}^1 \rightarrow \mathcal{C}^2)$, and

$\mathbf{g} \stackrel{\text{def}}{=} (g_0: E^1 \rightarrow E^2, g_1: \mathcal{C}^1 \rightarrow \mathcal{C}^2)$ be m-atlas (near) morphisms of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$. Then $\mathbf{f}$ is equivalent to $\mathbf{g}$ as an E^1 - E^2 m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$, abbreviated $\mathbf{f}: (\mathbf{A}^1, E^1, \mathcal{C}^1) \rightarrow (\mathbf{A}^2, E^2, \mathcal{C}^2)$ is equivalent to $\mathbf{g}: (\mathbf{A}^1, E^1, \mathcal{C}^1) \rightarrow (\mathbf{A}^2, E^2, \mathcal{C}^2)$, iff $f_0 = g_0$.

By abuse of language we write $\mathbf{f}$ is equivalent to $\mathbf{g}$ when the spaces and atlases are understood by context.

11.1.2 Definitions of semi-strict and strict

{def:M-ATLmorphsemi}

Definition 11.9 (Semi-strict m-atlas (near) morphisms). Let $\mathcal{E}^i$, $\mathcal{C}^i$, $i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i$, $\mathcal{C}^i \in \mathcal{C}^i$, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathcal{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathcal{C}^1 \rightarrow \mathcal{C}^2)$ an E^1 - E^2 m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$.

$\mathbf{f}$ is a semi-strict $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$, iff f_0 is a local m-morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$, f_1 is a local m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$ and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local $\mathbf{C}^1\text{-}\mathbf{C}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff each $E^i \in \mathbf{E}^i$, each $C^i \in \mathbf{C}^i$, $\mathbf{E}^1 \stackrel{\text{Ob}}{\subseteq} \mathbf{E}^2$ and $\mathbf{C}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathbf{C}^2$.

$\mathbf{f}$ is a semi-strict $\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff f_0 is a local m-morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$, f_1 is a local $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$ and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff each $E^i \in \mathbf{E}^i$, $\mathbf{E}^1 \stackrel{\text{Ob}}{\subseteq} \mathbf{E}^2$ and $\mathcal{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$.

$\mathbf{f}$ is a semi-strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$, f_1 is a local m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$ and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local $\mathbf{C}^1\text{-}\mathbf{C}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff each $C^i \in \mathbf{C}^i$, $\mathcal{E}^1 \stackrel{\text{Ob}}{\subseteq} \mathcal{E}^2$ and $\mathbf{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathbf{C}^2$.

$\mathbf{f}$ is a semi-strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$ and f_1 is a local $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$ and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ and $\mathcal{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$.

By abuse of language we will write $\mathbf{f}$ is coherent when it is a $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2$ ($\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$) m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$

where the model spaces, model categories and atlases are understood by context.

{def:M-ATLmorphstrict}

Definition 11.10 (Strict m-atlas (near) morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{E}^i$
3. $C^i \in \mathcal{C}^i$
4. A^i be an m-atlas of E^i in the coordinate space C^i
5. $f \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ be an m-atlas (near) morphism

f is a strict $E^1 - E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\text{strict})(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$, iff f_0 is an m-morphism of E^1 to E^2 , f_1 is an m-morphism of C^1 to C^2 and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow U^2$ is a morphism of $\text{Mod}(V^2, C^2)$.

It is also a coherent strict $E^1 - E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $E^i \in \mathcal{E}^i$, each $C^i \in \mathcal{C}^i$, $E^1 \stackrel{\text{Ob}}{\subseteq} E^2$ and $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$.

f is a strict $E^1 - E^2 - \mathcal{E}^1 - \mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\text{strict})(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}^1, \mathcal{C}^2)$, iff f_0 is an m-morphism of E^1 to E^2 , f_1 is an m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of $\mathcal{C}^2$.

It is also a coherent strict $E^1 - E^2 - \mathcal{E}^1 - \mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $E^i \in \mathcal{E}^i$, $E^1 \stackrel{\text{Ob}}{\subseteq} E^2$ and $\mathcal{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$.

f is a strict $\mathcal{E}^1 - \mathcal{C}^2 - E^1 - E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\text{strict})(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}^1, \mathcal{C}^2)$, iff f_0 is an m-morphism of $\mathcal{E}^1$ to $\mathcal{C}^2$, f_1 is an m-morphism of C^1 to C^2 and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow U^2$ is a morphism of $\text{Mod}(\phi^1[I], C^2)$.

It is also a coherent strict $\mathcal{E}^1 - \mathcal{C}^2 - E^1 - E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$, $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$ and $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$.

f is a strict $\mathcal{E}^1 - \mathcal{C}^2 - E^1 - E^2 - \mathcal{E}^1 - \mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{near})}(\text{strict})(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}^1, \mathcal{C}^2, \mathcal{E}^1, \mathcal{C}^2)$, iff f_0 is an m-morphism of

$\mathcal{E}^1$ to $\mathcal{E}^2$, f_1 is an m-morphism of $\mathcal{E}^1$ to $\mathcal{E}^2$ and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathcal{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of $\mathcal{E}^2$.

It is also a coherent strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ and $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$.

By abuse of language we will write $\mathbf{f}$ is coherent when it is a $(\mathcal{E}^1\text{-}\mathcal{E}^2)\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ $(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ where the model spaces, model categories and atlases are understood by context.

11.1.3 Abbreviated nomenclature for m-atlas morphisms

{def:M-ATLmorphabbrev}

Definition 11.11 (Abbreviated nomenclature for m-atlas (near) morphisms). Let $\mathcal{E}^i, \mathcal{E}^i, i = 1, 2$, be model categories, $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{E}^i, \mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ a pair of model functions.

$\mathbf{f}$ is a constrained semi-strict (strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const, const-near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$, iff it is a constrained $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-semi-strict (strict)

atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$ and it is a semi-strict (strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const, const-near, near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, (semi-strict, strict)

strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const, const-near, near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff it is a (constrained) (semi-strict, (semi-strict, strict)

strict, strict) $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const, const-near, near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{C}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, (semi-strict, strict)

strict) $\mathbf{E}^1\text{-}\mathbf{E}^1$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const, const-near, near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, (semi-strict, strict)

strict) $\mathbf{E}^1\text{-}\mathbf{E}^1$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{C}^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^1\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}\mathcal{C}^1$ m-atlas (near) morphism of A^1 to A^2 in the coordinate space $\mathbf{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, f_0, f_1, \mathcal{C}^1)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^1\text{-}\mathcal{C}^1\text{-}\mathcal{C}^1$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

Let $\mathcal{C}^i, i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i, E^i$ be a topological space, A^i be an m-atlas of E^i in the coordinate space $\mathbf{C}^i, f_0: E^1 \rightarrow E^2$ be continuous, $f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2$ be a model function and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and a (constrained) (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate model spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (constrained) (near) morphism of A^1 to A^2 in the coordinate space $\mathbf{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}E^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of A^1 to A^2 in the coordinate model spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $E^1\text{-}E^2\text{-}\mathcal{C}^1$ m-atlas (near) morphism of A^1 to A^2 in the coordinate model category $\mathcal{C}^1$, abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, f_0, f_1, \mathcal{C}^1)$, iff it is a (constrained) (semi-strict, strict) $E^1\text{-}E^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^1$ m-atlas (near) morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^1$.

Similar definitions apply with restrictions on the admissible atlases, i.e., $\mathbf{f}$ is a (constrained) (semi-strict, strict) full (semi-maximal, maximal, full semi-maximal, full maximal) $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1$, abbreviated as

$$\begin{aligned} \text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2) &\iff \\ &\quad \text{full (S-max, max, S-max-full, max-full)} \\ &\quad \text{(semi-strict, strict)} \\ \text{isAtl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2) &\wedge \\ &\quad \text{semi-strict (strict)} \\ \text{isAtl}_{\text{Ob}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1) &\wedge \\ \text{full (S-max, max, S-max-full, max-full)} & \\ \text{isAtl}_{\text{Ob}}(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) & \\ \text{full (S-max, max, S-max-full, max-full)} & \end{aligned}$$

11.1.4 Proclamations on m-atlas (near) morphisms

{lem:M-ATLmorph}

Lemma 11.12 (M-atlas (near) morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, \mathcal{E}'^i, \mathcal{C}'^i, i = 1, 2$, be model categories
2. $\mathcal{E}^i \xrightarrow{\text{full-cat}} \mathcal{E}'^i, \mathcal{C}^i \xrightarrow{\text{full-cat}} \mathcal{C}'^i$
3. $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i, \mathbf{C}'^i \in \mathcal{C}'^i, \mathbf{C}^i \xrightarrow{\text{mod}} \mathbf{C}'^i$
4. $f'_1: \mathbf{C}'^1 \rightarrow \mathbf{C}'^2$ be a model function such that $f'_1[\mathbf{C}^1] \subseteq \mathbf{C}^2$ and $f_1 \stackrel{\text{def}}{=} f'_1 \upharpoonright_{\mathbf{C}^1, \mathbf{C}^2}$ is a model function
5. $\mathbf{A}^i$ be an M-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$
6. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ be a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2$ $(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$
7. $\mathbf{f}' \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f'_1: \mathbf{C}^1 \rightarrow \mathbf{C}'^2)$,
 $\mathbf{f}'' \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f'_1: \mathbf{C}'^1 \rightarrow \mathbf{C}'^2)$ be pairs of model functions

Then

{M-ATLmorph:morphisnear}

1. If $\mathbf{f}$ is an m-atlas morphism then $\mathbf{f}$ is an m-atlas near morphism.

Proof. Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, be charts with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, u^1 \in I$ and $u^2 \stackrel{\text{def}}{=} f_0(u^1)$. Let $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$

and $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in \mathbf{A}^2$ be charts as in definition 11.1 (M-atlas morphisms for model spaces) on page 53. $U'^2 \subseteq I$ and eqs. (11.3) to (11.8) of definition 11.4 (M-atlas near morphisms for model spaces) on page 55 hold. $\square$

2. If $\mathbf{A}^i, i = 1, 2$, is semi-maximal and $\mathbf{f}$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2$ $(\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then it is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2(\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof.

$\mathbf{f}$ is an M-atlas morphism. Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$. Diagram (11.2) in definition 11.4 (M-atlas near morphisms for model spaces) on page 55 is M-locally nearly commutative in $\mathbf{C}^2$, i.e., for any $e^1 \in I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2]$ there are model neighborhoods $U'^1 \subseteq I, V'^1 \subseteq V^1, U'^2 \subseteq U^2, V'^2 \subseteq V^2, \hat{V}'^2 \subseteq \mathbf{C}^2$ and an isomorphism $\hat{f}: V'^2 \xrightarrow{\sim} \hat{V}'^2$ such that eqs. (11.3) to (11.8) in definition 11.4 hold, as shown in figs. 5 to 6. $f_0[U'^1] \subseteq U'^2$ by eq. (11.4) definition 11.4. $f_1[V'^1] \subseteq \hat{V}'^2$ by eq. (11.6) definition 11.4.

Since $\mathbf{A}^i$ is semi-maximal, $(U'^i, V'^i, \phi^i: U'^i \xrightarrow{\sim} V'^i) \in \mathbf{A}^i$ and $(U'^2, \hat{f}[V'^2], \hat{f} \circ \phi^2) \in \mathbf{A}^2$.

Define $\phi'^2: U'^2 \xrightarrow{\sim} \hat{f}[V'^2] \stackrel{\text{def}}{=} \hat{f} \circ \phi^2$. Then Diagram (11.1) is commutative.

$\mathbf{f}$ is (semi-strict, strict). The definitions of strict and semi-strict are the same for m-atlas near morphisms and for m-atlas morphisms. $\square$

3. Let $E^i \stackrel{\text{def}}{=} \text{Top}(\mathbf{E}^i), i = 1, 2$. If each $\mathbf{C}^i$ is fine grained then $\hat{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ is a (semi-strict, strict) $E^1\text{-}E^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof.

- (a) $\mathbf{A}^i$ is an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ by lemma 10.3 (M-atlases) on page 48.
- (b) $f_0: E^1 \longrightarrow E^2$ is a model function, hence continuous, hence by lemma 7.4 (Model functions and trivial model spaces) on page 28, $f_0: E^1_{\text{triv}} \longrightarrow E^2_{\text{triv}}$ is a model function.

- (c) $f_1: \mathcal{C}^1 \longrightarrow \mathcal{C}^2$ is a model function by hypothesis.
- (d) Any model neighborhood of E^i is open, hence a model neighborhood of $\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}$.
- (e) Let $(U^i, V^i, \phi^i) \in \mathcal{A}^i$, $i = 1, 2$. All of the model neighborhoods of E^i mentioned in definition 11.4 (M-atlas near morphisms for model spaces) on page 55 and definition 11.1 (M-atlas morphisms for model spaces) on page 53 are also model neighborhoods of $\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}$. {def:M-ATLmorphnear}
{def:M-ATLmorph}

□

{M-ATLmorph:istosup}

4. $\mathbf{f}'$ is an E^1 - E^2 m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}^1, \mathcal{C}'^2$ and $\mathbf{f}''$ is an E^1 - E^2 m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}'^1, \mathcal{C}'^2$.

Proof. $f_0: E^1 \longrightarrow E^2$ is a model function by definition 11.4 (M-atlas near morphisms for model spaces) on page 55 (definition 11.1 (M-atlas morphisms for model spaces) on page 53). {def:M-ATLmorphnear}
{def:M-ATLmorph}

$f'_1: \mathcal{C}'^1 \longrightarrow \mathcal{C}'^2$ and $f''_1: \mathcal{C}^1 \longrightarrow \mathcal{C}'^2$ are model functions by hypothesis.

$\mathcal{C}^i \stackrel{\text{mod}}{\subseteq} \mathcal{C}'^i$ by hypothesis,

$\mathcal{C}^i$ is an open subset of $\mathcal{C}'^i$ by lemma 2.9 (Model subspaces) on page 21.

{lem:modsub}

The expanded model space does not change the atlases or the commutation.

□

{M-ATLmorph:strictissemi}

5. If $\mathbf{f}$ is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}^1, \mathcal{C}^2$ then $\mathbf{f}$ is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}^1, \mathcal{C}^2$

Proof. There are four overlapping cases:

E^i without category: f_0 is an m-morphism of E^1 to E^2 by definition 11.10 (Strict m-atlas (near) morphisms) on page 60. f_0 is a local m-morphism of E^1 to E^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphstrict}
{lem:modMmorph}

$\mathcal{C}^i$ without category: f_1 is an m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by definition 11.10. f_1 is a local m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by lemma 7.13. {def:M-ATLmorphstrict}
{lem:modMmorph}

E^i with category: f_0 is an $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.10. f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by lemma 7.13. {def:M-ATLmorphstrict}
{lem:modMmorph}

$\mathcal{C}^i$ with category: f_1 is a $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by definition 11.10. f_1 is a local $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by lemma 7.13. {def:M-ATLmorphstrict}
{lem:modMmorph}

□

{M-ATLmorph:strictSstrictis

6. If f is a coherent semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , then f is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. There are four overlapping cases.

- E^i without category:** f_0 is a local m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $E^1 \xrightarrow{\text{strict-mod}} E^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\text{Set}(E^2)$ is a model neighborhood of E^2 by definition 2.6 (Model subspaces) on page 20. The result follows by item 6 (Restricted sheaf condition) of definition 2.6. {def:M-ATLmorphsemi}
{def:M-ATLmorphsemi}
{def:modsub}
{mod:sheaf}
- E^i with category:** f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^1 \xrightarrow{\text{strict-cat}} \mathcal{E}^2$ by definition 11.9. $\mathcal{E}^2$ satisfies the restricted sheaf condition by definition 7.7 (Model categories) on page 29 and the result follows. {def:M-ATLmorphsemi}
{def:M-ATLmorphsemi}
- C^i without category:** f_0 is a local m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $C^1 \xrightarrow{\text{strict-mod}} C^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\text{Set}(C^2)$ is a model neighborhood of C^2 by definition 2.6 (Model subspaces) on page 20. The result follows by item 6 (Restricted sheaf condition) of definition 2.6. {def:M-ATLmorphsemi}
{def:M-ATLmorphsemi}
{def:modsub}
{mod:sheaf}
- C^i with category:** f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^1 \xrightarrow{\text{strict-cat}} \mathcal{E}^2$ by definition 11.9. $\mathcal{E}^2$ satisfies the restricted sheaf condition by definition 7.7 (Model categories) on page 29 and the result follows. {def:M-ATLmorphsemi}
{def:M-ATLmorphsemi}

□

{M-ATLmorph:SstrictisSstrictis

7. If f is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 then f' is a semi-strict $(\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-})E^1\text{-}E^2(-\mathcal{E}'^1\text{-}\mathcal{E}'^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C'^2 .

Proof. There are four overlapping cases.

- E^i without category:** f_0 is a (strict) local m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}
- E^i with category:** f_0 is a (strict) local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^i \xrightarrow{\text{full-cat}}$ {def:M-ATLmorphsemi}

- $\mathcal{E}^i$ by hypothesis. f_0 is a (strict) local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {lem:modMmorph}
- C^i without category:** f_1 is a (strict) local m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $C^2 \stackrel{\text{mod}}{\subseteq} C'^2$ by hypothesis. f_1 is a (strict) local m-morphism of C^1 to C'^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}
- C^i with category:** f_0 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is a (strict) local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of C^1 to C^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}
- {M-ATLmorph:strictisstrict}
8. If f is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 then f' is a strict $(\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}'^1\text{-}\mathcal{E}'^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C'^2 .
- Proof.* There are four overlapping cases.
- E^i without category:** f_0 is an m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}
- E^i with category:** f_0 is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is an $\mathcal{E}'^1$ - $\mathcal{E}'^2$ m-morphism of E^1 to E^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}
- C^i without category:** f_1 is an m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $C^2 \stackrel{\text{mod}}{\subseteq} C'^2$ by hypothesis. f_1 is an m-morphism of C^1 to C'^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}
- C^i with category:** f_0 is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of C^1 to C^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is a $\mathcal{E}'^1$ - $\mathcal{E}'^2$ m-morphism of C^1 to C^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}
- {M-ATLmorph:Sstrictlocalis}
9. If f is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and f'_1 is a local m-morphism of C'^1 to C'^2 (a local $\mathcal{E}'^1$ - $\mathcal{E}'^2$ m-morphism) of C'^1 to C'^2 then f'' is a semi-strict $(\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}'^1\text{-}\mathcal{E}'^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C'^1, C'^2 .

Proof. There are four overlapping cases:

E^i without categories: f_0 is a local m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}

E^i with categories: f_0 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.9. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}'^i$ by hypothesis. f_0 is a local $\mathcal{E}'^1$ - $\mathcal{E}'^2$ m-morphism of E^1 to E^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorphsemi}
{lem:modMmorph}

C^i without categories: f'_1 is a local m-morphism of C'^1 to C'^2 by hypothesis.

C^i with categories: f'_1 is a local $\mathcal{C}'^1$ - $\mathcal{C}'^2$ m-morphism of C^1 to C^2 by hypothesis.

□

{M-ATLmorph:strictmorphiss}

10. If f is a strict $(\mathcal{E}^1$ - $\mathcal{E}^2$ -) E^1 - E^2 - $(\mathcal{C}^1$ - $\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1 , C^2 and f'_1 is an m-morphism of C'^1 to C'^2 (a $\mathcal{C}'^1$ - $\mathcal{C}'^2$ m-morphism) of C'^1 to C'^2 then f'' is a strict $(\mathcal{E}'^1$ - $\mathcal{E}'^2$ -) E^1 - E^2 - $(\mathcal{C}'^1$ - $\mathcal{C}'^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C'^1 , C'^2 .

Proof. There are four overlapping cases:

E^i without categories: f_0 is an m-morphism of E^1 to E^2 by definition 11.1 (M-atlas morphisms for model spaces) on page 53. {def:M-ATLmorph}

E^i with categories: f_0 is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.1. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}'^i$ by hypothesis. f_0 is an $\mathcal{E}'^1$ - $\mathcal{E}'^2$ m-morphism of E^1 to E^2 by lemma 7.13 ((Local) m-morphisms) on page 32. {def:M-ATLmorph}
{lem:modMmorph}

C^i without categories: f'_1 is an m-morphism of C'^1 to C'^2 by hypothesis.

C^i with categories: f'_1 is a $\mathcal{C}'^1$ - $\mathcal{C}'^2$ m-morphism of C^1 to C^2 by hypothesis.

□

{cor:M-ATLmorph}

Corollary 11.13 (M-atlas morphisms). Let

1. $\mathcal{E}^i, \mathcal{E}'^i, i = 1, 2$, be model categories
2. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}'^i$
3. $C^i \in \mathcal{E}^i$
4. $C'^i \in \mathcal{E}'^i$
5. $C^i \stackrel{\text{mod}}{\subseteq} C'^i$

6. $f'_1: \mathbf{C}'^1 \longrightarrow \mathbf{C}'^2$ be a model function such that $f'_1[\mathbf{C}^1] \subseteq \mathbf{C}^2$ and $f_1 \stackrel{\text{def}}{=} f'_1 \upharpoonright_{\mathbf{C}^1, \mathbf{C}^2}$ is a model function
7. E^i be a topological space
8. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$
9. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be a (semi-strict, strict) $(\mathcal{E}^1 \cdot \mathcal{E}^2 \cdot)E^1 \cdot E^2$ $(\cdot \mathcal{E}^1 \cdot \mathcal{E}^2)$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$

If $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ is a (semi-strict, strict) $E^1 \cdot E^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then it is a (semi-strict, strict) $E^1 \cdot E^2$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$

If $\mathbf{A}^i, i = 1, 2$, is semi-maximal and $\mathbf{f}$ is a (semi-strict, strict) $E^1 \cdot E^2$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then it is a (semi-strict, strict) $E^1 \cdot E^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

If $\mathbf{f}$ is a semi-strict (strict) $E^1 \cdot E^2 \cdot \mathcal{E}^1 \cdot \mathcal{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then $\mathbf{f}$ is a semi-strict (strict) $E^1 \cdot E^2 \cdot \mathcal{E}^1 \cdot \mathcal{E}^2$ m-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. The result follows from definition 11.11 (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61. $\square$

{def:M-ATLmorphabbrev}

{lem:M-ATLmorphequiv}

Lemma 11.14 (Equivalence of m-atlas (near) morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i$,
3. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$,
4. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, g_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be equivalent m-atlas near morphisms of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

If $\mathbf{f}$ is a (coherent) semi-strict $(\mathcal{E}^1 \cdot \mathcal{E}^2 \cdot)E^1 \cdot E^2 (\cdot \mathcal{E}^1 \cdot \mathcal{E}^2)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then $\mathbf{g}$ is a (coherent) semi-strict $(\mathcal{E}^1 \cdot \mathcal{E}^2 \cdot)E^1 \cdot E^2 (\cdot \mathcal{E}^1 \cdot \mathcal{E}^2)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. Let $v^1 \in \mathbf{C}^1$ be an arbitrary point. Let $(U^1, V^1, \phi^1) \in \mathbf{A}^1$ be an arbitrary chart such that $v^1 \in V^1, u^1 \stackrel{\text{def}}{=} \phi^{1-1}(v^1), (U^2, V^2, \phi^2) \in \mathbf{A}^2$ be an arbitrary chart at $u^2 \stackrel{\text{def}}{=} f_0(u^1), v^2 \stackrel{\text{def}}{=} \phi^2(u^2)$ and $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2]$, as shown in fig. 7 (Uncompleted equivalent m-atlas near morphisms).

{fig:At1E1}

Let $f U^1 \subseteq I, f V^1 \subseteq V^1 f U^2 \subseteq U^2, f V^2 \subseteq V^2$ and $f \hat{V}^2 \subseteq \mathbf{C}^2$ be model neighborhoods and $\hat{f}: f V^2 \xrightarrow{\cong} f \hat{V}^2$ an isomorphism of $\mathbf{C}^2 (\mathcal{C}^2)$ as in definition 11.4 (M-atlas near morphisms for model spaces) on page 55.

{def:M-ATLmorphnear}




Figure 7: Uncompleted equivalent m-atlas near morphisms

{fig:At1E1}

Let $gU'^1 \subseteq I$, $gV'^1 \subseteq V^1$, $gU'^2 \subseteq U^2$, $gV'^2 \subseteq V^2$ and $g\hat{V}'^2 \subseteq \mathcal{C}^2$ be model neighborhoods and $\hat{g}: gV'^2 \xrightarrow{\cong} g\hat{V}'^2$ an isomorphism of $\mathcal{C}^2(\mathcal{C}^2)$ as in definition 11.4. $f\hat{V}'^2 \cap g\hat{V}'^2$ is a model neighborhood of $\mathcal{C}^2$ by [Item 2](#) of definition 2.1 (Model spaces) on page 18.

{def:M-ATLmorphnear}
{def:model}

There are four overlapping cases.

E^i without category: $f_0 = g_0$ by definition 11.8 (Equivalence of m-atlas (near) morphisms) on page 58. f_0 is a local m-morphism of E^1 to E^2 by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59.

{def:M-ATLmorphequiv}
{def:M-ATLmorphsemi}

If f is coherent then $E^1 \xrightarrow{\text{strict-mod}} E^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59.

{def:M-ATLmorphsemi}

C^i without category: f_1 is a local m-morphism of C^1 to C^2 .

$\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \rightarrow V^2$ is a local m-morphism of $\phi^1[I]$ to $\mathcal{C}^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59.

{def:M-ATLmorphsemi}

Let $\hat{V}'^2 \subseteq f\hat{V}'^2 \cap g\hat{V}'^2$ be a model neighborhood of $\mathcal{C}^2$ at v^2 such that

$f_1: V'^1 \xrightarrow{\text{def}} f_1^{-1}[\hat{V}'^2] \rightarrow \hat{V}'^2$ is a morphism of $\mathcal{C}^2$.

The following, shown in [figs. 8 to 9](#), are model neighborhoods of $\mathcal{C}^2$ by

{fig:At1E2}
{def:modfunc}

definition 7.1 (Model functions) on page 27:

$$V'^2 \stackrel{\text{def}}{=} \hat{f}^{-1}[\hat{V}'^2]$$

$$U'^2 \stackrel{\text{def}}{=} \phi^{2-1}[V'^2]$$

$$U'^1 \stackrel{\text{def}}{=} f_0^{-1}[U'^2]$$




Figure 8: Completed equivalent m-atlas near morphisms

{fig:At1E2}




Figure 9: Completed equivalent m-atlas near morphisms

{fig:At1E3}

Then

$$\begin{aligned}
 g_1: V'^1 \longrightarrow \hat{V}'^2 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad \hat{f}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_1: V'^1 \longrightarrow \hat{V}'^2 \quad \circ
 \end{aligned}$$

is a composition of morphisms of $\mathcal{C}^2$, hence a morphism.

If $\mathbf{f}$ is coherent then $\mathcal{C}^1 \xrightarrow{\text{strict-mod}} \mathcal{C}^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}

E^i with category: $f_0 = g_0$ by definition 11.8. f_0 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 11.9. {def:M-ATLmorphequiv}
{def:M-ATLmorphsemi}

If $\mathbf{f}$ is coherent then $\mathcal{E}^1 \xrightarrow{\text{strict-cat}} \mathcal{E}^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}

$\mathcal{C}^i$ with category: f_1 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ and $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \rightarrow V^2$ is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of $\text{Mod}(\phi^1[I], \mathcal{C}^2)$ to $\text{Mod}(V^2, \mathcal{C}^2)$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}

Let $U'^2 \subseteq {}^f U'^2 \cap {}^g U'^2$ be a model neighborhood of $\mathcal{C}^2$ at u^2 such that

$f_0: U'^1 \xrightarrow{\text{def}} f_0^{-1}[U'^2] \rightarrow U'^2$ is a morphism of $\mathcal{E}^2$.

Let $V'^2 \xrightarrow{\text{def}} \phi^2[U'^2]$, $f\hat{V}'^2 \xrightarrow{\text{def}} f[fV'^2]$, $g\hat{V}'^2 \xrightarrow{\text{def}} g[gV'^2]$, $fV'^1 \xrightarrow{\text{def}} f_1^{-1}[f\hat{V}'^2]$ and

$gV'^1 \xrightarrow{\text{def}} g_1^{-1}[g\hat{V}'^2]$. All of these are model neighborhoods by definition 7.1 (Model functions) on page 27 and the corresponding relative model subspaces {def:modfunc}

of V^i are objects of $\mathcal{V}^i$ by item 5 of definition 7.7 (Model categories) on page 29. The restrictions of morphisms to them are morphisms by item 5 of definition 7.7. {modcat:rest}
{def:modcat}
{def:modcat}

Then

$$\begin{aligned} g_1: V'^1 \longrightarrow \hat{V}'^2 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\ &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\ &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\ &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\ &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\ &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\ &\quad \hat{f}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\ &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\ &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\ &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\ &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\ &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\ &\quad f_1: V'^1 \longrightarrow \hat{V}'^2 \quad \circ \end{aligned}$$

is a composition of morphisms of $\mathcal{E}^2$, hence a morphism.

If $\mathbf{f}$ is coherent then $\mathcal{E}^1 \xrightarrow{\text{strict-cat}} \mathcal{E}^2$ by definition 11.9 (Semi-strict m-atlas (near) morphisms) on page 59. {def:M-ATLmorphsemi}

□

{cor:M-ATLmorphequiv}

Corollary 11.15 (Equivalence of m-atlas (near) morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{E}^i$
3. $C^i \in \mathcal{C}^i$
4. A^i be an m-atlas of E^i in the coordinate space C^i
5. A^1 be full
6. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ be a coherent semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2$ $(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .
7. $\mathbf{g} \stackrel{\text{def}}{=} (g_0: E^1 \longrightarrow E^2, g_1: C^1 \longrightarrow C^2)$ be an m-atlas (near) morphisms of A^1 to A^2 in the coordinate spaces C^1, C^2 equivalent to $\mathbf{f}$

Then $\mathbf{g}$ is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. $\mathbf{g}$ is a coherent semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 by lemma 11.14 (Equivalence of m-atlas (near) morphisms) on page 68. The result follows from item 6 of lemma 11.12 (M-atlas (near) morphisms) on page 63. $\square$

`{lem:M-ATLmorphequiv}`
`{M-ATLmorph:strictSstricti}`
`{lem:M-ATLmorphcomp}`

Lemma 11.16 (Composition of m-atlas (near) morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2, 3$, be a model category, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$,
2. A^i an m-atlas of E^i in the coordinate space C^i
3. $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: C^i \longrightarrow C^{i+1})$ and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_0^i: E^i \longrightarrow E^{i+1}, g_1^i: C^i \longrightarrow C^{i+1})$ $i = 1, 2$, be equivalent (semi-strict, strict) $(\mathcal{E}^i\text{-}\mathcal{E}^{i+1}\text{-})E^i\text{-}E^{i+1}(\mathcal{C}^i\text{-}\mathcal{C}^{i+1})$ m-atlas (near) morphisms of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} .

Then:

`{M-morphcomp:nearcompmorph}`

1. If $\mathbf{f}^1$ is an $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and $\mathbf{f}^2$ is an $(\mathcal{E}^2\text{-}\mathcal{E}^3\text{-})E^2\text{-}E^3(\mathcal{C}^2\text{-}\mathcal{C}^3)$ m-atlas near morphism of A^2 to A^3 in the coordinate spaces C^2, C^3 then $\mathbf{f}^2 \circ \mathbf{f}^1$ is an $(\mathcal{E}^1\text{-}\mathcal{E}^3\text{-})E^1\text{-}E^3(\mathcal{C}^1\text{-}\mathcal{C}^3)$ m-atlas near morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof. $f_0^2 \circ f_0^1$ and $f_1^2 \circ f_1^1$ are model functions by lemma 7.5 (Composition of model functions) on page 28.

`{lem:modcomp}`

Let $(U^i, V^i, \phi^i) \in A^i, i = 1, 3$, be a chart such that $I \stackrel{\text{def}}{=} U^1 \cap (f_0^2 \circ f_0^1)^{-1}[U^3] \neq \emptyset$. It suffices to show that diagram (11.9) below is locally nearly commutative.

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} \phi^1: I \rightarrow V^1, f_1^2 \circ f_1^1: V^1 \longrightarrow \mathbf{C}^3, \\ f_0^2 \circ f_0^1: I \longrightarrow U^3, \phi^3: U^3 \xrightarrow{\cong} V^3 \end{array} \right\} \quad (11.9) \quad \{\text{eq:morphcompN1}\}$$

Let $u^1 \in I$, $u^2 \stackrel{\text{def}}{=} f_0^1(u^1)$, $u^3 \stackrel{\text{def}}{=} f_0^2(u^2)$. $(U^2, V^2, \phi^2) \in \mathbf{A}^2$ be a chart at u^2 .

Since $\mathbf{f}^1$ is a morphism, there exists a subchart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\cong} V'^1) \in \mathbf{A}^1$ at u^1 , a model neighborhood $U'^2 \subseteq U^2$ of u^2 and a chart

$(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\cong} \hat{V}'^2) \in \mathbf{A}^2$ at u^2 such that $f_0^1[U'^1] \subseteq U'^2$ and diagram (11.1) in definition 11.1 (M-atlas morphisms for model spaces) on page 54 is commutative, as shown in fig. 4 (Completed m-atlas morphism) on page 54. \{def:M-ATLmorph\}
\{fig:AtlM1\}

Let $I^{2,3} \stackrel{\text{def}}{=} U'^2 \cap f_0^{2-1}[U^3]$. Since $\mathbf{f}^2$ is a near morphism, diagram (11.10) below is locally nearly commutative at u^2 , and thus there exist model neighborhoods $U''^2 \subseteq I^{2,3}$, $U'^3 \subseteq U^3$, $V''^2 \subseteq V'^2$, $V'^3 \subseteq V^3$, $\hat{V}'^3 \subseteq \mathbf{C}^3$ and an isomorphism $\hat{f}: V'^3 \xrightarrow{\cong} \hat{V}'^3$ such that diagram (11.10) below is commutative, i.e., eqs. (11.11) to (11.16) below hold, as shown in figs. 5 to 6. \{eq:xin2\}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^2\} \rightarrow I^{2,3}, \phi'^2: I^{2,3} \rightarrow V'^2, f_1^2: V'^2 \longrightarrow \mathbf{C}^3, \\ f_0^2: I^{2,3} \longrightarrow U'^3, \phi^3: U'^3 \xrightarrow{\cong} V'^3 \end{array} \right\} \quad (11.10) \quad \{\text{eq:morphcompN2}\}$$

$$u^2 \in U''^2 \quad (11.11) \quad \{\text{eq:xin2}\}$$

$$f_0[U''^2] \subseteq U'^3 \quad (11.12) \quad \{\text{eq:xin2}\}$$

$$\phi^1[U''^2] = V''^2 \quad (11.13)$$

$$f_1[V''^2] \subseteq \hat{V}'^3 \quad (11.14) \quad \{\text{eq:fvplin2}\}$$

$$\phi^3[U'^3] = V'^3 \quad (11.15) \quad \{\text{eq:fvplin2}\}$$

$$\hat{f} \circ \phi^3 \circ f_0^2 = f_1^2 \circ \phi'^2 \quad (11.16) \quad \{\text{eq:fphigeqgphi2}\}$$

Define $U''^1 \stackrel{\text{def}}{=} f_0^1[U''^2]$ and $V''^1 \stackrel{\text{def}}{=} f_1^1[V''^2]$. Diagram (11.17) below is com-

mutative, i.e., eqs. $\{eq:xin3\}$ (11.18) to (11.23) below hold, as shown in figs. 10 and 11. {fig:morphcompN1,fig:morphcompN2}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^1\} \rightarrow U'^1, \phi^1: U'^1 \rightarrow V'^1, f_1^2 \circ f_1^1: V'^1 \rightarrow V'^3, \\ f_0^2 \circ f_0^1: U'^1 \rightarrow U'^3 \phi^3: U'^3 \xrightarrow{\cong} V'^3 \end{array} \right\} \quad (11.17) \quad \{eq:morphcompN3\}$$

$$u^1 \in U'^1 \subseteq U'^1 \quad (11.18) \quad \{eq:xin3\}$$

$$f_0^2 \circ f_0^1[U'^1] \subseteq U'^3 \quad (11.19) \quad \{eq:xin3\}$$

$$\phi^1[U'^3] = V'^3 \quad (11.20)$$

$$f_1^2 \circ f_1^1[V'^1] \subseteq V'^3 \quad (11.21) \quad \{eq:fvp1in3\}$$

$$\phi^3[U'^3] = V'^3 \quad (11.22) \quad \{eq:fvp1in3\}$$

$$\begin{aligned} \hat{f} \circ \phi^3 \circ f_0^2 \circ f_0^1 &= f_1^2 \circ \phi'^2 \circ f_0^1 \\ &= f_1^2 \circ f_1^1 \circ \phi^1 \end{aligned} \quad (11.23) \quad \{eq:fphigeqgphi3\}$$




Figure 10: Partially completed composition of M-atlas near morphisms

{fig:morphcompN1}

□

{M-morphcomp:morphcompmorph}

2. If each f^i is an E^1 - E^{i+1} m-atlas morphism of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} then $f^2 \circ f^1$ is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .




Figure 11: Completed composition of M-atlas near morphisms

{fig:morphcompN2}

Proof. $f_0^2 \circ f_0^1$ and $f_1^2 \circ f_1^1$ are model functions by lemma 7.5 (Composition of model functions) on page 28. {lem:modcomp}

Let $(U^i, V^i, \phi^i) \in \mathbf{A}^i, i = 1, 3$, be a chart, $I \stackrel{\text{def}}{=} U^1 \cap (f_0^2 \circ f_0^1)^{-1}[U^3] \neq \emptyset$.

Let $u^1 \in I, u^2 \stackrel{\text{def}}{=} f_0^1(u^1), u^3 \stackrel{\text{def}}{=} f_0^2(u^2)$ and $(U^2, V^2, \phi^2) \in \mathbf{A}^2$ be a chart at u^2 . Since f^1 is a morphism by hypothesis, there exists a subchart

$(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , a model neighborhood $U'^2 \subseteq U^2$ of

u^2 and a chart $(U'^2, V'^2, \phi'^2: U'^2 \xrightarrow{\sim} V'^2) \in \mathbf{A}^2$ at u^2 such that $f_0^1[U'^1] \subseteq U'^2$ and diagram (11.1) in definition 11.1 (M-atlas morphisms for model spaces) on page 54 is commutative, as shown in fig. 4 (Completed m-atlas morphism) on page 54. {def:M-ATLmorph}
{fig:AtlM1}

Since f^2 is a morphism by hypothesis, there exists a subchart

$(U''^2, V''^2, \phi^2: U''^2 \xrightarrow{\sim} V''^2) \in \mathbf{A}^2$ at u^2 , a model neighborhood $U'^3 \subseteq U^3$

of u^3 and a chart $(U'^3, V'^3, \phi'^3: U'^3 \xrightarrow{\sim} V'^3) \in \mathbf{A}^3$ at u^2 such that $f_0^2[U'^2] \subseteq$




Figure 12: Uncompleted composition of m-atlas morphisms

{fig:morphcompM1}

U'^3 and diagram (11.24) below is commutative, as shown in figs. 12 and 13.

{fig:morphcompM1,fig:morphcompM2}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^2\} \hookrightarrow U''^2, \phi^2: U''^2 \xrightarrow{\cong} V''^2, f_1: V''^2 \longrightarrow \hat{V}^3, \\ f_0: U'^2 \longrightarrow U^3, \phi'^3: U^3 \xrightarrow{\cong} \hat{V}^3 \end{array} \right. \quad (11.24) \quad \{\text{eq:AtlM2}\}$$

Define $U'''^1 \stackrel{\text{def}}{=} f_0^{1-1}[U''^2]$, $V'''^1 \stackrel{\text{def}}{=} f_1^{1-1}[V''^2] = \phi^1[U'''^1]$, Figure 14 is commutative. $\square$

{M-morphcomp:nearcompnear1}

3. If A^2 is semi-maximal then $\mathbf{f}^2 \circ \mathbf{f}^1 \stackrel{\text{def}}{=} (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is an E^1 - E^3 m-atlas near morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof. Since A^2 is semi-maximal and $\mathbf{f}^1$ is a (near) morphism, $\mathbf{f}^1$ is a morphism either by hypothesis or by item 2 of lemma 11.12 (M-atlas (near) morphisms) on page 64.

{M-ATLmorph:semimaxnearism}




Figure 13: Partially completed composition of m-atlas morphisms

{fig:morphcompM2}

The result follows from item 2 above,

□ {M-morphcomp:morphcompmorph}
{M-morphcomp:nearcompnear2}

4. If A^2 and A^3 are semi-maximal then $f^2 \circ f^1 \stackrel{\text{def}}{=} (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof. Since A^2 is semi-maximal and f^1 is a (near) morphism, f^1 is a morphism either by hypothesis or by item 2 of lemma 11.12 (M-atlas (near) morphisms) on page 64. Similarly, f^2 is a morphism.

{M-ATLmorph:semimaxnearismorph}

The result follows from item 2 above,

□ {M-morphcomp:morphcompmorph}
{M-morphcomp:semicompsemi}

5. If each f^i , $i = 1, 2$, is a semi-strict $(\mathcal{C}^1$ - $\mathcal{C}^2$)- E^1 - E^2 -($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas (near) morphism then $f^2 \circ f^1$ is a semi-strict $(\mathcal{C}^1$ - $\mathcal{C}^2$)- E^1 - E^2 -($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas (near) morphism.

Proof. There are four overlapping cases.




Figure 14: Completed composition of M-atlas morphisms

{fig:morphcompM3}

E^i without category: If each $E^i \subseteq E^{i+1} \text{ mod}$ and each f_0^i is a local m-morphism of E^i to E^{i+1} , then by lemma 7.17 (Composition of m-morphisms) on page 39 $f_0^2 \circ f_0^1$ is a local m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

C^i without category: If each $C^i \subseteq C^{i+1} \text{ mod}$ and each f_1^i is a local m-morphism of C^i to C^{i+1} , then by lemma 7.17 $f_1^2 \circ f_1^1$ is a local morphism of C^1 to C^3 . Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 3$, with $I \stackrel{\text{def}}{=} U^1 \cap (f_0^2 \circ f_0^1)^{-1}[U^3] \neq \emptyset$. For every $u^1 \in I$ let $u^2 \stackrel{\text{def}}{=} f_0^1(u^1)$, $u^3 \stackrel{\text{def}}{=} f_0^2(u^2)$ and $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in A^2$ be a chart at u^2 , $I^1 \stackrel{\text{def}}{=} U^1 \cap f_0^{1-1}[U^2] \neq \emptyset$ and $I^2 \stackrel{\text{def}}{=} U^2 \cap f_0^{2-1}[U^3] \neq \emptyset$. If $\phi^2 \circ f_0^1 \circ \phi^{1-1}: \phi^1[I^1] \rightarrow V^2$ is a local m-morphism of $\text{Mod}(\phi^1[I^1], C^1)$ to $\text{Mod}(V^2, C^2)$ and $\phi^3 \circ f_0^2 \circ \phi^{2-1}: \phi^2[I^2] \rightarrow V^3$ is a local m-morphism of $\text{Mod}(\phi^1[I^2], C^1)$ to $\text{Mod}(V^2, C^2)$ then by definition 7.11 there exist model neighborhoods $V'^i \subseteq V^i, i \in [1, 3]$, such that $f^i[V'^i] \subseteq V'^{i+1}$ and {def:modlocalMmorph}

$\phi^{i+1} \circ f_0^i \circ \phi^{i-1}: V'^i \longrightarrow V'^{i+1}$ is an m-morphism of $\mathbf{C}^{i+1}$, $i = 1, 2$. Then

$$\begin{aligned} \phi^3 \circ f_0^2 \circ f_0^1 \circ \phi^{1-1}: V'^1 \longrightarrow V'^3 &= \phi^3 \circ f_0^2 \circ \phi^{2-1}: V'^2 \longrightarrow V'^3 \circ \\ &\phi^2 \circ f_0^1 \circ \phi^{1-1}: V'^1 \longrightarrow V'^2 \end{aligned}$$

is an m-morphism of $\mathbf{C}^3$. Since $u^1 \in I$ is arbitray, $\phi^3 \circ f_0^2 \circ f_0^1 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3$ is a local m-morphism of $\text{Mod}(\phi^1[I], \mathbf{C}^3)$ to $\text{Mod}(V^3, \mathbf{C}^3)$.

E^i with category: If each $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^{i+1}$ and each f_0^i is a local $\mathcal{E}^i$ - $\mathcal{E}^{i+1}$ m-morphism of E^i to E^{i+1} , then by lemma 7.17 (Composition of m-morphisms) on page 39 $f_0^2 \circ f_0^1$ is a local $\mathcal{E}^1$ - $\mathcal{E}^3$ m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

C^i with category: If each $\mathcal{C}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^{i+1}$ and each f_1^i is a local $\mathcal{E}^i$ - $\mathcal{E}^{i+1}$ m-morphism of C^i to C^{i+1} , then by lemma 7.17 (Composition of m-morphisms) on page 39 $f_1^2 \circ f_1^1$ is a local $\mathcal{E}^1$ - $\mathcal{E}^3$ m-morphism of C^1 to C^3 . {lem:modMmorphcomp}

□

{M-morphcomp: strictcompstr}

6. If each f^i , $i = 1, 2$, is strict then $f^2 \circ f^1$ is strict.

Proof. There are four overlapping cases.

E^i without category: If each $E^i \stackrel{\text{mod}}{\subseteq} E^{i+1}$ and each f_0^i is an m-morphism of E^i to E^{i+1} , then by lemma 7.17 (Composition of m-morphisms) on page 39 $f_0^2 \circ f_0^1$ is an m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

C^i without category: If each $C^i \stackrel{\text{mod}}{\subseteq} C^{i+1}$ and each f_1^i is an m-morphism of C^i to C^{i+1} , then by lemma 7.17 (Composition of m-morphisms) on page 39 $f_1^2 \circ f_1^1$ is an m-morphism of C^1 to C^3 . {lem:modMmorphcomp}

E^i with category: If each $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^{i+1}$ and each f_0^i is a morphism of $\mathcal{E}^{i+1}$, then $f_0^2 \circ f_0^1$ is a morphism of $\mathcal{E}^3$.

C^i with category: If each $\mathcal{C}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^{i+1}$ and each f_1^i is a morphism of $\mathcal{C}^i$, then $f_1^2 \circ f_1^1$ is a morphism of $\mathcal{C}^3$.

□

{M-morphcomp: constcompconst}

7. If each f^i is constrained then $f^2 \circ f^1$ is constrained.

Proof. This is a corollary of lemma 7.5 (Composition of model functions) on page 28. {lem:modcomp}

□

{M-morphcomp: compequivcomp}

8. $\mathbf{f}^2 \circ^{\text{Ob}} \mathbf{f}^1$ is equivalent to $\mathbf{g}^2 \circ^{\text{Ob}} \mathbf{g}^1$.

Proof. $f_0^1 = g_0^1$ and $f_0^2 = g_0^2$, hence $f_0^2 \circ f_0^1 = g_0^2 \circ g_0^1$. $\square$

{cor:M-ATLmorphcomp}

Corollary 11.17 (Composition of m-atlas (near) morphisms). Let $\mathcal{E}^1$ be a model category, $\mathbf{E}^1 \in \mathcal{E}^1$, $\mathcal{E}^i$, $i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{E}^i$, $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^1$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^1 \longrightarrow \mathbf{E}^1, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$, $i = 1, 2$, be a (semi-strict, strict) $(\mathcal{E}^1\text{-})\mathbf{E}^1(-\mathcal{E}^i\text{-}\mathcal{E}^{i+1})$ m-atlas (near) morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i$, $\mathbf{C}^{i+1}$. Then

1. If $\mathbf{A}^2$ is semi-maximal then $\mathbf{f}^2 \circ^{\text{Ob}} \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-})\mathbf{E}^1(-\mathcal{E}^1\text{-}\mathcal{E}^3)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1$, $\mathbf{C}^3$.

Proof. Since each $\mathbf{f}^i$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^1\text{-})\mathbf{E}^1\text{-}\mathbf{E}^1(-\mathcal{E}^i\text{-}\mathcal{E}^{i+1})$ m-atlas (near) morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i$, $\mathbf{C}^{i+1}$ by definition 11.11 (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61, and $\mathbf{A}^2$ is semi-maximal, $\mathbf{f}^2 \circ^{\text{Ob}} \mathbf{f}^1$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^1\text{-})\mathbf{E}^1\text{-}\mathbf{E}^1(-\mathcal{E}^1\text{-}\mathcal{E}^3)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1$, $\mathbf{C}^3$ by item 3 of lemma 11.16 (Composition of m-atlas (near) morphisms) on page 73. Apply definition 11.11. $\square$

{def:M-ATLmorphabbrev}

{M-morphcomp: nearcompnear1}

{def:M-ATLmorphabbrev}

2. If each $\mathbf{f}^i$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-})\mathbf{E}^1\text{-}\mathbf{E}^1(-\mathcal{E}^i\text{-}\mathcal{E}^{i+1})$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i$, $\mathbf{C}^{i+1}$ then $\mathbf{f}^2 \circ^{\text{Ob}} \mathbf{f}^1$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathbf{E}^1\text{-})\mathbf{E}^1\text{-}\mathbf{E}^1(-\mathcal{E}^1\text{-}\mathcal{E}^3)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1$, $\mathbf{C}^3$.

Proof. Apply definition 11.11 as above. $\square$

{def:M-ATLmorphabbrev}

Let $\mathcal{E}^i$ $i = 1, 2, 3$, be a model category, $\mathbf{E}^i \in \mathcal{E}^i$, $\mathcal{E}^1$ be a model category, $\mathbf{C}^1 \in \mathcal{E}^1$, $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^1$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}, f_1^i: \mathbf{C}^1 \longrightarrow \mathbf{C}^1)$, $i = 1, 2$, be a (semi-strict, strict) $(\mathcal{E}^i\text{-}\mathcal{E}^{i+1}\text{-})\mathbf{E}^i\text{-}\mathbf{E}^{i+1}(-\mathcal{E}^1)$ m-atlas (near) morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate space $\mathbf{C}^1$.

3. If $\mathbf{A}^2$ is semi-maximal then $\mathbf{f}^2 \circ^{\text{Ob}} \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^3\text{-})\mathbf{E}^1\text{-}\mathbf{E}^3(-\mathcal{E}^1)$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\mathbf{C}^1$.

Proof. Apply definition 11.11 as above. $\square$

{def:M-ATLmorphabbrev}

4. If each $\mathbf{f}^i$ is a (semi-strict, strict) $(\mathcal{E}^i - \mathcal{E}^{i+1})\mathbf{E}^i - \mathbf{E}^{i+1}(-\mathcal{C}^1)$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$ then $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1$ is a (semi-strict, strict) $(\mathcal{E}^1 - \mathcal{E}^3)\mathbf{E}^1 - \mathbf{E}^3(-\mathcal{C}^1 - \mathcal{C}^1)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

Proof. Apply definition 11.11 as above. □ `{def:M-ATLmorphabbrev}`

Let $\mathcal{E}^1, \mathcal{C}^1$ be model categories, $\mathbf{E}^1 \in \mathcal{E}^1, \mathbf{C}^1 \in \mathcal{C}^1, \mathbf{A}^i, i = 1, 2, 3$, be an m-atlas of $\mathbf{E}^1$ in the coordinate space $\mathbf{C}^1$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^1 \longrightarrow \mathbf{E}^1, f_1^i: \mathbf{C}^1 \longrightarrow \mathbf{C}^1), i = 1, 2$, be a (semi-strict, strict) $\mathbf{E}^1$ m-atlas (near) morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate space $\mathbf{C}^1$.

5. If $\mathbf{A}^2$ is semi-maximal then $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $\mathbf{E}^1$ m-atlas m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\mathbf{C}^1$.

Proof. Apply definition 11.11 as above. □ `{def:M-ATLmorphabbrev}`

6. If each $\mathbf{f}^i$ is a (semi-strict, strict) $\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate space $\mathbf{C}^1$ then $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1$ is a (semi-strict, strict) $\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\mathbf{C}^1$.

Proof. Apply definition 11.11 as above. □ `{def:M-ATLmorphabbrev}`

Let $\mathcal{E}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{E}^i, \mathbf{E}^i$ be a topological space, $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i, \mathbf{A}^2$ be semi-maximal and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), i = 1, 2$, be a (semi-strict, strict) $\mathbf{E}^i - \mathbf{E}^{i+1}$ m-atlas near morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

7. $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $\mathbf{E}^1 - \mathbf{E}^3$ m-atlas near morphisms of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 11.11 as above. □ `{def:M-ATLmorphabbrev}`

Let $\mathcal{E}^i, \mathcal{C}^i, i \in [1, 3]$, be model categories, $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i, \mathbf{A}^i$ an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i, \mathbf{A}^2$ maximal and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ a (semi-strict, strict) $\mathbf{E}^i - \mathbf{E}^{i+1}$ m-atlas near morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

8. $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $\mathbf{E}^1 - \mathbf{E}^3$ m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 11.11 as above. □ {def:M-ATLmorphabbrev}

Let $\mathcal{C}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$, $\mathbf{A}^2$ maximal and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ be a (semi-strict, strict) E^i - E^{i+1} m-atlas near morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

9. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 - E^3 m-atlas near morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 11.11 as above. □ {def:M-ATLmorphabbrev}

Let $\mathcal{C}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$, $i = 1, 2$, be a (semi-strict, strict) E^i - E^{i+1} m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

10. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 - E^3 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 11.11 as above. □ {def:M-ATLmorphabbrev}

{lem:M-ATLid}

Lemma 11.18 (M-atlas identity). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$
3. $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$
4. $\mathbf{E}^i \in \mathcal{C}^i$
5. $\mathbf{E}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{E}^2$
6. $\mathbf{C}^i \in \mathcal{C}^i$
7. $\mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$
8. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$
9. $\mathbf{A}^1 \subseteq \mathbf{A}^2$

The inclusion morphism $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a semi-strict $(\mathcal{E}^1, \mathcal{E}^2) \cdot E^1 \cdot E^2 \cdot (-\mathcal{E}^1, \mathcal{E}^2)$ m-atlas near morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is an m-atlas near morphism: Let $(U^i, V^i, \phi^i) \in A^i, i = 1, 2$,

be charts such that $I \stackrel{\text{def}}{=} U^1 \cap U^2 \neq \emptyset$. To show that diagram D below is M-locally nearly commutative in C^2 , let $u^1 \in I$. Since $E^1 \stackrel{\text{mod}}{\subseteq} E^2$ by hypothesis, U^1 is a model neighborhood of E^2 and thus I is a model neighborhood of E^2 and the identity map $\text{Id}_{U^2}^I$ is a morphism of E^2 . Then define $U'^1 \stackrel{\text{def}}{=} I, U'^2 \stackrel{\text{def}}{=} I$, $V'^1 \stackrel{\text{def}}{=} \phi^1[I], V'^2 \stackrel{\text{def}}{=} \phi^2[I], \hat{V}'^2 \stackrel{\text{def}}{=} V'^1$ and $\hat{f} \stackrel{\text{def}}{=} \phi^1 \circ \phi^{2-1}: V'^2 \xrightarrow{\sim} V'^1$. Equations (11.3) to (11.8) in definition 11.4 (M-atlas near morphisms for model spaces) hold. {def:M-ATLmorphnear}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} \phi^1: I \stackrel{\text{def}}{=} U^1 \cap U^2 \rightarrow V^1, \text{Id}_{C^2}^{V^1}: V^1 \rightarrow C^2, \\ \text{Id}_{U^2}^{U^1}: U^1 \rightarrow U^2, \phi^2: U^2 \xrightarrow{\sim} V^2 \end{array} \right\}$$

$\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is semi-strict: Let $u^1 \in E^1$. Since the model neighborhoods of E^1 form an open cover, there is a model neighborhood U of E^1 at u^1 . Id_U is a morphism of E^1 and of $\mathcal{E}^1$; since $E^1 \stackrel{\text{mod}}{\subseteq} E^2$ and $\mathcal{E}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^2$, Id_U is also a morphism of E^2 and of $\mathcal{E}^2$. Thus $\text{Id}_{E^2}^{E^1}$ is a local m-morphism of E^1 to E^2 and a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism E^1 to E^2 .

Let $v^1 \in C^1$. Since the model neighborhoods of C^1 form an open cover, there is a model neighborhood V of C^1 at v^1 . Id_V is a morphism of C^1 and of $\mathcal{C}^1$; since $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ and $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$, Id_V is also a morphism of C^2 and of $\mathcal{C}^2$. Thus $\text{Id}_{C^2}^{C^1}$ is a local m-morphism of C^1 to C^2 and a local $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism C^1 to C^2 . □

If A^1 and A^2 are semi-maximal then $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is an m-atlas morphism.

Proof. Since $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is an m-atlas near morphism by the above and the atlases are semi-maximal by hypothesis, it is an m-atlas morphism by lemma 11.12 (M-atlas (near) morphisms) on page 63. {lem:M-ATLmorph} □

Proof. If $E^1 \in \mathcal{E}^2$ and $C^1 \in \mathcal{C}^2$ then $\text{Id}_{(A^2, E^2, C^2)}^{(A^1, E^1, C^1)}$ is strict:

$$\text{Id}_{E^1} \in \mathcal{E}^1 \stackrel{\text{Ar}}{\subseteq} \mathcal{E}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^2 \text{ and } \text{Id}_{C^1} \in \mathcal{C}^1 \stackrel{\text{Ar}}{\subseteq} \mathcal{C}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2. \quad \square$$

If each $\text{Top}E^i \in E^i$ ($E^i \in \mathcal{E}^i$) and each $\text{Top}C^i \in C^i$ ($C^i \in \mathcal{C}^i$) then $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2\text{-}(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. There are four overlapping cases

Each $\text{Top}E^i \in E^i$: $\text{Id}_{E^2}^{E^1}$ is a morphism of E^2 by item 5 of [definition 2.1](#) (Model spaces) on page 18, hence an m-morphism of E^1 to E^2 . {mod:inclusion}

Each $E^i \in \mathcal{E}^2$: $\text{Id}_{E^2}^{E^1}$ is a morphism of $\mathcal{E}^2$ by item 6 of [definition 7.7](#) (Model categories) on page 29, hence an $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 . {modcat:morph}

Each $\text{Top}C^i \in C^i$: $\text{Id}_{C^2}^{C^1}$ is a morphism of C^2 by item 5 of [definition 2.1](#), hence an m-morphism of C^1 to C^2 . {mod:inclusion}

Each $C^i \in \mathcal{C}^2$: $\text{Id}_{C^2}^{C^1}$ is a morphism of $\mathcal{C}^2$ by item 6 of [definition 7.7](#), hence a $\mathcal{C}^1\text{-}\mathcal{C}^2$ m-morphism of C^1 to C^2 . {modcat:morph}

□

11.2 Categories of m-atlases and functors

This subsection defines categories of m-atlases and functors among them, and proves some basic results. {sub:M-cat}

11.2.1 M-atlas categories

In the following, ... will refer to one of

$$A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1$$

$$A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1$$

depending on context.

Definition 11.19 (Sets of m-atlas morphisms). Let E^i and C^i . $i = 1, 2$, be model spaces. Then the set of (constrained) (semi-strict, strict) (full, semi-maximal, maximal, full semi-maximal, full maximal) $E^1\text{-}E^2$ m-atlas (near) morphisms in the coordinate spaces C^1, C^2 is {def:M-ATLmorphsets}

$$\begin{aligned}
& \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{E}^1, \mathbf{C}^1, \mathbf{E}^2, \mathbf{C}^2) \stackrel{\text{def}}{=} \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right) \\
& \quad \left\{ ((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\
& \quad \left. \text{isAtl}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1) \right\} \quad (11.25) \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right)
\end{aligned}$$

Let E^i , $i = 1, 2$, be a topological space and $\mathbf{C}^1 = (C^i, \mathcal{C}^i)$, $i = 1, 2$, be a model space. Then the set of (constrained) (semi-strict, strict) (full, semi-maximal, maximal, full semi-maximal, full maximal) E^1 - E^2 m-atlas (near) morphisms in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is

$$\begin{aligned}
& \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{E}^1, \mathbf{C}^1, \mathbf{E}^2, \mathbf{C}^2) \stackrel{\text{def}}{=} \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right) \\
& \quad \left\{ ((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\
& \quad \left. \text{isAtl}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1) \right\} \quad (11.26) \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right)
\end{aligned}$$

Let

1. $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces
2. $\mathbf{P} \stackrel{\text{def}}{=} \mathbf{E} \times \mathbf{C}$

$$\begin{aligned}
& \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right) \\
& \quad \left\{ ((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\
& \quad \left. (\mathbf{E}^i, \mathbf{C}^i) \in \mathbf{P} \wedge \text{isAtl}^{(\text{const}, \text{const-near}, \text{near})}(\dots) \right\} \quad (11.27) \\
& \quad \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right)
\end{aligned}$$

Let

1. $\mathcal{C}$ be a model category
2. $\mathbf{C}$ be a set of model spaces
3. $\mathbf{P} \stackrel{\text{def}}{=} \text{Ob}(\mathcal{C}) \times \mathbf{C}$

$$\begin{aligned} \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathbf{E}, \mathcal{C}) &\stackrel{\text{def}}{=} \\ & \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right) \\ & \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ & \left. (E^i, C^i) \in \mathbf{P} \wedge \text{isAtl}^{(\text{const}, \text{const-near}, \text{near})}_{\left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right)}(..., \mathcal{C}, \mathcal{C}) \right\} \quad (11.28) \end{aligned}$$

Let

1. $\mathbf{E}$ be a set of model spaces
2. $\mathbf{C}$ be a model category
3. $\mathbf{P} \stackrel{\text{def}}{=} \mathbf{E} \times \text{Ob}(\mathcal{C})$

$$\begin{aligned} \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const-near}, \text{near})}(\mathcal{C}, \mathcal{C}) &\stackrel{\text{def}}{=} \\ & \left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right) \\ & \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ & \left. (E^i, C^i) \in \mathbf{P} \wedge \text{isAtl}^{(\text{const}, \text{const-near}, \text{near})}_{\left(\begin{array}{c} \text{full, S-max, max, S-max-full, max-full} \\ \text{(semi-strict, strict)} \end{array} \right)}(..., \mathcal{C}, \mathcal{C}) \right\} \quad (11.29) \end{aligned}$$

Let

1. $\mathcal{C}$ and $\mathcal{C}$ be model categories
2. $\mathbf{P} \stackrel{\text{def}}{=} \text{Ob}(\mathcal{C}) \times \text{Ob}(\mathcal{C})$

$$\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. (E^i, C^i) \in P \wedge \text{isAtl}^{(\text{const}, \text{const}-\text{near}, \text{near})}_{(\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full})}(\dots, \mathcal{E}, \mathcal{E}, \mathcal{C}, \mathcal{C}) \right\} \quad (11.30)$$

(semi-strict, strict)

Lemma 11.20 (Sets of m-atlas morphisms). Let E and C be sets of model spaces.

{lem:M-ATLmorphsets}

$$\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{near})}(E, C).$$

(full, S-max, max, S-max-full, max-full)
(semi-strict, strict)

$$\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \\ \text{max-full} \quad \text{S-max-full} \\ \text{(semi-strict, strict)} \quad \text{(semi-strict, strict)} \\ \subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \\ \text{full} \\ \text{(semi-strict, strict)}$$

Proof. The result follows from [definition 11.11](#) (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61 and definition 11.19 above. $\square$

{def:M-ATLmorphsets}

$$\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{near})}(E, C) \\ \text{(full, S-max, max, S-max-full, max-full)} \quad \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \quad \text{(semi-strict, strict)} \\ \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) = \mathcal{Atl}_{\text{Ar}}^{(\text{near})}(E, C) \\ \text{(S-max, max)} \quad \text{(S-max, max)} \\ \text{(semi-strict, strict)} \quad \text{(semi-strict, strict)}$$

Proof. The result follows from [item 1](#) of [lemma 11.12](#) (M-atlas (near) morphisms) on page 63, [definition 11.11](#) (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61 and definition 11.19 above. $\square$

{M-ATLmorph:morphisnear}

{def:M-ATLmorphsets}

Proof. The result follows from [lemma 11.12](#) (M-atlas (near) morphisms) on page 63, [definition 11.11](#) (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61 and definition 11.19 above. $\square$

{lem:M-ATLmorph}

{def:M-ATLmorphsets}
{cor:M-ATLmorphsets}

Corollary 11.21 (Sets of m-atlas morphisms). Let E and C be sets of topological spaces.

$$\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(E, C) \subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{near})}(E, C) \\ \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)}$$

$$\begin{aligned}
\mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathbf{E}, \mathbf{C}) &\subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathbf{E}, \mathbf{C}) \\
&\quad \begin{array}{c} \text{max-full} \\ (\text{semi-strict}, \text{strict}) \end{array} \quad \begin{array}{c} \text{S-max-full} \\ (\text{semi-strict}, \text{strict}) \end{array} \\
&\subseteq \mathcal{Atl}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathbf{E}, \mathbf{C}) \\
&\quad \begin{array}{c} \text{full} \\ (\text{semi-strict}, \text{strict}) \end{array} \\
\mathcal{Atl}_{\text{Ar}}(\mathbf{E}, \mathbf{C}) &\subseteq \mathcal{Atl}_{\text{Ar}}^{\text{near}}(\mathbf{E}, \mathbf{C}) \\
&\quad \begin{array}{c} (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array} \quad \begin{array}{c} (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array}
\end{aligned}$$

Proof. The result follows from ^{def:M-ATLmorphabbrev}definition 11.11 (Abbreviated nomenclature for m-atlas (near) morphisms) on page 61 and definition 11.19 above. $\square$

`{def:M-ATLmorphsets}`
`{def:M-cat}`

Definition 11.22 (Categories $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Let $P \stackrel{\text{def}}{=} \mathbf{E} \times \mathbf{C}$.

$$\begin{aligned}
\mathcal{Atl}^{(\text{const})}(\mathbf{E}, \mathbf{C}) &\stackrel{\text{def}}{=} \\
&\quad \begin{array}{c} (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array} \\
&\quad \left(\begin{array}{c} \mathcal{Atl}_{\text{Ob}}(\mathbf{E}, \mathbf{C}) \\ (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \end{array}, \begin{array}{c} \mathcal{Atl}_{\text{Ar}}^{(\text{const})}(\mathbf{E}, \mathbf{C}) \\ (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array}, \overset{A}{\circ} \right) \quad (11.31)
\end{aligned}$$

Let $\mathbf{E}^i$ and $\mathbf{C}^i$, $i = 1, 2$, be model spaces, $\mathbf{A}^i$ be an atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be a (strict, semi-strict) $\mathbf{E}^1$ - $\mathbf{E}^2$ M-atlas (near) morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The identity functor $\mathcal{F}_{\text{M}, \text{Id}}$ is

$$\mathcal{F}_{\text{M}, \text{Id}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \quad (11.32)$$

This nomenclature will be justified below.

$$\begin{aligned}
\mathcal{F}_{\text{M}, \text{Id}} \left((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right) &\stackrel{\text{def}}{=} \\
&\quad ((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \quad (11.33)
\end{aligned}$$

These are fine grained iff $\mathbf{C}$ is fine grained.

`{lem:M-ATLiscat}`
`{Mcat:near}`

Lemma 11.23 ($\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ is a category). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Then

1. $\mathcal{Atl}^{(\text{const}-\text{near}, \text{near})}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ of $\mathcal{Atl}_{\text{Ob}}^{(\text{const}-\text{near}, \text{near})}(\mathbf{E}, \mathbf{C})$ is $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)}$.

{Mcat:morph}

2. $\mathcal{Atl}_{(full,S-max,max,S-max-full,max-full)}^{(const)}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{Atl}_{(full,S-max,max,S-max-full,max-full)}^{(const)}(\mathbf{E}, \mathbf{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$ be a morphism of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$. Then

Composition:

$m^2 \circ m^1 = ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3))$ is a morphism of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ by item 2 of lemma 11.16 (Composition of m-atlas (near morphisms)) on page 75.

{M-morphcomp:morphcomp}

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

{lem:labcomp}

Identity:

$\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{cor:M-ATLiscat}

Corollary 11.24 (Subcategories of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Then $\mathcal{Atl}_{(full,S-max,max,S-max-full,max-full)}^{(const)}(\mathbf{E}, \mathbf{C})$ is a subcategory of $\mathcal{Atl}^{(const)}(\mathbf{E}, \mathbf{C})$.

Remark 11.25. They are not, in general, full subcategories.

The identity functor $\mathcal{F}_{M, \text{Id}}$ is a functor from each of the subcategories to each of the containing categories.

{def:M-cat-cat}

Definition 11.26 (Categories $\mathcal{Atl}(\mathcal{E}, \mathcal{C})$). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Let $P \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \text{Ob}(\mathcal{C})$. Then

$$\mathcal{Atl}_{(full,S-max,max,S-max-full,max-full)}^{(const)}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E \in \mathcal{E} \\ C \in \mathcal{C}}} \mathcal{Atl}_{(full,S-max,max,S-max-full,max-full)}^{(const)}(E, C) \quad (11.34)$$

$$\begin{aligned} \mathcal{Att}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C}) &\stackrel{\text{def}}{=} \\ & \left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \left\{ \begin{array}{c} ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \\ (E^i, C^i) \in P \wedge \text{isAtI}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}, \mathcal{C}) \end{array} \right\} \right) \end{aligned} \quad (11.35)$$

$$\begin{aligned} \mathcal{Att}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C}) &\stackrel{\text{def}}{=} \\ & \left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \left(\mathcal{Att}_{\text{Ob}}(\mathcal{E}, \mathcal{C}), \mathcal{Att}_{\text{Ar}}^{(\text{const}, \text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C}), \circ^A \right) \right) \end{aligned} \quad (11.36)$$

{lem:M-cat-ATLiscat}

Lemma 11.27 ($\mathcal{Att}(\mathcal{E}, \mathcal{C})$ is a category). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Then

1. $\mathcal{Att}^{(\text{const})}(\mathcal{E}, \mathcal{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{Att}^{(\text{const})}(\mathcal{E}, \mathcal{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{Att}(\mathcal{E}, \mathcal{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$ be a morphism of $\mathcal{Att}(\mathcal{E}, \mathcal{C})$. Then

Composition:

$f_0^2 \circ f_0^1$ is a morphism of $\mathcal{E}$, $f_1^2 \circ f_1^1$ is a morphism of $\mathcal{C}$ and $f^2 \circ f^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^2 by item 2 of lemma 11.16 (Composition of m-atlas (near morphisms) on page 75).

{M-morphcomp:morphcompmorph}

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

2. $\mathcal{Att}^{(\text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{Att}^{(\text{const}-\text{near}, \text{near})}(\mathcal{E}, \mathcal{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Definition 11.28 (Categories $\mathcal{Att}(\mathbf{E}, \mathbf{C})$ of topological spaces). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Let $\mathbf{P} \stackrel{\text{def}}{=} \mathbf{E} \times \mathbf{C}$. Then {def:T-cat}

$$\begin{aligned} & \mathcal{Att}^{(\text{const})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \\ & \left(\begin{array}{c} \mathcal{Att}_{\text{Ob}}(\mathbf{E}, \mathbf{C}) \\ (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array}, \begin{array}{c} \mathcal{Att}_{\text{Ar}}^{(\text{const})}(\mathbf{E}, \mathbf{C}) \\ (\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{array}, \overset{A}{\circ} \right) \quad (11.37) \end{aligned}$$

Let $(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1) \in \mathcal{Att}_{\text{Ob}}(\mathbf{E}, \mathbf{C})$.

$$\text{Id}_{(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} ((\text{Id}_{\mathbf{E}^1}, \text{Id}_{\mathbf{C}^1}), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)) \quad (11.38)$$

Lemma 11.29 ($\mathcal{Att}(\mathbf{E}, \mathbf{C})$ of topological spaces is a category). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then $\mathcal{Att}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object $(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)$ is $\text{Id}_{(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)}$. {lem:T-ATLiscat}

Proof. Let $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Att}(\mathbf{E}, \mathbf{C})$. and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^{i+1}, \mathbf{E}^{i+1}, \mathbf{C}^{i+1}))$ be a morphism of $\mathcal{Att}(\mathbf{E}, \mathbf{C})$. Then

Composition:

$((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^3, \mathbf{E}^3, \mathbf{C}^3))$ is a morphism of $\mathcal{Att}(\mathbf{E}, \mathbf{C})$ by corollary 11.17 (Composition of m-atlas (near) morphisms) on page 80. {cor:M-ATLmorphcomp}

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)}$ is an identity morphism by lemma 1.19. {lem:labcomp}

□

Similar results follow with restrictions on the admissible atlases, the admissible morphisms, or both:

1. $\mathcal{Att}_{\text{full}}(\mathcal{E}, \mathcal{C})$
2. $\mathcal{Att}_{\text{S-max}}(\mathcal{E}, \mathcal{C})$
3. $\mathcal{Att}_{\text{max}}(\mathcal{E}, \mathcal{C})$
4. $\mathcal{Att}_{\text{S-max-full}}(\mathcal{E}, \mathcal{C})$
5. $\mathcal{Att}_{\text{max-full}}(\mathcal{E}, \mathcal{C})$

6. $\mathcal{Att}_{\text{full semi-strict}}(\mathcal{E}, \mathcal{C})$
7. $\mathcal{Att}_{\text{S-max semi-strict}}(\mathcal{E}, \mathcal{C})$
8. $\mathcal{Att}_{\text{max semi-strict}}(\mathcal{E}, \mathcal{C})$
9. $\mathcal{Att}_{\text{S-max-full semi-strict}}(\mathcal{E}, \mathcal{C})$
10. $\mathcal{Att}_{\text{max-full semi-strict}}(\mathcal{E}, \mathcal{C})$
11. $\mathcal{Att}_{\text{full strict}}(\mathcal{E}, \mathcal{C})$
12. $\mathcal{Att}_{\text{S-max strict}}(\mathcal{E}, \mathcal{C})$
13. $\mathcal{Att}_{\text{max strict}}(\mathcal{E}, \mathcal{C})$
14. $\mathcal{Att}_{\text{S-max-full strict}}(\mathcal{E}, \mathcal{C})$
15. $\mathcal{Att}_{\text{max-full strict}}(\mathcal{E}, \mathcal{C})$

Definition 11.30 ($\mathbf{E}_{\text{triv}}$ and $\mathcal{Att}_{\text{triv}}(\mathbf{E}, \mathbf{C})$). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then {def:M-triv}

$$\mathbf{E}_{\text{triv}} \stackrel{\text{def}}{=} \left\{ E^\mu \mid E^\mu \in \mathbf{E} \right\} \quad (11.39)$$

$$\mathcal{Att}_{\text{triv}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \mathcal{Att}\left(\mathbf{E}_{\text{triv}}, \mathbf{C}\right) \quad (11.40)$$

11.2.2 M-atlas functors

Definition 11.31 ($\mathcal{F}_{\text{M,Top}}$). Let $\mathbf{E}^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i)$, $i = 1, 2$, and $\mathbf{C}^i \stackrel{\text{def}}{=} (C^i, \mathcal{C}^i)$ be model spaces and $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$. Then {def:Func-M-Top}

$$\mathcal{F}_{\text{M,Top}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, E^i, \mathbf{C}^i) \quad (11.41)$$

Let $f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2$ and $f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2$ be model functions. Then

$$\begin{aligned} \mathcal{F}_{\text{M,Top}}((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} \\ ((f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2), (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2)) \end{aligned} \quad (11.42)$$

Lemma 11.32 ($\mathcal{F}_{M, \text{Top}}$ maps M-atlases to M-atlases if the coordinate model space is fine grained.). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C}$ be model spaces, $\mathbf{C}$ be fine grained and $\mathbf{A}$ be an M-atlas of $\mathbf{E}$ in the model space $\mathbf{C}$. Then $\mathbf{A}$ is an M-atlas of E in the model space $\mathbf{C}$. {lem:Func-M-Top}

Proof. Let (U, V, ϕ) be an M-chart in $\mathbf{A}$. Every open subset of V is a model neighborhood of $\mathbf{C}$ since it is fine grained by hypothesis. Then (U, V, ϕ) is an M-chart of E in the model space $\mathbf{C}$ by lemma 9.3 (M-charts) on page 42. {lem:M-chart}

The definition of mutually compatible charts neighborhood of $\mathbf{C}$. (U, V, ϕ) is also an M-chart of E in the coordinate space $\mathbf{C}$:

Since U is a model neighborhood of $\mathbf{E}$, it is open and hence a model neighborhood of E . {lem:M-chart}

V is a model neighborhood of $\mathbf{C}$ by hypothesis.

Since U' is a model neighborhood of E and hence open, and ϕ is a homeomorphism, $\phi[U']$ is open. Since V is a model neighborhood of $\mathbf{C}$, $\mathbf{C}$ is fine grained and $\phi[U'] \subseteq V$ is open, $\phi[U']$ is a model neighborhood of $\mathbf{C}$.

Since V' is a model neighborhood of $\mathbf{C}$, it is open, $\phi^{-1}[V']$ is open and hence a model neighborhood of E . □

Theorem 11.33 ($\mathcal{F}_{M, \text{Top}}$ is a functor if $\mathbf{C}$ is fine grained.). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces and $\mathbf{C}$ be fine grained. Then $\mathcal{F}_{M, \text{Top}}$ is a functor from $\mathcal{A}tl(\mathbf{E}, \mathbf{C})$ to $\mathcal{A}tl(\text{Top}(\mathbf{E}), \mathbf{C})$. {the:Func-M-Top}

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i), \mathbf{C}^i), i \in [1, 3]$, be an object of $\mathcal{A}tl(\mathbf{E}, \mathbf{C})$, $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{M, \text{Top}} o^i = (\mathbf{A}^i, E^i, \mathbf{C}^i)$ the corresponding object of $\mathcal{A}tl(\text{Top}(\mathbf{E}), \mathbf{C})$, $m^i \stackrel{\text{def}}{=} ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o^i, o^{i+1}), i = 1, 2$, be a morphism of $\mathcal{A}tl(\mathbf{E}, \mathbf{C})$ and $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{M, \text{Top}} m^i = ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o'^i, o'^{i+1})$ the corresponding morphism of $\mathcal{A}tl(\text{Top}(\mathbf{E}), \mathbf{C})$.

$\mathcal{F}_{M, \text{Top}}$ satisfies these criteria:

Preservation of objects:

$\mathbf{A}^i$ is an M-atlas of E^i in the coordinate space $\mathbf{C}^i$ by lemma 11.32 ($\mathcal{F}_{M, \text{Top}}$ maps M-atlases to M-atlases if the coordinate model space is fine grained) above. {lem:Func-M-Top}

Preservation of morphisms:

$f_0^i: E^i \longrightarrow E^{i+1}$ is continuous and $f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}$ is a model function.

Preservation of endpoints:

$$\mathcal{F}_{M, \text{Top}} o^i = o'^i \in \mathcal{A}tl(\text{Top}(\mathbf{E}), \mathbf{C})$$

$$\mathcal{F}_{M, \text{Top}} m^i = ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o'^i, o'^{i+1}) \stackrel{\text{Ar}}{\in} \mathcal{A}tl(\text{Top}(\mathbf{E}), \mathbf{C})$$

is a morphism from o'^i to o'^{i+1} .

Identity:

Let $((\text{Id}_{E^i}, \text{Id}_{C^i}), o^i, o^i)$ be an identity morphism of o^i in $\mathcal{A}tl(E, C)$. Then $((\text{Id}_{E^i}: \text{Top}(E^i) \longrightarrow \text{Top}(E^i), \text{Id}_{C^i}), o'^i, o'^i)$ is an identity morphism of o'^i in $\mathcal{A}tl(\text{Top}(E), C)$.

Composition:

$$\begin{aligned}
\mathcal{F}_{M, \text{Top}}(m^2 \overset{A}{\circ} m^1) &= \mathcal{F}_{M, \text{Top}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\
&= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o'^1, o'^3) \\
&= ((f_0^2, f_1^2), o'^1, o'^2) \overset{A}{\circ} ((f_0^1, f_1^1), o'^2, o'^3) \\
&= \mathcal{F}_{M, \text{Top}}((f_0^2, f_1^2), o^2, o^3) \overset{A}{\circ} \mathcal{F}_{M, \text{Top}}((f_0^1, f_1^1), o^1, o^2) \\
&= \mathcal{F}_{M, \text{Top}} m^2 \overset{A}{\circ} \mathcal{F}_{M, \text{Top}} m^1
\end{aligned}$$

□

{def:Func-Top-M}

Definition 11.34 ($\mathcal{F}_{\text{Top}, M}$). Let $E^i, i = 1, 2$, be a topological space, C^i a model space and A^i an m-atlas of E^i and A an m-atlas of E in the coordinate space C^i . Then

$$\mathcal{F}_{\text{Top}, M}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A, E^i, C^i)_{\text{triv}} \quad (11.43)$$

Let $f_0: E^1 \longrightarrow E^2$ be continuous and $f_1: C^1 \longrightarrow C^2$ be a model function. Then

$$\begin{aligned}
\mathcal{F}_{\text{Top}, M}((f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2), (A^1, E^1, C^1), (A^2, E^2, C^2)) &\stackrel{\text{def}}{=} \\
\left((f_0: E^1_{\text{triv}} \longrightarrow E^2_{\text{triv}}, f_1: C^1 \longrightarrow C^2), (A^1, E^1, C^1), (A^2, E^2, C^2) \right) &\quad (11.44)
\end{aligned}$$

{lem:Func-Top-M}

Lemma 11.35 ($\mathcal{F}_{\text{Top}, M}$ maps M-atlases to M-atlases). Let E be a topological space, C be a model space and A be an M-atlas of E in the model space C . Then A is an M-atlas of E_{triv} in the model space C .

Proof. Let (U, V, ϕ) be a M-chart in A , $U' \subseteq U$ a model neighborhood of E_{triv} and $V' \subseteq V$ a model neighborhood of C . (U, V, ϕ) is also an M-chart of E in the coordinate space C : □

{the:Func-Top-M}

Theorem 11.36 ($\mathcal{F}_{\text{Top}, M}$ is a functor). Let E be a set of topological spaces and C a set of model spaces, Then $\mathcal{F}_{\text{Top}, M}$ is a functor from $\mathcal{A}tl(E, C)$ to $\mathcal{A}tl(E_{\text{triv}}, C)$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i), i \in [1, 3]$, be an object of $\mathcal{A}tl(E, C)$, $o'^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)_{\text{triv}}$

the corresponding object of $\mathcal{A}tl(E_{\text{triv}}, C)$, $m^i \stackrel{\text{def}}{=} ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: C^i \longrightarrow C^{i+1}), o^i, o^{i+1})$,

$i = 1, 2$, be a morphism of $\mathcal{A}tl(\mathbf{E}, \mathbf{C})$ and $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$ the corresponding morphism of $\mathcal{A}tl(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$.

$\mathcal{F}_{\text{Top}, \mathbf{M}}$ satisfies these criteria:

Preservation of endpoints: A^i is an $\mathbf{M}$ -atlas of $\underset{\text{triv}}{E^i}$ in the coordinate space $\mathbf{C}^i$ by

lemma 11.32 ($\mathcal{F}_{\mathbf{M}, \text{Top}}$ maps $\mathbf{M}$ -atlases to $\mathbf{M}$ -atlases if the coordinate model space is fine grained) above. $f_0^i: E^i \rightarrow E^{i+1}$ is continuous and $f_1^i: \mathbf{C}^i \rightarrow \mathbf{C}^{i+1}$ is a model function. {lem:Func-M-Top}

$$\mathcal{F}_{\text{Top}, \mathbf{M}} o^i = o^i \stackrel{\text{Ob}}{\in} \mathcal{A}tl(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$$

$$\mathcal{F}_{\text{Top}, \mathbf{M}} m^i = m^i \stackrel{\text{Ar}}{\in} \mathcal{A}tl(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$$

is a morphism from o^i to o^{i+1} .

Identity: $\text{Id}_{o^i} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{\mathbf{C}^i}), o^i, o^i)$. Then $\mathcal{F}_{\text{Top}, \mathbf{M}} \text{Id}_{o^i} = ((\text{Id}_{\underset{\text{triv}}{E^i}}, \text{Id}_{\mathbf{C}^i}), o^i, o^i)$ is an identity morphism of o^i .

Composition:

$$\begin{aligned} \mathcal{F}_{\text{Top}, \mathbf{M}}(m^2 \stackrel{A}{\circ} m^1) &= \mathcal{F}_{\text{Top}, \mathbf{M}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o'^1, o'^3) \\ &= ((f_0^2, f_1^2), o'^1, o'^2) \stackrel{A}{\circ} ((f_0^1, f_1^1), o'^2, o'^3) \\ &= \mathcal{F}_{\text{Top}, \mathbf{M}}((f_0^2, f_1^2), o^2, o^3) \stackrel{A}{\circ} \mathcal{F}_{\text{Top}, \mathbf{M}}((f_0^1, f_1^1), o^1, o^2) \\ &= \mathcal{F}_{\text{Top}, \mathbf{M}} m^2 \stackrel{A}{\circ} \mathcal{F}_{\text{Top}, \mathbf{M}} m^1 \end{aligned}$$

□

$\mathcal{F}_{\mathbf{M}, \text{Top}} \circ \mathcal{F}_{\text{Top}, \mathbf{M}}$ is the identity functor on $\mathcal{A}tl(\mathbf{E}, \mathbf{C})$.

Proof.

$$\begin{aligned} \mathcal{F}_{\mathbf{M}, \text{Top}} \circ \mathcal{F}_{\text{Top}, \mathbf{M}} o^i &= \mathcal{F}_{\mathbf{M}, \text{Top}} o'^i \\ &= o^i \end{aligned}$$

$$\begin{aligned} \mathcal{F}_{\mathbf{M}, \text{Top}} \circ \mathcal{F}_{\text{Top}, \mathbf{M}} m^i &= \mathcal{F}_{\mathbf{M}, \text{Top}} m'^i \\ &= m^i \end{aligned}$$

□

12 Associated model spaces and functors

Definition 12.1 (Coordinate model spaces associated with m-atlases). sec:assmod {def:M-modC} Let $\mathbf{A}^i$, $i =$

$1, 2$, be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$, $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ be an E^1 - E^2 m-atlas (near) morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1$, Then:

The minimal coordinate model space with neighborhoods in the m-atlas $\mathbf{A}^i$ of E^i in the coordinate space $\mathbf{C}^i$ is

$$\mathcal{F}_2^{\text{M-min}}(\mathbf{A}^i, E^i, \mathbf{C}^i) \stackrel{\text{def}}{=} \text{Mod}_{\min} \left(\mathbf{C}^i, \pi_2[\mathbf{A}^i], \left\{ \phi' \circ \phi^{-1} \mid \left(\exists_{\substack{(U,V,\phi) \in \mathbf{A}^i \\ (U',V',\phi') \in \mathbf{A}^i}} U \cap U' \neq \emptyset \right) \right\} \right) \quad (12.1)$$

The coordinate mapping associated with the E^1 - E^2 m-atlas (near) morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is

$$\mathcal{F}_2^{\text{M-min}}((f_0, f_1), (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} f_1: \mathcal{F}_2^{\text{M-min}}(\mathbf{A}^1, E^1, \mathbf{C}^1) \longrightarrow \mathcal{F}_2^{\text{M-min}}(\mathbf{A}^2, E^2, \mathbf{C}^2) \quad (12.2)$$

If it is a model function then it is also the coordinate m-morphism associated with the E^1 - E^2 m-atlas (near) morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Lemma 12.2 (Coordinate model spaces associated with m-atlases). {lem:M-ismodC} Let $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$, $i = 1, 2$.

$\mathcal{F}_2^{\text{M-min}}(\mathbf{A}^i, E^i, \mathbf{C}^i)$ is a model space.

Proof. $\mathcal{F}_2^{\text{M-min}}(\mathbf{A}^i, E^i, \mathbf{C}^i)$ satisfies the conditions for a model space.

1. Since $\pi_2[\mathbf{A}^i]$ is an open cover of $\bigcup \pi_2[\mathbf{A}^i]$, the set of finite intersections is also an open cover.
2. Finite intersections of finite intersections are finite intersections
3. Restrictions of continuous functions are continuous
4. If $f: A \longrightarrow B$ is a morphism of $\mathcal{F}_2^{\text{M-min}}(\mathbf{A}^i, E^i, \mathbf{C}^i)$, A, A', B, B' model neighborhoods of $\mathcal{F}_2^{\text{M-min}}(\mathbf{A}^i, E^i, \mathbf{C}^i)$, $A' \subseteq A$, $B' \subseteq B$ and $f[A'] \subseteq B'$ then since $f: A \longrightarrow B$ is a morphism it is a restriction of a transition function between sets in $\pi_2[\mathbf{A}^i]$ and its restrictions are also, hence morphisms, and thus $f|_{A'}: A' \longrightarrow B'$ is a morphism.

5. If $(U, V, \phi) \in \mathbf{A}^i$ then $\text{Id}_V = \phi \circ \phi^{-1}$ is a transition function and hence a morphism of $\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$. If A, A' are objects of $\pi_2(\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i))$ and $A' \subseteq A$ then the inclusion map $i: A' \hookrightarrow A$ is a restriction of an identity morphism of $\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ and hence a morphism.

6. Restricted sheaf condition: let

(a) $U_\alpha, V_\alpha, \alpha < A$, be model neighborhoods of $\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$

(b) $f_\alpha: U_\alpha \rightarrow V_\alpha$ be a morphism of $\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$

(c) $U \stackrel{\text{def}}{=} \bigcup_{\alpha < A} U_\alpha$

(d) $V \stackrel{\text{def}}{=} \bigcup_{\alpha < A} V_\alpha$

(e) $f: U \rightarrow V$ be continuous and $\left(\bigvee_{\substack{\alpha < A \\ x \in U_\alpha}} \right) f(x) = f_\alpha(x)$

Then f is a morphism of $\mathbf{C}^i$ and hence a morphism of $\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$.

□

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ be an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$. If $\mathbf{f}$ is a morphism or each $\mathbf{A}^i$ is semi-maximal then

$f_1: \mathcal{F}_2^{\text{M}\min}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1) \rightarrow \mathcal{F}_2^{\text{M}\min}(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$ is well defined, i.e., $f_1 \left[\mathcal{F}_2^{\text{M}\min}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1) \right] \subseteq \mathcal{F}_2^{\text{M}\min}(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$.

Proof. $\mathbf{f}$ is a morphism either by hypothesis or by lemma 11.12 (M-atlas (near) morphisms) on page 63. Let $v^1 \in \mathcal{F}_2^{\text{M}\min}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)$, $(U^1, V^1, \phi^1) \in \mathbf{A}^1$ be a chart

with $v^1 \in V^1$, $u^1 \stackrel{\text{def}}{=} \phi^{1-1}(v^1)$, $u^2 \stackrel{\text{def}}{=} f_0(u^1)$ and $(U^2, V^2, \phi^2) \in \mathbf{A}^2$ be a chart at

u^2 . Let $(U'^1, V'^1, \phi'^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$, $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in \mathbf{A}^2$ be as in definition 11.1 (M-atlas morphisms for model spaces) on page 53. Then

$f_1(v^1) \in \hat{V}'^2 \subseteq \mathcal{F}_2^{\text{M}\min}(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$. □

{def:M-ATLmorph}

{def:M-modE}

Definition 12.3 (Model spaces associated with m-atlases). Let $\mathbf{A}^i, i = 1, 2$, be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ be an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas (near) morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1$, Then:

The minimal model space with neighborhoods in the m-atlas $\mathbf{A}^i$ of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ is

$$\mathcal{F}_1^{\text{M-min}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} \text{Mod}_{\min} \left(\mathbf{E}^i, \pi_1[\mathbf{A}^i], \left\{ \phi'^{-1} \circ \phi \mid \left(\exists_{\substack{(U,V,\phi) \in \mathbf{A}^i \\ (U',V',\phi') \in \mathbf{A}^i}} \right) U \cap U' \neq \emptyset \right\} \right) \quad (12.3)$$

The mapping associated with the $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is

$$\mathcal{F}_1^{\text{M-min}}((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} f_0: \mathcal{F}_1^{\text{M-min}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1) \longrightarrow \mathcal{F}_1^{\text{M-min}}(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \quad (12.4)$$

If it is a model function then the it is also the m-atlas morphism associated with the $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$. {lem:M-ismodE}

Lemma 12.4 (Model spaces associated with m-atlases). Let $\mathbf{A}$ be an m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$. Then $\mathcal{F}_1^{\text{M-min}}(\mathbf{A}, \mathbf{E}, \mathbf{C})$ is a model space.

Proof. The result follows from Lemma 5.4 (Minimal model spaces are model spaces) on page 26. □

Theorem 12.5 (Functors from m-atlases to model spaces). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. {the:M-isfunc}

$$\begin{aligned} \mathcal{F}_1^{\text{M-min}} & \text{ is a functor from } \mathcal{At}(\mathbf{E}, \mathbf{C}) \text{ to } \mathbf{E} \text{ full-submod-}\mathcal{T}riv \stackrel{\text{full-cat}}{\subseteq} \mathbf{E} \text{ submod-}\mathcal{T}riv \\ \mathcal{F}_1^{\text{M-min}} & \text{ is a functor from } \mathcal{At}_{\text{full}}(\mathbf{E}, \mathbf{C}) \text{ to } \mathbf{E} \text{ full-submod-}\mathcal{T}riv \stackrel{\text{full-cat}}{\subseteq} \mathbf{E} \text{ submod-}\mathcal{T}riv \\ \mathcal{F}_2^{\text{M-min}} & \text{ is a functor from } \mathcal{At}_{\text{S-max}}(\mathbf{E}, \mathbf{C}) \text{ to } \mathbf{C} \text{ submod-}\mathcal{T}riv \\ \mathcal{F}_2^{\text{M-min}} & \text{ is a functor from } \mathcal{At}_{\text{full}}(\mathbf{E}, \mathbf{C}) \text{ to } \mathbf{C} \text{ full-submod-}\mathcal{T}riv \stackrel{\text{full-cat}}{\subseteq} \mathbf{C} \text{ submod-}\mathcal{T}riv \end{aligned}$$

Proof. $\mathbf{E} \text{ full-submod-}\mathcal{T}riv \stackrel{\text{full-cat}}{\subseteq} \mathbf{E} \text{ submod-}\mathcal{T}riv$ and $\mathbf{C} \text{ full-submod-}\mathcal{T}riv \stackrel{\text{full-cat}}{\subseteq} \mathbf{C} \text{ submod-}\mathcal{T}riv$ by lemma 4.2 (Trivial model spaces) on page 23. {lem:trivmod}

Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, $i \in [1, 3]$, be objects in $\mathcal{At}(\mathbf{E}, \mathbf{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$ be a morphism in $\mathcal{At}(\mathbf{E}, \mathbf{C})$.

$$\mathcal{F}_1^{\text{M-min}}: \mathcal{At}(\mathbf{E}, \mathbf{C}) \longrightarrow \mathbf{E} \text{ submod-}\mathcal{T}riv :$$

Preservation of endpoints: $\mathcal{F}_1^{\text{M-min}}(m^i) = f_0^i: \mathcal{F}_1^{\text{M-min}} o^i \longrightarrow \mathcal{F}_1^{\text{M-min}} o^{i+1}$

Composition:

$$\begin{aligned}
\mathcal{F}_1^{\text{M}}(m^2 \circ^A m^1) &= \mathcal{F}_1^{\text{M}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)(\mathbf{A}^1, E^1, C^1), (\mathbf{A}^3, E^3, C^3)) \\
&= f_0^2 \circ f_0^1: \mathcal{F}_1^{\text{M}} o^1 \longrightarrow \mathcal{F}_1^{\text{M}} o^3 \\
&= (f_0^2: \mathcal{F}_1^{\text{M}} o^2 \longrightarrow \mathcal{F}_1^{\text{M}} o^3) \circ (f_0^1: \mathcal{F}_1^{\text{M}} o^1 \longrightarrow \mathcal{F}_1^{\text{M}} o^2) \\
&= \mathcal{F}_1^{\text{M}}((f_0^2, f_1^2)(\mathbf{A}^2, E^2, C^2), (\mathbf{A}^3, E^3, C^3)) \circ \\
&\quad \mathcal{F}_1^{\text{M}}((f_0^1, f_1^1)(\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \\
&= \mathcal{F}_1^{\text{M}}(m^2) \circ \mathcal{F}_1^{\text{M}}(m^1)
\end{aligned}$$

Identity:

$$\begin{aligned}
1. \mathcal{F}_1^{\text{M}}(\text{Id}_{oi}) &= \mathcal{F}_1^{\text{M}}((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) = \\
&\quad \text{Id}_{E^i}: \mathcal{F}_1^{\text{M}} o^i \longrightarrow \mathcal{F}_1^{\text{M}} o^i \\
2. \text{Id}_{\mathcal{F}_1^{\text{M}}} o^i &= \text{Id}_{E^i}: \mathcal{F}_1^{\text{M}} o^i \longrightarrow \mathcal{F}_1^{\text{M}} o^i
\end{aligned}$$

The proof for $\mathcal{F}_1^{\text{M}}: \mathcal{Att}(\mathbf{E}, \mathbf{C}) \xrightarrow{\text{full}} \mathbf{E} \xrightarrow{\text{full-submod-}\mathcal{F}riv} \mathbf{C} \xrightarrow{\text{full-cat}} \mathbf{C}$ is identical.

$$\mathcal{F}_2^{\text{M}}: \mathcal{Att}(\mathbf{E}, \mathbf{C}) \xrightarrow{\text{op-}\mathcal{F}riv} \mathbf{C}$$

Preservation of endpoints:

$$\mathcal{F}_2^{\text{M}}(m^i) = f_1^i: \mathcal{F}_2^{\text{M}} o^i \longrightarrow \mathcal{F}_2^{\text{M}} o^{i+1}$$

$$\begin{aligned}
\mathcal{F}(g \circ f) &= \mathcal{F}(g) \circ \mathcal{F}(f): \mathcal{F}_2^{\text{M}}(m^2 \circ^A m^1) = \\
&\quad \mathcal{F}_2^{\text{M}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)(\mathbf{A}^1, E^1, C^1), (\mathbf{A}^3, E^3, C^3)) = \\
&\quad f_0^2 \circ f_0^1: \mathcal{F}_2^{\text{M}} o^1 \longrightarrow \mathcal{F}_2^{\text{M}} o^3 = \\
&\quad (f_1^2: \mathcal{F}_2^{\text{M}} o^2 \longrightarrow \mathcal{F}_2^{\text{M}} o^3) \circ (f_1^1: \mathcal{F}_2^{\text{M}} o^1 \longrightarrow \mathcal{F}_2^{\text{M}} o^2) = \\
&\quad \mathcal{F}_2^{\text{M}}((f_0^2, f_1^2)(\mathbf{A}^2, E^2, C^2), (\mathbf{A}^3, E^3, C^3)) \circ \\
&\quad \mathcal{F}_2^{\text{M}}((f_0^1, f_1^1)(\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) = \\
&\quad \mathcal{F}_2^{\text{M}}(m^2) \circ \mathcal{F}_2^{\text{M}}(m^1)
\end{aligned}$$

$$\mathcal{F}(\text{Id}_A) = \text{Id}_{\mathcal{F}(A)}:$$

$$\begin{aligned}
1. \mathcal{F}_2^{\text{M}}(\text{Id}_{oi}) &= \mathcal{F}_2^{\text{M}}((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) = \\
&\quad \text{Id}_{C^i}: \mathcal{F}_2^{\text{M}} o^i \longrightarrow \mathcal{F}_2^{\text{M}} o^i
\end{aligned}$$

$$2. \text{Id}_{\mathcal{F}_2^{\text{M}}} o^i = \text{Id}_{C^i}: \mathcal{F}_2^{\text{M}} o^i \longrightarrow \mathcal{F}_2^{\text{M}} o^i$$

The proof for $\mathcal{F}_2^{\text{M}}: \mathcal{A}tl_{\text{full}}^{\text{M}}(E, C) \longrightarrow \mathbf{C}_{\text{full-submod-}\mathcal{F}riv} \stackrel{\text{full-cat}}{\subseteq}$ is identical. $\square$

13 Classic m-atlas morphisms and functors

{sec:M-ATLcmorph}

This subsection introduces an alternate definition of and taxonomy for morphisms between m-atlases, defines categories of m-atlases, defines functors among them and proves some basic results. It introduces the notion of classic m-atlas morphisms, which is very similar to the conventional definitions for a manifold, although it does not require the space to be locally Euclidean. The reader can safely skip it, but it does help to establish a context for other notions that are used in the exposition of local coordinate spaces.

While it is easy to define satisfactory notions of classic m-morphisms, there are technical difficulties in defining a satisfactory notion of classic near m-morphisms.

13.1 Classic m-atlas morphisms

{sub:M-ATLcmorph}

This subsection introduces the notion of classic m-atlas morphisms, and proves some basic results.

13.1.1 Definitions of classic m-atlas morphisms

{def:M-ATLcmorph}

Definition 13.1 (Classic m-atlas morphisms for model spaces). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

$f: E^1 \longrightarrow E^2$ is a classic E^1 - E^2 ($\mathcal{E}^1$ - $\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$. and a classic $\mathcal{E}^1$ - $\mathcal{E}^2$ ($\mathcal{E}^1$ - $\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{E}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, \mathcal{E}^1, C^1, A^2, \mathcal{E}^2, C^2, f)$, iff $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ ($\mathcal{E}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{E}^2$) and for any $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset, \phi^2 \circ f \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of C^2 ($\mathcal{E}^2$).

f is also a constrained classic E^1 - E^2 ($\mathcal{E}^1$ - $\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{constrained,classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$, and a constrained classic $\mathcal{E}^1$ - $\mathcal{E}^2$ ($\mathcal{E}^1$ - $\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{E}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{constrained,classic}}(A^1, \mathcal{E}^1, C^1, A^2, \mathcal{E}^2, C^2, f_0, f_1)$, iff f is constrained, i.e., $f[E]$ is contained in a model neighborhood of E^2 .

Let f be a classic E^1 - E^2 ($\mathcal{E}^1$ - $\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The triple $((f), (A^1, E^1, C^1), (A^2, E^2, C^2))$ will refer to f considered as a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ and $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{def:M-ATLcmorph^{top}}

Definition 13.2 (Classic m-atlas morphisms for topological spaces). Let $\mathcal{C}^i, i = 1, 2$, be a model category, $\mathcal{E}^i$ be a topological category, $E^i \in \mathcal{E}^i$, $C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

$f: E^1 \longrightarrow E^2$ is a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$. and a classic $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f)$, iff f is a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The triple $((f), (A^1, E^1, C^1), (A^2, E^2, C^2))$ will refer to f considered as a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ and $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{def:M-ATLcid}

Definition 13.3 (Classic m-atlas inclusion and identity morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{E}^i$
3. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
4. $C^i \in \mathcal{C}^i$
5. $C^1 \stackrel{\text{mod}}{\subseteq} C^2 (\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2)$
6. A^i be an m-atlas of E^i in the coordinate space C^i
7. $A^1 \subseteq A^2$

The classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas inclusion morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 is $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)} \stackrel{\text{def}}{=} \text{Id}_{E^2}^{E^1}$.

The classic m-atlas inclusion morphism of (A^1, E^1, C^1) to (A^2, E^2, C^2) is $\text{Id}_{(A^2, E^2, C^2)}^{(A^1, E^1, C^1)} \stackrel{\text{def}}{=} \left(\text{Id}_{(E^2)}^{(E^1)}, (A^1, E^1, C^1), (A^2, E^2, C^2) \right), \text{Id}_{E^2}^{E^1}$ considered as a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The classic $E^i(-\mathcal{C}^i)$ m-atlas identity morphism of A^i in the coordinate space C^i is $\text{Id}_{(E^i, C^i)} \stackrel{\text{def}}{=} \text{Id}_{E^i}$.

The classic m-atlas identity morphism of (A^i, E^i, C^i) is $\text{Id}_{(A^i, E^i, C^i)} \stackrel{\text{def}}{=} (\text{Id}_{(E^i)}, (A^i, E^i, C^i), (A^i, E^i, C^i)), \text{Id}_{E^i}$ considered as a classic $E^i(-\mathcal{C}^i)$ m-atlas morphism of A^i in the coordinate space C^i .

Let

1. E^i be a topological space, $i = 1, 2$,
2. $E^1 \stackrel{\text{subsp}}{\subseteq} E^2$
3. $\mathcal{C}^i$ be a model category
4. $\mathbf{C}^i \in \mathcal{C}^i$
5. $\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2$
6. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$
7. $\mathbf{A}^1 \subseteq \mathbf{A}^2$

The classic E^1 - E^2 ($-\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\text{Id}_{E^2}^{E^1}$.

The classic m-atlas inclusion morphism of $(\mathbf{A}^1, E^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, E^2, \mathbf{C}^2)$ is $\text{Id}_{(\mathbf{A}^2, E^2, \mathbf{C}^2)}^{(\mathbf{A}^1, E^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} (\text{Id}_{(E^2)}^{(E^1)}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2))$, $\text{Id}_{E^2}^{E^1}$ considered as a classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The classic E^i m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$ is $\text{Id}_{(\mathbf{A}^i, \mathbf{C}^i)}^{E^i}$.

The classic m-atlas identity morphism of $(\mathbf{A}^i, E^i, \mathbf{C}^i)$ is

$\text{Id}_{(\mathbf{A}^i, E^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} (\text{Id}_{(E^i)}^{E^i}, (\mathbf{A}^i, E^i, \mathbf{C}^i), (\mathbf{A}^i, E^i, \mathbf{C}^i))$, $\text{Id}_{E^i}^{E^i}$ considered as an E^i m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

13.1.2 Definitions of semi-strict and strict

{def:M-ATLcmorphsemi}

Definition 13.4 (Semi-strict classic m-atlas morphisms). Let $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{C}^i, \mathbf{C}^i \in \mathcal{C}^i$ and $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$.

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

f is a semi-strict classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtI}_{\text{Ar}}^{\text{classic}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f)$, iff f is a local m-morphism of E^1 to E^2 .

Remark 13.5. Definitions of semi-strict classic m-atlas morphisms for topological spaces would be pointless, as any semi-strict classic m-atlas morphism would be strict.

It is a semi-strict classic $\mathcal{C}^1$ - $\mathcal{C}^2$ - E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtI}_{\text{Ar}}^{\text{classic}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f, \mathcal{C}^1, \mathcal{C}^2)$, iff f is a local $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism of E^1 to E^2 .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a semi-strict classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{C}^1, \mathcal{C}^2)$, iff f is a local m-morphism of E^1 to E^2 .

It is a semi-strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

{def:M-ATLcmorphstrict}

Definition 13.6 (Strict classic m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a strict classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff f is an m-morphism of E^1 to E^2 .

It is a strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2)$, iff f is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a strict classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{C}^1, \mathcal{C}^2)$, iff f is an m-morphism of E^1 to E^2 .

It is a strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAt}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

13.1.3 Abbreviated nomenclature for classic m-atlas morphisms

{def:M-ATLcmorphabbrev}

Definition 13.7 (Abbreviated nomenclature for classic m-atlas morphisms). Let $\mathcal{E}^i,$

$\mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i and $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas

morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{constrained,})\text{classic}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f)$, iff it is a (full) (maximal) (constrained) (semi-strict, strict) classic $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{constrained,})\text{classic}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f, \mathcal{E}^1, \mathcal{E}^2)$, iff it is a (full) (maximal) (constrained) (semi-strict, strict) classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

Let $\mathcal{E}^i, i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{E}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $f: E^1 \rightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{constrained,})\text{classic}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f)$, iff it is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{constrained,})\text{classic}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, f)$, iff it is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

13.1.4 Proclamations on classic m-atlas morphisms

{lem:M-ATLcmorph}

Lemma 13.8 (Classic m-atlas morphisms). Let E be a set of topological spaces, $E^i \in E, i = 1, 2$, $\mathbf{C}^i$ be a model space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$.

1. Let $f: E^1 \rightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ and $\mathbf{C}^1 \subseteq^{\text{mod}} \mathbf{C}^2$. Then f is a strict constrained classic E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof.

f is **constrained**: $f[E^1] \subseteq E^2$ by hypothesis. E^2 is a model neighborhood of E^2 by definition 4.1 (Trivial model spaces) on page 23.

{def:trivmod}

f is **strict**: $E_{\mathcal{T}riv}^{full-cat} \subseteq E_{\mathcal{T}riv}$. f is continuous by definition 7.1 (Model functions) on page 27. $E^i \in E$ by hypothesis, $E^i \in E_{\mathcal{T}riv}^{Ob}$ and $f \in E_{\mathcal{T}riv}^{Ar}$ by

{def:modfunc}

{def:trivmod}

definition 4.1. $C^1 \subseteq^{mod} C^2$ by definition 13.1 (Classic m-atlas morphisms for model spaces) on page 101.

{def:M-ATLcmorph}

□

2. If E^1 is an open subspace of E^2 then $f: E_{\mathcal{T}riv}^1 \longrightarrow E_{\mathcal{T}riv}^2$ is also a strict constrained classic $E^1 - E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. Each $E^i \in E$ by hypothesis and each $E_{\mathcal{T}riv}^i \in E_{\mathcal{T}riv}$ by definition 4.1.

{def:trivmod}

$E_{\mathcal{T}riv}^1 \subseteq^{open-subsp} E^2$ by hypothesis.

$E_{\mathcal{T}riv}^1 \subseteq^{mod} E_{\mathcal{T}riv}^2$ by lemma 4.2 (Trivial model spaces) on page 23.

{lem:trivmod}

$f \in E_{\mathcal{T}riv}^{Ar}$ by definition 4.1.

□

{def:trivmod}

Let

1. $\mathcal{C}^i, \mathcal{C}'^i, \mathcal{C}''^i, \mathcal{C}'^i, i = 1, 2$, be model categories
2. $\mathcal{C}^i \subseteq^{full-cat} \mathcal{C}'^i$
3. $\mathcal{C}^i \subseteq^{full-cat} \mathcal{C}''^i$
4. $\mathcal{C}'^1 \subseteq^{full-cat} \mathcal{C}'^2$
5. $E^i \in \mathcal{C}^i$
6. $C^i \in \mathcal{C}^i$
7. $C''^i \in \mathcal{C}''^i$
8. $C^i \subseteq^{mod} C''^i$
9. A^i be an m-atlas of E^i in the coordinate space C^i

10. $f: E^1 \longrightarrow E^2$ be a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2)\text{-}E^1\text{-}E^2$ ($\mathcal{E}^1\text{-}\mathcal{E}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 (model function?)

If f is a (semi-strict, strict) $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}E^1\text{-}E^2$ ($E^1\text{-}E^2$) classic m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , then f is a (semi-strict, strict) $\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-}E^1\text{-}E^2$ ($E^1\text{-}E^2$) classic m-atlas morphism of A^1 to A^2 in the coordinate spaces C'^1, C'^2 .

Proof.

f remains a classic m-atlas morphism with the expanded categories. $\mathcal{E}'^1 \overset{\text{full-cat}}{\subseteq} \mathcal{E}'^2$ by hypothesis.

Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \overset{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset$.

$\phi^2 \circ f \circ \phi^{-1}$ is a morphism of $\mathcal{E}^2 \overset{\text{full-cat}}{\subseteq} \mathcal{E}'^2$ by definition 13.1.

{def:M-ATLcmorph}

f remains semi-strict (strict). If f is a local morphism of $\mathcal{E}^2$ then f is a local morphism of $\mathcal{E}'^2$. If f is a local morphism of E^2 then f is a local morphism of E'^2 . If $f \in \mathcal{E}^2$ then $f \in \mathcal{E}'^2$. If f is a morphism of E^2 then f is a morphism of E'^2 .

□

{lem:M-ATLcmorphcomp}

Lemma 13.9 (Composition of classic m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i \in [1, 3]$, be a model category, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i and $f^i: E^i \longrightarrow E^{i+1}, i = 1, 2$, be a (semi-strict, strict) classic $E^i\text{-}E^{i+1}$ m-atlas morphism of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} .

$f^2 \circ f^1: E^1 \longrightarrow E^3$ is a (semi-strict, strict) classic $E^1\text{-}E^3$ m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof.

$f^2 \circ f^1$ is a classic $E^1\text{-}E^3$ m-atlas morphism. If $\mathcal{E}^1 \overset{\text{cat}}{\subseteq} \mathcal{E}^2$ and $\mathcal{E}^2 \overset{\text{cat}}{\subseteq} \mathcal{E}^3$ then

$\mathcal{E}^1 \overset{\text{cat}}{\subseteq} \mathcal{E}^3$. Let $(U^1, V^1, \phi^1: U^1 \xrightarrow{\sim} V^1) \in A^1, (U^3, V^3, \phi^3: U^3 \xrightarrow{\sim} V^3) \in A^3$

with $I \overset{\text{def}}{=} U^1 \cap (f^2 \circ f^1)^{-1}[U^3] \neq \emptyset, (f^2 \circ f^1)^{-1}[U^3]$ is a model neighborhood by definition 7.1 (Model functions) on page 27 and I is a model neighborhood by item 2 of definition 2.1 (Model spaces) on page 18.

{def:modfunc}

{mod:closed}

For any $u^1 \in I$ and any chart $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in A^2$ at $f^1(u^1)$, define

$I^{1,2} \overset{\text{def}}{=} U^1 \cap f^{1-1}[U^2], I^{2,3} \overset{\text{def}}{=} U^2 \cap f^{2-1}[U^3]$ and $I_{u^1}^{1,3} \overset{\text{def}}{=} U^1 \cap f^{1-1}[I^{2,3}] \subseteq I$.

$I_{u^1}^{1,3}$ is a model neighborhood by definition 7.1 and item 2 of definition 2.1.

{mod:closed}

$\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I^{1,2}] \longrightarrow V^2$ is a morphism of $\mathcal{E}^2 \overset{\text{cat}}{\subseteq} \mathcal{E}^3$ by hypothesis

and $\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^2$ is a morphism of $\mathcal{E}^2 \overset{\text{cat}}{\subseteq} \mathcal{E}^3$ by item 4. mod:restriction

of definition 2.1. $\phi^3 \circ f^2 \circ \phi^{2-1}: \phi^2[I^{2,3}] \longrightarrow V^3$ is a morphism of $\mathcal{C}^3$ by hypothesis and thus $(\phi^3 \circ f^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^3) = (\phi^3 \circ f^2 \circ \phi^{2-1}: \phi^2[I^{2,3}] \longrightarrow V^3) \circ (\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^2)$ is a morphism of $\mathcal{C}^3$. $(\phi^3 \circ f^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^3)$ agrees with $(\phi^3 \circ f^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3)$ on $\phi^1[I_{u^1}^{1,3}]$ and $\phi^1[I] = \bigcup_{\text{mod! sheaf}} \phi^1[I_{u^1}^{1,3}]$, thus $(\phi^3 \circ f^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3)$ is a morphism of $\mathcal{C}^3$ by item 6 of definition 2.1. {def:model}

$f^2 \circ f^1$ is **(semi-strict, strict)**. If $E^1 \subseteq^{\text{mod}} E^2$ and $E^2 \subseteq^{\text{mod}} E^3$ then $E^1 \subseteq^{\text{mod}} E^3$. If $C^1 \subseteq^{\text{mod}} C^2$ and $C^2 \subseteq^{\text{mod}} C^3$ then $C^1 \subseteq^{\text{mod}} C^3$. If $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^2$ and $\mathcal{C}^2 \subseteq^{\text{cat}} \mathcal{C}^3$ then $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^3$.

If f^1 is a local morphism of E^2 and f^2 is a local morphism of E^3 then $f^2 \circ f^1$ is a local morphism of E^3 by lemma 7.15 (Composition of local m-morphisms) on page 38. {lem:modlocalMmorphcomp}

If f^1 is a local $\mathcal{C}^1$ - $\mathcal{C}^2$ morphism of E^2 and f^2 is a local $\mathcal{C}^2$ - $\mathcal{C}^3$ morphism of E^3 then $f^2 \circ f^1$ is a local $\mathcal{C}^1$ - $\mathcal{C}^3$ morphism of E^3 by lemma 7.15. {lem:modlocalMmorphcomp}

If f^1 is a morphism of $E^2 E^3$ and f^2 is a morphism of E^3 then $f^2 \circ f^1$ is a morphism of E^3 .

□

Similar results apply with restrictions on the admissible atlases.

{lem:M-ATLcid}

Lemma 13.10 (Classic m-atlas inclusion and identity). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathcal{C}^1 \subseteq^{\text{full-cat}} \mathcal{C}^2$
3. $E^i \in \mathcal{C}^i$
4. $E^1 \subseteq^{\text{mod}} E^2$
5. $C^i \in \mathcal{C}^i$
6. $C^1 \subseteq^{\text{mod}} C^2$ ($\mathcal{C}^1 \subseteq^{\text{full-cat}} \mathcal{C}^2$)
7. A^i be a maximal m-atlas of E^i in the coordinate space C^i
8. $A^1 \subseteq A^2$

The classic m-atlas inclusion morphism $\text{Id}_{(E^1, C^1)}^{(E^1, C^1)} = \text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a semi-strict classic E^1 - E^2 -($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and a classic $\mathcal{C}^1$ - $\mathcal{C}^2$ -($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof.

$\text{Id}_{E^2}^{E^1}$ is a classic m-atlas morphism:

Let $(U^i, V^i, \phi^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap U^2 \neq \emptyset$.

Since $\mathbf{A}^1 \subseteq \mathbf{A}^2$, (U^1, V^1, ϕ^1) is compatible with (U^2, V^2, ϕ^2) ,

$\phi^2 \circ \text{Id}_{E^2}^{E^1} \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2 = \phi^2 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of $\mathbf{C}^2$ and a morphism of $\mathcal{C}^2$.

$\text{Id}_{E^2}^{E^1}$ is semi-strict:

□

If each $\text{Top}(E^i) \in E^i$ ($E^i \in \mathcal{E}^i$) and $\mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$ ($\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2$) then the classic m-atlas inclusion morphism $\text{Id}_{(E^2, \mathbf{C}^2)}^{(E^1, \mathbf{C}^1)}$ is a strict classic $(\mathcal{E}^1 - \mathcal{E}^2 -)E^1 - E^2(-\mathcal{C}^1 - \mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof.

$\text{Id}_{(E^2, \mathbf{C}^2)}^{(E^1, \mathbf{C}^1)}$ is a classic m-atlas morphism $\text{Id}_{(E^2, \mathbf{C}^2)}^{(E^1, \mathbf{C}^1)}$ is a classic $E^1 - E^2(-\mathcal{C}^1 - \mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ and a classic $\mathcal{E}^1 - \mathcal{E}^2(-\mathcal{C}^1 - \mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$. by the above proof.

$\text{Id}_{(E^2, \mathbf{C}^2)}^{(E^1, \mathbf{C}^1)}$ is strict: If each $\text{Top}(E^i) \in E^i$ then $\text{Id}_{E^2}^{E^1}$ is a morphism of E^2 by item 5 of definition 2.1 (Model spaces) on page 18, hence an m-morphism of E^1 to E^2 . {mod:inclusion}

If each $E^i \in \mathcal{E}^i$ $\text{Id}_{E^2}^{E^1}$ is a morphism of $\mathcal{E}^2$ by item 6 of definition 7.7 (Model categories) on page 29, hence an $\mathcal{E}^1 - \mathcal{E}^2$ m-morphism of E^1 to E^2 . {modcat:morph}

□

{cor:M-ATLcid}

Corollary 13.11 (Classic m-atlas identity). Let $\mathcal{E}, \mathcal{C}$, be model categories, $E \in \mathcal{E}$, $\mathbf{C} \in \mathcal{C}$, $\mathbf{A}$ an m-atlas of E in the coordinate spaces $\mathbf{C}$.

The identity morphism $\mathbf{f} \stackrel{\text{def}}{=} \text{Id}_{(\mathbf{A}, E, \mathbf{C})}$ is a strict m-atlas morphism of $\mathbf{A}$ to $\mathbf{A}$ in the coordinate spaces $\mathbf{C}, \mathbf{C}$.

13.2 Categories of classic m-atlases and functors

{sub:C-cat}

This subsection defines categories of m-atlases with classic m-atlas morphisms and functors among them, and proves some basic results.

13.2.1 Classic m-atlas categories

Definition 13.12 (Sets of classic m-atlas morphisms). Let E^i and C^i . $i = 1, 2$, be model spaces. Then

$$\begin{aligned} \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(E^1, C^1, E^2, C^2) &\stackrel{\text{def}}{=} \\ &\left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \right) \\ &\left\{ ((f), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \begin{array}{c} \text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f) \\ \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \right\} \end{aligned} \quad (13.1)$$

{def:C-cat}

Definition 13.13 (Categories $\mathcal{Atl}^{\text{Classic}}(E, C)$). Let E and C be sets of model spaces.

Let $P \stackrel{\text{def}}{=} E \times C$. Then

$$\begin{aligned} \mathcal{Atl}_{\text{Ob}}^{\text{classic}}(E, C) &\stackrel{\text{def}}{=} \bigcup_{(E^\mu, C^\mu) \in P} \mathcal{Atl}_{\text{Ob}}^{\text{classic}}(E^\mu, C^\mu) \\ &\left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \right) \end{aligned} \quad (13.2)$$

$$\begin{aligned} \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(E, C) &\stackrel{\text{def}}{=} \bigcup_{\substack{(E^\mu, C^\mu) \in P \\ (E^\nu, C^\nu) \in P}} \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(E^\mu, C^\mu, E^\nu, C^\nu) \\ &\left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \right) \end{aligned} \quad (13.3)$$

$$\begin{aligned} \mathcal{Atl}^{\text{classic}}(E, C) &\stackrel{\text{def}}{=} \left(\mathcal{Atl}_{\text{Ob}}^{\text{classic}}(E, C), \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(E, C), \overset{A}{\circ} \right) \\ &\left(\begin{array}{c} \text{(full, S-max, max, S-max-full, max-full)} \\ \text{(semi-strict, strict)} \end{array} \right) \end{aligned} \quad (13.4)$$

Let E^i and C^i , $i = 1, 2$, be model spaces, A^i be an atlas of E^i in the coordinate space C^i and $f: E^1 \longrightarrow E^2$ be a (strict, semi-strict) E^1 - E^2 classic M-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The identity functor $\mathcal{F}_{\text{Classic, Id}}$ is

$$\mathcal{F}_{\text{Classic, Id}}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A^i, E^i, C^i) \quad (13.5)$$

$$\begin{aligned} \mathcal{F}_{\text{Classic, Id}}((f: E^1 \longrightarrow E^2), (A^1, E^1, C^1), (A^2, E^2, C^2)) &\stackrel{\text{def}}{=} \\ ((f: E^1 \longrightarrow E^2), (A^1, E^1, C^1), (A^2, E^2, C^2)) \end{aligned} \quad (13.6)$$

This nomenclature will be justified below.

{lem:M-ATLCiscat}

Lemma 13.14 ($\mathcal{Atl}^{\text{Classic}}(E, C)$ is a category). Let E and C be sets of model spaces. Then $\mathcal{Atl}^{\text{Classic}}(E, C)$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{Atl}_{\text{Ob}}^{\text{Classic}}(E, C)$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{A}tl^{\text{Classic}}(E, C)$ and let $m^i \stackrel{\text{def}}{=} ((f^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$ be a morphism of $\mathcal{A}tl^{\text{Classic}}(E, C)$. Then

Composition: $f^2 \circ f^1$ is a classic E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 by lemma 13.9 (Composition of classic m-atlas morphisms) on page 106 and $(f^2) \circ (f^1) = (f^2 \circ f^1)$. {lem:M-ATLcmorphcomp}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19. {lem:labcomp}

□

{cor:ATLCiscat}

Corollary 13.15 (Subcategories of $\mathcal{A}tl^{\text{Classic}}(E, C)$). Let E and C be sets of model spaces. Then all of the following are subcategories of $\mathcal{A}tl^{\text{Classic}}(E, C)$.

1. $\mathcal{A}tl_{\text{full}}^{\text{Classic}}(E, C)$
2. $\mathcal{A}tl_{\text{S-max}}^{\text{Classic}}(E, C)$
3. $\mathcal{A}tl_{\text{max}}^{\text{Classic}}(E, C)$
4. $\mathcal{A}tl_{\text{S-max-full}}^{\text{Classic}}(E, C)$
5. $\mathcal{A}tl_{\text{max-full}}^{\text{Classic}}(E, C)$
6. $\mathcal{A}tl_{\text{semi-strict}}^{\text{Classic}}(E, C)$
7. $\mathcal{A}tl_{\text{full semi-strict}}^{\text{Classic}}(E, C)$
8. $\mathcal{A}tl_{\text{S-max semi-strict}}^{\text{Classic}}(E, C)$
9. $\mathcal{A}tl_{\text{max semi-strict}}^{\text{Classic}}(E, C)$
10. $\mathcal{A}tl_{\text{S-max-full semi-strict}}^{\text{Classic}}(E, C)$
11. $\mathcal{A}tl_{\text{max-full semi-strict}}^{\text{Classic}}(E, C)$

$$12. \mathcal{Att}_{\text{strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

$$13. \mathcal{Att}_{\text{full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

$$14. \mathcal{Att}_{\text{S-max strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

$$15. \mathcal{Att}_{\text{max strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

$$16. \mathcal{Att}_{\text{S-max-full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

$$17. \mathcal{Att}_{\text{max-full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$$

Remark 13.16. They are not, in general, full subcategories.

The identity functor $\mathcal{F}_{\text{Classic}, \text{Id}}$ is a functor from each of the subcategories to each of the containing categories.

{def:C-cat-cat}

Definition 13.17 (Categories $\mathcal{Att}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories.

Let $\mathbf{P} \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \text{Ob}(\mathcal{C})$. Then

$$\mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E \in \mathcal{E} \\ C \in \mathcal{C}}} \mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathbf{E}, \mathbf{C}) \quad (13.7)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left\{ ((f), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. (E^i, C^i) \in \mathbf{P} \wedge \text{isAtl}_{\text{Ar}}^{\text{strict}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}, \mathcal{C}) \right\} \quad (13.8)$$

$$\mathcal{Att}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} (\mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}), \mathcal{Att}_{\text{Ar}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}), \overset{A}{\circ}) \quad (13.9)$$

Similar definitions apply with restrictions on the admissible atlases, the admissible morphisms, or both:

$$1. \mathcal{Att}_{\text{full}}^{\text{Classic}}$$

$$2. \mathcal{Att}_{\text{S-max}}^{\text{Classic}}$$

$$3. \mathcal{Att}_{\text{max}}^{\text{Classic}}$$

4. $\mathcal{Att}_{\text{S-max-full}}^{\text{Classic}}$
5. $\mathcal{Att}_{\text{max-full}}^{\text{Classic}}$
6. $\mathcal{Att}_{\text{full semi-strict}}^{\text{Classic}}$
7. $\mathcal{Att}_{\text{S-max semi-strict}}^{\text{Classic}}$
8. $\mathcal{Att}_{\text{max semi-strict}}^{\text{Classic}}$
9. $\mathcal{Att}_{\text{S-max-full semi-strict}}^{\text{Classic}}$
10. $\mathcal{Att}_{\text{max-full semi-strict}}^{\text{Classic}}$
11. $\mathcal{Att}_{\text{full strict}}^{\text{Classic}}$
12. $\mathcal{Att}_{\text{S-max strict}}^{\text{Classic}}$
13. $\mathcal{Att}_{\text{max strict}}^{\text{Classic}}$
14. $\mathcal{Att}_{\text{S-max-full strict}}^{\text{Classic}}$
15. $\mathcal{Att}_{\text{max-full strict}}^{\text{Classic}}$

e.g.,

$$\mathcal{Att}_{\text{Ob S-max-full}}^{\text{Classic}} l(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{(E^\mu, C^\mu) \in \mathbf{P}} \mathcal{Att}_{\text{Ob S-max-full}}^{\text{Classic}}(E^\mu, C^\mu) \quad (13.10)$$

$$\mathcal{Att}_{\text{Ar S-max-full semi-strict}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^\mu, C^\mu) \in \mathbf{P} \\ (E^\nu, C^\nu) \in \mathbf{P}}} \mathcal{Att}_{\text{Ar S-max-full semi-strict}}^{\text{Classic}}(E^\mu, C^\mu, E^\nu, C^\nu) \quad (13.11)$$

$$\mathcal{Att}_{\text{S-max-full semi-strict}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob S-max-full}}^{\text{Classic}} l(\mathcal{E}, \mathcal{C}), \mathcal{Att}_{\text{Ar S-max-full semi-strict}}^{\text{Classic}}(E, C), \circ^A \right) \quad (13.12)$$

Let $(A^1, E^1, C^1) \in \mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$.

$$\text{Id}_{(A^1, E^1, C^1)} \stackrel{\text{def}}{=} ((\text{Id}_{E^1}, \text{Id}_{C^1}), (A^1, E^1, C^1), (A^1, E^1, C^1)) \quad (13.13)$$

{lem:C-cat-ATLCiscat}

Lemma 13.18 ($\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is a category). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Then $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ and let $m^i \stackrel{\text{def}}{=} ((f^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$, $i = 1, 2$, be a morphism of $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$. Then

Composition: $f^2 \circ f^1$ is a morphism of $\mathcal{E}$ and is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 by lemma 13.9 (Composition of classic m-atlas morphisms) on page 106. {lem:M-ATLcmorphcomp}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

Similar results follow with restrictions on the admissible atlases, the admissible morphisms, or both.

13.2.2 Classic m-atlas functors

{def:Func-M-C}

Definition 13.19 ($\mathcal{F}_{\text{M,Classic}}$). Let E^i , $i = 1, 2$, C^i be model spaces and A^i an m-atlas of E^i in the coordinate model space C^i . Then

$$\mathcal{F}_{\text{M,Classic}}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A^i, E^i, C^i) \quad (13.14)$$

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2,)$ be an M-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then

$$\begin{aligned} \mathcal{F}_{\text{M,Classic}}((f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2,), (A^1, E^1, C^1), (A^2, E^2, C^2)) &\stackrel{\text{def}}{=} \\ ((f_0: E^1 \longrightarrow E^2), (A^1, E^1, C^1), (A^2, E^2, C^2)) &\quad (13.15) \end{aligned}$$

{the:Func-M-C}

Theorem 13.20 ($\mathcal{F}_{\text{M,Classic}}$ is a functor). Let E and C be sets of model spaces, Then $\mathcal{F}_{\text{M,Classic}}$ is a functor from $\mathcal{A}tl(E, C)$ to $\mathcal{A}tl^{\text{Classic}}(E, C)$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $i \in [1, 3]$, be an object of $\mathcal{A}tl(E, C)$ and $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$, $i = 1, 2$, be a morphism of $\mathcal{A}tl(E, C)$. $\mathcal{F}_{\text{M,Classic}}$ satisfies these criteria:

Preservation of endpoints:

$$\begin{aligned}\mathcal{F}_{\mathbf{M}, \text{Classic}} o^i &= o^i \in \text{Ob} \mathcal{Att}^{\text{classic}}(\mathbf{E}, \mathbf{C}) \\ \mathcal{F}_{\text{Top}, \mathbf{M}} m^i &= ((f_0: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^{i+1}, \mathbf{E}^{i+1}, \mathbf{C}^{i+1})) \\ &\text{is a morphism from } o^i \text{ to } o^{i+1}.\end{aligned}$$

Identity: Let $((\text{Id}_{\mathbf{E}^i}, \text{Id}_{\mathbf{C}^i}), o^i, o^i)$ be an identity morphism of o^i in $\mathcal{Att}(\mathbf{E}, \mathbf{C})$. Then $((\text{Id}_{\mathbf{E}^i}: \mathbf{E}^i \longrightarrow \mathbf{E}^i), o^i, o^i)$ is an identity morphism of o^i in $\mathcal{Att}^{\text{classic}}(\mathbf{E}, \mathbf{C})$.

Composition:

$$\begin{aligned}\mathcal{F}_{\mathbf{M}, \text{Classic}}(m^2 \overset{A}{\circ} m^i) &= \mathcal{F}_{\mathbf{M}, \text{Classic}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ &= ((f_0^2 \circ f_0^1), o^1, o^3) \\ &= ((f_0^2), o^1, o^2) \overset{A}{\circ} ((f_0^1), o^2, o^3) \\ &= \mathcal{F}_{\mathbf{M}, \text{Classic}}((f_0^2, f_1^2), o^2, o^3) \overset{A}{\circ} \\ &\quad \mathcal{F}_{\mathbf{M}, \text{Classic}}((f_0^1, f_1^1), o^1, o^2) \\ &= \mathcal{F}_{\mathbf{M}, \text{Classic}} m^2 \overset{A}{\circ} \mathcal{F}_{\mathbf{M}, \text{Classic}} m^1\end{aligned}$$

□

Part VII

Equivalence of manifolds

{part:man}

For a manifold¹¹ the coordinate category is open subsets of a Banach space or more generally a Fréchet space, with an appropriate choice of morphisms. Choosing a separating hyperplane and half space, with open sets in the chosen half space, allows manifolds with boundary. Similarly, choosing a ball with tails allows a manifold with tails.

For differentiable manifolds the coordinate category is similar, but the morphisms are limited to those sufficiently differentiable, in order to impose a differentiability constraint on the transition functions $t_\beta^\alpha = \phi_\beta \circ \phi_\alpha^{-1}$.

This part of the paper defines C^k -atlases, C^k -manifolds, local coordinate spaces equivalent to C^k -manifolds, categories of them and functors, and gives some basic results.

¹¹The literature has several different definitions of a manifold. This paper uses one chosen for ease of exposition.

14 Linear spaces and linear model spaces

This subsection defines spaces and related model spaces suitable for use as the coordinate spaces of generalized C^k manifolds. sec:lin

Definition 14.1 (Linear spaces). Let C be a locally arcwise connected topological subspace of a real (complex) Banach or Fréchet space. Then C is a real (complex) linear space. If C has a non-void interior, i.e., contains a ball, then it is non-degenerate. def:lin

Remark 14.2. This paper uses the term *ball* to refer to balls of the underlying space but uses the terms *open set* and *neighborhoods* to refer to the relative topology.

Let $\mathcal{C}$ be a small category whose objects are real (complex) linear spaces and whose morphisms are C^k functions. Then $\mathcal{C}$ is a C^k linear category. If each object of $\mathcal{C}$ is non-degenerate then $\mathcal{C}$ is non-degenerate.

Let C be a real (complex) linear space. Then $\mathcal{C}_{C^k\text{-}Triv}$, the category of all C^k functions between nonvoid open subspaces of C , is the trivial C^k linear category of C .

Lemma 14.3 (Open subsets of linear spaces are locally arcwise connected). Let U be an open subset of the real (complex) linear space S . Then U is locally arcwise connected.

Proof. An open subset of a locally arcwise connected space is locally arcwise connected. □

Definition 14.4 (Linear model spaces). Let S be a real (complex) linear space and $\mathcal{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ a model space for S . Then $\mathcal{S}$ is a real (complex) linear model space.

Let $\mathcal{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a real (complex) linear model space such that every morphism of $\mathcal{S}$ is a C^k function. Then $\mathcal{S}$ is a real (complex) C^k linear model space.

Let $\mathcal{S}$ be a small category whose objects are real (complex) C^k linear model spaces and whose morphisms are C^k model functions. Then $\mathcal{S}$ is a C^k linear model category.

Definition 14.5 (Trivial C^k linear model spaces). Let C be a real (complex) linear space and $\mathcal{C}$ the category of all C^k functions between open sets of C . Then $\mathcal{C}_{C^k\text{-}triv} \stackrel{\text{def}}{=} (C, \mathcal{C})$ is the trivial C^k linear model space of C and $\mathcal{C}_{C^k\text{-}triv}$ is a real (complex) trivial C^k linear model space. {def:trivck}

Let $\mathcal{C}$ be a set of real (complex) linear spaces.

$\mathcal{C}_{C^k\text{-}triv} \stackrel{\text{def}}{=} \left\{ \mathcal{C}' \mid \mathcal{C}' \in \mathcal{C} \right\}$ is the set of trivial C^k linear model spaces of $\mathcal{C}$.

The category of trivial C^k linear model spaces of $\mathcal{C}$, abbreviated $\mathcal{C}_{C^k\text{-}Triv}$, is the category whose objects are $\mathcal{C}_{C^k\text{-}triv}$ and whose morphisms are all the C^k functions among them.

$\mathcal{C}_{\text{C}^k\text{-op-triv}}$, the category of open trivial C^k model spaces in $\mathcal{C}$, is the category

whose objects are $\left\{ U_{\text{C}^k\text{-triv}} \mid U \in \mathcal{C}_{\text{op}} \right\}$, the trivial C^k linear model spaces of non-null open sets of spaces in $\mathcal{C}$, and whose morphisms are all the C^k functions among them.

{lem:trivck}

Lemma 14.6 (Trivial C^k linear model spaces). Let C^i , $i = 1, 2$, be a real (complex) linear space and $\mathcal{C}^i \stackrel{\text{def}}{=} \text{Cat}_{\text{C}^k\text{-triv}}(C^i)$.

1. $\mathcal{C}_{\text{C}^k\text{-triv}}^i$ is a C^k linear model space.

Proof.

- (a) $\text{Ob}(\mathcal{C}^i)$ is an open cover for C^i .
- (b) $\text{Ob}(\mathcal{C}^i)$ is closed under finite intersections.
- (c) The morphisms of $\mathcal{C}^i$ are C^k , hence continuous.
- (d) If $f: A \rightarrow B$ is a morphism, $A' \in \mathcal{C}^i \subseteq A \in \mathcal{C}^i$, $B' \in \mathcal{C}^i \subseteq B \in \mathcal{C}^i$ and $f[A'] \subseteq B'$ then $f|_{A'}: A' \rightarrow B'$ is C^k and thus a morphism.
- (e) If $A' \in \mathcal{C}^i \subseteq A \in \mathcal{C}^i$ then the inclusion map $i: A' \hookrightarrow A$ is C^k and thus a morphism.
- (f) Restricted sheaf condition: Whenever
 - i. U_α and V_α , $\alpha < A$, are objects of $\mathcal{C}^i$.
 - ii. $f_\alpha: U_\alpha \rightarrow V_\alpha$ are morphisms of $\mathcal{C}^i$.
 - iii. $U \stackrel{\text{def}}{=} \bigcup_{\alpha < A} U_\alpha \in \mathcal{C}^i$,
 - iv. $V \stackrel{\text{def}}{=} \bigcup_{\alpha < A} V_\alpha \in \mathcal{C}^i$
 - v. $f: U \rightarrow V$ is a continuous function and for every $\alpha < A$, f agrees with f_α on U_α
then f is C^k and thus a morphism of $\mathcal{C}^i$.

□

2. Let $f: C^1 \rightarrow C^2$ be C^k . Then f is a model function from $\mathcal{C}_{\text{C}^k\text{-triv}}^1$ to $\mathcal{C}_{\text{C}^k\text{-triv}}^2$.

Proof. Let U^i be a model neighborhood of $\mathcal{C}_{\text{C}^k\text{-triv}}^i$, $i = 1, 2$.

- (a) Since U^2 is a model neighborhood of $\mathcal{C}_{\text{C}^k\text{-triv}}^2$, U^2 is open, $f^{-1}[U^2]$ is open and thus a model neighborhood of $\mathcal{C}_{\text{C}^k\text{-triv}}^1$.

(b) $f[U^1] \subseteq C^2$. Since C^2 is open, it is a model neighborhood of $C^2_{\mathbf{C}^k\text{-triv}}$.

□

3. Let C^1 be an open subspace of C^2 . Then $C^1_{\mathbf{C}^k\text{-triv}} \stackrel{\text{mod}}{\subseteq} C^2_{\mathbf{C}^k\text{-triv}}$.

Proof. By definition 14.5 above, $C^i_{\mathbf{C}^k\text{-triv}} = (C^i, \mathcal{C}^i)$, where $\mathcal{C}^i$ is the category of all $\mathbf{C}^k$ functions between open sets of C^i . {def:trivck}

$$C^1_{\mathbf{C}^k\text{-triv}} \stackrel{\text{mod}}{\subseteq} C^2_{\mathbf{C}^k\text{-triv}} :$$

- (a) C^1 is an open subspace of C^2 by hypothesis.
- (b) Every object of $\mathcal{C}^1$ is open in C^1 , thus open in C^2 and thus an object of $\mathcal{C}^2$. If $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^1$ then U and V are open in C^1 and f is $\mathbf{C}^k$ in C^1 , thus U and V are open in C^2 and f is $\mathbf{C}^k$ in C^2 and $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^2$. If $f: U \xrightarrow{\text{Ob}} V$ is a morphism of $\mathcal{C}^2$, $U \in \mathcal{C}^1$ and $V \in \mathcal{C}^1$, then U and V are open in C^1 and f is $\mathbf{C}^k$ in C^2 , thus $\mathbf{C}^k$ in C^1 , and $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^1$.
- (c) If $U \in \mathcal{C}^2$ then U is open in C^2 , hence $U \cap C^1$ is open in C^2 , hence $U \cap C^1$ is open in C^1 and $U \in \mathcal{C}^1$.
- (d) If $U \in \mathcal{C}^2$ and $U \subseteq C^1$ then U is open in C^2 , hence open in C^1 , hence $U \in \mathcal{C}^1$.

□

{def:singCk}

Definition 14.7 ($\mathbf{C}^k$ singleton categories). Let $\mathbf{C}$ be a $\mathbf{C}^k$ linear model space. Then the $\mathbf{C}^k$ singleton category of $\mathbf{C}$, abbreviated $\mathbf{C}_{\mathbf{C}^k\text{-Sing}}$, is the category whose sole object is $\mathbf{C}$ and whose morphisms are all of the $\mathbf{C}^k$ model functions from $\mathbf{C}$ to itself.

15 $\mathbf{C}^k$ -nearly commutative diagrams

{sec:Ck-NCD}

Let C be a linear space, $\mathcal{C} \stackrel{\text{def}}{=} S_{\mathbf{C}^k\text{-Triv}}$, $\mathbf{C} \stackrel{\text{def}}{=} C_{\mathbf{C}^k\text{-triv}}$ and D a tree with two branches, whose nodes are topological spaces U_i and V_j and whose links are continuous functions $f_i: U_i \longrightarrow U_{i+1}$ and $f'_j: U_j \longrightarrow U_{j+1}$ between the spaces:

$$D = \{f_0: U_0 = V_0 \longrightarrow U_1, \dots, f_{m-1}: U_{m-1} \longrightarrow U_m, \\ f'_0: U_0 = V_0 \longrightarrow V_1, \dots, f'_{m-1}: V_{m-1} \longrightarrow V_n\}$$

with $U_0 = V_0$, $U_m \subseteq C$ and $V_n \subseteq C$ open, as shown in fig. 1 (Uncompleted nearly commutative diagram) on page 15. {fig:NCDb}

Definition 15.1 (C^k -nearly commutative diagrams). D is (left,right,strongly) C^k -nearly commutative in linear space C iff D is (left,right,strongly) M -nearly commutative in model space $\mathcal{C}$. {def:Ck-NCD}

Definition 15.2 (C^k -nearly commutative diagrams at a point). Let $C, \mathcal{C}, \mathbf{C}$ and D be as above and x be an element of the initial node. D is (left,right,strongly) C^k -nearly commutative in C at x iff D is (left,right,strongly) M -nearly commutative in model space $\mathbf{C}$ at x . {def:Ck-NCDp}

Definition 15.3 (C^k -locally nearly commutative diagrams). Let $C, \mathcal{C}, \mathbf{C}$ and D be as above. D is (left,right,strongly) C^k -locally nearly commutative in C iff D is (left,right,strongly) M -locally nearly commutative in model space $\mathbf{C}$. {def:Ck-NCDl}

Lemma 15.4. Let $C, \mathcal{C}, \mathbf{C}$ and D be as above. D is C^k -nearly commutative in linear space C iff There is a C^k diffeomorphism $\hat{f}: U_m \xrightarrow{\cong} V_n$ making the graph a commutative diagram, as shown in fig. 2. {lem:Ck-NCD}

Proof. A set is a model neighborhood of $\mathbf{C}$ (an object of $\mathcal{C}^{k-\text{Triv}}$) iff it is an open subset of A . A function is a morphism of $\mathbf{C}$ (a morphism of $\mathcal{C}^{k-\text{Triv}}$) iff it is a C^k function between open subsets of S . {fig:NCDa}

16 C^k charts

Definition 16.1 (C^k charts). Let C be a linear space and E a topological space. A C^{k12} chart (U, V, ϕ) of E in the coordinate space C consists of sec:ckcharts {def:Ck-chart}

1. A nonvoid open subset $U \subseteq E$, known as a coordinate patch
2. An open subset $V \subseteq C$
3. A homeomorphism $\phi: U \xrightarrow{\cong} V$, known as a coordinate function

Remark 16.2. I consider it clearer to explicate the range, rather than the conventional usage of specifying only the domain and function or the minimalist usage of specifying only the function.

A chart (U, V, ϕ) is non-degenerate iff V contains a ball of the underlying Banach or Fréchet space. {lem:Ck-chart}

Lemma 16.3 (C^k charts). Let C be a linear space and E a topological space. The triple (U, V, ϕ) is a C^k chart of E in the coordinate space C iff it is an m -chart of E in the coordinate space $\mathcal{C}^{k-\text{triv}}$. {triv}

¹²With $k \in \mathbb{N} \cup \{\infty, \omega\}$.

Proof. U is a model neighborhood of E_{triv} iff it is a nonvoid open set of E . V is a model neighborhood of $C_{\text{Ck-triv}}$ iff it is a nonvoid open set of C . ϕ is required to be a homeomorphism from U to V in either case. $\square$

{def:Ck-subchart}

Definition 16.4 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and U' be a nonvoid open subset of U . Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

By abuse of language we will write (U', V', ϕ) for (U', V', ϕ') .

{lem:Ck-subchart}

Lemma 16.5 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and (U', V', ϕ') be a triple. Then (U', V', ϕ') is a subchart of (U, V, ϕ) iff it is a subchart of (U, V, ϕ) considered as a chart of E_{triv} in the coordinate space $C_{\text{Ck-triv}}$.

Proof. (U', V', ϕ') satisfies the conditions of definition 16.1;

{def:Ck-chart}

1. U is open by definition 16.1
2. U' is required to be a nonvoid open subset of U in either case, and thus a model neighborhood of E_{triv} .
3. ϕ is a homeomorphism, so $V' = \phi[U']$ is open, and thus a model neighborhood of $C_{\text{Ck-triv}}$.
4. U' is a model neighborhood of E_{triv} iff it is an open set of E .
5. ϕ is a homeomorphism, so $\phi|_{U'}: U' \xrightarrow{\cong} \phi[U']$ is also.

$\square$

{cor:Ck-subchart}

Corollary 16.6 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and (U', V', ϕ') be a subchart of (U, V, ϕ) . Then (U', V', ϕ') is a C^k chart of E in the coordinate space C .

Proof. The result follows from lemma 16.3 (C^k charts) above, lemma 9.3 (M -charts) on page 42 and definition 9.5 (Subcharts) on page 43. $\square$

{lem:Ck-chart}

{def:M-subchart}
{Ck-compat}

Definition 16.7 (C^k compatibility). Let (U, V, ϕ) and (U', V', ϕ') be C^k charts of E in the coordinate space C . Then (U, V, ϕ) is C^k compatible with (U', V', ϕ') iff either

1. U and U' are disjoint
2. The transition function $t = \phi' \circ \phi^{-1}|_{\phi[U \cap U']}$ is a C^k diffeomorphism.

{lem:Ck-compatism}

Lemma 16.8 (Symmetry of C^k compatibility). Let (U, V, ϕ) and (U', V', ϕ') be C^k charts of E in the coordinate space C . Then (U, V, ϕ) is C^k compatible with (U', V', ϕ') iff (U', V', ϕ') is C^k compatible with (U, V, ϕ) .

Proof. It suffices to prove the implication in only one direction.

1. $U \cap U' = U' \cap U$.
2. Since the transition function $t = \phi' \circ \phi^{-1} \upharpoonright_{\phi[U \cap U']}$ is a C^k diffeomorphism of C , so is $t^{-1} = \phi \circ \phi'^{-1} \upharpoonright_{\phi'[U \cap U']}$.

□

Lemma 16.9 (C^k compatibility of subcharts). Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be C^k charts of E in the coordinate space C , (U'_i, V'_i, ϕ'_i) be subcharts and (U_1, V_1, ϕ_1) be C^k compatible with (U_2, V_2, ϕ_2) . Then (U'_1, V'_1, ϕ'_1) is C^k compatible with (U'_2, V'_2, ϕ'_2) .

Proof. If $U_1 \cap U_2 = \emptyset$ then $U'_1 \cap U'_2 = \emptyset$. If $U'_1 \cap U'_2 = \emptyset$ then (U'_1, V'_1, ϕ'_1) is C^k compatible with (U'_2, V'_2, ϕ'_2) . Otherwise, the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism and hence $t_2^1 \upharpoonright_{\phi_1[U'_1 \cap U'_2]}: \phi_1[U'_1 \cap U'_2] \xrightarrow{\cong} \phi_2[U'_1 \cap U'_2]$ is a C^k diffeomorphism. □

Corollary 16.10 (C^k compatibility with subcharts). Let (U, V, ϕ) be a C^k charts of E in the coordinate space C and (U', V', ϕ') a subchart. Then (U', V', ϕ') is C^k compatible with (U, V, ϕ) .

Proof. (U, V, ϕ) is C^k compatible with itself and is a subchart of itself. □

Definition 16.11 (Covering by C^k charts). Let $\mathbf{A}$ be a set of charts of E in the coordinate space C . $\mathbf{A}$ covers E iff $\pi_1[\mathbf{A}]$ covers E , i.e., $E = \bigcup \pi_1[\mathbf{A}]$.

17 C^k -atlases

{sec:ckatlases}

A set of charts can be atlases for different coordinate spaces even if it is for the same total space. In order to aggregate them into categories, there must be a way to distinguish them. Including the two¹³ spaces in the definitions of the categories serves the purpose.

{def:Ck-atlas}

Definition 17.1 (C^k -atlases). Let $\mathbf{A}$ be a set of mutually C^k compatible charts of E in the coordinate space C . $\mathbf{A}$ is a C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C)$, iff $\mathbf{A}$ covers E .

$\mathbf{A}$ is a full C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C)$,
iff

1. $\pi_1[\mathbf{A}]$ covers E
2. $\pi_2[\mathbf{A}]$ covers C .

¹³The total space is redundant, but convenient.

$\mathbf{A}$ is non-degenerate iff it contains a non-degenerate chart.

By abuse of language we write $U \in \mathbf{A}$ for $U \in \pi_1[\mathbf{A}]$.

Let E be a topological space and C a linear space. Then

$$\mathcal{Atl}_{\text{Ob}}^{\text{C}^k}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{\text{C}^k}(\mathbf{A}, E, C) \right\} \quad (17.1)$$

$$\mathcal{Atl}_{\text{Ob}}^{\text{C}^k}{}_{\text{full}}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{\text{C}^k}{}_{\text{full}}(\mathbf{A}, E, C) \right\} \quad (17.2)$$

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then

$$\mathcal{Atl}_{\text{Ob}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E_\mu \in \mathbf{E} \\ C_\mu \in \mathbf{C}}} \mathcal{Atl}_{\text{Ob}}^{\text{C}^k}(E_\mu, C_\mu) \quad (17.3)$$

$$\mathcal{Atl}_{\text{Ob}}^{\text{C}^k}{}_{\text{full}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{\text{C}^k}{}_{\text{full}}(\mathbf{A}, E, C) \right\} \quad (17.4)$$

{lem:Ck-atlas}

Lemma 17.2 (C^k -atlases). Let E be a topological space, C be a linear space and $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C .

If $\mathbf{A}$ is non-degenerate then C is non-degenerate.

Proof. Let (U, V, ϕ) be a non-degenerate chart of $\mathbf{A}$. V contains a ball and is contained in C . $\square$

If C is non-degenerate and $\mathbf{A}$ is full then $\mathbf{A}$ is non-degenerate.

Proof. Let B be a ball in C with center v . Since $\mathbf{A}$ is full, it contains a chart (U, V, ϕ) with $v \in V$. Then V contains a ball with center v . $\square$

{def:CkCompat}

Definition 17.3 (Compatibility of charts with C^k -atlases). A chart (U, V, ϕ) of E in the coordinate space C is C^k compatible with a C^k -atlas $\mathbf{A}$ iff it is C^k compatible with every chart in the atlas.

{lem:CkCompat}

Lemma 17.4 (Compatibility of subcharts with C^k -atlases). Let $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C and $\mathbf{C}_1 = (U_1, V_1, \phi_1)$ a C^k chart in $\mathbf{A}$. Then any subchart of $\mathbf{C}_1$ is C^k compatible with $\mathbf{A}$.

Proof. Let $\mathbf{C}' = (U', V', \phi')$ be a subchart of $\mathbf{C}_1$ and $\mathbf{C}_2 = (U_2, V_2, \phi_2)$ another chart in $\mathbf{A}$.

1. If $U_1 \cap U_2 = \emptyset$, then $U' \cap U_2 = \emptyset$.
2. If $U' \cap U_2 = \emptyset$ then $\mathbf{C}'$ is C^k compatible with $\mathbf{C}_2$.
3. Otherwise the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism and thus $t_2^1 \upharpoonright_{\phi_1[U' \cap U_2]}$ is a C^k diffeomorphism.

$\square$

{Ck-atl:extensions}

Lemma 17.5 (Extensions of C^k -atlases). Let $\mathbf{A}$ be a C^k atlas of E in the coordinate space C and (U_i, V_i, ϕ_i) , $i = 1, 2$, be a C^k chart of E in the coordinate space C compatible with $\mathbf{A}$ in the coordinate space C . Then (U_1, V_1, ϕ_1) is C^k compatible with (U_2, V_2, ϕ_2) in the coordinate space C .

Proof. If $U_1 \cap U_2 = \emptyset$ then (U_1, V_1, ϕ_1) is C^k compatible with (U_2, V_2, ϕ_2) . Otherwise, $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}: \phi_1[U_1 \cap U_2] \xrightarrow{\cong} \phi_2[U_1 \cap U_2]$ is a homeomorphism. It remains to show that $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism. Let $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\alpha < A$, be charts in $\mathbf{A}$ such that $U_1 \cap U_2 \subseteq \bigcup_{\alpha < A} U'_\alpha$ and $U_1 \cap U_2 \cap U'_\alpha \neq \emptyset$, $\alpha < A$. Since the charts are C^k compatible with $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\phi_2 \circ \phi'^{-1}_\alpha \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ and $\phi'_\alpha \circ \phi_1^{-1} \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ are C^k diffeomorphisms and thus $\phi_2 \circ \phi_1^{-1} = \phi_2 \circ \phi'^{-1}_\alpha \circ \phi'_\alpha \circ \phi_1^{-1}$ is a C^k diffeomorphism. $\square$

{def:Ck-ATLmax}

Definition 17.6 (Maximal C^k -atlases). Let E be a topological space, C be a linear space and $\mathbf{A}$ be a non-degenerate C^k -atlas of E in the coordinate space C .

$\mathbf{A}$ is a maximal C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C)$, iff $\mathbf{A}$ is a C^k -atlas that cannot be extended by adding an additional C^k compatible chart.

$$\text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \wedge \text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \quad (17.5)$$

max-full full max

$\mathbf{A}$ is a semi-maximal C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C)$,
iff whenever $(U, V, \phi) \in \mathbf{A}$, $U' \subseteq U$, $V' \subseteq V$ and $V'' \subseteq C$ are open, $\phi[U'] = V'$ and
 $\phi': V' \xrightarrow{\cong} V''$ is a C^k diffeomorphism then $(U', V'', \phi' \circ \phi) \in \mathbf{A}$.

$$\text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \wedge \text{isAtl}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, E, C) \quad (17.6)$$

S-max-full full S-max

Lemma 17.7 (Maximal C^k -atlases are semi-maximal C^k -atlases). Let E be a topological space, C a C^k -linear space and $\mathbf{A}$ a maximal C^k -atlas of E in the coordinate space C . Then $\mathbf{A}$ is a semi-maximal C^k -atlas of E in the coordinate space C .

Proof. Let $(U, V, \phi) \in \mathbf{A}$, $U' \subseteq U$, $V' \subseteq V$ and $V'' \subseteq C$ be open, $\phi[U'] = V'$ and $\phi': V' \xrightarrow{\cong} V''$ be a C^k diffeomorphism. (U', V', ϕ) is a subchart of (U, V, ϕ) and by lemma 17.4 (Compatibility of subcharts with C^k -atlases) on page 121 is C^k compatible with the charts of $\mathbf{A}$. Since ϕ' is a C^k diffeomorphism, $(U', V'', \phi' \circ \phi)$ is C^k compatible with the charts of $\mathbf{A}$. Since $\mathbf{A}$ is maximal, $(U', V'', \phi' \circ \phi)$ is a chart of $\mathbf{A}$. $\square$

{lem:CkCompat}

Theorem 17.8 (Existence and uniqueness of maximal C^k -atlases). Let $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C . Then there exists a unique maximal C^k -atlas $\text{Atlas}_{\max}^{C^k}(\mathbf{A}, E, C)$ of E in the coordinate space C compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all C^k -atlases of E in the coordinate space C containing $\mathbf{A}$ and C^k compatible in the coordinate space C with all of the C^k charts in $\mathbf{A}$. Let $\mathbf{P}$ be a maximal chain of $\mathbf{A}$. Then $\mathbf{A}' = \bigcup_{\max} \mathbf{P}$ is a maximal C^k atlas of E in the coordinate space C C^k compatible with $\mathbf{A}$. Uniqueness follows from lemma 17.5 (Extensions of C^k -atlases) on page 122. $\square$

{Ck-atl:extensions}

{def:Ck-ATLsets}

Definition 17.9 (Sets of C^k -atlases). Let E be a topological space and C be a linear space.

$$\text{Att}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \begin{array}{c} \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \\ \text{(full,S-max,max,S-max-full,max-full)} \end{array} \right\} \quad (17.7)$$

Let E be a set of topological spaces and C a set of linear spaces. Then

$$\text{Att}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \bigcup_{\substack{E \in E \\ C \in C(\text{full,S-max,max,S-max-full,max-full})}} \text{Att}_{\text{Ob}}^{C^k}(E, C) \quad (17.8)$$

18 C^k -atlas (near) morphisms and functors

{sec:ckmorph}

This section introduces a taxonomy for morphisms between C^k -atlases, defines categories of C^k -atlases, defines functors among them, defines functors between them and categories of m-atlases, defines inverse functors and proves some basic results.

18.1 C^k -atlas (near) morphisms

This subsection introduces the notions of C^k -atlas near morphisms and of C^k -atlas morphisms, and proves some basic results. C^k -atlas near morphisms between semi-maximal atlases will be proven to be C^k -atlas morphisms.

18.1.1 Definitions of C^k -atlas (near) morphisms

{def:Ck-ATLmorphnear}

Definition 18.1 (C^k -atlas near morphisms). Let E^i , $i = 1, 2$, be a topological space, C^i a linear space, $\mathbf{A}^i$ a C^k -atlases of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ a pair¹⁴ of functions.

¹⁴The conventional definition uses only the first of the two functions and a slightly different compatibility condition.

$\mathbf{f}$ is an E^1 - E^2 C^k near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{C^k, \text{near}}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1)$, iff

1. $f_0: E^1 \longrightarrow E^2$ is a continuous function.
2. $f_1: C^1 \longrightarrow C^2$ is a C^k function.
3. for any $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, diagram (11.2) in def: M-ATLmorphnear
definition 11.4 (M-atlas near morphisms for model spaces) on page 55 is C^k -
locally nearly commutative in C^2 , i.e., for any $u^1 \in I$ there are open sets
 $U'^1 \subseteq I, V'^1 \subseteq V^1, U'^2 \subseteq U^2, V'^2 \subseteq V^2, \hat{V}'^2 \subseteq C^2$ and a C^k diffeomor-
phism $\hat{f}: \hat{V}'^2 \xrightarrow{\sim} V'^2$ such that eqs. (11.3) to (11.8) on pages 56 to 56 in {eq: xin}
definition 11.1 (M-atlas morphisms for model spaces) on page 53 hold.

$\mathbf{f}$ is also a full (semi-maximal, maximal, full semi-maximal, full maximal) E^1 - E^2 C^k near morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{C^k, \text{near}}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1)$, iff

$$\text{isAtl}_{\text{Ob}}^{C^k, \text{near}}(\mathbf{A}^1, E^1, C^1) \wedge \text{isAtl}_{\text{Ob}}^{C^k, \text{near}}(\mathbf{A}^2, E^2, C^2)$$

full (S-max, max, S-max-full, max-full) full (S-max, max, S-max-full, max-full)

The identity morphism of $(\mathbf{A}^i, E^i, C^i)$ is

$$\text{Id}_{(\mathbf{A}^i, E^i, C^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) \quad (18.1)$$

This nomenclature will be justified later.

Let $E^i, i = 1, 2$, be topological spaces and C^i be linear spaces. Then

$$\begin{aligned} \text{Atl}_{\text{Ar}}^{C^k, \text{near}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \right\} \\ \text{isAtl}_{\text{Ar}}^{C^k, \text{near}}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \end{aligned} \quad (18.2)$$

(full, S-max, max, S-max-full, max-full) (full, S-max, max, S-max-full, max-full)

The identity morphism of $(\mathbf{A}^i, E^i, C^i)$ is

$$\text{Id}_{(\mathbf{A}^i, E^i, C^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) \quad (18.3)$$

This nomenclature will be justified later.

Definition 18.2 (C^k -atlas morphisms). Let $E^i, i = 1, 2$, be a topological space, C^i a linear space, $\mathbf{A}^i$ a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ a pair¹⁵ of functions.

$\mathbf{f}$ is an E^1 - E^2 C^k -morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1)$, iff

¹⁵The conventional definition uses only the first of the two functions and a slightly different compatibility condition.

1. $f_0: E^1 \longrightarrow E^2$ is a continuous function.
2. $f_1: C^1 \longrightarrow C^2$ is a C^k function.
3. for any $u^1 \in E^1$, any chart $(U^1, V^1, \phi^1: U^1 \xrightarrow{\sim} V^1) \in \mathbf{A}^1$ at u^1 and any chart $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in \mathbf{A}^2$ at $f_0(u^1)$ there exists a subchart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , an open set $U'^2 \subseteq U^2$ and a chart $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in \mathbf{A}^2$ at $f_0(u^1)$ such that $f_0[U'^1] \subseteq U'^2$ and diagram (11.1) in definition 11.1 (**M-atlas morphisms for model spaces**) on page 53 is commutative, as shown in fig. 4 (**Completed m-atlas morphism**) on page 54. {def:M-ATLmorph}
{fig:AtlM1}

$\mathbf{f}$ is also a full (semi-maximal, maximal, full semi-maximal, full maximal) E^1 - E^2 C^k morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1)$, iff

$$\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}^1, E^1, C^1) \wedge \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}^2, E^2, C^2) \\ \text{full (S-max,max,S-max-full,max-full)} \quad \text{full (S-max,max,S-max-full,max-full)}$$

The identity morphism of $(\mathbf{A}^i, E^i, C^i)$ is

$$\text{Id}_{(\mathbf{A}^i, E^i, C^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) \quad (18.4)$$

This nomenclature will be justified later.

Let $E^i, i = 1, 2$, be topological spaces and C^i be linear spaces. Then

$$\mathcal{Atl}_{\text{Ar}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (18.5)$$

$$\mathcal{Atl}_{\text{full}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{full}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (18.6)$$

$$\mathcal{Atl}_{\text{max}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{max}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (18.7)$$

$$\mathcal{Atl}_{\text{S-max}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{S-max}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (18.8)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{C}^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (18.9)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{C}^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (18.10)$$

{def:Ck-ATLmorphequiv}

Definition 18.3 (Equivalence of C^k -atlas (near) morphisms). Let E^i , $i = 1, 2$, be a topological space, C^i a linear space, A^i a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_0, g_1)$ E^1 - E^2 C^k (near) morphisms of A^1 to A^2 in the coordinate spaces C^1, C^2 .

$\mathbf{f}$ is C^k -equivalent to $\mathbf{g}$ iff $f_0 = g_0$.

18.1.2 Proclamations on C^k -atlas (near) morphisms

{lem:Ck-ATLmorph}

Lemma 18.4 (C^k -atlas (near) morphisms). Let $\mathbf{E}$ be a set of topological spaces, $\mathbf{C}$ a set of C^k linear spaces, $E^i \in \mathbf{E}$, $i = 1, 2$, $C^i \in \mathbf{C}$, A^i a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$ a pair of functions.

$\mathbf{f}$ is an E^1 - E^2 C^k (near) morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff $\mathbf{f}$ is a strict $\mathbf{E} - \mathbf{E} - E^1 - E^2 - \mathbf{C} - \mathbf{C}$ m-atlas (near) morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. The model neighborhoods of $\mathbf{C}^i$ are the open sets of $\mathbf{C}^i$, $\mathbf{E} \stackrel{\text{full-cat}}{\subseteq} \mathbf{E}$, the morphisms of $\mathbf{E}$ are the continuous functions between spaces in $\mathbf{E}$, $\mathbf{C} \stackrel{\text{full-cat}}{\subseteq} \mathbf{C}$, and the morphisms of $\mathbf{C}$ are the C^k functions between spaces in $\mathbf{C}$. $\square$

$\mathbf{f}$ is an E^1 - E^2 C^k -morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff $\mathbf{f}$ is a strict E^1 - E^2 $\mathbf{C} - \mathbf{C}$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. The model neighborhoods of $\mathbf{C}^i$ are the open sets of $\mathbf{C}^i$, $\mathbf{E} \stackrel{\text{full-cat}}{\subseteq} \mathbf{E}$, $\mathbf{C} \stackrel{\text{full-cat}}{\subseteq} \mathbf{C}$, and the morphisms of $\mathbf{C}$ are the C^k functions between spaces in $\mathbf{C}$. $\square$

Corollary 18.5 (C^k -atlas (near) morphisms). Let $E^i, i = 1, 2$, be a topological space, C^i a linear space, A^i a semi-maximal C^k -atlas of E^i in the coordinate space C^i and $f \stackrel{\text{def}}{=} (f_0, f_1)$ an E^1 - E^2 C^k near morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then f is an E^1 - E^2 C^k morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{cor:Ck-ATLmorph}

Proof. The result follows from lemma 11.12 (M-atlas (near) morphisms) on page 63

{lem:M-ATLmorph}

Let $E^i, i = 1, 2, 3$, be a topological space, C^i a linear space, A^i a C^k -atlas of E^i in the coordinate space C^i and (f_0^i, f_1^i) an E^i - E^{i+1} C^k -morphism of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} . Then $(f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a E^1 - E^3 C^k -morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof. The result follows from item 2 of lemma 11.16 (Composition of m-atlas (near) morphisms) on page 75.

{lem:M-ATLmorphcomp}

{M-morphcomp:morphcompmorph}

Lemma 18.6 (Composition of equivalent C^k -atlas (near) morphisms). Let $E^i, i = 1, 2, 3$, be a topological space, C^i be a C^k linear space, A^i be a C^k -atlas of E^i in the coordinate space C^i and $f^i \stackrel{\text{def}}{=} (f_0^i: E^i \rightarrow E^{i+1}, f_1^i: C^i \rightarrow C^{i+1})$ and $g^i \stackrel{\text{def}}{=} (g_0^i: E^i \rightarrow E^{i+1}, g_1^i: C^i \rightarrow C^{i+1})$ be C^k -equivalent E^i - E^{i+1} m-atlas (near) morphisms of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} .

{lem:Ck-ATLmorphequiv}

Then $f^2 \circ f^1$ is C^k -equivalent to $g^2 \circ g^1$.

Proof. $f_0^1 = g_0^1$ and $f_1^1 = g_1^1$, hence $f_0^2 \circ f_1^1 = g_0^2 \circ g_1^1$.

□

18.2 Categories of C^k atlases and functors

18.2.1 Categories of C^k atlases

This subsection defines categories of C^k -atlases with C^k -atlas morphisms as morphisms. It does not define categories of C^k -atlases with C^k -atlas near morphisms as morphisms because the composition of two C^k -atlas near morphisms is not in general a C^k -atlas near morphism, and requiring the atlases to be semi-maximal would cause all C^k -atlas near morphisms to be C^k -atlas morphisms.

{def:Ck-ATLcat}

Definition 18.7 (Categories of C^k atlases). Let E be a set of topological spaces and C a set of linear spaces. Let $P \stackrel{\text{def}}{=} E \times C$. Then

$$\mathcal{Att}_{\text{Ar}}^{C^k}(E, C) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^\mu, C^\mu) \in P \\ (E^\nu, C^\nu) \in P}} \mathcal{Att}_{\text{Ar}}^{C^k}(E^\mu, C^\mu, E^\nu, C^\nu) \quad (18.11)$$

$$\mathcal{Att}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob}}^{C^k}(E, C), \mathcal{Att}_{\text{Ar}}^{C^k}(E, C), \circ^A \right) \quad (18.12)$$

Remark 18.8. It is pointless to define categories of near morphisms, since

$$\text{is}\mathcal{A}tl_{\text{Ar}}^{\text{C}^k, \text{near}}(E^\mu, C^\mu, E^\nu, C^\nu) \iff \text{is}\mathcal{A}tl_{\text{Ar}}^{\text{C}^k}(E^\mu, C^\mu, E^\nu, C^\nu).$$

{lem:Ck-ATLiscat}

Lemma 18.9 ($\mathcal{A}tl^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ is a category). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then each of $\mathcal{A}tl_{(\text{full}, \text{S-max}, \text{max}, \text{S-max-full}, \text{max-full})}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ is a category.

Let $(A^i, E^i, C^i) \in \mathcal{A}tl_{\text{Ob}}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$. Then $\text{Id}_{(A^i, E^i, C^i)}$ is the identity morphism for (A^i, E^i, C^i) .

Proof. Let $(A^i, E^i, C^i), i \in [1, 3]$, be an object of $\mathcal{A}tl^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$, $i = 1, 2$, be a morphism of $\mathcal{A}tl^{\text{C}^k}(\mathbf{E}, \mathbf{C})$. Then

Composition: $((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3))$ is a morphism of

$\mathcal{A}tl^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ by corollary 18.5 (**C^k-atlas (near) morphisms**) on page 127.

{cor:Ck-ATLmorph}

Associativity: Composition is associative by lemma 1.19 (**Tuple composition for labeled morphisms**) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

18.2.2 C^k atlas functors

{def:Func-Ck-M}

Definition 18.10 (Functors from C^k atlases to m-atlases). Let $E^i, i = 1, 2$, be a topological space, C^i a linear space, A^i C^k-atlases of E^i in the coordinate space C^i , $f_0: E^1 \longrightarrow E^2$ continuous and $f_1: C^1 \longrightarrow C^2$ C^k. Then

$$\mathcal{F}_{\text{C}^k, \text{M}}^{\text{C}^k}(A^i, E^i, C^i) \stackrel{\text{def}}{=} \left(A^i, E^i, C^i \right)_{\text{triv } \text{C}^k\text{-triv}} \quad (18.13)$$

$$\mathcal{F}_{\text{C}^k, \text{M}}^{\text{C}^k}((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \stackrel{\text{def}}{=} \left((f_0, f_1), \left(A^1, E^1, C^1 \right)_{\text{triv } \text{C}^k\text{-triv}}, \left(A^2, E^2, C^2 \right)_{\text{triv } \text{C}^k\text{-triv}} \right) \quad (18.14)$$

{the:Func-Ck-M}

Theorem 18.11 (Functors from C^k atlases to m-atlases). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then $\mathcal{F}_{\text{C}^k, \text{M}}^{\text{C}^k}$ is a functor from $\mathcal{A}tl^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ to

$$\mathcal{A}tl_{\text{triv } \text{C}^k\text{-triv}}(\mathbf{E}, \mathbf{C})$$

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $i \in [1, 3]$, be an object of $\mathcal{At}\ell^{\text{Ck}}(E, C)$,
 $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i = (A^i, E^i, \underset{\text{triv Ck-triv}}{C^i})$, $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$, $i = 1, 2$, be a mor-
 phism from o^i to o^{i+1} and $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} m^i = ((f_0^i, f_1^i).o'^i, o'^{i+1})$, be the corre-
 sponding morphism in $\mathcal{At}\ell(\underset{\text{triv Ck-triv}}{E}, \underset{\text{triv Ck-triv}}{C})$.

Preservation of endpoints:

$$\begin{aligned} \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i &= (A^i, E^i, \underset{\text{triv Ck-triv}}{C^i}) \\ &= o'^i \\ \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} m^i &= m'^i \\ &= ((f_0^i, f_1^i), o'^i, o'^{i+1}) \end{aligned}$$

$\mathcal{F}_{\text{Ck}, M}^{\text{Ck}} m^i$ is a morphism from $\mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i$ to $\mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^{i+1}$:

Identity:

$$\begin{aligned} \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} \text{Id}_{o^i} &= \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} ((\text{Id}_{E^1}, \text{Id}_{C^1}), o^i, o^i) \\ &= ((\text{Id}_{\underset{\text{triv}}{E^i}}, \text{Id}_{C^i}), o'^i, o'^i) \\ &= ((\text{Id}_{\underset{\text{triv}}{E^i}}, \text{Id}_{C^i}), \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i, \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i) \\ &= \text{Id}_{\mathcal{F}_{\text{Ck}, M}^{\text{Ck}} o^i} \end{aligned}$$

Composition:

$$\begin{aligned} m^2 \circ^A m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} (m^2 \circ^A m^1) &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o'^1, o'^3) \\ &= ((f_0^2, f_1^2), o'^2, o'^3) \circ^A ((f_0^1, f_1^1), o'^1, o'^2) \\ &= \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} ((f_0^2, f_1^2), o^2, o^3) \circ^A \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} ((f_0^1, f_1^1), o^1, o^2) \\ &= \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} m^2 \circ^A \mathcal{F}_{\text{Ck}, M}^{\text{Ck}} m^1 \end{aligned}$$

□

{def:Func-M-Ck}

Definition 18.12 (Functors from m-atlases to Ck atlases). Let E^i , $i = 1, 2$, be model spaces, C^i be linear model spaces, A^i a maximal m-atlas of E^i in the coordinate space C^i and (f_0, f_1) an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , Then

$$\mathcal{F}_{M, \text{Ck}}^{\text{Ck}}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A^i, \pi_1(E^i), \pi_1(C^i)) \quad (18.15)$$

$$\begin{aligned} \mathcal{F}_{M, C^k}^{C^k}((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) &\stackrel{\text{def}}{=} \\ &\left((f_0, f_1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^2, \pi_1(E^2), \pi_1(C^2))\right) \end{aligned} \quad (18.16)$$

{the:Func-M-Ck}

Theorem 18.13 (Functors from m-atlases to C^k atlases). Let E be a set of model spaces and C a set of C^k linear model spaces. Then $\mathcal{F}_{M, C^k}^{C^k}$ is a functor from $\mathcal{Atl}(E, C)$ to $\mathcal{Atl}^{C^k}(\pi_1[E], \pi_1[C])$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, (E^i, \mathcal{E}^i), (C^i, \mathcal{C}^i))$, $i \in [1, 3]$, be an object of $\mathcal{Atl}^{C^k}(E, C)$ and

$m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i).o^i, o^{i+1})$, $i = 1, 2$, be a morphism from o^i to o^{i+1} .

$\mathcal{F}_{M, C^k}^{C^k}(m^1)$ is a morphism from $\mathcal{F}_{M, C^k}^{C^k} o^1$ to $\mathcal{F}_{M, C^k}^{C^k} o^2$:

$$\begin{aligned} \mathcal{F}_{M, C^k}^{C^k}(m^1) &= \\ \mathcal{F}_{M, C^k}^{C^k}((f_0^1, f_1^1), (A^1, E^1, C^1), (A^2, E^2, C^2)) &= \\ ((f_0^1, f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^2, \pi_1(E^2), \pi_1(C^2))) \end{aligned}$$

$\mathcal{F}_{M, C^k}^{C^k}$ maps identity functions to identity functions:

$$\begin{aligned} \mathcal{F}_{M, C^k}^{C^k} \text{Id}_{(A^i, E^i, C^i)} &= \\ \mathcal{F}_{M, C^k}^{C^k}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) &= \\ ((\text{Id}_{\pi_1(E^i)}, \text{Id}_{\pi_1(C^i)}), (A^i, \pi_1(E^i), \pi_1(C^i)), (A^i, \pi_1(E^i), \pi_1(C^i))) &= \\ ((\text{Id}_{E^i}, \text{Id}_{C^i}), \mathcal{F}_{M, C^k}^{C^k}(A^i, E^i, C^i), \mathcal{F}_{M, C^k}^{C^k}(A^i, E^i, C^i)) &= \\ \text{Id}_{\mathcal{F}_{M, C^k}^{C^k}(A^i, E^i, C^i)} \end{aligned}$$

$$\mathcal{F}_{M, C^k}^{C^k} m^2 \circ^A \mathcal{F}_{M, C^k}^{C^k} m^1 = \mathcal{F}_{M, C^k}^{C^k} (m^2 \circ^A m^1):$$

$$\begin{aligned} m^2 \circ^A m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3)) \\ \mathcal{F}_{M, C^k}^{C^k}((A^i, E^i, C^i)) &= (A^i, \pi_1(E^i), \pi_1(C^i)) \\ \mathcal{F}_{M, C^k}^{C^k}(m^i) &= ((f_0^i, f_1^i), (A^i, \pi_1(E^i), \pi_1(C^i)), (A^{i+1}, \pi_1(E^{i+1}), \pi_1(C^{i+1}))) \\ \mathcal{F}_{M, C^k}^{C^k} m^2 \circ^A \mathcal{F}_{M, C^k}^{C^k} m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^3, \pi_1(E^3), \pi_1(C^3))) \\ \mathcal{F}_{M, C^k}^{C^k}(m^2 \circ m^1) &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^3, \pi_1(E^3), \pi_1(C^3))) \end{aligned}$$

□

19 Associated model spaces and functors

$\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k}$ is an obvious functor of $\mathbf{C}^k$ -atlases to $\mathbf{m}$ -atlases, but $\mathbf{E}$ and $\mathbf{C}$ may have more model neighborhoods or more morphisms than needed for consistency with the atlas. There exist, however, model spaces with the minimum model neighborhoods and morphisms needed.

Definition 19.1 (Coordinate model spaces associated with $\mathbf{C}^k$ -atlases). Let $\mathbf{A}^i$, $i = 1, 2$, be a $\mathbf{C}^k$ -atlas of E^i in the coordinate space C^i , $f_0: E^1 \rightarrow E^2$ a continuous function and $f_1: C^1 \rightarrow C^2$ a $\mathbf{C}^k$ function. Then

$$\mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^i, E^i, C^i) \stackrel{\text{def}}{=} \text{Mod}_{\min} \left(C^i, \pi_2[\mathbf{A}^i], \left\{ \phi' \circ \phi^{-1} \mid \left(\exists_{\substack{(U, V, \phi) \in \mathbf{A}^i \\ (U', V', \phi') \in \mathbf{A}^i}} \right) U \cap U' \neq \emptyset \right\} \right) \quad (19.1)$$

$$\mathcal{F}_2^{\mathbf{C}^k, \min}((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \stackrel{\text{def}}{=} f_1: \mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^1, E^1, C^1) \rightarrow \mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^2, E^2, C^2) \quad (19.2)$$

The minimal coordinate $\mathbf{C}^k$ model space with neighborhoods in the $\mathbf{C}^k$ -atlas $\mathbf{A}^i$ of E^i in the coordinate space C^i is $\mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^i, E^i, C^i)$.

The coordinate mapping associated with the $\mathbf{E}^1$ - $\mathbf{E}^2$ $\mathbf{C}^k$ -atlas morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $f_1: \mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^1, E^1, C^1) \rightarrow \mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^2, E^2, C^2)$. If it is a model function then it is also the coordinate $\mathbf{m}$ -atlas morphism associated with the $\mathbf{E}^1$ - $\mathbf{E}^2$ $\mathbf{C}^k$ -atlas morphism (f_0, f_1) of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Lemma 19.2 (Coordinate model spaces associated with $\mathbf{C}^k$ -atlases). Let $\mathbf{A}^i$ be a $\mathbf{C}^k$ -atlas of E^i in the coordinate space C^i , $i = 1, 2$. Then $\mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^i, E^i, C^i)$ is a $\mathbf{C}^k$ linear model space.

Proof. $\mathcal{F}_2^{\mathbf{C}^k, \min}(\mathbf{A}^i, E^i, C^i)$ satisfies the conditions for a model space.

1. Since $\pi_2[\mathbf{A}^i]$ is an open cover of $\bigcup \pi_2[\mathbf{A}^i]$, the set of finite intersections is also an open cover.
2. Finite intersections of finite intersections are finite intersections
3. Restrictions of continuous functions are continuous

4. If $f: A \longrightarrow B$ is a morphism of $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$, A, A', B, B' model neighborhoods of $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$, $A' \subseteq A$, $B' \subseteq B$ and $f[A'] \subseteq B'$ then since $f: A \longrightarrow B$ is a morphism it is a restriction of a transition function between its restrictions to sets in $\pi_2[\mathbf{A}^i]$ and its restrictions are also, hence morphisms, and thus $f|_{A'}: A' \longrightarrow B'$ is a morphism.
5. If $(U, V, \phi) \in \mathbf{A}^i$ then $\text{Id}_V = \phi \circ \phi^{-1}A$ is a transition function and hence a morphism of $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$. If A, A' are objects of $\pi_2(\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i))$ and $A' \subseteq A$ then the inclusion map $i: A' \hookrightarrow A$ is a restriction of an identity morphism of $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$ and hence a morphism.
6. Restricted sheaf condition: let

- (a) $U_\alpha, V_\alpha, \alpha < A$, be an object of $\pi_2(\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i))$
- (b) $f_\alpha: U_\alpha \longrightarrow V_\alpha$ be a morphism of $\pi_2(\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i))$
- (c) $U \stackrel{\text{def}}{=} \bigcup_{\alpha < A} U_\alpha$
- (d) $V \stackrel{\text{def}}{=} \bigcup_{\alpha < A} V_\alpha$
- (e) $f: U \longrightarrow V$ be continuous and $\left(\bigvee_{\substack{\alpha < A \\ x \in U_\alpha}} \right) f(x) = f_\alpha(x)$

Then f is C^k and hence a morphism of $\pi_2(\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i))$.

□

$\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$ is non-degenerate iff $\mathbf{A}^i$ is non-degenerate.

Proof. If $\mathbf{A}^i$ is non-degenerate then by definition 17.1 (**C^k-atlases**) on page 121 there is at least one chart $(U, V, \phi) \in \mathbf{A}$ such that V contains a ball of the underlying Banach or Fréchet space. $U \subseteq \bigcup \pi_2[\mathbf{A}^i]$, so by definition 19.1 (**Coordinate model spaces associated with C^k-atlases**) on page 131 that ball is contained in $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$. {def:Ck-atlas}

Conversly, if $\mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$ is non-degenerate then by definition 14.1 (**Linear spaces**) on page 115 it contains a ball B at $v \in \mathcal{F}_2^{\text{C}^k, \text{min}}(\mathbf{A}^i, E^i, C^i)$ with radius r . By definition 19.1 there is a chart $(U, V, \phi) \in \mathbf{A}^i$ at v . Since V is open in C^i , there is a ball B' at v with radius $r' > 0$ such that $B' \cap C^i \subseteq V$. Then the ball at v with radius $\min(r, r')$ is contained in V . {def:lin}
{def:Ck-modC}

□

Let $A^i, i = 1, 2$, be a C^k -atlas of E^i in the coordinate space C^i , $f_0: E^1 \longrightarrow E^2$ a continuous function, $f_1: C^1 \longrightarrow C^2$ a C^k function and $f \stackrel{\text{def}}{=} (f_0, f_1)$ either a semi-maximal C^k -atlas near morphism or a C^k -atlas morphism from A^1 to A^2 . Then $f_1: \mathcal{F}_2^{\min, C^k}(A^1, E^1, C^1) \longrightarrow \mathcal{F}_2^{\min, C^k}(A^2, E^2, C^2)$ is well defined, i.e., $f_1 \left[\mathcal{F}_2^{\min, C^k}(A^1, E^1, C^1) \right] \subseteq \mathcal{F}_2^{\min, C^k}(A^2, E^2, C^2)$.

Proof. f is a morphism either by corollary 18.5 (C^k -atlas (near) morphisms) on page 127 or by hypothesis. By definition 18.2 (C^k -atlas morphisms) on page 125

{cor:Ck-ATLmorph}
{def:Ck-ATLmorph}

there exists a subchart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in A^1$ at u^1 , an open set $U'^2 \subseteq U^2$

and a chart $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in A^2$ at u^2 such that $f_0[U'^1] \subseteq U'^2$ and diagram (11.1) in definition 11.1 (M-atlas morphisms for model spaces) on page 53 is commutative, as shown in fig. 4 (Completed m-atlas morphism) on page 54. Then $f_1(v^1) \in \hat{V}'^2$. $\square$

{def:M-ATLmorph}
{fig:AtLM1}
{def:Ck-modE}

Definition 19.3 (Model spaces associated with C^k -atlases). Let $A^i, i = 1, 2$, be a C^k -atlas of E^i in the coordinate space C^i and $f \stackrel{\text{def}}{=} (f_0, f_1)$ an E^1 - E^2 C^k -atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then:

The minimal C^k model space with neighborhoods in the C^k -atlas A^i of E^i in the coordinate space C^i is

$$\mathcal{F}_1^{\min, C^k}(A^i, E^i, C^i) \stackrel{\text{def}}{=} \text{Mod}_{\min} \left(E^i, \pi_1[A^i], \left\{ \phi'^{-1} \circ \phi \mid \left(\exists_{\substack{(U, V, \phi) \in A^i \\ (U', V', \phi') \in A^i}} U \cap U' \neq \emptyset \right) \right\} \right) \quad (19.3)$$

The mapping associated with the E^1 - E^2 C^k -atlas morphism (f_0, f_1) of A^1 to A^2 in the coordinate spaces C^1, C^2 is

$$\mathcal{F}_1^{\min, C^k}((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \stackrel{\text{def}}{=} f_0: \mathcal{F}_1^{\min, C^k}(A^1, E^1, C^1) \longrightarrow \mathcal{F}_1^{\min, C^k}(A^2, E^2, C^2) \quad (19.4)$$

If it is a model function then it is also the m-atlas morphism associated with the E^1 - E^2 C^k -atlas morphism (f_0, f_1) of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{lem:Ck-ismodE}

Lemma 19.4 (Model spaces associated with C^k -atlases). Let A be a C^k -atlas of E in the coordinate space C . Then $\mathcal{F}_1^{\min, C^k}(A, E, C)$ is a model space.

Proof. The result follows from Lemma 5.4 (Minimal model spaces are model spaces) on page 26. $\square$

Theorem 19.5 (Functors from C^k atlases to model spaces). Let E be a set of topological spaces and C a set of linear spaces. Then $\mathcal{F}_1^{C^k}$ is a functor from $\mathcal{Atl}^{C^k}(E, C)$ to $\mathcal{E}_{\mathcal{T}riv}$, $\mathcal{F}_1^{C^k}$ is a functor from $\mathcal{Atl}_{full}^{C^k}(E, C)$ to $\mathcal{E}_{\mathcal{T}riv}$, $\mathcal{F}_2^{C^k}$ is a functor from $\mathcal{Atl}_{S-max}^{C^k}(E, C)$ to $\mathcal{C}_{C^k-op-\mathcal{T}riv}$ and $\mathcal{F}_2^{C^k}$ is a functor from $\mathcal{Atl}_{full}^{C^k}(E, C)$ to $\mathcal{C}_{C^k-triv}$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $i \in [1, 3]$, be objects in $\mathcal{Atl}^{C^k}(E, C)$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$ be a morphism in $\mathcal{Atl}^{C^k}(E, C)$.

$$\mathcal{F}_1^{C^k}: \mathcal{Atl}^{C^k}(E, C) \longrightarrow \mathcal{E}_{\mathcal{T}riv}$$

Preservation of endpoints: $\mathcal{F}_1^{C^k}(m^i) = f_0^i: \mathcal{F}_1^{C^k} o^i \longrightarrow \mathcal{F}_1^{C^k} o^{i+1}$

Composition:

$$\begin{aligned} \mathcal{F}_1^{C^k}(m^2 \circ^A m^1) &= \mathcal{F}_1^{C^k}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)(A^1, E^1, C^1), (A^3, E^3, C^3)) \\ &= f_0^2 \circ f_0^1: \mathcal{F}_1^{C^k} o^1 \longrightarrow \mathcal{F}_1^{C^k} o^3 \\ &= (f_0^2: \mathcal{F}_1^{C^k} o^2 \longrightarrow \mathcal{F}_1^{C^k} o^3) \circ (f_0^1: \mathcal{F}_1^{C^k} o^1 \longrightarrow \mathcal{F}_1^{C^k} o^2) \\ &= \mathcal{F}_1^{C^k}((f_0^2, f_1^2)(A^2, E^2, C^2), (A^3, E^3, C^3)) \circ \\ &\quad \mathcal{F}_1^{C^k}((f_0^1, f_1^1)(A^1, E^1, C^1), (A^2, E^2, C^2)) \\ &= \mathcal{F}_1^{C^k}(m^2) \circ \mathcal{F}_1^{C^k}(m^1) \end{aligned}$$

Identity:

$$\begin{aligned} 1. \mathcal{F}_1^{C^k}(\text{Id}_{o^i}) &= \mathcal{F}_1^{C^k}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) = \\ &\quad \text{Id}_{E^i}: \mathcal{F}_1^{C^k} o^i \longrightarrow \mathcal{F}_1^{C^k} o^i \\ 2. \text{Id}_{\mathcal{F}_1^{C^k}} o^i &= \text{Id}_{E^i}: \mathcal{F}_1^{C^k} o^i \longrightarrow \mathcal{F}_1^{C^k} o^i \end{aligned}$$

The proof for $\mathcal{F}_1^{C^k}: \mathcal{Atl}_{full}^{C^k}(E, C) \longrightarrow \mathcal{E}_{\mathcal{T}riv}$ is identical.

$$\mathcal{F}_2^{C^k}: \mathcal{Atl}^{C^k}(E, C) \longrightarrow \mathcal{C}_{C^k-op-\mathcal{T}riv}$$

Preservation of endpoints:

$$\mathcal{F}_2^{\min}(m^i) = f_1^i: \mathcal{F}_2^{\min} o^i \longrightarrow \mathcal{F}_2^{\min} o^{i+1}$$

$$\begin{aligned} \mathcal{F}(g \circ f) &= \mathcal{F}(g) \circ \mathcal{F}(f): \mathcal{F}_2^{\min}(m^2 \circ m^1) = \\ &= \mathcal{F}_2^{\min}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)(A^1, E^1, C^1), (A^3, E^3, C^3)) = \\ &= f_0^2 \circ f_0^1: \mathcal{F}_2^{\min} o^1 \longrightarrow \mathcal{F}_2^{\min} o^3 = \\ &= (f_1^2: \mathcal{F}_2^{\min} o^2 \longrightarrow \mathcal{F}_2^{\min} o^3) \circ (f_1^1: \mathcal{F}_2^{\min} o^1 \longrightarrow \mathcal{F}_2^{\min} o^2) = \\ &= \mathcal{F}_2^{\min}((f_0^2, f_1^2)(A^2, E^2, C^2), (A^3, E^3, C^3)) \circ \\ &= \mathcal{F}_2^{\min}((f_0^1, f_1^1)(A^1, E^1, C^1), (A^2, E^2, C^2)) = \\ &= \mathcal{F}_2^{\min}(m^2) \circ \mathcal{F}_2^{\min}(m^1) \end{aligned}$$

$$\mathcal{F}(\text{Id}_A) = \text{Id}_{\mathcal{F}(A)}:$$

$$\begin{aligned} 1. \quad \mathcal{F}_2^{\min}(\text{Id}_{o^i}) &= \mathcal{F}_2^{\min}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) = \\ &= \text{Id}_{C^i}: \mathcal{F}_2^{\min} o^i \longrightarrow \mathcal{F}_2^{\min} o^i \\ 2. \quad \text{Id}_{\mathcal{F}_2^{\min} o^i} &= \text{Id}_{C^i}: \mathcal{F}_2^{\min} o^i \longrightarrow \mathcal{F}_2^{\min} o^i \end{aligned}$$

The proof for $\mathcal{F}_2^{\min}: \mathcal{A}t\ell_{\text{full}}^{C^k}(E, C) \longrightarrow C_{\text{ck-triv}}$ is identical. $\square$

20 Classic C^k -atlas morphisms and functors

{sec:Ck-ATLcmorph}

This section introduces an alternate definition of and taxonomy for morphisms between C^k -atlases, defines categories of C^k -atlases, defines functors among them and proves some basic results. It introduces the notion of classic C^k -atlas morphisms, which is equivalent to the conventional definitions for a manifold.

20.1 Classic C^k -atlas morphisms

{sub:Ck-ATLcmorph}

This subsection introduces the notion of classic C^k -atlas morphisms, and proves some basic results.

20.1.1 Definitions of classic C^k -atlas morphisms

{def:Ck-ATLcmorph}

Definition 20.1 (Classic C^k -atlas morphisms). Let $E^i, i = 1, 2$, be a topological space, C^i be a linear space, A^i be a C^k -atlases of E^i in the coordinate space C^i and f be a continuous function.

f is a E^1 - E^2 classic C^k morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}, C^k}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff for any $(U^i, V^i, \phi^i: U^i \xrightarrow{\cong} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset, \phi^2 \circ f \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a C^k function.

Remark 20.2. Definitions of constrained, semistrict and strict classic C^k -atlas morphisms would be pointless, as any classic C^k -atlas morphism would be constrained and strict.

20.1.2 Proclamations on classic C^k -atlas morphisms

{lem:Ck-ATLcmorph}

Lemma 20.3 (Classic C^k -atlas morphisms). Let E be a set of topological spaces, C a set of C^k linear spaces, $E^i \in E, i = 1, 2, C^i \in C, A^i$ a C^k -atlas of E^i in the coordinate space C^i and $f: E^1 \longrightarrow E^2$ a function.

f is an E^1 - E^2 classic C^k -atlas near morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff f is an E^1 - E^2 - $\begin{smallmatrix} C \\ C^k\text{-op-}\mathcal{T}riv \end{smallmatrix}$ - $\begin{smallmatrix} C \\ C^k\text{-op-}\mathcal{T}riv \end{smallmatrix}$ classic m-atlas near morphism of A^1 to A^2 in the coordinate spaces $\begin{smallmatrix} C^1 \\ C^k\text{-triv} \end{smallmatrix}, \begin{smallmatrix} C^2 \\ C^k\text{-triv} \end{smallmatrix}$.

Let E^1 be an open subspace of E^2 . Then f is an E^1 - E^2 classic C^k -atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff f is a strict E^1 - E^2 classic m-atlas morphism of A^1 to A^2 in the coordinate spaces $\begin{smallmatrix} C^1 \\ C^k\text{-triv} \end{smallmatrix}, \begin{smallmatrix} C^2 \\ C^k\text{-triv} \end{smallmatrix}$.

f is an E^1 - E^2 classic C^k -atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff f is a strict E^1 - E^2 - $\begin{smallmatrix} C \\ C^k\text{-op-}\mathcal{T}riv \end{smallmatrix}$ - $\begin{smallmatrix} C \\ C^k\text{-op-}\mathcal{T}riv \end{smallmatrix}$ classic m-atlas morphism of A^1 to A^2 in the coordinate spaces $\begin{smallmatrix} C^1 \\ C^k\text{-triv} \end{smallmatrix}, \begin{smallmatrix} C^2 \\ C^k\text{-triv} \end{smallmatrix}$.

21 C^k manifolds

{sec:ckman}

Conventionally a manifold is different from its atlases, but $\mathcal{Att}_{\text{max}}^{C^k}(E, C)$ in definition 18.7 (Categories of C^k atlases) on page 128 encourages treating them on an equal footing. All of the results for maximal atlases carry directly over to results for manifolds.

{def:Ck-ATLcat}

{def:man}

Definition 21.1 (C^k manifolds). Let E be a topological space, C a linear space and A a maximal¹⁶ C^k -atlas of E in the coordinate space C . Then (E, C, A) is a C^k manifold.

Let E be a set of topological spaces and C be a set of linear spaces. Then

$$\text{Man}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{max}}^{C^k}(E, C) \quad (21.1)$$

Remark 21.2. The manifold (E, C, A) corresponds to the object (A, E, C) .

{def:manmorph}

Definition 21.3 (C^k manifold morphisms). Let $S^i \stackrel{\text{def}}{=} (E^i, C^i, A^i), i = 1, 2$, be C^k manifolds and $(f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ be an E^1 - E^2 C^k -morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then (f_0, f_1) is a C^k morphism of S^1 to S^2 .

¹⁶Requiring that the atlas be full would eliminate some pathologies.

Let $E^i, i = 1, 2$, be a topological space and C^i be a linear space. Then

$$\text{Man}_{\text{Ar}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \mathcal{A}t\ell_{\text{Ar}}^{C^k}(E^1, C^1, E^2, C^2) \quad (21.2)$$

Let E be a set of topological spaces and C a set of linear spaces. Then

$$\text{Man}_{\text{Ar}}^{C^k}(E, C) \stackrel{\text{def}}{=} \mathcal{A}t\ell_{\text{Ar}}^{C^k}(E, C) \quad (21.3)$$

$$\text{Man}^{C^k}(E, C) \stackrel{\text{def}}{=} \mathcal{A}t\ell_{\text{max}}^{C^k}(E, C) \quad (21.4)$$

Theorem 21.4 (Categories of C^k manifolds). Let E be a set of topological spaces and C a set of linear spaces. Then $\text{Man}^{C^k}(E, C)$ is a category and the identity morphism of (A, E, C) is an identity morphism.

Proof. The result follows directly from $\stackrel{\text{def:man}, \text{def:manmorph}}{\text{definitions 21.1 and 21.3}}$ above and $\stackrel{\text{lem: Ck-ATLiscat}}{\text{lemma 18.9}}$ ($\mathcal{A}t\ell^{C^k}(E, C)$ is a category) on page 128. $\square$

Part VIII

Equivalence of fiber bundles

{part:bun}

For fiber bundles¹⁷, the adjunct spaces are the base space $X = (X, \mathcal{X})$, the fiber $Y = (Y, \mathcal{Y})$ and the group G ; the category of the coordinate space is the category of Cartesian products $\{U \times Y \mid U \in \mathcal{X}\}$ of model neighborhoods in the base space with the entire fiber, with morphisms $t: U \times Y \longrightarrow U \times Y$ that preserve the fibers, i.e., $\pi_1 \circ t = \pi_1$, and are generated by the group action on the fiber (Equation (24.1)).

The sole adjunct functions are the projection $\pi: E \rightarrow X$, the group operation and the group action on the fiber.

This part of the paper defines bundle atlases, fiber bundles, local coordinate spaces equivalent to fiber bundles, categories of them and functors, and gives basic results.

Definition 21.5 (Trivial group category of groups). Let G be a set of topological groups. The trivial group category of G , abbreviated $\mathbf{G}_{\text{group-}\mathcal{T}riv}$, is the category of all continuous homomorphisms between groups of G . By abuse of language it will be shortened to $\mathbf{G}_{\mathcal{T}riv}$ when the meaning is clear from context.

Definition 21.6 (Group actions). Let Y be a topological space, G a topological group, ρ an effective group action of G on Y , $y \in Y$ and $g \in G$. Then $y \star g \stackrel{\text{def}}{=} \rho(y, g)$.

¹⁷The literature has several definitions of fiber bundle. This paper uses one chosen for clarity of exposition. It differs from [Steenrod, 1999, p. 8] in that, e.g., it uses the machinery of maximal atlases rather than equivalence classes of coordinate bundles, the nomenclature differs in several minor regards. {TopFib}

Let X be a topological space and $x \in X$. Then $(x, y) \star g \stackrel{\text{def}}{=} (x, y \star g)$.

A $\star$ with a subscript, superset, underset or overset refers to the group action ρ with the same subscript, superset, underset or overset.

Remark 21.7. This notation is only used when it is clear from context what the group action is.

{def:ProtoBun}

Definition 21.8 (Protobundles). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$, where E , X and Y are topological spaces, G a topological group, $\pi: E \rightarrow X$ a continuous surjection and ρ an effective group action of G on Y . Then $\mathbf{B}$ is a protobundle.

Remark 21.9. While this definition does not itself require E to have a local product structure nor require π to have the Covering Homotopy Property, only those protobundles having an atlas are of interest, and for them [definition 25.1 \(Bundle atlases\)](#) on page 148 imposes additional constraints.

{def:BunAtl}

22 G -rho model spaces

def:Grhomod {sec:grhomod}

Definition 22.1 (G - ρ -model spaces). Let $\mathbf{XY} \stackrel{\text{def}}{=} (X \times Y, \mathcal{XY})$ be a model space, G a topological group and ρ an effective group action of G on Y such that the objects of $\mathcal{XY}$ are products of open sets with Y and the morphisms are fiber-preserving automorphisms generated by the group action, i.e.,

$$\left(\bigvee_{U \in \mathcal{XY}}^{\text{Ob}} \right) \left(\exists_{V \in X}^{\text{op}} \right) U = V \times Y \quad (22.1)$$

$$\left(\bigvee_{f \in \mathcal{XY}: V \times Y \xrightarrow{\sim} V \times Y}^{\text{Ar}} \right) \left(\exists_{g \in G^V} \right) \left(\bigvee_{(x,y) \in V \times Y} \right) f(x, y) = (x, y \star g(x)) \quad (22.2) \quad \{\text{eq:fGrho}\}$$

Then $\mathbf{XY}$ is a G - ρ model space of $X \times Y$, abbreviated is $G\rho(\mathbf{XY}, Y, G, \rho)$, $G_{\mathbf{XY},f} \stackrel{\text{def}}{=} \text{range}(g)$ is the set of group elements associated with f and $G_{\mathbf{XY}} \stackrel{\text{def}}{=} \bigcup_{f \in \mathcal{XY}}^{\text{Ar}} G_{\mathbf{XY},f}$ is the set of group elements associated with $\mathbf{XY}$ morphisms.

{lem:Grhomod}

Lemma 22.2 (G - ρ -model spaces). Let $\mathbf{XY}$ be a G - ρ model space of $X \times Y$ and $f \in \mathcal{XY}: V \times Y \xrightarrow{\sim} V \times Y$.

The function g in eq. (22.2) is unique.

{eq:fGrho}

Proof. The group action is effective. $\square$

$G_{\mathbf{XY}}$ is unique.

Proof. The function g is unique. $\square$

$$\begin{array}{ccc}
((x^1, y^1), g^1) & \xrightarrow{f_C \times f_G} & ((x^2, y^2), g^2) \\
\downarrow \star^1 & & \downarrow \star^2 \\
(x^1, y^1 \star g^1) & \xrightarrow{f_C} & (x^2, y^2 \star g^2)
\end{array}$$

Figure 15: Preserving group action

{fig:Grho}

Definition 22.3 (Morphisms of G - ρ -model spaces). Let $X^i, Y^i, i = 1, 2$, be topological spaces, G^i a topological group, ρ^i an effective group action on Y^i and $\mathbf{XY}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$. Then a model function $f_C: \mathbf{XY}^1 \rightarrow \mathbf{XY}^2$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$, abbreviated is $G\rho\text{morph}(\mathbf{XY}^1, G^1, \rho^1, \mathbf{XY}^2, G^2, \rho^2, f_C)$ iff it preserves the group action, i.e., there is a continuous homomorphism $f_G: G^1 \rightarrow G^2$ such that [fig. 15 \(Preserving group action\)](#) is commutative, i.e., [eq. \(22.3\)](#) below holds.

{def:Grhomorph}

{fig:Grho}

$$\left(\exists_{f_G: G^1 \rightarrow G^2} \right) \left(\bigvee_{\substack{(x,y) \in X^1 \times Y^1 \\ g \in G^1}} f_C((x, y) \star^1 g) = f_C((x, y)) \star^2 f_G(g) \right) \stackrel{\text{eq:Grho}}{(22.3)}$$

{lem:Grhomorph}

Lemma 22.4 (Morphisms of G - ρ -model spaces). Let $X^i, Y^i, i = 1, 2, 3$, be topological spaces, G^i a topological group, ρ^i an effective group action on Y^i , $\mathbf{XY}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$ and $f_C^i: X^i \times Y^i \rightarrow X^{i+1} \times Y^{i+1}$ a G^i - G^{i+1} - ρ^i - ρ^{i+1} morphism of $X^i \times Y^i$ to $X^{i+1} \times Y^{i+1}$.

{grhomorph:fgunique}

1. The function f_G in [eq. \(22.3\)](#) is unique.

{eq:Grho}

Proof. The group action is effective. $\square$

{grhomorph:fiber}

2. f_C^i preserves fibers, i.e., $\pi_1(f_C^i(x, y)) = \pi_1(f_C^i(x, y'))$ for x in X^i and y, y' in Y^i ,

Proof. Since ρ^i is effective, there is a $g \in G^i$ such that $y' = y \star^i g$ and thus by [definition 22.3](#)

{def:Grhomorph}

$$\begin{aligned}
\pi_1(f_C^i(x, y')) &= \\
\pi_1(f_C^i(x, y \star^i g)) &= \\
\pi_1(f_C^i(x, y) \star^{i+1} f_G^i(g)) &= \\
\pi_1(f_C^i(x, y)) &
\end{aligned} \tag{22.4}$$




Figure 16: Preserving group actions

{fig:Grho2}

□

{grhomorph:fxunique}

3. There exists a unique function $f_X^i: X^i \longrightarrow X^{i+1}$ such that $f_X^i \circ \pi_1 = \pi_1 \circ f_C^i$.

Proof. Define $f_X^i(x) = \pi_1(f_C^i(x, y))$, where y is an arbitrary point of Y^i . It does not depend on the choice of y because f_C^i preserves fibers. □

Remark 22.5. f_C^i may be a twisted product: there need not exist $f_Y^i: Y^i \longrightarrow Y^{i+1}$ such that $f_C^i = f_X^i \times f_Y^i$.

{grhomorph:comp}

4. $f_C^2 \circ f_C^1$ is a G^1 - G^3 - ρ^1 - ρ^3 morphism of $X^1 \times Y^1$ to $X^3 \times Y^3$.

Proof. Let $f_G^i: G^i \longrightarrow G^{i+1}$ be a continuous homomorphism such that

$$\left(\bigvee_{\substack{(x^i, y^i) \in X^i \times Y^i \\ g^i \in G^i}} \right) f_C^i((x^i, y^i) \star g^i) = f_C^i((x^i, y^i)) \star f_G^i(g^i) \quad (22.5) \quad \{\text{eq:Grho2}\}$$

Let $(x^1, y^1) \in X^1 \times Y^1$ and $g^1 \in G^1$. Then fig. 16 (Preserving group actions) is commutative and {fig:Grho2}

$$f_C^2 \circ f_C^1((x^1, y^1) \star g^1) = \quad (22.6)$$

$$= f_C^2(f_C^1((x^1, y^1)) \star f_G^1(g^1)) \quad (22.7)$$

$$= f_C^2(f_C^1((x^1, y^1))) \star f_G^2 \circ f_G^1(g^1) = \quad (22.8)$$

$$(22.9)$$

$$= f_C^2 \circ f_C^1((x^1, y^1)) \star f_G^2 \circ f_G^1(g^1) \quad (22.10)$$

□

{def:Grhomodcat}

Definition 22.6 (Categories of G - ρ -model spaces). Let $X^\alpha, Y^\alpha, \alpha < A$, be topological spaces, G^α a topological group, ρ^α an effective group action on Y^α ,

$\mathbf{XY}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{XY}^\alpha)$ a G^α - ρ^α model space of $X^\alpha \times Y^\alpha$, $\mathcal{XY}\mathcal{E}\rho_{\text{Ob}} \stackrel{\text{def}}{=} \{\mathbf{XY}^\alpha | \alpha < A\}$, $\mathcal{XY}\mathcal{E}\rho_{\text{Ar}} \stackrel{\text{def}}{=} \{f_C: X^\alpha \times Y^\alpha \longrightarrow X^\beta \times Y^\beta |$

is $G\rho$ morph($\mathbf{XY}^\alpha, G^\alpha, \rho^\alpha, \mathbf{XY}^\beta, G^\beta, \rho^\beta, f_C) \wedge \alpha < A \wedge \beta < A\}$ and $\mathcal{XY}\mathcal{E}\rho \stackrel{\text{def}}{=} (\mathcal{XY}\mathcal{E}\rho_{\text{Ob}}, \mathcal{XY}\mathcal{E}\rho_{\text{Ar}})$. Then any subcategory of $\mathcal{XY}\mathcal{E}\rho$ is a G - ρ model category.

{def:Grhomodtriv}

Definition 22.7 (Trivial G - ρ -model spaces). Let X and Y be topological spaces, G a topological group, ρ an effective group action of G on Y and $\mathcal{XY}$ the category of all products of open subsets of X with Y and all homeomorphisms induced by the group action, i.e.,

$$\text{Ob}(\mathcal{XY}) \stackrel{\text{def}}{=} \left\{ V \times Y \mid V \in X_{\text{op}} \right\} \quad (22.11)$$

$$\text{Ar}(\mathcal{XY}) \stackrel{\text{def}}{=} \left\{ f: V \times Y \xrightarrow{\sim} V \times Y \mid \begin{array}{l} V \in X_{\text{op}} \wedge \pi_1 \circ f = \pi_1 \wedge \left(\exists_{g \in G^V} \left(\bigvee_{\substack{x \in V \\ y \in Y}} f(x, y) = (x, y \star g(x)) \right) \right) \end{array} \right\} \quad (22.12)$$

Then the trivial G - ρ model space of X, Y , abbreviated $X, Y_{G-\rho-\text{triv}}$, is $(X \times Y, \mathcal{XY})$ and

$X, Y_{G-\rho-\text{triv}}$ is a trivial G - ρ model space of X, Y .

Remark 22.8. Let G' be a topological group and ρ' an effective group action of G' on Y such that $X, Y_{G-\rho-\text{triv}} = X, Y_{G'-\rho'-\text{triv}}$. Although G' must be isomorphic to G , it need not have the same topology.

The identity morphism of $X, Y_{G-\rho-\text{triv}}$ is $\text{Id}_{X, Y_{G-\rho-\text{triv}}} \stackrel{\text{def}}{=} \text{Id}_{X \times Y}$.

{lem:Grhomodtriv}

Lemma 22.9 (The trivial G - ρ model space of X, Y is a G - ρ model space of X, Y). Let X and Y be topological spaces, G a topological group and ρ an effective group action on Y . Then $X, Y_{G-\rho-\text{triv}}$ is a G - ρ model space of $X \times Y$.

Proof. $X, Y_{G-\rho-\text{triv}}$ satisfies the conditions of definition 22.1 (G - ρ -model spaces) on page 138:

{def:Grhomod}

1. Product with fiber:

$$\left(\bigvee_{U \in \pi_2 \left(X, Y_{G-\rho-\text{triv}} \right)}^{\text{Ob}} \right) \left(\exists_{V \in X_{\text{op}}} U = V \times Y \right)$$

by definition 22.7.

{def:Grhomodtriv}

2. Generated by ρ : Let $f \in \pi_2^{\text{Ar}} \left(\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix} \right): V \times Y \xrightarrow{\sim} V \times Y$. Then

$$\left(\exists_{g \in G^V} \right) \left(\forall_{(x,y) \in V \times Y} \right) f(x,y) = (x, y \star g(x))$$

by definition 22.7.

{def:Grhomodtriv}

□

{lem:Grhomodtrivmorph}

Lemma 22.10 (Morphisms of trivial G - ρ -model spaces). Let X^i and Y^i , $i = 1, 2$, be topological spaces, G^i a topological group, ρ^i an effective group action of G^i on Y^i and $\mathbf{f} \stackrel{\text{def}}{=} f_X \times f_Y: \begin{smallmatrix} X^1, Y^1 \\ G^1-\rho^1-\text{triv} \end{smallmatrix} \longrightarrow \begin{smallmatrix} X^2, Y^2 \\ G^2-\rho^2-\text{triv} \end{smallmatrix}$ a pair of continuous functions. Then $\mathbf{f}$ is a model function from $\begin{smallmatrix} X^1, Y^1 \\ G^1-\rho^1-\text{triv} \end{smallmatrix}$ to $\begin{smallmatrix} X^2, Y^3 \\ G^2-\rho^3-\text{triv} \end{smallmatrix}$.

Proof.

1. $\mathbf{f}$ is a Cartesian product of continuous functions, hence continuous.
2. Let $m^2 \stackrel{\text{def}}{=} U^2 \times Y^2$ be a model neighborhood of $\begin{smallmatrix} X^2, Y^2 \\ G^2-\rho^2-\text{triv} \end{smallmatrix}$. Then $U^1 \stackrel{\text{def}}{=} f_X^{-1}(U^2)$ is open and $\mathbf{f}^{-1}(m^2) = U^1 \times Y^1$ is a model function.
3. Let $m^1 \stackrel{\text{def}}{=} U^1 \times Y^1$ be a model neighborhood of $\begin{smallmatrix} X^1, Y^1 \\ G^1-\rho^1-\text{triv} \end{smallmatrix}$. $X^2 \times Y^2$ is a model neighborhood of $\begin{smallmatrix} X^2, Y^2 \\ G^2-\rho^2-\text{triv} \end{smallmatrix}$ and $\mathbf{f}(m^1) \subseteq X^2 \times Y^2$.

□

{def:Grhomodtrivcat}

Definition 22.11 (Categories of trivial G - ρ -model spaces). Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles.

The trivial coordinate model category of $\mathbf{B}$, $\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}$, is the category with objects all trivial G_α - ρ_α model spaces of X_α, Y_α , $\alpha < A$ and morphisms all continuous functions compatible with the group action:

$$\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ob}}} \stackrel{\text{def}}{=} \left\{ \begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix} \mid (E, X, Y, \pi, G, \rho) \in \mathbf{B} \right\} \quad (22.13)$$

$$\begin{aligned} \mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ar}}} &\stackrel{\text{def}}{=} \left\{ f_C: X^1 \times Y^1 \longrightarrow X^2 \times Y^2 \mid \right. \\ &\left. \left(\exists_{\substack{(E^i, X^i, Y^i, \pi^i, G^i, \rho^i) \in \mathbf{B} \\ f_G: G^1 \longrightarrow G^2}} \right) \text{isG}\rho\text{morph}_{G-\rho-\text{triv}}(X^1, Y^1, G^1, \rho^1, X^2, Y^2, G^2, \rho^2, f_C) \right\} \end{aligned} \quad (22.14)$$

$$\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}} \stackrel{\text{def}}{=} \left(\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ob}}}, \mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ar}}} \right) \quad (22.15)$$

Remark 22.12. The morphisms f_C may be twisted products: there need not exist $f_Y: Y^1 \longrightarrow Y^2$ such that $f_C = f_X \times f_Y$.

The trivial product coordinate model category of $\mathbf{B}$, $\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$, is the category with objects all trivial G^α - ρ^α model spaces of $X^\alpha, Y^\alpha, \alpha < A$ and morphisms all products of continuous functions compatible with the group action:

$$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}_{\text{Ob}}} \stackrel{\text{def}}{=} \mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ob}}} \quad (22.16)$$

$$\begin{aligned} \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}_{\text{Ar}}} &\stackrel{\text{def}}{=} \left\{ f_C \stackrel{\text{def}}{=} f_X \times f_Y: X^1 \times Y^1 \longrightarrow X^2 \times Y^2 \mid \right. \\ &\left. \left(\exists_{\substack{(E^i, X^i, Y^i, \pi^i, G^i, \rho^i) \in \mathbf{B} \\ f_G: G^1 \longrightarrow G^2}} \right) \text{isG}\rho\text{morph}_{G-\rho-\text{triv}}(X^1, Y^1, G^1, \rho^1, X^2, Y^2, G^2, \rho^2, f_C) \right\} \end{aligned} \quad (22.17)$$

$$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}} \stackrel{\text{def}}{=} \left(\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}_{\text{Ob}}}, \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}_{\text{Ar}}} \right) \quad (22.18)$$

Any subcategory of $\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}$ is a trivial coordinate model category and any subcategory of $\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$ is a trivial product coordinate model category.

Lemma 22.13 (The trivial coordinate model category of $\mathbf{B}$ is a category). Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha), \alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha \mid \alpha < A\}$ be a set of protobundles.

Then $\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}$ is a category and the identity morphism for object $\mathbf{B}^\alpha$ is $\text{Id}_{X^\alpha \times Y^\alpha}$.

Proof.

Composition: The composition of G - ρ morphisms is a G - ρ morphism by lemma 22.4_{lem:Grhomorph} (Morphisms of G - ρ -model spaces) on page 139.

Associativity: Morphisms are simply functions and composition of morphisms is simply composition of functions.

Unit: The identity morphisms are simply identity functions and composition of morphisms is simply composition of functions.

□

$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$ is a category and the identity morphism for object $\mathbf{B}^\alpha$ is $\text{Id}_{X^\alpha \times Y^\alpha}$.

Proof.

Composition: The composition of G - ρ morphisms is a G - ρ morphism by lemma 22.4_{lem:Grhomorph}

(Morphisms of G - ρ -model spaces) on page 139. Let $X^i, Y^i \xrightarrow{G^i-\rho^i-\text{triv}} \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$, $i = 1, 2, 3$, $f_C^i = f_X^i \times f_Y^i: X^i \times Y^i \longrightarrow X^{i+1} \times Y^{i+1} \xrightarrow{\text{Ar}} \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$. Then $f_C^{i+1} \circ f_C^i = (f_X^{i+1} \circ f_X^i) \times (f_Y^{i+1} \circ f_Y^i)$.

Associativity: Morphisms are simply functions and composition of morphisms is simply composition of functions.

Unit: The identity morphisms are simply identity functions and composition of morphisms is simply composition of functions.

□

23 G -rho-nearly commutative diagrams

_{sec:grhoncd}

Let

1. X and Y be topological spaces
2. G be a topological group,
3. ρ an effective group action on Y
4. $\mathbf{C} = (C, \mathcal{C}) \stackrel{\text{def}}{=} \mathbf{X}, Y_{G-\rho-\text{triv}}$
5. D be, as in diagram (1.20) of 15, a tree with two branches, whose nodes are topological spaces U_i and V^j and whose links are continuous functions $f_i: U_i \longrightarrow U_{i+1}$ and $f'_j: U_j \longrightarrow U_{j+1}$ between the spaces, with $U_0 = V_0$, $U_m \subseteq C$ open and $V_n \subseteq C$ open, as shown in fig. 1 (Uncompleted nearly commutative diagram) on page 15. _{fig:NCDbb}

Definition 23.1 (G - ρ -nearly commutative diagrams). D is nearly commutative in X, Y, π, ρ iff D is nearly commutative in category $\mathcal{C}$.

_{def:grhoNCD}

Definition 23.2 (G - ρ -nearly commutative diagrams at a point). Let $\mathcal{C}$ and D be as above and x be an element of the initial node. D is nearly commutative in X, Y, π, ρ at x iff D is nearly commutative in $\mathcal{C}$ at x .

Definition 23.3 (G - ρ -locally nearly commutative diagrams). Let $\mathcal{C}$ and D be as above. D is locally nearly commutative in X, Y, π, ρ iff D is locally nearly commutative in $\mathcal{C}$.

24 Bundle charts

sec:buncharts {def:Ypichart}

Definition 24.1 (Y - π -bundle charts). Let E, X, Y be topological spaces and $\pi: E \rightarrow X$. A Y - π -bundle chart $C \stackrel{\text{def}}{=} (U, V \times Y, \phi)$ of E in the coordinate space $X \times Y$ consists of




Figure 17: ϕ preserves fibers

{fig:Ypichart}

1. a nonvoid open set $U \subseteq E$, known as a coordinate patch
2. An open set $V \times Y \subseteq X \times Y$
3. a homeomorphism $\phi: U \xrightarrow{\cong} V \times Y$, known as a coordinate function, that preserves fibers. i.e., $\pi_1 \circ \phi = \pi$.

C covers u iff $u \in U$.

{lem:Ypichart}

Lemma 24.2 (Y - π -bundle charts). Let E, X, Y be topological spaces, G a topological group, ρ an effective group action of G on Y and $\mathbf{XY} \stackrel{\text{def}}{=} (X \times Y, \mathcal{XY})$ a G - ρ model space of $X \times Y$ such that for every $e \in E$ there is a Y - π -bundle chart of E in the coordinate space $X \times Y$ in $\mathbf{A}$ that covers e .

Then every chart in $\mathbf{A}$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$.

Proof. Let $e \in E$, (U, V, ϕ) be an arbitrary Y - π -bundle chart of E in the coordinate space $X \times Y$ in $\mathbf{A}$ that covers e and (U', V', ϕ') be an arbitrary m-chart in $\mathbf{A}$ that covers e .

By definition 22.7 (Trivial G - ρ -model spaces) on page 141 the transition function is of the form $t(x, y) = (x, y \star g(x))$. Then $\pi_1 \circ t = \pi_1, \pi_1 \circ \phi' = \pi_1 \circ \phi = \pi$. $\square$

{def:Grhomodtriv}

Lemma 24.3 (Y - π -bundle charts in (X, Y)). Let X and Y be topological spaces, G

{lem:GrhomodtrivYpichart}

a topological group, ρ an effective group action of G on Y and $\pi \stackrel{\text{def}}{=} \pi_1: X \times Y \longrightarrow Y$.
 $C \stackrel{\text{def}}{=} (U, V \times Y, \phi)$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$ iff C is an m -chart of E in the coordinate space (X, Y) .

G - ρ -triv

Lemma 24.4 (Properties of projection). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $v \in V$. Then $\pi|_U$ is a surjection.

{def:piepi}

Proof. Let $v \in V$, $y \in Y$ and $u \stackrel{\text{def}}{=} \phi^{-1}((v, y)) \in U$. Then $\pi(u) = v$. $\square$

$\pi^{-1}[\{v\}]$ is homeomorphic to Y .

Proof. ϕ and ϕ^{-1} are homeomorphisms, so their restrictions are homeomorphisms and thus $\phi^{-1}[\{v\}]$ is homeomorphic to $\{v\} \times Y$, which is homeomorphic to Y . $\square$

{def:Ypibsub}

Definition 24.5 (Y - π subcharts). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and U' be a nonvoid open subset of U . Then $(U', V' \times Y, \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U'})$ is a subchart of $(U, V \times Y, \phi)$.

Lemma 24.6 (Y - π subcharts). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $(U', V' \times Y, \phi')$ a subchart of $(U, V \times Y, \phi)$. Then $(U', V' \times Y, \phi')$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$.

Proof. $(U', V' \times Y, \phi')$ satisfies the conditions of definition 24.1:

{def:Ypichart}

1. $U' \subseteq E$ is open.
2. $\phi[U']$ is open since ϕ is a homeomorphism.
3. $\phi|_{U'}: U' \longrightarrow V' \times Y$ is the restriction of a homeomorphism and thus a homeomorphism. $\phi|_{U'}$ preserves fibers because ϕ does.

$\square$

{def:Grho-compat}

Definition 24.7 (G - ρ -compatibility). Let E, X, Y be topological spaces, G a topological group, $\rho: Y \times G \longrightarrow Y$ an effective right action of G on Y , $\pi: E \twoheadrightarrow X$ surjective and $(U^\mu, V^\mu \times Y, \phi^\mu)$, $(U^\nu, V^\nu \times Y, \phi^\nu)$ Y - π -bundle charts. Then $(U^\mu, V^\mu \times Y, \phi^\mu)$ and $(U^\nu, V^\nu \times Y, \phi^\nu)$ are G - ρ -compatible iff either

1. U^μ and U^ν are disjoint
2. The transition function $t_\nu^\mu \stackrel{\text{def}}{=} \phi^\mu \circ \phi^{\nu-1}|_{\phi^\nu[U^\mu \cap U^\nu]}$ is generated by the group action, i.e., there is a continuous function $g_\nu^\mu: \pi_1[\phi^\nu[U^\mu \cap U^\nu]] \longrightarrow G$ such that

$$\left(\bigvee_{(x,y) \in \phi^\nu[U^\mu \cap U^\nu]} \right) t_\nu^\mu(x, y) = (x, y \star g_\nu^\mu(x)) \quad (24.1)$$

{lem:Grho-compat}

Lemma 24.8 (G - ρ -compatibility). Let E, X, Y be topological spaces, G a topological group, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y , $\pi: E \rightarrow X$ surjective and $(U^i, V^i \times Y, \phi^i)$, $i = 1, 2$ Y - π -bundle charts. Then $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are G - ρ -compatible iff $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are compatible m-charts of E in the coordinate space X, Y .

Proof.

Each U^i is a model neighborhood of E and each $V^i \times Y$ is a model neighborhood of X, Y .

2. If $U^1 \cap U^2 = \emptyset$ then the lemma is trivially true.
3. If the charts are G - ρ -compatible then the transition function $t_2^1 \stackrel{\text{def}}{=} \phi^1 \circ \phi^{2^{-1}}|_{\phi^2[U^1 \cap U^2]}$ is generated by the group action, i.e., there is a continuous function $g_2^1: \pi_1[\phi^2[U^1 \cap U^2]] \rightarrow G$ such that $\left(\bigvee_{(x,y) \in \phi^2[U^1 \cap U^2]} t_2^1(x,y) = (x,y \star g_2^1(x)) \right)$. Then $\pi_1 \circ t_2^1 = \pi_1$, t_2^1 is a morphism of X, Y and the charts are compatible m-charts of X, Y .
4. If the charts are compatible m-charts of X, Y then the transition function is an isomorphism of X, Y , hence generated by the group action, and the charts are G - ρ -compatible.

□

{cor:Grho-compatsym}

Corollary 24.9 (Symmetry of G - ρ compatibility). Let $(U^\mu, V^\mu \times Y, \phi^\mu)$ and $(U^\nu, V^\nu \times Y, \phi^\nu)$ be Y - π -bundle charts of E in the coordinate space $X \times Y$. Then $(U^\mu, V^\mu \times Y, \phi^\mu)$ is G - ρ -compatible with $(U^\nu, V^\nu \times Y, \phi^\nu)$ iff $(U^\nu, V^\nu \times Y, \phi^\nu)$ is G - ρ -compatible with $(U^\mu, V^\mu \times Y, \phi^\mu)$.

Proof. The result follows from lemma 9.11 (Symmetry of m-compatibility) on page 46.

□

Lemma 24.10 (G - ρ -compatibility of subcharts). Let $(U^i, V^i \times Y, \phi^i)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$, $(U'^i, V'^i \times Y, \phi'^i)$ a subchart and (U^1, V^1, ϕ^1) be G - ρ -compatible with (U^2, V^2, ϕ^2) . Then $(U'^1, V'^1 \times Y, \phi'^1)$ is G - ρ -compatible with $(U'^2, V'^2 \times Y, \phi'^2)$.

Proof. If $U^1 \cap U^2 = \emptyset$ then $U'^1 \cap U'^2 = \emptyset$. If $U'^1 \cap U'^2 = \emptyset$ then $(U'^1, V'^1 \times Y, \phi'^1)$ is G - ρ -compatible with $(U'^2, V'^2 \times Y, \phi'^2)$. Otherwise, the transition function $t_2'^1 \stackrel{\text{def}}{=} \phi'^1 \circ \phi'^{2^{-1}}|_{\phi'^2[U'^1 \cap U'^2]}$ is generated by the group action and hence $t_2'^1|_{\phi'^2[U'^1 \cap U'^2]}: \phi'^2[U'^1 \cap U'^2] \xrightarrow{\cong} \phi^1[U'^1 \cap U'^2]$ is generated by the group action.

□

Corollary 24.11 (G - ρ -compatibility with subcharts). Let $(U, V \times Y, \phi)$, be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $(U', V' \times Y, \phi')$ a subchart. Then $(U', V' \times Y, \phi')$ is G - ρ -compatible with $(U, V \times Y, \phi)$.

Proof. $(U, V \times Y, \phi)$ is G - ρ -compatible with itself and is a subchart of itself, $\square$

Definition 24.12 (Covering by Y - π -bundle charts). Let $\mathbf{A}$ be a set of Y - π -bundle charts of E in the coordinate space $X \times Y$. $\mathbf{A}$ covers E iff $E = \bigcup \pi_1[\mathbf{A}]$.

Lemma 24.13. Let $\mathbf{A}$ be a set of Y - π -bundle charts of E in the coordinate space $X \times Y$ that covers E and $x \in X$. Then $\pi^{-1}[\{x\}]$ is homeomorphic to Y .

Proof. Since $\mathbf{A}$ covers E , there is a chart (U, V, ϕ) in $\mathbf{A}$ containing x . Then $\pi^{-1}[\{x\}]$ is homeomorphic to Y by lemma 24.4 (Properties of projection) on page 146. $\square$ {def:piepi}

25 Bundle atlases

{sec:bunatlases}

A set of charts can be atlases for different fiber bundles even if it is for the same total model space, base space and fiber. In order to aggregate atlases into categories, there must be a way to distinguish them. Including the spaces¹⁸, group and group action in the definitions of the categories serves the purpose.

{def:BunAtl}

Definition 25.1 (Bundle atlases). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$, be a protobundle. Then $(\mathbf{A}, \mathbf{B})$ is a bundle atlas, $\mathbf{A}$ is a bundle atlas of $\mathbf{B}$, abbreviated is $\text{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ and $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$, abbreviated is $\text{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E, X, Y, \pi, G, \rho)$, iff $\mathbf{A}$ consists of a set of mutually G - ρ -compatible Y - π -bundle charts of E in the coordinate space $X \times Y$ that covers E ¹⁹.

By abuse of language we write $U \in \mathbf{A}$ for $U \in \pi_1[\mathbf{A}]$.

Remark 25.2. The definition of a π - G - ρ -bundle atlas is by design similar to the definition of a coordinate bundle in [Steenrod, 1999, p. 7], but there are significant differences. This paper will use the term bundle atlas to avoid confusion. {TopFib}

Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\mathcal{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \{(\mathbf{A}, \mathbf{B}^\alpha) | \text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha)\} \quad (25.1)$$

$$\mathcal{Atl}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\alpha < A} \mathcal{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \quad (25.2)$$

¹⁸The spaces are redundant, but convenient.

¹⁹There is no need to introduce the concept of a full π - G - ρ -bundle atlas because a π - G - ρ -bundle atlas is automatically full.

{lem:BunAtl}

Lemma 25.3 (Bundle atlases). Let E, X, Y be topological spaces, G a topological group, $\pi: E \rightarrow X$ surjective and $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y . Then $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ iff it is an m-atlas of E in the coordinate space X, Y and every coordinate function preserves fibers.

Proof. If $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ then

1. Every chart in $\mathbf{A}$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$, and hence its coordinate function preserves fibers.
2. The charts in $\mathbf{A}$ are mutually G - ρ -compatible; hence the transition functions are generated by the group action and are morphisms of X, Y .

If $\mathbf{A}$ is an m-atlas of E in the coordinate space X, Y then the transition functions are generated by the group action and thus the charts are mutually G - ρ -compatible.

If every coordinate function preserves fibers then the m-charts of $\mathbf{A}$ are Y - π -bundle charts. $\square$

Definition 25.4 (Compatibility of charts with bundle atlases). A Y - π -bundle chart $(U, V \times Y, \phi)$ of E in the coordinate space $X \times Y$ is G - ρ -compatible with a π - G - ρ -bundle atlas $\mathbf{A}$ iff it is G - ρ -compatible with every chart in the atlas.

{lem:BunCompat}

Lemma 25.5 (Compatibility of subcharts with bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ and $\mathbf{C}^1 = (U, V \times Y, \phi)$ a Y - π -bundle chart in $\mathbf{A}$. Then any subchart of $\mathbf{C}^1$ is G - ρ -compatible with $\mathbf{A}$.

Proof. Let $\mathbf{C}'^1 = (U'^1, V'^1 \times Y, \phi'^1)$ be a subchart of $\mathbf{C}^1$ and $\mathbf{C}^2 = (U^2, V^2 \times Y, \phi^2)$ another chart in $\mathbf{A}$.

1. If $U^1 \cap U^2 = \emptyset$, then $U'^1 \cap U^2 = \emptyset$.
2. If $U'^1 \cap U^2 = \emptyset$ then $\mathbf{C}'^1$ is G - ρ -compatible with $\mathbf{C}^2$.
3. Otherwise the transition function $t_2^1 \stackrel{\text{def}}{=} \phi^1 \circ \phi^{2-1} \upharpoonright_{\phi^1[U^1 \cap U^2]}$ is generated by the group action and thus $t_2^1 \upharpoonright_{\phi^2[U'^1 \cap U^2]}$ is generated by the group action.

$\square$

{def:piGrho-atl:extensions}

Lemma 25.6 (Extensions of bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -atlas of E in the coordinate space $X \times Y$ and (U_i, V_i, ϕ_i) , $i = 1, 2$, be a π - G - ρ chart of E in the coordinate space $X \times Y$ G - ρ -compatible with $\mathbf{A}$ in the coordinate space $X \times Y$. Then (U_1, V_1, ϕ_1) is G - ρ -compatible with (U_2, V_2, ϕ_2) in the coordinate space $X \times Y$.

Proof. If $U_1 \cap U_2 = \emptyset$ then (U_1, V_1, ϕ_1) is G - ρ -compatible with (U_2, V_2, ϕ_2) . Otherwise, $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}: \phi_1[U_1 \cap U_2] \xrightarrow{\cong} \phi_2[U_1 \cap U_2]$ is a homeomorphism. It remains to show that $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is generated by the group action, Let $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\alpha < A$, be charts in $\mathbf{A}$ such that $U_1 \cap U_2 \subseteq \bigcup_{\alpha < A} U'_\alpha$ and $U_1 \cap U_2 \cap U'_\alpha \neq \emptyset$, $\alpha < A$. Since the charts are G - ρ -compatible with $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\phi_2 \circ \phi'_\alpha \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ and $\phi'_\alpha \circ \phi_1^{-1} \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ are generated by the group action and thus $\phi_2 \circ \phi_1^{-1} = \phi_2 \circ \phi'_\alpha \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha} \circ \phi'_\alpha \circ \phi_1^{-1}$ is generated by the group action. $\square$

{def:Bun-ATLmax}

Definition 25.7 (Maximal bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. $\mathbf{A}$ is a maximal π - G - ρ -bundle atlas, abbreviated $\text{isAt}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E, X, Y, G, \pi, \rho)$, iff it cannot be extended by adding an additional G - ρ -compatible Y - π -bundle chart.

$\mathbf{A}$ is a semi-maximal π - G - ρ -bundle atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAt}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E, X, Y, G, \pi, \rho)$, iff whenever $(U, V \times Y, \phi) \in \mathbf{A}$, $U' \subseteq U$, $V' \times Y \subseteq V \times Y$ and $V' \times Y$ and $V'' \times Y \subseteq X \times Y$ are open sets, $\phi[U'] = V' \times Y$, $\phi': V' \times Y \xrightarrow{\cong} V'' \times Y$ is a fiber preserving homeomorphism generated by the group action then $(U', V'' \times Y, \phi' \circ \phi) \in \mathbf{A}$.

Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\text{At}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, \mathbf{B}^\alpha) \mid \text{isAt}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \right\} \quad (25.3)$$

$$\text{At}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\alpha < A} \text{At}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \quad (25.4)$$

Lemma 25.8 (Maximal π - G - ρ -bundle atlases are semi-maximal π - G - ρ -bundle atlases). Let E, X, Y be topological spaces, G a topological group, $\pi: E \rightarrow X$ surjective, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y and $\mathbf{A}$ a maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. Then $\mathbf{A}$ is a semi-maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$.

Proof. Let $(U, V \times Y, \phi) \in \mathbf{A}$, $U' \subseteq U$, $V' \times Y \subseteq V \times Y$ and $V'' \times Y \subseteq X \times Y$ be open sets and $\phi[U'] = V' \times Y$, $\phi': V' \times Y \xrightarrow{\cong} V'' \times Y$ be a fiber preserving homeomorphism generated by the group action. (U', V', ϕ) is a subchart of (U, V, ϕ) and by lemma 25.5 (Compatibility of subcharts with bundle atlases) on page 149 is G - ρ -compatible with the charts of $\mathbf{A}$. Since ϕ' is a fiber preserving homeomorphism generated by the group action, $(U', V'' \times Y, \phi' \circ \phi)$ is G - ρ -compatible with the charts of $\mathbf{A}$. Since $\mathbf{A}$ is maximal, $(U', V'', \phi' \circ \phi)$ is a chart of $\mathbf{A}$. $\square$

{lem:BunCompat}

Theorem 25.9 (Existence and uniqueness of maximal π - G - ρ -bundle atlases). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$ be a protobundle and $\mathbf{A}$ a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. Then there exists a unique maximal π - G - ρ -bundle atlas $\text{Atlas}_{\max}^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ of E in the coordinate space $X \times Y$ G - ρ -compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all π - G - ρ -bundle atlases E in the coordinate space $X \times Y$ containing $\mathbf{A}$ and G - ρ compatible in the coordinate space $X \times Y$ with $\mathbf{A}$. Let $\mathbf{P}_{\max}$ be a maximal chain of $\mathbf{P}$. Then $\mathbf{A}' = \bigcup_{\max} \mathbf{P}$ is a maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ G - ρ compatible with $\mathbf{A}$. Uniqueness follows from lemma 25.6 (Extensions of bundle atlases) on page 149. $\square$

{def:piGrho-atl:extensions}
{lem:mGrho}

Lemma 25.10 (Existence and uniqueness of projection for atlases in G - ρ -model spaces). Let E, X and Y be topological spaces, G a topological group, ρ an effective action of G on Y , $\mathbf{C} \stackrel{\text{def}}{=} (X \times Y, \mathcal{X}\mathcal{Y})$ a G - ρ model space of $X \times Y$ and $\mathbf{A}$ an m-atlas of E in the coordinate model space $\mathbf{C}$. Then there exists a unique function $\pi: E \rightarrow X$ such that for any chart (U, V, ϕ) in $\mathbf{A}$, $\pi|_U = \pi_1 \circ \phi$. If $\mathbf{A}$ is full then π is surjective.

Proof. Let (U, V, ϕ) be an arbitrary chart in $\mathbf{A}$ and define $\pi(e \in U) = \pi_1 \circ \phi(e)$. $\pi(e)$ does not depend on the choice of chart because the morphisms of a G - ρ model space preserve fibers. π is continuous because it is continuous on each coordinate patch.

Let $x \in X$. If $\mathbf{A}$ is full then there exists a chart (U, V, ϕ) in $\mathbf{A}$ such that $x \in \pi_1[V]$. Let u be an arbitrary point in $\phi^{-1}[\{x\} \times Y]$. Then $\pi(u) = \pi_1(\phi(u)) = x$. $\square$

26 Bundle-atlas (near) morphisms and functors

{sec:bunatlmorph}

This section introduces a taxonomy for morphisms between bundle atlases, defines categories of bundle atlases and constructs functors from them to categories of m-atlases. It only constructs reverse functors for subcategories of $\mathbf{B}_{\text{Bun-prod-Triv}}$.

26.1 Bundle-atlas (near) morphisms

26.1.1 Definitions of bundle-atlas (near) morphisms

{def:Bun-ATLmorphnear}

Definition 26.1 (Bundle-atlas near morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle and let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space

$C^i = X^i \times Y^i$. Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$

is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated

$\text{isAtl}_{\text{Ar}}^{\text{Bun,near}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$, iff

1. all four functions are continuous
2. f_G is a homomorphism

3. $\mathbf{f}$ commutes with π^i and ρ^i , i.e.,

$$(a) \quad \pi^2 \circ f_E = f_X \circ \pi^1$$

$$(b) \quad \left(\bigvee_{\substack{y \in Y^1 \\ g \in G^1}} \right) f_Y(y) \star^2 f_G(g) = f_Y(y \star^1 g)$$

4. For any charts $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i$, $i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_E^{-1}[U^2] \neq \emptyset$, diagram (26.1) below is locally nearly commutative in X, Y, π, ρ , i.e., for any $u^1 \in I$ there are open sets $U'^1 \subseteq I$, $V'^1 \subseteq V^1$, $U'^2 \subseteq U^2$, $V'^2 \subseteq V^2$,

$\hat{V}'^2 \subseteq X^2 \times Y^2$ and a homeomorphism $\hat{f}: V'^2 \xrightarrow{\sim} \hat{V}'^2$ ²⁰ such that eqs. (26.2) to (26.7) below hold, as shown in figs. 18 and 19. {eq:Bun-xin}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} f_E: I \stackrel{\text{def}}{=} U^1 \cap f_E^{-1}[U^2] \longrightarrow U^2, \phi^2: U^2 \xrightarrow{\sim} V^2, \\ \phi^1: I \hookrightarrow V^1, f_X \times f_Y: V^1 \longrightarrow X^2 \times Y^2 \end{array} \right. \quad (26.1) \quad \{\text{eq:Bun-At1N1}\}$$

$$u^1 \in U'^1 \quad (26.2) \quad \{\text{eq:Bun-xin}\}$$

$$f_E[U'^1] \subseteq U'^2 \quad (26.3) \quad \{\text{eq:Bun-fup1in}\}$$

$$\phi^1[U'^1] = V'^1 \quad (26.4) \quad \{\text{eq:Bun-fup1in}\}$$

$$f_X \times f_Y[V'^1] \subseteq \hat{V}'^2 \quad (26.5) \quad \{\text{eq:Bun-fvp1in}\}$$

$$\phi^2[U'^2] = V'^2 \quad (26.6) \quad \{\text{eq:Bun-fvp1in}\}$$

$$\hat{f} \circ \phi^2 \circ f_E = f_X \times f_Y \circ \phi^1 \quad (26.7) \quad \{\text{eq:Bun-fphigeqgphi}\}$$

Remark 26.2. $(U'^i, V'^i, \phi^i|_{U'^i, V'^i})$ need not be a chart of $\mathbf{A}^i$.

If $\mathbf{A}^1$ and $\mathbf{A}^2$ are semi-maximal (maximal) atlases then $\mathbf{f}$ is also a semi-maximal (maximal) $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas near morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\text{isAtl}_{\text{Ar}}^{\text{Bun, near}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$ {sec:grhoncd}

²⁰This reverses the direction of the arrow $\hat{f}: U_m \xrightarrow{\sim} V_n$ from section 23 (*G-rho-nearly commutative diagrams*) on page 144. This is permissible since $\hat{f}$ is a homeomorphism

$$\begin{array}{ccc}
I = U^1 \cap f_0^{-1}[U^2] & \xrightarrow{f_E} & U^2 \\
\downarrow \phi^1 & & \downarrow \tilde{\phi}^2 \\
V^1 & \xrightarrow{f_X \times f_Y} & C^2 \\
& & \downarrow \\
& & V^2
\end{array}$$

Figure 18: Uncompleted m-atlas near morphism

{fig:Bun-AtlN1}

$$\begin{array}{ccccccc}
& & u^1 & & & & \\
& & \downarrow i & & & & \\
I = U^1 \cap f_0^{-1}[U^2] & \xleftarrow{i} & U^{n1} & \xrightarrow{f_E} & U^{n2} & \xrightarrow{i} & U^2 \\
\downarrow \phi^1 & & \downarrow \tilde{\phi}^1 & & \downarrow \tilde{\phi}^2 & & \downarrow \tilde{\phi}^2 \\
V^1 & \xleftarrow{i} & V^{n1} & \xrightarrow{f_X \times f_Y} & V^{n2} & \xrightarrow{\tilde{f}} & V^2 \\
& & & & \downarrow \tilde{\phi} & & \\
& & & & V^2 & &
\end{array}$$

Figure 19: Completed m-atlas near morphism

{fig:Bun-AtlN2}

The identity morphism of (A^i, B^i) is

$$\text{Id}_{(A^i, B^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (A^i, B^i), (A^i, B^i)) \quad (26.8) \quad \{\text{eq:Bun-fphigeqgphi}\}$$

This nomenclature will be justified later.

{def:Bun-ATLmorph}

Definition 26.3 (Bundle-atlas morphisms). Let $B^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle and let A^i be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$.

Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\text{isAtl}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$, iff

1. all four functions are continuous
2. f_G is a homomorphism
3. $\mathbf{f}$ commutes with π^i and ρ^i , i.e.,
 - (a) $\pi^2 \circ f_E = f_X \circ \pi^1$

$$(b) \left(\bigvee_{\substack{y \in Y^1 \\ g \in G^1}} \right) f_Y(y) \star^2 f_G(g) = f_Y(y \star^1 g)$$

4. For any charts $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i$, $i = 1, 2$, and any $u^1 \in I \stackrel{\text{def}}{=} U^1 \cap f_E^{-1}[U^2]$, there exists a subchart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , an open set $U'^2 \subseteq U^2$ and a chart $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in \mathbf{A}^2$ at $f_E(u^1)$ such that $f_E[U'^1] \subseteq U'^2$ and diagram (26.9) below is commutative, as shown in fig. 20.

{fig:Bun-At1M1}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^1\} \rightarrow U'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1, f_X \times f_Y: V'^1 \rightarrow \hat{V}'^2, \\ f_E: U'^1 \rightarrow U'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2 \end{array} \right. \quad (26.9) \quad \{\text{eq:Bun-At1M1}\}$$

$$\begin{array}{ccccccc} & & u^1 & \xrightarrow{f_E} & u^2 & & \\ & & \downarrow i & & \downarrow i & & \\ U^1 & \xleftarrow{i} & U'^1 & \xrightarrow{f_E} & U'^2 & \xrightarrow{i} & U^2 \\ \downarrow \phi^1 & & \downarrow \phi^1 & & \downarrow \phi'^2 & & \downarrow \phi^2 \\ V^1 & \xleftarrow{i} & V'^1 & \xrightarrow{f_X \times f_Y} & \hat{V}'^2 & & V^2 \end{array}$$

Figure 20: Completed Bundle-atlas morphism

{fig:Bun-At1M1}

If $\mathbf{A}^1$ and $\mathbf{A}^2$ are semi-maximal (maximal) atlases then $\mathbf{f}$ is also a semi-maximal (maximal) $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated

$\text{isAtl}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$
 $\text{S-max}(\text{max})$

The identity morphism of $(\mathbf{A}^i, \mathbf{B}^i)$ is

$$\text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) \quad (26.10) \quad \{\text{eq:Bun-At1M1}\}$$

This nomenclature will be justified later.

{def:Bun-ATLmorphequiv}

Definition 26.4 (Equivalence of bundle-atlas (near) morphisms).

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is bundle equivalent to $\mathbf{g} \stackrel{\text{def}}{=} (g_E: E^1 \rightarrow E^2, g_X: X^1 \rightarrow X^2, g_Y: Y^1 \rightarrow Y^2, g_G: G^1 \rightarrow G^2)$ as a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\mathbf{f}$ and $\mathbf{g}$ are bundle equivalent $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphisms from $\mathbf{A}^1$ to $\mathbf{A}^2$, iff $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, is a protobundle, $\mathbf{A}^i$ is a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$, $\mathbf{f}$ and $\mathbf{g}$ are $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphisms from $\mathbf{A}^1$ to $\mathbf{A}^2$, $f_E = g_E, f_X = g_X$ and $f_G = g_G$,

By abuse of language we shall write $\mathbf{f}$ is bundle equivalent to $\mathbf{g}$ when the protobundles and bundle atlases are understood by context.

26.1.2 Proclamations on bundle-atlas (near) morphisms

{lem:Bun-ATLmorph}

Lemma 26.5 (Bundle-atlas (near) morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle and let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$. Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ iff $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 and $(f_E, f_X \times f_Y)$ is a E^1 - E^2 m-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 .

Proof.

1. If $\mathbf{f}$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism then $f_X \times f_Y$ is a model function from X^1, Y^1 to X^2, Y^2 by lemma 22.10 (Morphisms of trivial G - ρ -model spaces) on page 142 and f_G is the function asserted to exist in eq. (22.3) (preservation of group action) of definition 22.3 (Morphisms of G - ρ -model spaces) on page 139, so $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 .

A diagram is locally nearly commutative in X^2, Y^2, π^2, ρ^2 iff it is m-locally nearly commutative in $\pi_2 \left(X^2, Y^2 \right)_{G^2-\rho^2-\text{triv}}$, thus $(f_E, f_X \times f_Y)$ is an m-atlas morphism in the coordinate spaces X^1, Y^1 , X^2, Y^2 .

If $f_X \times f_Y$ is a G^1 - G^2 - π^1 - π^2 morphism from X^1, Y^1 to X^2, Y^2 then $\mathbf{f}$ commutes with ρ^i .

2. If $(f_E, f_X \times f_Y)$ is a E^1 - E^2 m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 then $\mathbf{f}$ commutes with π^i .

A diagram is locally nearly commutative in X^2, Y^2, π^2, ρ^2 iff it is m-locally nearly commutative in $\pi_2 \left(\begin{smallmatrix} X^2, Y^2 \\ G^2-\rho^2-\text{triv} \end{smallmatrix} \right)$ thus $(f_E, f_X \times f_Y)$ is a m-atlas morphism in the coordinate space $\begin{smallmatrix} X^2, Y^2 \\ G^2-\rho^2-\text{triv} \end{smallmatrix}$, so $\mathbf{f}$ is a $\mathbf{B}^1\text{-}\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$.

□

{cor:Bun-ATLmorph}

Corollary 26.6 (Bundle-atlas morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2, 3$, be a protobundle, $\mathbf{A}^i$ be a $\pi^i\text{-}G^i\text{-}\rho^i$ -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_E^i: E^i \longrightarrow E^{i+1}, f_X^i: X^i \longrightarrow X^{i+1}, f_Y^i: Y^i \longrightarrow Y^{i+1}, f_G^i: G^i \longrightarrow G^{i+1})$ a $\mathbf{B}^i\text{-}\mathbf{B}^{i+1}$ bundle-atlas (near) morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$.

If each $\mathbf{A}^i$ is semi-maximal or if each $\mathbf{f}^i$ is a morphism then $\mathbf{f}^2 \circ \mathbf{f}^1$ is a bundle-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^3$.

Proof. $f_X^1 \times f_Y^1 \circ f_X^2 \times f_Y^2$ is a $G^1\text{-}G^3\text{-}\pi^1\text{-}\pi^3$ morphism from $\begin{smallmatrix} X^1, Y^1 \\ G^1-\rho^1-\text{triv} \end{smallmatrix}$ to $\begin{smallmatrix} X^3, Y^3 \\ G^3-\rho^3-\text{triv} \end{smallmatrix}$ by item 4 of lemma 22.4 (Morphisms of $G\text{-}\rho$ -model spaces) on page 140 and $(f_E^1, f_X^1 \times f_Y^1) \circ (f_E^2, f_X^2 \times f_Y^2)$ is a $\begin{smallmatrix} E^1-E^3 \\ \text{triv triv} \end{smallmatrix}$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\begin{smallmatrix} X^3, Y^3 \\ G^3-\rho^3-\text{triv} \end{smallmatrix}$ by item 2 of lemma 11.16 (Composition of m-atlas (near) morphisms) on page 75.

{grhomorph: comp}

{M-morphcomp: morphcompmorph}

{lem:Bun-ATLmorphequiv}

Lemma 26.7 (Composition of equivalent bundle-atlas (near) morphisms). Let

1. $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2,$, be a protobundle
2. $\mathbf{A}^i$ be a $\pi^i\text{-}G^i\text{-}\rho^i$ -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$
3. $\mathbf{f}^i \stackrel{\text{def}}{=} (f_E^i: E^i \longrightarrow E^{i+1}, f_X^i: X^i \longrightarrow X^{i+1}, f_Y^i: Y^i \longrightarrow Y^{i+1}, f_G^i: G^i \longrightarrow G^{i+1})$ and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_E^i: E^i \longrightarrow E^{i+1}, g_X^i: X^i \longrightarrow X^{i+1}, g_Y^i: Y^i \longrightarrow Y^{i+1}, g_G^i: G^i \longrightarrow G^{i+1})$ be bundle equivalent $\mathbf{B}^i\text{-}\mathbf{B}^{i+1}$ bundle-atlas (near) morphisms from $\mathbf{A}^1$ to $\mathbf{A}^2$

Then $\mathbf{f}^{i+1} \circ \mathbf{f}^1$ is $\mathbf{C}^k$ -equivalent to $\mathbf{g}^2 \circ \mathbf{g}^1$.

Proof. $f_E^1 = g_E^1$, and $f_E^2 = g_E^2$, hence $f_E^2 \circ f_E^1 = g_E^2 \circ g_E^1$. $f_X^1 = g_X^1$, and $f_X^2 = g_X^2$, hence $f_X^2 \circ f_X^1 = g_X^2 \circ g_X^1$. $f_Y^1 = g_Y^1$, and $f_Y^2 = g_Y^2$, hence $f_Y^2 \circ f_Y^1 = g_Y^2 \circ g_Y^1$. $f_G^1 = g_G^1$, and $f_G^2 = g_G^2$, hence $f_G^2 \circ f_G^1 = g_G^2 \circ g_G^1$. □

26.2 Categories of bundle atlases and functors

This subsection only defines categories whose morphisms are bundle atlas morphisms; bundle atlas near morphisms do not satisfy the requirements for forming a category.

{def:BunATLsets}

Definition 26.8 (Sets of bundle atlas morphisms). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles.

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\alpha, \mathbf{B}^\beta) \stackrel{\text{def}}{=} \left\{ (f, (A^\alpha, \mathbf{B}^\alpha), (A^\beta, \mathbf{B}^\beta)) \mid \right. \\ \left. \text{isAtI}_{\text{Ar}}^{\text{Bun}}(A^\alpha, \mathbf{B}^\alpha, A^\beta, \mathbf{B}^\beta, f) \right\} \quad (26.11)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \left\{ (f, (A^\alpha, \mathbf{B}^\alpha), (A^\alpha, \mathbf{B}^\alpha)) \mid \right. \\ \left. \text{isAtI}_{\text{Ar}}^{\text{Bun}}(A^\alpha, \mathbf{B}^\alpha, A^\alpha, \mathbf{B}^\alpha, f) \right\} \quad (26.12)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\substack{\mathbf{B}^\mu \in \mathbf{B} \\ \mathbf{B}^\nu \in \mathbf{B}(\text{S-max, max})}} \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\mu, \mathbf{B}^\nu) \quad (26.13)$$

{def:Bun-AtIcat}

Definition 26.9 (Categories of bundle atlases). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha), \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\alpha, \mathbf{B}^\alpha), \circ^A \right) \quad (26.14)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}), \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}), \circ^A \right) \quad (26.15)$$

{lem:BunATLiscat}

Lemma 26.10 ($\mathcal{Att}^{\text{Bun}} \mathbf{B}$ is a category). Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ is a category.

Let $(A^\alpha, \mathbf{B}^\alpha) \in \mathcal{Att}_{\text{Ob}}^{\text{Bun}} \mathbf{B}$. Then $\text{Id}_{(A^\alpha, \mathbf{B}^\alpha)}$ is the identity morphism for $(A^\alpha, \mathbf{B}^\alpha)$.

Proof. Let $(A^i, \mathbf{B}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ and let

$m^i \stackrel{\text{def}}{=} (f^i, (A^i, \mathbf{B}^i), (A^{i+1}, \mathbf{B}^{i+1}))$ be a morphism of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$. Then

Composition: $(f^2 \circ f^1, (A^1, \mathbf{B}^1, \mathbf{C}^1), (A^3, \mathbf{B}^3, \mathbf{C}^3))$ is a morphism of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ by corollary 26.6 (Bundle-atlas morphisms) on page 156.

{cor:Bun-ATLmorph}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{def:FuncBuntoM}

Definition 26.11 (Functor from Bundle atlases to m-atlases). Let

$\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle, let A^i be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$ and let

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \longrightarrow E^2, f_X: X^1 \longrightarrow X^2, f_Y: Y^1 \longrightarrow Y^2, f_G: G^1 \longrightarrow G^2)$ be a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from A^1 to A^2 . Then

$$\mathcal{F}_{\text{Bun}, M}^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i) \stackrel{\text{def}}{=} \left(\mathbf{A}^i, \begin{matrix} E^i \\ \text{triv} \end{matrix}, \begin{matrix} X^i, Y^i \\ G^i - \rho^i - \text{triv} \end{matrix} \right) \quad (26.16)$$

$$\begin{aligned} \mathcal{F}_{\text{Bun}, M}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) &\stackrel{\text{def}}{=} \\ &\left((f_E, f_X \times f_Y), \left(\mathbf{A}^1, \begin{matrix} E^1 \\ \text{triv} \end{matrix}, \begin{matrix} C^1 \\ G^1 - \rho^1 - \text{triv} \end{matrix} \right), \left(\mathbf{A}^2, \begin{matrix} E^2 \\ \text{triv} \end{matrix}, \begin{matrix} C^2 \\ G^2 - \rho^2 - \text{triv} \end{matrix} \right) \right) \end{aligned} \quad (26.17)$$

Theorem 26.12 (Functor from Bundle atlases to m-atlases). Let $\mathbf{E}$ be a set of topo-

{the:FuncBuntoM}

logical spaces, $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha \in \mathbf{E}, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle,

$\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles and $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha, Y^\alpha)$. Then $\mathcal{F}_{\text{Bun}, M}^{\text{Bun}}$ is a

functor from $\mathcal{Att}_{(S-\text{max}, \text{max})}^{\text{Bun}} \mathbf{B}$ to $\mathcal{Att}_{(S-\text{max}, \text{max})} \left(\begin{matrix} \mathbf{E} \\ \mathcal{F}^{\text{triv}} \end{matrix}, \begin{matrix} \mathbf{B} \\ \text{Bun} - \mathcal{F}^{\text{triv}} \end{matrix} \right)$ and a functor from

$\mathcal{Att}_{(S-\text{max}, \text{max})}^{\text{Bun}} \mathbf{B}$ to $\mathcal{Att}_{(S-\text{max}, \text{max})} \left(\begin{matrix} \mathbf{E} \\ \mathcal{F}^{\text{triv}} \end{matrix}, \begin{matrix} \mathbf{B} \\ \text{Bun} - \text{prod} - \mathcal{F}^{\text{triv}} \end{matrix} \right)$.

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Att}^{\text{Bun}}(\mathbf{B})$, $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Bun}, M}^{\text{Bun}} o^i =$

$\left(\mathbf{A}^i, \begin{matrix} E^i \\ \text{triv} \end{matrix}, \begin{matrix} C^i \\ G^i - \rho^i - \text{triv} \end{matrix} \right)$ be the corresponding object of $\mathcal{Att} \left(\begin{matrix} \mathbf{E} \\ \mathcal{F}^{\text{triv}} \end{matrix}, \begin{matrix} \mathbf{B} \\ \text{Bun} - \mathcal{F}^{\text{triv}} \end{matrix} \right)$,

$m^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o^i, o^{i+1})$, $i = 1, 2$, be a morphism of $\mathcal{Att}^{\text{Bun}}(\mathbf{B})$ from o^i to o^{i+1} and

$$\begin{aligned} m'^i &\stackrel{\text{def}}{=} \mathcal{F}_{\text{Bun}, M}^{\text{Bun}} m^i \\ &= \left((f_E^i, f_X^i \times f_Y^i), \left(\mathbf{A}^i, \begin{matrix} E^i \\ \text{triv} \end{matrix}, \begin{matrix} C^i \\ G^i - \rho^i - \text{triv} \end{matrix} \right), \left(\mathbf{A}^{i+1}, \begin{matrix} E^{i+1} \\ \text{triv} \end{matrix}, \begin{matrix} C^{i+1} \\ G^{i+1} - \rho^{i+1} - \text{triv} \end{matrix} \right) \right) \\ &= ((f_E^i, f_X^i \times f_Y^i), o'^i, o'^{i+1}) \end{aligned}$$

be the corresponding morphism of $\mathcal{Att} \left(\begin{matrix} \mathbf{E} \\ \mathcal{F}^{\text{triv}} \end{matrix}, \begin{matrix} \mathbf{B} \\ \text{Bun} - \mathcal{F}^{\text{triv}} \end{matrix} \right)$.

Preservation of endpoints: $\begin{matrix} E^i \\ \text{triv} \end{matrix} \overset{\text{Ob}}{\in} \begin{matrix} E \\ \mathcal{T}riv \end{matrix} \cdot \begin{matrix} C^i \\ G^i-\rho^i-\text{triv} \end{matrix} \overset{\text{Ar}}{\in} \begin{matrix} B \\ \text{Bun-prod-}\mathcal{T}riv \end{matrix}.$

By theorem 26.12 (Functor from Bundle atlases to m-atlases) on page 158, $\{the : FuncBuntoM\}$
 $(f_E, f_X \times f_Y)$ is a $\begin{matrix} E^i-E^{i+1} \\ \text{triv} \end{matrix}$ m-atlas morphism from A^i to A^{i+1} in the coordinate
spaces $\begin{matrix} X^i, Y^i \\ G^i-\rho^i-\text{triv} \end{matrix}, \begin{matrix} X^{i+1}, Y^{i+1} \\ G^{i+1}-\rho^{i+1}-\text{triv} \end{matrix}.$

$$\begin{aligned} \mathcal{F}_{\text{Bun},M}^{\text{Bun}} m^i &= \mathcal{F}_{\text{Bun},M}^{\text{Bun}} (f^i, o^i, o^{i+1}) \\ &= ((f_E^i, f_X^i \times f_Y^i), o^i, o^{i+1}) \end{aligned}$$

Identity:

$$\begin{aligned} \mathcal{F}_{\text{Bun},M}^{\text{Bun}} \text{Id}_{(A^i, B^i)} &= \mathcal{F}_{\text{Bun},M}^{\text{Bun}} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), o^i, o^i) \\ &= ((\text{Id}_{\begin{matrix} E^i \\ \text{triv} \end{matrix}}, \text{Id}_{C^i}), o^i, o^i) \\ &= ((\text{Id}_{\begin{matrix} E^i \\ \text{triv} \end{matrix}}, \text{Id}_{C^i}), \mathcal{F}_{\text{Bun},M}^{\text{Bun}} o^i, \mathcal{F}_{\text{Bun},M}^{\text{Bun}} o^i) \\ &= \text{Id}_{\mathcal{F}_{\text{Bun},M}^{\text{Bun}} o^i} \end{aligned}$$

Composition:

$$\begin{aligned} m^2 \circ^A m^1 &= ((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), o^1, o^3) \\ \mathcal{F}_{\text{Bun},M}^{\text{Bun}} (m^2 \circ m^1) &= ((f_E^2 \circ f_E^1, (f_X^2 \circ f_X^1) \times (f_Y^2 \circ f_Y^1)), o'^1, o'^3) \\ &= ((f_E^2 \circ f_E^1, (f_X^2 \times f_Y^2) \circ (f_X^1 \times f_Y^1)), o'^1, o'^3) \\ &= \mathcal{F}_{\text{Bun},M}^{\text{Bun}} m^2 \circ^A \mathcal{F}_{\text{Bun},M}^{\text{Bun}} m^1 \end{aligned}$$

□

$\{lem: \text{BunAtlbasemorph}\}$

Lemma 26.13 (Base space functions derived from bundle-atlas morphisms). Let $E^i, i = 1, 2$, be a model space, X^i, Y^i topological spaces, G^i a topological group, ρ^i an effective action of G^i on Y^i , $C^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$, $f_C: C^1 \rightarrow C^2$ a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$, i.e., a model function that preserves group action, A^i an m-atlas of E^i in the coordinate model space C^i and $f \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_C: C^1 \rightarrow C^2)$ an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Then there exists a unique function $f_X: X^1 \rightarrow X^2$ such that $f_X \circ \pi_1 = \pi_1 \circ f_C$.

Proof. Let x in X^1, y, y' in Y^1 , Then f_C preserves fibers, i.e.,

$\pi_1(f_C(x, y)) = \pi_1(f_C(x, y'))$. by lemma 22.4 (Morphisms of G - ρ -model spaces)

$\{lem: \text{Grhomorph}\}$

on page 139. Define $f_X(x) \stackrel{\text{def}}{=} \pi_1(f_C(x, y))$

□

Remark 26.14. There need not exist $f_Y: Y^1 \rightarrow Y^2$ such that $f_C = f_X \times f_Y$.

Definition 26.15 (Functor from m-atlases to Bundle atlases). Let $\mathcal{E}$ be a trivial model category, $\mathcal{C}$ be a trivial product coordinate model category with objects {def:FuncMtoBun}

$\{\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}\mathcal{Y}^\alpha) \mid \alpha < A\}$, G a group valued function on $\text{Ob}(\mathbf{C})$ and ρ a function

valued function on $\text{Ob}(\mathbf{C})$ such that for every $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}\mathcal{Y}^\alpha) \in \mathcal{C}$, $\rho(\mathbf{C}^\alpha)$ is an effective action of $G(\mathbf{C}^\alpha)$ on Y^α and $\frac{X^\alpha, Y^\alpha}{G(\mathbf{C}^\alpha) - \rho(\mathbf{C}^\alpha) - \text{triv}} = \mathbf{C}^\alpha$.

Let $\mathbf{E}^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i) \in \mathcal{E}$, $i = 1, 2$, be a trivial model space, $\mathbf{C}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i) \in \mathcal{C}$ {lem:mGrho}
 $G^i \stackrel{\text{def}}{=} G(\mathbf{C}^i)$, $\rho^i \stackrel{\text{def}}{=} \rho(\mathbf{C}^i)$, A^i a full m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$, $\pi^i: E^i \rightarrow X^i$ the unique function asserted in lemma 25.10 and $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$. Then define

$$\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i) \quad (26.18)$$

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_E: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f_C \stackrel{\text{def}}{=} f_X \times f_Y: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ be an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ that preserves the group action, $f_G: G^1 \rightarrow G^2$ the unique function asserted to exist in eq. (22.3) (preservation of group action) of definition 22.3 (Morphisms of G - ρ -model spaces) on page 139. {eq:Grho}
Then {def:Grhomorph}

$$\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} ((f_E, f_X, f_Y, f_G), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \quad (26.19)$$

Theorem 26.16 (Functor from m-atlases to bundle atlases). Let {the:FuncMtoBun}

1. $\mathcal{E}$ be a trivial product coordinate model category,
2. $\mathcal{E}$ be a model category,
3. G be a group valued function on $\text{Ob}(\mathcal{E})$
4. ρ be a function valued function on $\text{Ob}(\mathbf{C})$ such that for every $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}\mathcal{Y}^\alpha) \in \mathcal{C}$, $\rho(\mathbf{C}^\alpha)$ is an effective action of $G(\mathbf{C}^\alpha)$ on Y^α and $\mathbf{C}^\alpha = \frac{X^\alpha, Y^\alpha}{G(\mathbf{C}^\alpha) - \rho(\mathbf{C}^\alpha) - \text{triv}}$.
5. π be the unique function valued function on $\mathcal{A}t\ell_{\text{Ob}(\mathcal{E}, \mathcal{C})}^{\text{full}}$ such that for every $(\mathbf{A}^\alpha, (E^\alpha, \mathcal{E}^\alpha), (C^\alpha, \mathcal{C}^\alpha)) \in \mathcal{A}t\ell_{\text{Ob}(\mathcal{E}, \mathcal{C})}^{\text{full}}$ and every $(U, V, \phi) \in \mathbf{A}^\alpha$, $\pi_1 \circ \phi = \pi(\mathbf{A}^\alpha) \upharpoonright_U$
6. $\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, G(\mathbf{C}^\alpha), \pi(\mathbf{A}^\alpha), \rho(\mathbf{C}^\alpha)) \mid \right.$
 $\left. (\mathbf{A}^\alpha, (E^\alpha, \mathcal{E}^\alpha), (C^\alpha \stackrel{\text{def}}{=} X^\alpha \times Y^\alpha, \mathcal{C}^\alpha)) \in \mathcal{A}t\ell_{\text{Ob}(\mathcal{E}, \mathcal{C})}^{\text{full}} \right\}$

Then $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}$ is a functor from $\mathcal{Att}\left(\frac{\mathbf{E}}{\mathcal{T}^{\text{triv}}}, \text{Bun-prod-}\frac{\mathbf{B}}{\mathcal{T}^{\text{triv}}}\right)$ to $\mathcal{Att}^{\text{Bun}}(\mathbf{B})$.

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}^i \mathcal{Y}^i)) \in \mathcal{M}$, $i \in [1, 3]$, $G^i \stackrel{\text{def}}{=} G(o^i)$,
 $\pi^i \stackrel{\text{def}}{=} \pi(o^i)$, $\rho^i \stackrel{\text{def}}{=} \rho(o^i)$, $\mathbf{A}^i$ an m -atlas of E^i in the coordinate model space $\mathbf{C}^i$,
 $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$, $m^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o^i, o^{i+1}) \stackrel{\text{Ar}}{\in} \mathcal{M}$, $i = 1, 2$, an E^i -
 E^{i+1} m -atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$ that preserves
the group action, $f_G: G^i \rightarrow G^{i+1}$ the unique function asserted to exist in eq. (22.3)
(preservation of group action) of definition 22.3 (Morphisms of G - ρ -model spaces)
on page 139 and $\pi^i: E^i \rightarrow X^i$ the unique function asserted in lemma 25.10.

Let $f_G^i: G^i \rightarrow G^{i+1}$ be the function asserted by eq. (22.3) (Morphisms of G - ρ -
model spaces) on page 139; let $f_1^i = f_X^i \times f_Y^i$ be the decomposition given by defini-
tion 22.7 (Trivial G - ρ -model spaces) on page 141, and let $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^i =$
 $(\mathbf{f}^i, (f_0^i f_X^i f_Y^i f_G^i), o^i, o^{i+1})$.

{eq:Grho}
{def:Grhomorph}
{lem:mGrho}
{eq:Grho}
{def:Grhomodtriv}

Preservation of endpoints: E^i, X^i and Y^i are topological spaces.

G^i is a topological group,

$\pi^i: E^i \rightarrow X^i$ is a continuous surjection.

$\rho^i \stackrel{\text{def}}{=} \rho(\mathbf{C}^i)$ is an effective action of $G^i \stackrel{\text{def}}{=} G(\mathbf{C}^i)$ on Y^α .

Hence B^i is a protobundle.

Identity:

$$\begin{aligned} \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} &= \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)}((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), o^i, o^i) \\ &= ((\text{Id}_{\frac{E^i}{\text{triv}}}, \text{Id}_{C^i}), o^i, o^i) \\ &= ((\text{Id}_{\frac{E^i}{\text{triv}}}, \text{Id}_{C^i}), \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i) \\ &= \text{Id}_{\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i} \end{aligned}$$

Composition:

$\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^i$ is a morphism from $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i$ to $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^{i+1}$:

$$\begin{aligned} \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}(m^i) &= \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}(\mathbf{f}^i, o^i, o^{i+1}) \\ &= ((f_0^i, f_X^i, f_Y^i, f_G^i), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^{i+1}, \mathbf{B}^{i+1})) \\ &= ((f_0^i, f_X^i, f_Y^i, f_G^i), \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^{i+1}) \end{aligned}$$

$\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}$ maps identity functions to identity functions:

$$\begin{aligned}
& \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} \text{Id}_{o^i} = \\
& \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} ((\text{Id}_{E^i}, \text{Id}_{C^i}), o^i, o^i) = \\
& ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) = \\
& ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i) = \\
& \text{Id}_{\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} o^i}
\end{aligned} \tag{26.20}$$

$$\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^2 \circ^A \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^1 = \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (m^2 \circ^A m^1):$$

1. $m^2 \circ^A m^1 = ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3)$
2. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (o^i) = (\mathbf{A}^i, \mathbf{B}^i)$
3. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (m^i) = ((f_0^i, f_X^i, f_Y^i, f_G^i), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^{i+1}, \mathbf{B}^{i+1}))$
4. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^2 \circ^A \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^1 = ((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^3, \mathbf{B}^3))$
5. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (m^2 \circ m^1) = ((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^3, \mathbf{B}^3))$

□

$$\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} \circ \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} = \text{Id}.$$

Proof. Expanding the definitions, we have

1. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) = (\mathbf{A}^i, \mathbf{B}^i)$
2. $\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} (\mathbf{A}^i, \mathbf{B}^i) = (\mathbf{A}^i, \underset{\text{triv } G^i-\rho^i-\text{triv}}{E^i, X^i, Y^i})(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, since by definition 26.15, $\mathbf{E}^i$ and $\mathbf{C}^i$ are trivial. {def:FuncMtoBun}
3. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} ((f_E^1, f_X^1 \times f_Y^1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) = ((f_E^1, f_X^1, f_Y^1, f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2))$
4. $\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} ((f_E^1, f_X^1, f_Y^1, f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) = ((f_E^1, f_X^1 \times f_Y^1), (\mathbf{A}^1, \underset{\text{triv } G^1-\rho^1-\text{triv}}{E^1, C^1}), (\mathbf{A}^2, \underset{\text{triv } G^2-\rho^2-\text{triv}}{E^2, C^2})) = ((f_E^1, f_X^1 \times f_Y^1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2))$, since by definition 26.15, $\mathbf{E}^i$ and $\mathbf{C}^i$ are trivial. {def:FuncMtoBun}

□

27 Associated model spaces and functors

Definition 27.1 (Coordinate model spaces associated with bundle atlases). Let $\text{sec:assmodB} \quad \{\text{def:ismodBF}\}$

$\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$ be a protobundle $\mathbf{A}^i$ be a π^i - G^i - ρ^i -bundle atlas of E^i in the coordinate space $C^i = X^i \times Y^i$ and $\mathbf{f}$

$\stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ be a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$. Then:

The minimal G - ρ coordinate model space with neighborhoods in $\mathbf{A}^i$ is $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i)$.

$$\mathcal{F}_2^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i) \stackrel{\text{def}}{=} \text{Mod}_{\min} \left(C^i, \pi_2[\mathbf{A}^i], \left\{ \phi' \circ \phi^{-1} \mid \left(\exists_{\substack{(U,V,\phi) \in \mathbf{A}^i \\ (U',V',\phi') \in \mathbf{A}^i}} \right) U \cap U' \neq \emptyset \right\} \right) \quad (27.1)$$

The coordinate mapping associated with the $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism $\mathbf{f}$ from $\mathbf{A}^1$ to $\mathbf{A}^2$ is

$$\mathcal{F}_2^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} f_X \times f_Y: \mathcal{F}_2^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1) \rightarrow \mathcal{F}_2^{\text{Bun}}(\mathbf{A}^2, \mathbf{B}^2) \quad (27.2)$$

If it is a model function then it is also the coordinate G^1 - G^2 - ρ^1 - ρ^2 (near) morphism associated with the $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism $\mathbf{f}$ from $\mathbf{A}^1$ to $\mathbf{A}^2$ lem:ismodBF

Lemma 27.2 (Coordinate model spaces associated with bundle atlases). Let

$\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, G, \pi, \rho)$ and let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $C = X \times Y$. Then $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ is a model space.

Proof. $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ satisfies the conditions for a model space. for a model space. Let

$$\mathcal{C} \stackrel{\text{def}}{=} \text{Cat}(\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})).$$

1. Since $\pi_2[\mathbf{A}]$ is an open cover of $\bigcup \pi_2[\mathbf{A}]$, the set of finite intersections is also an open cover.
2. Finite intersections of finite intersections are finite intersections
3. Restrictions of continuous functions are continuous
4. If $f: A \rightarrow B$ is a morphism of $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ A, A', B, B' objects of $\mathcal{C}$ $A' \subseteq A$, $B' \subseteq B$ and $f[A'] \subseteq B'$ then since $f: A \rightarrow B$ is a morphism it is a restriction of a transition function between its restrictions to sets in $\pi_2[\mathbf{A}]$ and its restrictions are also, hence morphisms, and thus $f|_{A'}: A' \rightarrow B'$ is a morphism.
5. If $(U, V, \phi) \in \mathbf{A}$ then $\text{Id}_V = \phi \circ \phi^{-1}$ is a transition function and hence a morphism of $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})$. If A, A' objects of $\mathcal{C}$ and $A' \subseteq A$ then the inclusion map $i: A' \hookrightarrow A$ is a restriction of an identity morphism of $\mathcal{F}_2^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ and hence a morphism.

6. Restricted sheaf condition: let

- (a) $U_\alpha, V_\alpha, \alpha < A$, be an object of $\mathcal{C}$
- (b) $f_\alpha: U_\alpha \longrightarrow V_\alpha$ be a morphism of $\mathcal{C}$
- (c) $U \stackrel{\text{def}}{=} \bigcup_{\alpha < A} U_\alpha$
- (d) $V \stackrel{\text{def}}{=} \bigcup_{\alpha < A} V_\alpha$
- (e) $f: U \longrightarrow V$ be continuous and $\left(\bigvee_{\substack{\alpha < A \\ x \in U_\alpha}} \right) f(x) = f_\alpha(x)$

Then f is generated by the group action and hence a morphism of $\mathcal{C}$

□

{def:ismodT}

Definition 27.3 (Model spaces associated with bundle atlases). Let

$\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, and $\mathbf{A}^i$ be a π^i - G^i - ρ^i -bundle atlas of E^i in the coordinate space $C^i = X^i \times Y^i$. Then

The minimal model space with neighborhoods in the atlas $\mathbf{A}^i$ is

$$\mathcal{F}_1^{\text{Bun}}(\mathbf{B}^i, \mathbf{A}^i) \stackrel{\text{def}}{=} \text{Mod}_{\min}(E^i, \pi_1[\mathbf{A}^i], \left\{ \phi'^{-1} \circ \phi \mid \left(\exists_{\substack{(U, V, \phi) \in \mathbf{A}^i \\ (U', V', \phi') \in \mathbf{A}^i}} \right) U \cap U' \neq \emptyset \right\} \right) \quad (27.3)$$

The mapping associated with the $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism $\mathbf{f}$ from $\mathbf{A}^1$ to $\mathbf{A}^2$ is

$$\mathcal{F}_1^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} f_E: \mathcal{F}_1^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1) \longrightarrow \mathcal{F}_1^{\text{Bun}}(\mathbf{A}^2, \mathbf{B}^2) \quad (27.4)$$

If it is a model function then it is also the m-atlas (near) morphism associated with the $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas (near) morphism $\mathbf{f}$ from $\mathbf{A}^1$ to $\mathbf{A}^2$.

{lem:ismodT}

Lemma 27.4 (Model spaces associated with bundle atlases). Let

$\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, G, \pi, \rho)$ and let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $C = X \times Y$. Then $\mathcal{F}_1^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ is a model space.

Proof. Lemma 5.4 (Minimal model spaces are model spaces) on page 26

□

{the:FuncBuntoMod}

Theorem 27.5 (Functors from bundle atlases to model spaces). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle, and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha \mid \alpha < A\}$ be a set of protobundles. Then $\mathcal{F}_1^{\text{Bun}}$ is a functor from $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ to $\mathbf{E}_{\text{triv}}$ and $\mathcal{F}_2^{\text{Bun}}$ is a functor from $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ to $\mathbf{B}_{\text{Bun-}\mathcal{F}\text{riv}}$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $i \in [1, 3]$, be objects in $\text{Bun}(E, X, Y, \pi, G, \rho)$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$ be a morphism in $\mathcal{A}t\ell^{\text{Bun}}(E, \mathbf{C})$.

$\mathcal{F}_1^{\text{Bun}}$ is a functor from $\mathcal{A}t\ell^{\text{Bun}} \mathbf{B}$ to $\mathbf{E}_{\text{triv}}$.

Preservation of endpoints:

$$\begin{aligned} \mathcal{F}_2^{\text{Bun}} m^i &= \mathcal{F}_1^{\text{Bun}}(\mathbf{f}^i, o^i, o^{i+1}) \\ &= f_E^i: \mathcal{F}_1^{\text{Bun}} o^i \longrightarrow o^{i+1} \end{aligned}$$

by definition 27.3 (Model spaces associated with bundle atlases) on page 164 `{def:ismodT}`

Composition:

$$\begin{aligned} \mathcal{F}_1^{\text{Bun}}(m^2 \circ^A m^1) &= \mathcal{F}_1^{\text{Bun}}(\mathbf{f}^2 \circ \mathbf{f}^1(A^1, E^1, C^1), (A^3, E^3, C^3)) \\ &= f_E^2 \circ f_E^1: \mathcal{F}_1^{\text{Bun}} o^1 \longrightarrow \mathcal{F}_1^{\text{Bun}} o^3 \\ &= (f_E^2: \mathcal{F}_1^{\text{Bun}} o^2 \longrightarrow \mathcal{F}_1^{\text{Bun}} o^3) \circ \\ &\quad (f_E^1: \mathcal{F}_1^{\text{Bun}} o^1 \longrightarrow \mathcal{F}_1^{\text{Bun}} o^2) \\ &= \mathcal{F}_1^{\text{Bun}}(\mathbf{f}^2, (A^2, E^2, C^2), (A^3, E^3, C^3)) \circ \\ &\quad \mathcal{F}_1^{\text{Bun}}(\mathbf{f}^1, (A^1, E^1, C^1), (A^2, E^2, C^2)) \\ &= \mathcal{F}_1^{\text{Bun}} m^2 \circ \mathcal{F}_1^{\text{Bun}} m^1 \end{aligned}$$

Identity:

1. $\mathcal{F}_1^{\text{Bun}}(\text{Id}_{o^i}) = \mathcal{F}_1^{\text{Bun}}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) =$
 $\text{Id}_{E^i}: \mathcal{F}_1^{\text{Bun}} o^i \longrightarrow \mathcal{F}_1^{\text{Bun}} o^i$
2. $\text{Id}_{\mathcal{F}_1^{\text{Bun}} o^i} = \text{Id}_{E^i}: \mathcal{F}_1^{\text{Bun}} o^i \longrightarrow \mathcal{F}_1^{\text{Bun}} o^i$

$\mathcal{F}_2^{\text{Bun}}$ is a functor from $\mathcal{A}t\ell^{\text{Bun}} \mathbf{B}$ to $\mathbf{B}_{\text{Bun-}\mathcal{F}\text{triv}}$.

Preservation of endpoints: $\mathcal{F}_2^{\text{Bun}}(m^i) = f_1^i: \mathcal{F}_2^{\text{Bun}} o^i \longrightarrow \mathcal{F}_2^{\text{Bun}} o^{i+1}$

Composition:

$$\begin{aligned} \mathcal{F}_2^{\text{Bun}}(m^2 \circ^A m^1) &= \mathcal{F}_2^{\text{Bun}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)(A^1, E^1, C^1), (A^3, E^3, C^3)) \\ &= f_0^2 \circ f_0^1: \mathcal{F}_2^{\text{Bun}} o^1 \longrightarrow \mathcal{F}_2^{\text{Bun}} o^3 \\ &= (f_1^2: \mathcal{F}_2^{\text{Bun}} o^2 \longrightarrow \mathcal{F}_2^{\text{Bun}} o^3) \circ (f_1^1: \mathcal{F}_2^{\text{Bun}} o^1 \longrightarrow \mathcal{F}_2^{\text{Bun}} o^2) \\ &= \mathcal{F}_2^{\text{Bun}}((f_0^2, f_1^2)(A^2, E^2, C^2), (A^3, E^3, C^3)) \circ \\ &\quad \mathcal{F}_2^{\text{Bun}}((f_0^1, f_1^1)(A^1, E^1, C^1), (A^2, E^2, C^2)) \\ &= \mathcal{F}_2^{\text{Bun}}(m^2) \circ \mathcal{F}_2^{\text{Bun}}(m^1) \end{aligned}$$

Identity:

1. $\mathcal{F}_2^{\text{Bun}} \text{Id}_{o^i} = \mathcal{F}_2^{\text{Bun}}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) =$
 $\text{Id}_{C^i}: \mathcal{F}_2^{\text{Bun}} o^i \longrightarrow \mathcal{F}_2^{\text{Bun}} o^i$
2. $\text{Id}_{\mathcal{F}_2^{\text{Bun}} o^i} = \text{Id}_{C^i}: \mathcal{F}_2^{\text{Bun}} o^i \longrightarrow \mathcal{F}_2^{\text{Bun}} o^i$

□

28 Fiber bundles

{sec:fib}

Conventionally a fiber bundle is different from its atlases, but definition 26.9 (Categories of bundle atlases) on page 157 encourages treating them on an equal footing. All of the results for maximal bundle atlases carry directly over to results for fiber bundles.

{def:Bun-Atlcat}

28.1 Definitions

{def:Bun}

Definition 28.1 (fiber bundles). Let E, X and Y be topological spaces, $\pi: E \rightarrow X$ surjective, G a topological group, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y and $\mathbf{A}$ a maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. Then $(E, X, Y, \pi, G, \rho, \mathbf{A})$ is a fiber bundle.

Let $\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha) \mid \alpha < A \right\}$, where $E^\alpha, X^\alpha, Y^\alpha$ are topological spaces, G^α a topological group, $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ surjective and $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ an effective right action of G^α on Y^α . Then

$$\text{Bun}_{\text{Ob}} \mathbf{B} \stackrel{\text{def}}{=} \mathcal{Att}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \quad (28.1)$$

{def:Bunmorph}

Definition 28.2 (Bundle maps). Let

$\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha) \mid \alpha < A \right\}$, where $E^\alpha, X^\alpha, Y^\alpha$ are topological spaces, G^α a topological group, $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ surjective and $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ an effective right action of G^α on Y^α . Then

$$\text{Bun}_{\text{Ar}} \mathbf{B} \stackrel{\text{def}}{=} \mathcal{Att}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \quad (28.2)$$

$$\text{Bun} \mathbf{B} \stackrel{\text{def}}{=} (\text{Bun}_{\text{Ob}} \mathbf{B}, \text{Bun}_{\text{Ar}} \mathbf{B}, \circ^A) \quad (28.3)$$

Let $(A^i, \mathbf{B}^i) \in \text{Bun}_{\text{Ob}} \mathbf{B}, i = 1, 2$. Then

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a bundle map from $(A^1, \mathbf{B}^1)$ to $(A^2, \mathbf{B}^2)$ iff it is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from A^1 to A^2 . The identity morphism for $(A^i, \mathbf{B}^i)$ is

$$\text{Id}_{(A^i, \mathbf{B}^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (A^i, \mathbf{B}^i), (A^i, \mathbf{B}^i)) \quad (28.4)$$

28.2 Categories of fiber bundles

{the:Bun}

Theorem 28.3 (Categories of fiber bundles). Let

$\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \mid \alpha < A \right\}$, where $E^\alpha, X^\alpha, Y^\alpha$ are topological spaces, G^α a topological group, $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ surjective and $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ an effective right action of G^α on Y^α . Then $\text{Bun}\mathbf{B}$ is a category and $Id_{\mathbf{B}^\alpha}$ is the identity morphism for $\mathbf{B}^\alpha$.

Proof. The result follows directly from [definition 28.1 \(fiber bundles\)](#), [definition 28.2 \(Bundle maps\)](#) and [lemma 26.10 \(\$\mathcal{A}tl^{\text{Bun}}\mathbf{B}\$ is a category\)](#) on page 157. $\square$

{def:Bunmorph}
{lem:BunATLiscat}

Part IX

Future directions

{part:fut}

If this paradigm proves useful, it can be extended to include a set of admissible functions on the model neighborhoods of the charts, possibly using the language of sheaves. That might be desirable for coordinate spaces more general than Fréchet spaces.

Further work is needed to determine whether there is a productive way to define a notion of classic m-atlas near morphisms.

Further work is needed to determine whether it is productive to define (semi-)strict m-atlas morphisms in terms of concrete categories over a category of model spaces rather than directly in terms of a model category.

The extension of paracompactness to model spaces is intended to be useful for partitions of unity on fiber bundles. Further work is needed to determine whether that is actually the case.

The definitions given here include some fairly strong conditions, e.g., AOC. Further work is needed to determine whether they should be relaxed for applications beyond manifolds and fiber bundles.

Further work is needed to determine whether the concept of category-based atlases²¹ of model spaces has general utility.

If the concept of nearly commutative diagrams proves useful, further work is needed to determine whether a more general definition has utility.

Further work is needed to determine where it is most fruitful to require full subcategories and where just subcategories.

Further work is needed to determine conditions for mappings associated with atlas morphisms to be model functions.

This paper uses the language of category theory as an organizing principle, but defines various notions concretely with sets. It may be desirable to abstract away some of the details, in the spirit of, e.g., topoi.

²¹As opposed to pseudogroup based

References

- [Adámek, Herrlich, Strecker, 1990] Jiří Adámek, Horst Herrlich, George E. Strecker. *Abstract and Concrete Categories The Joy of Cats*, John Wiley and Sons, Inc., 1990.
- [Kelley, 1955] John L. Kelley, *General Topology*, D. Van Nostrand Company (first edition), 1955.
- [Kobayashi, 1996] Shoshichi Kobayashi, Katsumi Nomizu, *Foundations of Differential Geometry, Volume I*, ISBN 0-471-15733-3, John Wiley and Sons, Inc., 1996.
- [Mac Lane, 1998] Saunders Mac Lane, *Categories for the Working Mathematician, 2nd edition*, ISBN 0-387-98403-8, Springer-Verlag, 1998.
- [Metz, 2018] Shmuel (Seymour J.) Metz, *Local Coordinate Spaces: a proposed unification of manifolds with fiber bundles, and associated machinery*, [arXiv:1801.05775, 2018](#)
- [Steenrod, 1999] Norman Steenrod, *The Topology Of Fibre Bundles*, ISBN 0-691-00548-6, Princeton University Press (seventh printing), 1999.