

Local Coordinate Spaces: a proposed unification of manifolds and fiber bundles, and associated machinery*

Shmuel (Seymour J.) Metz

September 15, 2026

Abstract

Manifolds and fiber bundles, while superficially different, have strong parallels; in particular, they are both defined in terms of equivalence classes of atlases or in terms of maximal atlases, with the atlases treated as mere adjuncts. This paper introduces the notions of an m-atlas and of a local coordinate space, and shows that special cases are equivalent to fiber bundles and manifolds. It defines categories of, e.g., atlases, and constructs functors among them.

Part I

Introduction

Historically, manifolds were formalized in terms of atlases based on Euclidean spaces and fiber bundles were formalized in terms of atlases based on product spaces, using either equivalence classes of atlases or maximal atlases. The concept of pseudogroups allowed unifying manifolds and manifolds with boundary. The definitions of fiber bundles and manifolds have strong parallels, and can be unified in a similar fashion; there are several ways to do so. The central part of this paper, [part VIII \(Local Coordinate Spaces\)](#) on page [113](#) defines an approach using categories and commutative diagrams that is designed for easy exposition at the possible expense of abstractness and generality. In particular, I have chosen to assume the Axiom of Choice (AOC).

This paper treats atlases as objects of interest in their own right, although it does not give them primacy. It introduces notions that are convenient for use here and others that, while not used here, may be useful for future work. [Section 1](#) below lists the more important of these.

*I wish to gratefully thank Walter Hoffman (z"l), Milton Parnes, Dr. Stanley H. Levy (z"l), the Mathematics department of Wayne State University, the Mathematics department of the State University of New York at Buffalo and others who guided my education.

Although this paper incidentally defines partial equivalents to manifolds and fiber bundles using model spaces and model atlases, it proposes the more general **Local Coordinate Space (LCS)** in order to explicitly reflect the role of the group in fiber bundles. [5] further develops some of the notions used here.

{Taxonomy}

A local coordinate space (LCS) is a space (total space) with some additional structure, including a coordinate model space and an atlas whose transition functions are restricted to morphisms of the coordinate model space; one can impose, e.g., differentiability restrictions, by appropriate choice of the coordinate category. There is an equivalent paradigm that avoids explicit mention of the total space by imposing a cocycle (compatibility) condition on the transition functions, but that approach is beyond the scope of this paper.

This paper defines functors among categories of atlases, categories of model spaces, categories of local coordinate spaces, categories of manifolds and categories of fiber bundles; it constructs more machinery than is customary in order to facilitate the presentation of those categories and functors.

Remark 0.1. The definitions of morphisms given here are nonstandard, but there are also definitions of classical morphisms that are much closer to the standard ones.

Parts II to VI present nomenclature and give basic results.

Part VII defines m-atlases, m-atlas morphisms and categories of them; lemma 13.20 proves that the defined categories are indeed categories.

lem:M-ATLiscat

Part VIII defines local coordinate spaces and categories of them; theorem 16.7 (LCS($\mathcal{M}, \Sigma$) is a category) on page 120 proves that the defined categories are indeed categories. Section 17 (Examples) on page 120 gives some examples of structures that can be represented as local coordinate spaces.

{the:LCSiscat}

part IX (Equivalence of manifolds) on page 122 and part X (Equivalence of fiber bundles) on page 146 present two examples of an LCS in detail, showing the equivalence of manifolds and fiber bundles with special cases of local coordinate spaces by explicitly exhibiting functors to and from local coordinate spaces.

{part:man}

{part:bun}

Remark 0.2. The unconventional definitions of manifold and fiber bundle morphisms are intended to make their relationship to local coordinate spaces more natural.

Many of the lemmata, theorems and corollaries in this paper are substantially identical to results that may be familiar to the reader. What is novel is the perspective and the material directly related to local coordinate spaces. The presentation assumes only a basic knowledge of Category Theory, such as may be found in the first chapter of [4] or [1].

JoyCat

{CftWM}

Many of the proofs are obvious and may be omitted from other versions of this paper.

1 New concepts and notation

{sec:new}

This paper introduces a significant number of new concepts and some modifications of the definitions for some conventional concepts. It also introduces some

notation of lesser importance. The following are the most important.

1. **Model space** and related concepts. A **model space**¹ is a topological space with a category of permissible open sets and mappings satisfying specified conditions. It is similar to a pseudo-group, with some important differences.
2. **Model topology** and **M-paracompactness**
3. **Signature, Σ -commutation** and related concepts. A signature is a means of characterising the domains and range of a multi-variable function. Σ -commutation is a naturality condition for a sequence of functions.
4. An **m-atlas** is a generalization of atlases in manifolds.
5. An **m-atlas morphism** is a generalization of a C^k map between differentiable manifolds. It differs in using both a function between the manifolds and a function between the coordinate spaces.
6. **Local Coordinate Space (LCS)** and related concepts
7. **Linear space and related concepts** A linear space is a subspace of a Banach space or of a Fréchet space that is suitable for defining coordinate patches in a manifold.
8. **Trivial C^k linear model space** and related concepts. A trivial C^k linear model space is a maximal model space with morphisms C^k maps between open subsets of a linear space.
9. G - ρ bundle atlases² and related concepts

Part II

Conventions

{part:conv}

An arrow with an Equal-Tilde ($A \xrightarrow[\phi]{\approx} B$) represents an isomorphism. One with a hook ($A \hookrightarrow_i B$) represents an inclusion map. One with a tail ($A \rightarrowtail_i B$) represents a monomorphism. One with a double head ($A \twoheadrightarrow_{\pi} B$) represents a surjection.

When a superscript ends in -1 , e.g., f^{i-1} , it is to be taken as function inverse rather than subtraction of 1.

All diagrams shown are commutative; none are exact.

A definition of a base term and several related terms may specify modifiers in parentheses in the base definition and then give the definitions for each modifier; usually the modifiers will be restrictions and the base definition will not be repeated. E.g., “ **f** is a (semi-strict, strict) prestructure morphism of P^1 to P^2 iff” is

¹The phrase has been used before, but with a different meaning.

²Similar to coordinate bundles

When a definition defines a base propositional function and variant propositional functions with a qualifier given as a superscript or underset, then the form $\text{base}^{(qualifiers)}$ will refer to either $\text{base}^{qualifier}$ or $\text{base}_{(qualifiers)}$ and the form $\text{base}_{(qualifiers)}$ will

refer to either $\underset{qualifier}{base}$ or $base$, e.g., $\text{isAtI}_{Ar}^{(classic)}$ refers to either isAtI_{Ar} , $\text{isAtI}_{Ar}^{classic}$,
 isAtI_{Ar} , $\text{isAtI}_{Ar}^{classic}$, $\text{isAtI}_{Ar}^{(semi-strict,strict)}$ or $\text{isAtI}_{Ar}^{classic}$.
 semi-strict semi-strict strict strict

The form $\text{base}^{\text{qualifier}(\text{qualifiers})}$ will refer to either $\text{base}^{\text{qualifier}}$ or to the variant $\text{base}^{\text{classic}(\text{proper})}$ with one or more of the additional qualifiers in the superscript, e.g., $\text{isAt}^{\text{strict}}$

refers to either $\text{isAtI}_{\text{Ar}}^{\text{classic}}$ or $\text{isAtI}_{\text{Ar}}^{\text{classic,proper}}$.

Alternatively, a definition may specify a numbered list of alternatives, and subsequently specify additional numbered lists with items corresponding to those in the first list, e.g., $\mathbf{f}$ is also a

1. full
2. semi-maximal
3. maximal
4. full semi-maximal
5. full maximal

E^1 - E^2 C^k morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

1. abbreviation for full
2. abbreviation for semi-maximal
3. abbreviation for maximal
4. abbreviation for full semi-maximal
5. abbreviation for full maximal

iff

1. definition for full
2. definition for semi-maximal
3. definition for maximal
4. definition for full semi-maximal
5. definition for full maximal

A Corollary, Lemma or Theorem that applies to related terms defined with the above convention will specify the modifiers in parentheses to indicate that it applies to the base term and to the variant terms, e.g., “a (semi-strict, strict) E^i - E^{i+1} m-atlas morphism” If it applies only to more restrictive terms then it will specify the first relevant modifier followed by the remaining relevant modifiers in parentheses, e.g.,

“If $\mathcal{S}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}'^i$ and f^i is a semi-strict (strict) prestructure morphism”.

Blackboard bold upper case will denote specific sets, e.g., the Naturals.

Bold lower case italic letters will refer to sets, sequences and tuples of functions, e.g., $f \stackrel{\text{def}}{=} (f_1, f_2)$.

Bold lower case Latin letters will refer to sequence valued functions of sequences and tuple valued functions of tuples, e.g., **range** yields the sequence of ranges of a sequence of functions.

Bold upper case italic letters will refer to sequences or tuples, e.g., $A = (x, y, z)$, to sets of them, to sets of topological spaces or to sets of open sets.

Bold upper case script letters will refer to sequences of categories, e.g.,

$\mathcal{A} \stackrel{\text{def}}{=} (\mathcal{A}_\alpha, \alpha \in A)$.

Fraktur will refer to topologies and to topology-valued functions, e.g., $\mathfrak{Top}$.

Functions have a range, domain and relation, not just a relation. Unless otherwise stated, they are assumed or proven to be continuous.

Groups are assumed to be topological groups. The ambiguous notation x^{-1} will be used when it is obvious from context what the group operation $\star$ and the group identity 1_G are.

Lower case Greek letters other than π, ρ, σ, ϕ and ψ will refer to ordinals, possibly transfinite, and to formal labels. A letter with a Greek superscript and a letter with a Latin or numeric superscript always refer to distinct variables.

Lower case π will refer to a projection operator

Lower case ρ will refer to a continuous effective group action, i.e., a continuous representation of a group in a homeomorphism group.

Lower case σ will refer to a sequence of ordinals, referred to as a signature.

Lower case ϕ will refer to a coordinate function.

Lower case italic and Latin letters will refer to

1. elements of a set or sequence

2. formal labels

When the letters u and v , with or without a superscript, refer to an element of a set then they may also be used as formal labels for sets associated with that element.

3. functions

4. morphisms of a category

5. natural numbers

6. objects of a category

7. ordinal numbers

Upper case Greek letters other than Σ may refer to

1. ordinal used as the limit of a sequence of consecutive ordinals, e.g., $x_\alpha, \alpha \leq A$
2. ordinal used as the order type of a sequence of consecutive ordinals, e.g., $x_\alpha, \alpha < A$

Upper case Σ will refer to a sequence of signatures

Upper case Latin letters will refer to

1. Natural numbers
2. Topological spaces
3. Open sets
4. Formal labels for elements of a sequence or tuple of functions, e.g., f_E might be $f_0: E_1 \longrightarrow E_2$.

Upper case Script Latin letters will refer to categories and functors.

Upright Latin letters will be used for long names. In particular,

1. Ar , Hom , Mor and Ob are the conventional set-valued functions on small categories:

$\text{Ar}(\mathcal{C})$ All morphisms of $\mathcal{C}$

$\text{Hom}_{\mathcal{C}}(a, b)$ All morphisms of $\mathcal{C}$ from a to b

$\text{Mor}(\mathcal{C})$ All morphisms of $\mathcal{C}$. Synonymous with $\text{Ar}(\mathcal{C})$.

$\text{Ob}(\mathcal{C})$ All objects of $\mathcal{C}$

They will never be applied to large categories in this paper.

2. **Set** and **Top** are the conventional categories of sets and topological spaces.
3. The letters E , G , X and Y will be used for formal labels.

The term C^k includes C^∞ (smooth) and C^ω (analytic).

This paper uses the term morphism in preference to arrow, but uses the conventional Ar .

This paper uses the terms empty, null and void interchangeably.

The term sequence without an explicit reference to $\mathbb{N}$ will refer to a general ordinal sequence, possibly transfinite.

Sequence numbering, unlike tuple numbering, starts at 0 and the exposition assumes a von Neumann definition of ordinals, so that $\alpha \in \beta \equiv \alpha < \beta$.

Except where explicitly stated otherwise, all categories mentioned are small categories with underlying sets, but the morphisms will often not be set functions between the objects and there will not always be a forgetful functor to **Set** or **Top**. By abuse of language no distinction will consistently be made between a category

$\mathcal{A}$ of topological spaces and the concrete category $(\mathcal{A}, \mathcal{U})$ over **Top**. Similarly, no distinction will consistently be made among the object $U \in \text{Ob}(\mathcal{A})$, the topological space $\mathcal{U}(U)$ and the underlying set.

The notation G^V will refer only to the set of continuous functions from V to G , never to the set of all functions from V to G .

When defining a category, the ordered pair $(\mathbf{O}, \mathbf{M})$ refers to the smallest concrete category over **Set** or **Top** whose objects are the elements in $\mathbf{O}$, whose morphisms include all of the elements of $\mathbf{M}$ and whose morphisms from $o^1 \in \mathbf{O}$ to $o^2 \in \mathbf{O}$ are functions $f: o^1 \longrightarrow o^2$ and whose composition is function composition.

When defining a category, the ordered triple $(\mathbf{O}, \mathbf{M}, C)$ refers to the small category whose objects are in $\mathbf{O}$, whose morphisms are in $\mathbf{M}$, whose Hom is

$$\text{Hom}_{(\mathbf{O}, \mathbf{M}, C)}(o_1 \in \mathbf{O}, o_2 \in \mathbf{O}) \stackrel{\text{def}}{=} \{f, o_1, o_2\} \in \mathbf{M} \quad (1.1)$$

and whose composition is C .

By abuse of language I may write “ $\mathcal{S}$ ” for $\text{Ob}(\mathcal{S})$, “ $A \in \mathcal{A}$ ” for $A \in \text{Ob}(\mathcal{A})$, “ $A \subset \mathcal{A}$ ” for $A \subset \text{Ob}(\mathcal{A})$, “ $A \in \mathcal{A} \subset B \in \mathcal{B}$ ” for “the underlying set of A is contained in the underlying set of B and the inclusion $i: x \in A \hookrightarrow x \in B$ is a morphism” and “ $f: A \longrightarrow B$ ” for $f \in \text{Hom}_{\mathcal{C}}(A, B)$, where $\mathcal{C}$ is understood by context.

By abuse of language I shall use the same nomenclature for sequences and tuples, and will use parentheses around a single expression both for grouping and for a tuple with a single element; the intent should be clear from context.

By abuse of language I will omit parentheses around the operands of Functors when they can be assumed by context.

By abuse of language I shall use the $\times$ and $\times$ symbols for both Cartesian products of sets and Cartesian products of functions on those sets.

By abuse of language, and assuming AOC, I shall refer to some sets as ordinal sequences, e.g., “ $(C_\alpha, \alpha \in A)$ ” for “ $\{C_\alpha \mid \alpha \in A\}$ ”, in cases where the order is irrelevant.

Where a proof requires proving intermediate propositions, a bold leadin to a paragraph will indicate such a proposition, e.g.

Intermediate proposition 1: Intermediate proof 1.

Intermediate proposition 2: Intermediate proof 2.

By abuse of language, I may omit universal quantifiers in cases where the intent is clear.

In some cases I define a notion similar to a conventional notion and also need to refer to the conventional notion. In those cases I prefix a letter or phrase to the term, e.g., m-paracompact versus paracompact.

Part III

General notions

{part: notions}

This part of the paper describes nomenclature used throughout the paper. In some cases this reflects new nomenclature or notions, in others it simply makes a choice from among the various conventions in the literature.

{def: catprop}

Definition 1.1 (Operations on categories). If $\mathcal{C}$ is a category then

1. $x \in^{\text{Ob}} \mathcal{C} \equiv x \in \text{Ob}(\mathcal{C})$
2. $x \in^{\text{Ar}} \mathcal{C} \equiv x \in \text{Ar}(\mathcal{C})$
3. If $\mathcal{C}$ is a concrete category over **Set** or **Top** and $o \in^{\text{Ob}} \mathcal{C}$, then $\mathbf{Set}_{\mathcal{C}}(o)$ is the underlying set of o . By abuse of language we will write $\mathbf{Set}(o)$ when the category is understood from context.
4. If $\mathcal{C}$ is a concrete category over **Top** and $o \in^{\text{Ob}} \mathcal{C}$, then $\text{Top}_{\mathcal{C}}(o)$ is the underlying topological space of o . By abuse of language we will write $\text{Top}(o)$ when the category is understood from context.

If $\mathcal{S}$ and $\mathcal{T}$ are categories then $\mathcal{S} \subseteq^{\text{cat}} \mathcal{T}$ iff $\mathcal{S}$ is a subcategory of $\mathcal{T}$ and $\mathcal{S} \subseteq^{\text{full-cat}} \mathcal{T}$ iff $\mathcal{S}$ is a full subcategory of $\mathcal{T}$.

If $\mathcal{S}$ and $\mathcal{T}$ are categories then the category union of $\mathcal{S}$ and $\mathcal{T}$, abbreviated $\mathcal{S} \cup^{\text{cat}} \mathcal{T}$, is the category whose objects are in $\mathcal{S}$ or in $\mathcal{T}$ and whose morphisms are in $\mathcal{S}$ or in $\mathcal{T}$.

{def: Id}

Definition 1.2 (Identity). $\text{Id}_{\mathcal{S}}$ is the identity function on the space $\mathcal{S}$,

Id_o is the identity morphism for the object o ³,

Id_V^U , for $U \subseteq V$, is the inclusion map⁴. $\text{Id}_{U,V}$ is synonymous with Id_V^U .

$\text{Id}_{\mathcal{C}}$ is the identity functor on the category $\mathcal{C}$.

$\mathbf{Id}_{\mathcal{S}^i}$, $i = 1, 2$, is the sequence of identity functions for the elements of the sequence $\mathcal{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_{\alpha}^i, \alpha < A)$. Let $\mathcal{S}^1 \subseteq^{\text{O}} \mathcal{S}^2$. Then $\mathbf{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}$ is the sequence of inclusion maps $(\text{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}, \alpha < A)$ for the elements of the sequences $\mathcal{S}^i$. $\mathbf{Id}_{\mathcal{S}^1, \mathcal{S}^2}$ is synonymous with $\mathbf{Id}_{\mathcal{S}^2}^{\mathcal{S}^1}$. A similar definition applies for tuples, except in a context where they are treated as objects of a category.

$\mathbf{Id}_{(A, \dots, Z)}$ is the tuple of identity maps $(\text{Id}_A, \dots, \text{Id}_Z)$. $\mathbf{Id}_{(A^1, \dots, Z^1)}^{(A^2, \dots, Z^2)}$ is the tuple of inclusion maps $(\text{Id}_{A^2}^{A^1}, \dots, \text{Id}_{Z^2}^{Z^1})$.

³The object is often expressed as a tuple, e.g., $\text{Id}_{(A,B)}$ is the identity morphism for the object (A, B)

⁴ U need not have the subspace topology.

Definition 1.3 (Images). $f[U] \stackrel{\text{def}}{=} \{f(x) | x \in U\}$ is the image of U under f and $f^{-1}[V] \stackrel{\text{def}}{=} \{x | f(x) \in V\}$ is the inverse image of V under f .

Remark 1.4. This notation, adopted from [2], avoids the ambiguity in the traditional $f(U)$ and $f^{-1}(V)$. {GenTop}

Definition 1.5 (Projections). π_α is the projection function that maps a sequence into element α of the sequence. π_i is also the projection function that maps a tuple into element i of the tuple. {projections}

Definition 1.6 (Topological category). A topological category is a small subcategory of **Top** or its concrete category over **Set**. {def:topcat}

$\mathcal{T}$ is a full topological category iff it is a topological category and whenever $U^i, V^i \in \mathcal{T}, i = 1, 2, V^i \stackrel{\text{subsp}}{\subseteq} U^i, f: U^1 \longrightarrow U^2 \in \mathcal{T}$ and $f[V^1] \subseteq V^2$ then $f: V^1 \longrightarrow V^2 \in \mathcal{T}$. {lem:topInc}

Lemma 1.7 (Inclusions in topological categories are morphisms). Let $\mathcal{T}$ be a full topological category, $S^i \in \mathcal{T}, i = 1, 2$, and $S^1 \stackrel{\text{subsp}}{\subseteq} S^2$. Then $\text{Id}_{S^2}^{S^1}$ is a morphism of $\mathcal{T}$.

Proof. 1. $U^i \stackrel{\text{def}}{=} S^2 \in \mathcal{T}, i = 1, 2$, by hypothesis.

2. $V^1 \stackrel{\text{def}}{=} S^1 \in \mathcal{T}$ and $V^2 \stackrel{\text{def}}{=} S^2 \in \mathcal{T}$ by hypothesis.

3. $V^i \stackrel{\text{subsp}}{\subseteq} U^i$,

4. $f \stackrel{\text{def}}{=} \text{Id}_{S^2}: U^1 \longrightarrow U^2 \in \mathcal{T}$.

5. $f[V^1] = V^1 \stackrel{\text{subsp}}{\subseteq} V^2$.

Thus $\text{Id}_{S^2}^{S^1} = f: V^1 \longrightarrow V^2 \in \mathcal{T}$ by definition 1.6. □

{def:topcat}
{def:topLocal}

Definition 1.8 (Local morphisms). Let $\mathcal{S}^i, i = 1, 2$, be a full topological category and $S^i \in \mathcal{S}^i$. A continuous function $f: S^1 \longrightarrow S^2$ is a local $\mathcal{S}^1$ - $\mathcal{S}^2$ morphism of S^1 to S^2 iff $\mathcal{S}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}^2$ and for every $u \in S^1$ there is an open neighborhood U_u for u and an open neighborhood V_u for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathcal{S}^2$.

Remark 1.9. The condition $f: U_u \longrightarrow V_u \in \mathcal{S}^2$ ensures that $U_u \in \mathcal{S}^1$ and $V_u \in \mathcal{S}^2$.

Let $\mathcal{T}$ be a full topological category and $S^i \in \mathcal{T}, i = 1, 2$. A continuous function $f: S^1 \longrightarrow S^2$ is a local $\mathcal{T}$ morphism of S^1 to S^2 iff it is a local $\mathcal{T}$ - $\mathcal{T}$ morphism of S^1 to S^2 .

{lem:topLocal}

Lemma 1.10 (Local morphisms). Let $\mathcal{T}^i, i \in [1, 3]$, be a full topological category, $S^i \in^{\text{Ob}} \mathcal{T}^i$ and $\mathcal{T}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^{i+1}$.

If $f^i: S^i \longrightarrow S^{i+1} \in^{\text{Ar}} \mathcal{T}^{i+1}$ then f^i is a local $\mathcal{T}^i$ - $\mathcal{T}^{i+1}$ morphism of S^i to S^{i+1} .

Proof. Let $u \in S^i$ and $v \stackrel{\text{def}}{=} f^i(u) \in S^{i+1}$. S^i is an open for u , S^{i+1} is an open neighborhood of v and $f^i: S^i \longrightarrow S^{i+1} \in^{\text{Ar}} \mathcal{T}^{i+1}$ by hypothesis. $\square$

If each $f^i: S^i \longrightarrow S^{i+1}$, is a local $\mathcal{T}^i$ - $\mathcal{T}^{i+1}$ morphism of S^i to S^{i+1} then $f^2 \circ f^1: S^1 \longrightarrow S^3$ is a local $\mathcal{T}^1$ - $\mathcal{T}^3$ morphism of S^1 to S^3 .

Proof. Since $\mathcal{T}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^2$ and $\mathcal{T}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^3$, $\mathcal{T}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^3$. Let $u \in S^1$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist an open neighborhood U_u for u , open neighborhoods V_u, V'_v for v and an open neighborhood W_v of w such that $f^1[U_u] \subseteq V_u$, $f^1: U_u \longrightarrow V_u$ is a morphism of $\mathcal{T}^2$, $f^2[V'_v] \subseteq W_v$ and $f^2: V'_v \longrightarrow W_v$ is a morphism of $\mathcal{T}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_u \cap V'_v \neq \emptyset$, $\hat{V}_u$ is an open neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{i-1}[\hat{V}_u]$ is an open neighborhood of u . $f^1: \hat{U}_u \longrightarrow \hat{V}_u$ and $f^2: \hat{V}_u \longrightarrow W_v$ are morphisms of $\mathcal{T}^3$ by definition 1.6 (Topological category) on page 9 and thus $f^2 \circ f^1: \hat{U}_u \longrightarrow W_v$ is a morphism of $\mathcal{T}^3$. $\square$

{def:topcat}

{cor:topLocal}

Corollary 1.11 (Local morphisms). Let $\mathcal{T}^i, i = 1, 2$, be a full topological category, $\mathcal{T}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{T}^{i+1}$, $S^i \in^{\text{Ob}} \mathcal{T}^i$ and $S^1 \subseteq S^2$. Then $\text{Id}_{S^2}^{S^1}$ is a local $\mathcal{T}^1$ - $\mathcal{T}^2$ morphism of S^1 to S^2 and Id_{S^i} is a local $\mathcal{T}$ morphism of S^i to S^i .

Proof. $S^1 \in^{\text{Ob}} \mathcal{T}^2$ because $S^1 \in^{\text{Ob}} \mathcal{T}^1$ and $\mathcal{T}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{T}^2$, $S^2 \in^{\text{Ob}} \mathcal{T}^2$ by hypothesis and $S^1 \subseteq S^2$ by hypothesis, so $\text{Id}_{S^2}^{S^1} \in^{\text{Ar}} \mathcal{T}^2$ by lemma 1.7.

{lem:topInc}

$\text{Id}_{S^i} \stackrel{\text{def}}{=} \text{Id}_{S^i}^{S^i}$. $\square$

Definition 1.12 (Sequence functions). Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\alpha, \alpha < A)$ be a sequence of functions. Then

$$\mathbf{domain}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{domain}(s_\alpha), \alpha < A) \quad (1.2)$$

$$\mathbf{range}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{range}(s_\alpha), \alpha < A) \quad (1.3)$$

Let $\mathbf{T} \stackrel{\text{def}}{=} (t_\alpha, \alpha < A)$ be a sequence of functions with $\mathbf{range}(\mathbf{S}) = \mathbf{domain}(\mathbf{T})$. Then their composition is the sequence $\mathbf{T} \circ \mathbf{S} \stackrel{\text{def}}{=} (t_\alpha \circ s_\alpha, \alpha < A)$,

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\gamma, \gamma \leq \Gamma)$, then these functions extract information about the sequence:

$$\mathbf{head}(\mathbf{S}, \Omega) \stackrel{\text{def}}{=} (s_\gamma, \gamma < \Omega) \quad (1.4)$$

$$\text{head}(\mathbf{S}) \stackrel{\text{def}}{=} \text{head}(\mathbf{S}, \Gamma) \quad (1.5)$$

$$\text{length0}(\mathbf{S}) \stackrel{\text{def}}{=} \Gamma \quad (1.6)$$

$$\text{tail}(\mathbf{S}) \stackrel{\text{def}}{=} S_\Gamma \quad (1.7)$$

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_\gamma, \gamma < \Gamma)$, then

$$\text{length}(\mathbf{S}) \stackrel{\text{def}}{=} \Gamma \quad (1.8)$$

Remark 1.13. If $\text{length0}(\mathbf{S})$ is defined then $\text{length}(\mathbf{S}) = \text{length0}(\mathbf{S}) + 1$. $\text{length0}(\mathbf{S})$ is the ordinal type of $\text{head}(\mathbf{S})$, not the ordinal type of $\mathbf{S}$.

Let $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$ and $\mathcal{T} \stackrel{\text{def}}{=} (\mathcal{T}_\alpha, \alpha < A)$ be sequences of categories. Then $\mathcal{S}$ is a subcategory sequence of $\mathcal{T}$, abbreviated $\mathcal{S} \subseteq^{\text{cat}} \mathcal{T}$, iff every category in $\mathcal{S}$ is a subcategory of the corresponding category in $\mathcal{T}$, i.e., $\left(\bigvee_{\alpha < A}\right) \mathcal{S}_\alpha \subseteq^{\text{cat}} \mathcal{T}_\alpha$, and $\mathcal{S}$ is a full subcategory sequence of $\mathcal{T}$, abbreviated $\mathcal{S} \subseteq^{\text{full-cat}} \mathcal{T}$, iff every category in $\mathcal{S}$ is a full subcategory of the corresponding category in $\mathcal{T}$, i.e., $\left(\bigvee_{\alpha < A}\right) \mathcal{S}_\alpha \subseteq^{\text{full-cat}} \mathcal{T}_\alpha$.

The category sequence union of $\mathcal{S}$ and $\mathcal{T}$, abbreviated $\mathcal{S} \cup^{\text{cat}} \mathcal{T}$, is the sequence of category unions of corresponding categories in $\mathcal{S}$ and $\mathcal{T}$, i.e., $(\mathcal{S}_\alpha \cup^{\text{cat}} \mathcal{T}_\alpha)$.

{lem:seqfunc}

Lemma 1.14 (Sequence functions). Let $\mathbf{f}^i \stackrel{\text{def}}{=} (f_\alpha^i, \alpha < A)$, $i \in [1, 3]$, be sequences of functions with $\mathbf{domain}(\mathbf{f}^2) = \mathbf{range}(\mathbf{f}^1)$ and $\mathbf{domain}(\mathbf{f}^3) = \mathbf{range}(\mathbf{f}^2)$. Then $(\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 = \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1)$.

Proof.

$$\begin{aligned} (\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 &= ((f_\alpha^3 \circ f_\alpha^2) \circ f_\alpha^1, \alpha < A) \\ &= (f_\alpha^3 \circ (f_\alpha^2 \circ f_\alpha^1), \alpha < A) \\ &= \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1) \end{aligned}$$

□

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha, \alpha < A)$ be a sequence of functions, $\mathbf{D} = \mathbf{domain}(\mathbf{f})$ and $\mathbf{R} = \mathbf{range}(\mathbf{f})$. Then $\mathbf{Id}_\mathbf{R}$ is a left $\circ$ identity for $\mathbf{f}$ and $\mathbf{Id}_\mathbf{D}$ is a right $\circ$ identity for $\mathbf{f}$.

Proof.

$$\begin{aligned} \mathbf{Id}_\mathbf{R} \circ \mathbf{f} &= (\mathbf{Id}_{\mathbf{range}(f_\alpha)} \circ f_\alpha, \alpha < A) \\ &= (f_\alpha, \alpha < A) \\ &= \mathbf{f} \end{aligned}$$

$$\begin{aligned}
\mathbf{f} \overset{\circ}{\circ} \mathbf{Id}_D &= (f_\alpha \circ \text{Id}_{\text{domain}(f_\alpha)}, \alpha < A) \\
&= (f_\alpha, \alpha < A) \\
&= \mathbf{f}
\end{aligned}$$

□

Definition 1.15 (Tuple functions). Let $\mathbf{S} \stackrel{\text{def}}{=} (s_n, n \in [1, N])$ be a tuple of functions. Then

$$\mathbf{domain}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{domain}(s_n), n \in [1, N]) \quad (1.9)$$

$$\mathbf{range}(\mathbf{S}) \stackrel{\text{def}}{=} (\text{range}(s_n), n \in [1, N]) \quad (1.10)$$

Let $\mathbf{T} \stackrel{\text{def}}{=} (t_n, n \in [1, N])$ be a tuple of functions with $\mathbf{range}(\mathbf{S}) = \mathbf{domain}(\mathbf{T})$. Then their composition is the tuple $\mathbf{T} \overset{\circ}{\circ} \mathbf{S} \stackrel{\text{def}}{=} (t_n \circ s_n, n \in [1, N])$

Let $\mathbf{S} \stackrel{\text{def}}{=} (s_m, m \in [1, M])$ and $\mathbf{T} \stackrel{\text{def}}{=} (t_n, n \in [1, N])$ be tuples. Then the following are tuple functions

$$\text{head}(\mathbf{S}, I) \stackrel{\text{def}}{=} (s_m, m \in [1, I]) \quad (1.11)$$

$$\text{head}(\mathbf{S}) \stackrel{\text{def}}{=} \text{head}(\mathbf{S}, M - 1) \quad (1.12)$$

$$\text{tail}(\mathbf{T}, I) \stackrel{\text{def}}{=} (t_n, n \in [I, N]) \quad (1.13)$$

$$\text{tail}(\mathbf{T}) \stackrel{\text{def}}{=} t_N \quad (1.14)$$

$$\text{join}(\mathbf{S}, \mathbf{T}) \stackrel{\text{def}}{=} (s_1, \dots, s_M, t_1, \dots, t_N) \quad (1.15)$$

{lem:tupfunc}

Lemma 1.16 (Tuple functions). Let $\mathbf{f} \stackrel{\text{def}}{=} (f_n, n \in [1, N])$ be a tuple of functions, $\mathbf{D} \stackrel{\text{def}}{=} \mathbf{domain}(\mathbf{f})$ and $\mathbf{R} \stackrel{\text{def}}{=} \mathbf{range}(\mathbf{f})$. Then $\mathbf{Id}_R$ is a left $\overset{\circ}{\circ}$ identity for $\mathbf{f}$ and $\mathbf{Id}_D$ is a right $\overset{\circ}{\circ}$ identity for $\mathbf{f}$.

Proof.

$$\begin{aligned}
\mathbf{Id}_R \overset{\circ}{\circ} \mathbf{f} &= (\text{Id}_{\text{range}(f_\alpha)} \circ f_n, n \in [1, N]) \\
&= (f_n, n \in [1, N]) \\
&= \mathbf{f}
\end{aligned}$$

$$\begin{aligned}
\mathbf{f} \overset{\circ}{\circ} \mathbf{Id}_D &= (f_n \circ \text{Id}_{\text{domain}(f_n)}, n \in [1, N]) \\
&= (f_n, n \in [1, N]) \\
&= \mathbf{f}
\end{aligned}$$

□

{lem:tupcomp}

Lemma 1.17 (Tuple composition for unlabeled morphisms). Let

$\mathbf{f}^i \stackrel{\text{def}}{=} (f_n^i, n \in [1, N]), i \in [1, 3]$, be tuples of functions with $\mathbf{domain}(\mathbf{f}^2) = \mathbf{range}(\mathbf{f}^1)$ and $\mathbf{domain}(\mathbf{f}^3) = \mathbf{range}(\mathbf{f}^2)$. Then $(\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 = \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1)$.

Proof.

$$\begin{aligned} (\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1 &= ((f_n^3 \circ f_n^2) \circ f_n^1, n \in [1, N]) \\ &= (f_n^3 \circ (f_n^2 \circ f_n^1), n \in [1, N]) \\ &= \mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1) \end{aligned}$$

□

{def:labcomp}

Definition 1.18 (Tuple composition for labeled morphisms). Let $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o_1^i, o_2^i)$, $i = 1, 2$, be tuples such that each $\mathbf{f}^i$ is a sequence of functions or each $\mathbf{f}^i$ is a tuple of functions, $\mathbf{range}(\mathbf{f}^1) = \mathbf{domain}(\mathbf{f}^2)$ and $o_2^1 = o_1^2$. Then

$$\mathbf{M}^2 \overset{A}{\circ} \mathbf{M}^1 \stackrel{\text{def}}{=} (\mathbf{f}^2 \circ \mathbf{f}^1, o_1^1, o_2^2) \quad (1.16)$$

{lem:labcomp}

Lemma 1.19 (Tuple composition for labeled morphisms). Let $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o_1^i, o_2^i)$, $i \in [1, 3]$, be a tuple such that $\mathbf{f}^i$ is a sequence or tuple of functions, $\mathbf{range}(\mathbf{f}^i) = \mathbf{domain}(\mathbf{f}^{i+1})$ and $o_2^i = o_1^{i+1}$, $i = 1, 2$. Then

$$\mathbf{M}^3 \overset{A}{\circ} (\mathbf{M}^2 \overset{A}{\circ} \mathbf{M}^1) = (\mathbf{M}^3 \overset{A}{\circ} \mathbf{M}^2) \overset{A}{\circ} \mathbf{M}^1 \quad (1.17)$$

Proof. From definition 1.18 (Tuple composition for labeled morphisms),

lemma 1.14 (Sequence functions) on page 11 and lemma 1.16 (Tuple functions), we have

{def:labcomp}

{lem:tupfunc}

$$\begin{aligned} \mathbf{M}^3 \overset{A}{\circ} (\mathbf{M}^2 \overset{A}{\circ} \mathbf{M}^1) &= \mathbf{M}^3 \overset{A}{\circ} (\mathbf{f}^2 \circ \mathbf{f}^1, o_1^1, o_2^2) \\ &= (\mathbf{f}^3 \circ \mathbf{f}^2 \circ \mathbf{f}^1, o_1^1, o_2^3) \\ &= (\mathbf{f}^3 \circ \mathbf{f}^2, o_1^2, o_2^3) \overset{A}{\circ} \mathbf{M}^1 \\ &= (\mathbf{M}^3 \overset{A}{\circ} \mathbf{M}^2) \overset{A}{\circ} \mathbf{M}^1 \end{aligned}$$

□

Let $\mathbf{D}^i \stackrel{\text{def}}{=} \mathbf{domain}(\mathbf{f}^i)$ and $\mathbf{R}^i \stackrel{\text{def}}{=} \mathbf{range}(\mathbf{f}^i)$. Then

$$(\mathbf{Id}_{\mathbf{R}^i}, o_2^i, o_2^i) \overset{A}{\circ} \mathbf{M}^i = \mathbf{M}^i \quad (1.18)$$

$$\mathbf{M}^i \overset{A}{\circ} (\mathbf{Id}_{\mathbf{D}^i}, o_1^i, o_1^i) = \mathbf{M}^i \quad (1.19)$$

Proof.

$$\begin{aligned} (\mathbf{Id}_{R^i}, o_2^i, o_2^i) \overset{A}{\circ} \mathbf{M}^i &= (\mathbf{Id}_{R^i} \overset{\circ}{\circ} \mathbf{f}^i, o_1^i, o_2^i) \\ &= (\mathbf{f}^i, o_1^i, o_2^i) \\ &= \mathbf{M}^i \end{aligned}$$

$$\begin{aligned} \mathbf{M}^i \overset{A}{\circ} (\mathbf{Id}_{D^i}, o_1^i, o_1^i) &= (\mathbf{f}^i \overset{\circ}{\circ} \mathbf{Id}_{D^i}, o_1^i, o_2^i) \\ &= (\mathbf{f}^i, o_1^i, o_2^i) \\ &= \mathbf{M}^i \end{aligned}$$

□

{def:Cart}

Definition 1.20 (Cartesian product of sequence). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence and $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha < A)$ be a sequence of functions, then $\bigtimes \mathbf{S}^i \stackrel{\text{def}}{=} \bigtimes_{\alpha < A} S_\alpha^i$ is the generalized Cartesian product of the sequence $\mathbf{S}^i$ and $\bigtimes \mathbf{f}: \mathbf{S}^1 \longrightarrow \mathbf{S}^2 \stackrel{\text{def}}{=} \bigtimes_{\alpha < A} f_\alpha$ is the generalized Cartesian product of the function sequence $\mathbf{f}$.

Definition 1.21 (underline). Let $\mathbf{S}^1 \stackrel{\text{def}}{=} (S_\alpha^1, \alpha \leq A)$, $\mathbf{S}^2 \stackrel{\text{def}}{=} (S_\alpha^2, \alpha \leq A)$ be sequences and $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha \leq A)$ be a sequence of functions, then

$$\underline{f}: \text{head}(S_1) \longrightarrow \text{head}(S_2) \stackrel{\text{def}}{=} \bigtimes \text{head}(\mathbf{f}) = \bigtimes_{\alpha < A} f_\alpha \quad (1.20)$$

is the function mapping $(s_\alpha \in S_\alpha^1, \alpha < A)$ into $(f_\alpha(s_\alpha), \alpha < A)$.

Definition 1.22 (Head and tail compositions). Let

$f^1: (S_\alpha^1, \alpha < A) \longrightarrow S_A^1$, $f^2: (S_\alpha^2, \alpha < A) \longrightarrow S_A^2$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha \leq A)$. Then (see figs. 1 and 2)

{fig:fg,fig:gf}

$$f^2 \odot \mathbf{g} \stackrel{\text{def}}{=} f^2 \circ \underline{\mathbf{g}} \quad (1.21a)$$

$$\mathbf{g} \cdot f^1 \stackrel{\text{def}}{=} \text{tail}(\mathbf{g}) \circ f^1 \quad (1.21b)$$

Definition 1.23 (Topology functions). Let S and T be a topological spaces and Y a subset of S . Then

1. $S \overset{\text{subsp}}{\subseteq} T$ iff S is a subspace of T .
2. $S \overset{\text{open-subsp}}{\subseteq} T$ iff S is an open subspace of T .
3. **Set**(S) is the underlying set of S .




Figure 1: $f^2 \odot g$

{fig:fg}




Figure 2: $g \cdot f^1$

{fig:gf}

4. $\mathfrak{Top}(S)$ is the topology of S .
5. $\mathfrak{Top}(Y, S) \stackrel{\text{def}}{=} \{U \cap Y \mid U \in \mathfrak{Top}(S)\}$ is the relative topology of Y .
6. $\text{Top}(Y, S) \stackrel{\text{def}}{=} (Y, \mathfrak{Top}(Y, S))$ is Y with the relative topology.
7. $S \stackrel{\text{def}}{=} \{(U, \mathfrak{Top}(U, S)) \mid U \in \mathfrak{Top}(S) \setminus \{\emptyset\}\}$ is the set of all non-null open subspaces of S .

Let $\mathbf{S}$ be a set of topological spaces. Then $\mathbf{S} \stackrel{\text{def}}{=} \bigcup_{S \in \mathbf{S}} S$ is the set of open subspaces in $\mathbf{S}$.

Let S and T' be spaces, $T \subseteq T'$ be a subspace and $f: S \longrightarrow T$ a function. Then $f: S \longrightarrow T' \stackrel{\text{def}}{=} \text{Id}_{T'}^T \circ f$ is f considered as a function from S to T' .

Let S' and T' be spaces, $S \subseteq S'$, $T \subseteq T'$ be subspaces and $f': S' \longrightarrow T'$ a function such that $f'[S] \subseteq T$. Then $f': S \longrightarrow T$, also written $f' \upharpoonright_{S,T}$, is $f' \upharpoonright_S$ considered as a function from S to T .

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha \leq A)$, $i = 1, 2$, be a sequence of spaces, $\mathbf{S}^1 \subseteq \mathbf{S}^2$ and $f^2: \text{head}(\mathbf{S}^2) \longrightarrow \text{tail}(\mathbf{S}^2)$ a function. $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)} \stackrel{\text{def}}{=} f^2 \upharpoonright_{\times \text{head}(\mathbf{S}^1)}$. If $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)} [\times \text{head}(\mathbf{S}^1)] \subseteq \text{tail}(\mathbf{S}^1)$ then $f^2: \text{head}(\mathbf{S}^1) \longrightarrow \text{tail}(\mathbf{S}^1)$, also written $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1), \text{tail}(\mathbf{S}^1)}$, is $f^2 \upharpoonright_{\text{head}(\mathbf{S}^1)}$ considered as a function from $\times \text{head}(\mathbf{S}^1)$ to $\text{tail}(\mathbf{S}^1)$.

Definition 1.24 (Truth space). $\text{False} \stackrel{\text{def}}{=} \emptyset$, $\text{True} \stackrel{\text{def}}{=} \{\emptyset\}$, the truth set is $\mathbb{T} \stackrel{\text{def}}{=} \{\text{False}, \text{True}\}$, $\mathfrak{Truthtop} \stackrel{\text{def}}{=} \{\emptyset, \{\text{True}\}, \mathbb{T}\}$ and the truth space $\text{Truthspace} \stackrel{\text{def}}{=} (\mathbb{T}, \mathfrak{Truthtop})$ is $\mathbb{T}$ with $\{\text{True}\}$ and $\mathbb{T}$ as the only nonvoid open sets

Definition 1.25 (Truth category). Let $f_0 = \text{Id}_{\text{Truthspace}}$ and $f_1: \text{Truthspace} \longrightarrow \text{Truthspace}$

be the constant true function, i.e., value True . The truth category is $\mathcal{T}ruth \stackrel{\text{def}}{=} (\text{Truthspace}, \{f_0, f_1\})$. The truth model space is

Truthspace $\stackrel{\text{def}}{=} (\text{Truthspace}, \mathcal{T}ruth)$.

Definition 1.26 (Constraint functions). A constraint function is a continuous function with range Truthspace or a model function with range **Truthspace**.

Definition 1.27 (Sequence inclusion). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ and $\mathbf{T} \stackrel{\text{def}}{=} (T_\alpha, \alpha < A)$ be sequences. $\mathbf{S} \stackrel{\circ}{\in} \mathbf{T}$ iff $(\bigvee_{\alpha < A} S_\alpha \in T_\alpha \text{ or } (\bigvee_{\alpha < A} S_\alpha \stackrel{\text{Ob}}{\in} T_\alpha$.

Lemma 1.28 (Sequence inclusion). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ be a sequence,

$\mathcal{T}^i \stackrel{\text{def}}{=} (\mathcal{T}_\alpha^i, \alpha < A)$, $i = 1, 2$, a sequence of categories, $\mathcal{T}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{T}^2$ and $\mathbf{S} \stackrel{\circ}{\in} \mathcal{T}^1$ Then $\mathbf{S} \stackrel{\circ}{\in} \mathcal{T}^2$.

Proof. If $(\bigvee_{\alpha < A} S_\alpha \stackrel{\text{Ob}}{\in} \mathcal{T}_\alpha^1$ then $(\bigvee_{\alpha < A} S_\alpha \stackrel{\text{Ob}}{\in} \mathcal{T}_\alpha^2$. □

Part IV

Model spaces and allied notions

Let S be a topological space. We need to formalize the notions of an open cover by sets that are “well behaved” in some sense, e.g., convex, sufficiently small, and of “well behaved” functions among those sets, e.g., preserving fibers, smooth. We do this by associating a category of acceptable sets and functions.

{part:modeldef}

Remark 1.29. Using pseudo-groups, as in [3, p. 1], would not allow restricting model neighborhoods to, e.g., convex sets.

{FoundDiffGeo1}

2 Model spaces

Definition 2.1 (Model spaces). Let S be a topological space and $\mathcal{S}$ a small category whose objects are open subsets of S and whose morphisms are continuous functions. $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is a model space for S iff

{def:model}

1. $\text{Ob}(\mathcal{S})$ is an open cover for S . Note that it need not be a basis for S . {mod:cover}
 2. $\text{Ob}(\mathcal{S})$ is closed under finite intersections. {mod:closed}
 3. The morphisms of $\mathcal{S}$ are continuous functions in S . {mod:continuous}
 4. If $f: A \rightarrow B$ is a morphism, $A' \in \mathcal{S} \subseteq A \in \mathcal{S}$, $B' \in \mathcal{S} \subseteq B \in \mathcal{S}$ and $f[A'] \subseteq B'$ then $f|_{A'}: A' \rightarrow B'$ is a morphism. {mod:restriction}
 5. If $A' \in \mathcal{S} \subseteq A \in \mathcal{S}$ then the inclusion map $\text{Id}_A^{A'}: A' \hookrightarrow A$ is a morphism. {mod:inclusion}
- Remark 2.2.* This is actually a consequence of item 4, but it is convenient to give it here. {mod:restriction}
6. Restricted sheaf condition: informally, consistent morphisms can be glued together. Whenever {mod:sheaf}
 - (a) U_α and V_α , $\alpha < A$, are objects of $\mathcal{S}$.
 - (b) $f_\alpha: U_\alpha \rightarrow V_\alpha$ are morphisms of $\mathcal{S}$.
 - (c) $U = \bigcup_{\alpha < A} U_\alpha$ is an object of $\mathcal{S}$.
 - (d) $V = \bigcup_{\alpha < A} V_\alpha$ is an object of $\mathcal{S}$.
 - (e) $f: U \rightarrow V$ is a continuous function and for every $\alpha < A$, f agrees with f_α on U_α

then f is a morphism of $\mathcal{S}$.

By abuse of language we write “ $U \in \mathbf{S}$ ” for “ $U \in \text{Cat}\mathbf{S}$ ” and “ $f \in \mathbf{S}$ ” for “ $f \in \text{Cat}\mathbf{S}$ ”.

Remark 2.3. A model space is similar to a pseudo-group, but has some important differences:

1. Open subsets of objects of $\mathcal{S}$ need not be objects of $\mathcal{S}$.
2. The morphisms of $\mathcal{S}$ need not be homeomorphisms or even 1-1, thus they need not have inverses.
3. The morphisms of $\mathcal{S}$ need not satisfy the sheaf condition, but only the restricted sheaf condition in item 6.

{mod:sheaf}

$\mathbf{S}$ is fine grained iff every open subset of an object of $\mathcal{S}$ is an object of $\mathcal{S}$. Note that it is not sufficient that $\text{Ob}(\mathcal{S})$ be a basis for S .

$\text{Top}(\mathbf{S}) \stackrel{\text{def}}{=} S = \pi_1(\mathbf{S})$ is the topological space of $\mathbf{S}$.

Let $\mathbf{S}^i$, $i = 1, 2$, be a model space. $\text{Cat}(\mathbf{S}^i) \stackrel{\text{def}}{=} \mathcal{S}^i = \pi_2(\mathbf{S}^i)$ is the category of model space $\mathbf{S}^i$. By abuse of language we write $\mathbf{S}^1 \stackrel{\text{subsp}}{\subseteq} \mathbf{S}^2$ ($\mathbf{S}^1 \stackrel{\text{open-subsp}}{\subseteq} \mathbf{S}^2$) for $\text{Top}(\mathbf{S}^1) \stackrel{\text{subsp}}{\subseteq} \text{Top}(\mathbf{S}^2)$ ($\text{Top}(\mathbf{S}^1) \stackrel{\text{open-subsp}}{\subseteq} \text{Top}(\mathbf{S}^2)$).

Let $U \in \mathcal{S}$. Then $\text{Top}(U, \mathbf{S}) \stackrel{\text{def}}{=} \text{Top}(U, S)$ is U with the relative topology.

$\mathcal{T}op(\mathbf{S}) \stackrel{\text{def}}{=} \left(\left\{ \text{Top}(U, S) \mid U \in \mathcal{S} \right\}, \text{Ar}(\mathcal{S}) \right)$ is the topological category of $\mathbf{S}$.

By abuse of language we write $U \subseteq \mathbf{S}$ for $U \in \mathcal{S}$, $\mathbf{Top}(\mathbf{S})$ for $\mathbf{Top}(\text{Top}(\mathbf{S}))$, f is a morphism of $\mathbf{S}$ for f is a morphism of $\mathcal{S}$ and f is an isomorphism of $\mathbf{S}$ for f is an isomorphism of $\mathcal{S}$.

Let $\mathbf{C}$ be a set of model spaces. $\mathbf{C}$ is fine grained iff every $(C, \mathcal{C}) \in \mathbf{C}$ is fine grained.

{lem:modtopcat}

Lemma 2.4 (The topological category of a model space is a full topological category).

Let $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then $\mathcal{T}op(\mathbf{M})$ is a full topological category.

Proof. $\mathcal{T}op(\mathbf{M})$ is a small subcategory of $\mathbf{Top}$ by construction. $\mathcal{T}op(\mathbf{M})$ is a full topological category by item 4 of definition 2.1. $\square$

{mod:restriction}

Definition 2.5 (Model neighborhoods). Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then the objects of $\mathcal{S}$ are model neighborhoods of $\mathbf{S}$. If $u \in U \in \mathcal{S}$ then U is a model neighborhood of $\mathbf{S}$ for u . If $U_i \in \mathcal{S}$, $i = 1, 2$, and $U_1 \subseteq U_2$ then U_1 is a model subneighborhood of U_2 ; if $u^1 \in U_1$ then U_1 is also a model subneighborhood for u^1 .

If $\emptyset \in \mathcal{S}$ then $\emptyset$ is a degenerate model neighborhood of $\mathbf{S}$; any nonvoid model neighborhood of $\mathbf{S}$ is a nondegenerate model neighborhood of $\mathbf{S}$.

{def:modsub}

Definition 2.6 (Model subspaces). $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is a model subspace of $\mathbf{T} \stackrel{\text{def}}{=} (T, \mathcal{T})$, abbreviated $\mathbf{S} \stackrel{\text{mod}}{\subseteq} \mathbf{T}$, iff

1. $\mathbf{S}$ and $\mathbf{T}$ are model spaces

2. S is a subspace of T

{modsub:sub}

3. $\mathcal{S} \subseteq^{\text{cat}} \mathcal{T}$

{modsub:subcat}

4. every intersection of S with an object of $\mathcal{T}$ is an object of $\mathcal{S}$

{modsub:rest}

$$\left(\bigvee_{U \in \mathcal{T}}^{\text{Ob}} \right) U \cap S \in \mathcal{S} \quad (2.1)$$

{modsub:full}

5. Every object of $\mathcal{T}$ contained in S is an object of $\mathcal{S}$

$$\left(\bigvee_{U \in \mathcal{T}: U \subseteq S}^{\text{Ob}} \right) U \in \mathcal{S} \quad (2.2)$$

Remark 2.7. This is actually a consequence of item 4, but it is convenient to state it here.

{modsub:rest}

$\mathbf{S}$ is also a full model subspace of $\mathbf{T}$, abbreviated $\mathbf{S} \subseteq^{\text{full-mod}} \mathbf{T}$, iff $S = T$.

$\mathbf{S}$ is also a strict model subspace of $\mathbf{T}$, abbreviated $\mathbf{S} \subseteq^{\text{strict-mod}} \mathbf{T}$, iff S is a model neighborhood of T .

Remark 2.8. S is also a model neighborhood of $\mathbf{S}$ by item 5.

{modsub:full}

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space and T a model neighborhood of $\mathbf{S}$. Then $\text{Mod}(T, \mathbf{S})$, the relative model space of T , is $(T, \mathcal{T})$, where $\mathcal{T}$ is the full subcategory of $\mathcal{S}$ containing all model subneighborhoods of T .

By abuse of language we write T for $\text{Mod}(T, \mathbf{S})$ when $\mathbf{S}$ is understood by context.

{lem:modsub}

Lemma 2.9 (Model subspaces). 1. $\subseteq^{\text{mod}}$, $\subseteq^{\text{full-mod}}$ and $\subseteq^{\text{strict-mod}}$ are transitive.

Proof. Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$ $i = 1, 2, 3$, be a model space. If $\mathbf{S}^i \subseteq^{\text{mod}} \mathbf{S}^{i+1}$, $i = 1, 2$, then

(a) $\mathbf{S}^1$ and $\mathbf{S}^3$ are model spaces by hypothesis.

(b) Since S^i is a subspace of S^{i+1} , $i = 1, 2$, then S^1 is a subspace of S^3 .

(c) Each $\mathcal{S}^i \subseteq^{\text{cat}} \mathcal{S}^{i+1}$ by item 3 of definition 2.6 (Model subspaces) above. Then $\mathcal{S}^1 \subseteq^{\text{cat}} \mathcal{S}^3$.

{def:modsub}

(d) Each intersection of S^i with an object of $\mathcal{S}^{i+1}$ is an object of $\mathcal{S}^i$ by item 4 of definition 2.6. Then every intersection of S^1 with an object of $\mathcal{S}^3$ is an object of $\mathcal{S}^1$.

{modsub:rest}

{def:modsub}

(e) Let $U \in^{\text{Ob}} \mathbf{S}^3$ and $U \subseteq S^1$. $U \cap S^1 = U \in^{\text{Ob}} \mathbf{S}^1$.

If each $\mathbf{S}^i \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}^{i+1}$, then each $S^i = S^{i+1}$ by definition 2.6; then $S^1 = S^3$ and thus $\mathbf{S}^1 \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}^3$. {def:modsub}

If each $\mathbf{S}^i \stackrel{\text{strict-mod}}{\subseteq} \mathbf{S}^{i+1}$, then each S^i is a model neighborhood of $\mathbf{S}^{i+1}$ by definition 2.6, and S^1 is a model neighborhood of $\mathbf{S}^3$ by item 5 of definition 2.6. {def:modsub} {def:modsub} $\square$

2. Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space and $T \subseteq \mathbf{S}$. Then $\text{Mod}(T, \mathbf{S}) \stackrel{\text{mod}}{\subseteq} \mathbf{S}$

Proof. Let $\mathbf{T} = ((T, \mathcal{T}), \mathcal{T}) \stackrel{\text{def}}{=} \text{Mod}(T, \mathbf{S})$.

(a) $\mathbf{S}$ and $\mathbf{T}$ are model spaces.

(b) T is a subspace of S .

(c) $\mathcal{T} \stackrel{\text{cat}}{\subseteq} \mathcal{S}$ by construction.

(d) Let $U \stackrel{\text{Ob}}{\in} \mathbf{S}$. $U \cap T \stackrel{\text{Ob}}{\in} \mathbf{S}$. Since $U \cap T \subseteq T$ is a model neighborhood of $\mathbf{S}$, it is a model neighborhood of $\text{Mod}(T, \mathbf{S})$ by construction.

(e) Let $U \stackrel{\text{Ob}}{\in} \mathbf{S}$ be contained in T . Then U is a model neighborhood of $\text{Mod}(T, \mathbf{S})$ by construction. \(\square\)

3. Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, and $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$. Then S^1 is an open subspace of S^2 .

Proof. The model neighborhoods of $\mathbf{S}^1$ are model neighborhoods of $\mathbf{S}^2$, hence open in S^2 . Their union, which is open in S^2 , is S^1 by item 1 of definition 2.6 (Model subspaces) on page 18. {def:modsub} {mod:cover} $\square$

4. Let S^i , $i = 1, 2$, be a topological space and $S^1 \stackrel{\text{open-subsp}}{\subseteq} S^2$. Then $\text{Id}_{S^2}^{S^1}$ is open and continuous.

Proof. Let U be open in S^1 . Since $S^1 \stackrel{\text{open-subsp}}{\subseteq} S^2$, there is a V open in S^2 such that $U = S^1 \cap V$. Since V and S^1 are open in S^2 , so is $U = S^1 \cap V$.

Let V be open in S^2 . Since $S^1 \stackrel{\text{subsp}}{\subseteq} S^2$, $\text{Id}_{S^2}^{S^1-1}[V] = V \cap S^1$ is open in S^1 . \(\square\)

{cor:modsub}

Corollary 2.10 (Model subspaces). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, $\mathbf{S}'^2 \stackrel{\text{def}}{=} (S'^2, \mathcal{S}'^2)$ be model spaces, and $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^2$.

1. $S^1 \stackrel{\text{open-subsp}}{\subseteq} S'^2$.

Proof. $S^1 \xrightarrow{\text{open-subsp}} \subseteq S^2 \xrightarrow{\text{open-subsp}} \subseteq S'^2$. □

2. Let $f: S^1 \longrightarrow S^2$ be continuous. Then $f: S^1 \longrightarrow S'^2$ is continuous.

Proof. $S^1 \xrightarrow{\text{open-subsp}} \subseteq S'^2$, Let $V \subseteq S'^2$ be an arbitrary open set. Since $S^2 \xrightarrow{\text{mod}} \subseteq S'^2$ by hypothesis, $V \cap S^2$ is open in S^2 . Since $f: S^1 \longrightarrow S^2$ is continuous by hypothesis, $f^{-1}[V] = f^{-1}[V \cap S^2]$ is open in S^1 . □

3 Trivial model spaces

Informally, a trivial model space of a specific type is one that does not restrict the potential objects and morphisms of its type. {sec:triv}

Definition 3.1 (Trivial model spaces). Let S be a topological space and $\mathcal{S}$ the category of all continuous functions between open sets of S . {def:trivmod}

$\mathcal{S}_{\text{triv}} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is the trivial model space of S and $\mathcal{S}_{\text{triv}}$ is a trivial model space.

Let $\mathcal{S}$ be a set of topological spaces.

$\mathcal{S}_{\text{triv}} \stackrel{\text{def}}{=} \left\{ \mathcal{S}'_{\text{triv}} \mid \mathcal{S}' \in \mathcal{S} \right\}$ is the set of all trivial model spaces in $\mathcal{S}$.

$\mathcal{S}_{\text{triv}}$, the category of all continuous functions between elements of $\mathcal{S}_{\text{triv}}$, is the trivial model category of $\mathcal{S}$.

$\mathcal{S}_{\text{op-triv}}$, the category of open trivial model spaces in $\mathcal{S}$, is the category whose

objects are $\left\{ U_{\text{triv}} \mid U \in \mathcal{S}_{\text{op}} \right\}$, the trivial model spaces of non-null open sets of spaces

in $\mathcal{S}$, and whose morphisms are all the continuous functions among them. {lem:trivmod}

Lemma 3.2 (Trivial model spaces). Let $S^i, i = 1, 2$, be a topological space.

1. $\mathcal{S}_{\text{triv}}^i$ is a model space.

Proof. Let $\mathcal{S}^i \stackrel{\text{def}}{=} \text{Cat} \left(\mathcal{S}_{\text{triv}}^i \right)$, $f: A \longrightarrow B$ be a morphism of $\mathcal{S}^i$, $A' \in \mathcal{S}^i \subseteq A$ and $B' \in \mathcal{S}^i \subseteq B$. Then the definitions of continuity and of $\mathcal{S}_{\text{triv}}^i$ imply each of the following.

- (a) $\text{Ob}(\mathcal{S}^i) = \mathfrak{Top}(S^i)$ and thus is an open cover of S^i .
- (b) $\text{Ob}(\mathcal{S}^i) = \mathfrak{Top}(S^i)$ and thus is closed under finite intersections.
- (c) All morphisms in $\text{Ar}(\mathcal{S}^i)$ are continuous.
- (d) Since $f: A \longrightarrow B$ is a morphism of $\mathcal{S}^i$, f is continuous.
- (e) Since $A' \in \mathcal{S}^i$ then A' is open.

- (f) Since $B' \in \mathcal{S}^i$ then B' is open.
- (g) Since $f: A \rightarrow B$ is continuous then $f|_{A'}: A' \rightarrow B$ is continuous.
- (h) Since $f[A'] \subseteq B'$ then $f|_{A', B'}: A' \rightarrow B'$ is well defined and continuous, hence a morphism.
- (i) If $A' \in \mathcal{S}^i \subseteq A \in \mathcal{S}^i$ then
 - i. A and A' are open.
 - ii. The inclusion map $\text{Id}_A^{A'}$ is continuous.
 - iii. The inclusion map $\text{Id}_A^{A'}$ is a morphism of $\mathcal{S}^i$ by the definition of $\mathcal{S}_{\text{triv}}^i$.
 - iv. $\mathcal{S}_{\text{triv}}^i$ trivially satisfies the restricted sheaf condition because every continuous function between open sets of S^i is a morphism of $\mathcal{S}^i$.

□

2. Let S^1 be an open subspace of S^2 . Then $\mathcal{S}_{\text{triv}}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathcal{S}_{\text{triv}}^2$.

Proof. Let $\mathcal{S}^i \stackrel{\text{def}}{=} \text{Cat}(\mathcal{S}_{\text{triv}}^i)$, $i = 1, 2$. By definition 3.1 (Trivial model spaces) on page 21, $\mathcal{S}^i$ is the category of all continuous functions between open sets of S^i ,

{def:trivmod}

$\mathcal{S}_{\text{triv}}^1$ and $\mathcal{S}_{\text{triv}}^2$ satisfy the six conditions of definition 2.6 (Model subspaces) on page 18:

{def:modsub}

$\mathcal{S}_{\text{triv}}^1$ and $\mathcal{S}_{\text{triv}}^2$ are model spaces: Proved above.

$\mathcal{S}_{\text{triv}}^1$ is a subspace of $\mathcal{S}_{\text{triv}}^2$:

$\text{Top}(\mathcal{S}_{\text{triv}}^i) = S^i$ by definition 3.1 (Trivial model spaces) on page 21. S^1 is an open subspace of S^2 by hypothesis.

{def:trivmod}

$\mathcal{S}_{\text{triv}}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}_{\text{triv}}^2$:

If $U \in \mathcal{S}_{\text{triv}}^1$ then U is open in S^1 . Since S^1 is an open subspace of S^2 , U is open in S^2 and thus $U \in \mathcal{S}_{\text{triv}}^2$.

If $f: U \rightarrow V \in \mathcal{S}_{\text{triv}}^1$ then f is a continuous function between open sets of S^1 . Since S^1 is an open subspace of S^2 , f is a continuous function between open sets of S^2 and thus a morphism of $\mathcal{S}_{\text{triv}}^2$.

If $f: U \rightarrow V \in \mathcal{S}_{\text{triv}}^2$ then f is a continuous function between open sets of S^2 . If $U \in \mathcal{S}_{\text{triv}}^1$ and $V \in \mathcal{S}_{\text{triv}}^1$, then U and V are open sets of S^1 . Since S^1 is an open subspace of S^2 , f is a continuous function between open sets of S^1 and thus a morphism of $\mathcal{S}_{\text{triv}}^1$.

Every intersection of S^1 with an object of $\mathcal{S}^2$ is an object of $\mathcal{S}^1$:

Let $U \in \mathcal{S}^2$. U and S^1 are open in S^2 . $U \cup \text{Set}(S^1)$ is open in S^2 and contained in S^1 . Since S^1 is an open subspace of S^2 , $U \cup \text{Set}(S^1)$ is open in S^1 and thus an object of $\mathcal{S}^1$.

Every object of $\mathcal{S}^2$ contained in S^1 is an object of $\mathcal{S}^1$:

Let $U \in \mathcal{S}^2$. U is open in S^2 . If U is contained in S^1 then it is open in S^1 and thus an object of $\mathcal{S}^1$.

Since S^1 is an open subspace of S^2 , $\text{Set}(S^1)$ is a model neighborhood of S^2_{triv} by definition 3.1 above □ {def:trivmod}

$$S^1_{\text{triv}} \overset{\text{mod}}{\subseteq} S^2_{\text{triv}}.$$

Proof. $S^1 \overset{\text{open-subsp}}{\subseteq} S^2$ by lemma 2.9 (Model subspaces) on page 19. □ {lem:modsub}

3. Let $f: S^1 \rightarrow S^2$ be continuous. Then f is a model function from S^1_{triv} to S^2_{triv} .

Proof. Let U^i be a model neighborhood of S^i_{triv} , $i = 1, 2$.

(a) Since U^2 is a model neighborhood of S^2_{triv} , U^2 is open, $f^{-1}[U^2]$ is open and thus a model neighborhood of S^1_{triv} .

(b) $f[U^1] \subseteq S^2$. Since S^2 is open, it is a model neighborhood of S^2_{triv} .

□

4 Minimal model spaces

Even if a set of open sets fails one of items 1 and 2 or a set of functions among them fails one of items 3 to 6 in the definition of a model space, there is a minimal model space containing them.

{sec:min}
{mod:cover,mod:closed}
{mod:continuous}
{def:minmod}

Definition 4.1 (Minimal model spaces). Let S be a topological space, $\mathcal{O}$ a set of open sets in S , $\mathcal{f}$ a set of continuous functions between elements of $\mathcal{O}$ and $\mathcal{S}$ the smallest concrete category over **Top** having all sets in $\mathcal{O}$ as objects, having all functions in $\mathcal{f}$ as morphisms and satisfying items 3 to 6 of definition 2.1 (Model spaces) on page 17.

{def:model}
{mod:continuous}

Then

$$\text{Mod}_{\min}(S, \mathcal{O}, \mathcal{f}) \stackrel{\text{def}}{=} \left(\text{Top} \left(\bigcup \mathcal{O}, S \right), \mathcal{S} \right) \quad (4.1)$$

is the minimal model space of S with neighborhoods $\mathcal{O}$ and neighborhood mappings $\mathcal{f}$.

Remark 4.2. The trivial model space $\underset{\text{triv}}{S}$ is a special case.

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space, $\mathbf{O}$ a set of model neighborhoods of S and $\mathbf{f}$ a set of morphisms between elements of $\mathbf{O}$.

Then

$$\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f}) \stackrel{\text{def}}{=} \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f}) \quad (4.2)$$

is the minimal model space of S with neighborhoods $\mathbf{O}$ and neighborhood mappings $\mathbf{f}$.

Remark 4.3. $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is not, in general, a model subspace of $\mathbf{S}$, since modsub:rest, modsub:full items 4 and 5 of definition 2.6 (Model subspaces) on page 18 may fail. {def:modsub}

While every model neighborhood of $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is a model neighborhood of $\underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$, the converse need not be true. {lem:minmod}

Lemma 4.4 (Minimal model spaces are model spaces). Let S be a topological space, $\mathbf{O}$ a set of open sets in S and $\mathbf{f}$ a set of continuous functions between elements of $\mathbf{O}$. Then $(C, \mathcal{C}) \stackrel{\text{def}}{=} \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ is a model space.

Proof.

1. Finite intersections of open sets are open and $C = \bigcup_{U \in \mathbf{O}} U$ by construction
2. $\text{Ob}(\mathcal{C})$ is closed under finite intersections by construction
3. Compositions of continuous functions, inclusion maps and restrictions of continuous functions are continuous
4. Restrictions of morphisms are morphisms by construction
5. Inclusion maps are morphisms by construction
6. The restricted sheaf condition holds by construction

□

Let $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space, $\mathbf{O}$ a set of model neighborhoods of S and $\mathbf{f}$ a set of morphisms between elements of $\mathbf{O}$. Then $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f})$ is a model space.

Proof. $\underset{\text{min}}{\text{Mod}}(\mathbf{S}, \mathbf{O}, \mathbf{f}) = \underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ and $\underset{\text{min}}{\text{Mod}}(S, \mathbf{O}, \mathbf{f})$ is a model space. □

5 M-paracompact model spaces

Paracompactness is an important property for topological spaces because of partitions of unity. There is an analogous property for model spaces.

{sec:m-para}

Definition 5.1 (Model topology). Let $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a model space for S . Then the model topology $\mathfrak{M}^*$ for M is the topology generated by $\text{Ob}(\mathcal{S})$.

{def:ModTop}

Remark 5.2. $\mathfrak{M}^*$ is not guaranteed to be T0 even if S is T4. However, $\mathfrak{M}^*$ may be normal or regular even if S is not.

{def:M-para}

Definition 5.3 (m-paracompactness). A model space $\mathbf{M} \stackrel{\text{def}}{=} (S, \mathcal{S})$ is m-paracompact iff $\mathfrak{M}^*$ is regular and every cover of S by model neighborhoods has a locally finite refinement by model neighborhoods.

Remark 5.4. This is a stronger condition than merely requiring $\mathfrak{M}^*$ to be paracompact.

Theorem 5.5 (m-paracompactness and paracompactness). If $\mathbf{M} = (S, \mathcal{S})$ is m-paracompact and the model neighborhoods form a basis for S then S is paracompact.

Proof. Let R be an open cover of S . Since the model neighborhoods form a basis, every set in R is a union of model neighborhoods and thus there is a refinement R_1 by model neighborhoods. Since by hypothesis $\mathbf{M}$ is m-paracompact, R_1 has a locally finite refinement R_2 by model neighborhoods. Since model neighborhoods are open sets, R_2 is a locally finite refinement of R in the conventional sense. $\square$

6 Model functions and model categories

It is convenient to have a notion of mappings between model spaces that are well behaved in some sense, e.g., fiber preserving; using that notion it is then possible to group model spaces into categories.

{sec:modcat}

6.1 Model functions

Definition 6.1 (Model functions). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: S^1 \rightarrow S^2$ be a continuous function. f is a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff the inverse images of model neighborhoods are model neighborhoods.

{def:modfunc}

$$\left(\bigvee_{V \in \mathcal{S}^2}^{\text{Ob}} \right) f^{-1}[V] \in \mathcal{S}^1 \quad (6.1)$$

f is also a constrained model function of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff $f[S^1]$ is contained in a model neighborhood of $\mathbf{S}^2$.

$$\left(\bigvee_{V \in \mathcal{S}^2}^{\text{Ob}} \right) f[S^1] \subseteq V \quad (6.2)$$

Remark 6.2. When $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is constrained, S^1 is a model neighborhood of $\mathbf{S}^1$.

By abuse of language we write $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ both for f considered as a model function and for f considered as a continuous function.

Let $\mathbf{S}^1 \stackrel{\text{def}}{=} (S^1, \mathcal{S}^1)$ and $\mathbf{S}'^2 \stackrel{\text{def}}{=} (S'^2, \mathcal{S}'^2)$ be model spaces, $\mathbf{S}^2 \stackrel{\text{def}}{=} (S^2, \mathcal{S}^2) \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^2$ be a model subspace and $f: S^1 \longrightarrow S^2$ be a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$. Then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}'^2 \stackrel{\text{def}}{=} \text{Id}_{\mathbf{S}'^2} \circ f$ is f considered as a model function from $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Let $\mathbf{S}'^i \stackrel{\text{def}}{=} (S'^i, \mathcal{S}'^i)$, $i = 1, 2$, be a model space, $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^i$ be a model subspace and f' be a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$ such that $f'[S^1] \subseteq S^2$. Then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$, also written $f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$, is $f' \upharpoonright_{\mathbf{S}^1}$ considered as a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$. {lem:modfunc}

Lemma 6.3 (Model functions). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$ and $\mathbf{S}'^i \stackrel{\text{def}}{=} (S'^i, \mathcal{S}'^i)$, $i = 1, 2$, be model spaces, $\mathbf{S}^i \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^i$ and $f': S'^1 \longrightarrow S'^2$ be a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$.

$f \stackrel{\text{def}}{=} f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$ is a model function of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. f' is a model function by hypothesis. f' is continuous by definition 6.1 above, so $f \stackrel{\text{def}}{=} f' \upharpoonright_{\mathbf{S}^1, \mathbf{S}^2}$ is continuous. {def:modfunc}

Let U be a model neighborhood of $\mathbf{S}^2$. $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}'^2$ by hypothesis, so U is a model neighborhood of $\mathbf{S}'^2$. f' is a model function of $\mathbf{S}'^1$ to $\mathbf{S}'^2$ by hypothesis, so $f'^{-1}[U] \in \mathbf{S}'^1$ by definition 6.1. Then $f^{-1}[U] = f'^{-1}[U] \cap S^1 \in \mathbf{S}^1$ by definition 6.1. {def:modfunc}
□ {lem:modfunctriv}

Lemma 6.4 (Model functions and trivial model spaces). Let S^i , $i = 1, 2$, be a topological space and $f: S^1 \longrightarrow S^2$ be a continuous function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ is a model function.

Proof. Let U^i be a model neighborhood of $\underset{\text{triv}}{S^i}$, $i = 1, 2$.

$f^{-1}[U^2]$ is open, hence a model neighborhood of $\underset{\text{triv}}{S^1}$.

S^2 is a model neighborhood of $\underset{\text{triv}}{S^2}$ and $f[U^1] \subseteq U^2$. □

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ be a model function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \underset{\text{triv}}{S^2}$ is a model function.

Proof. $f: S^1 \longrightarrow S^2$ is continuous. □

Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ be a model function. Then $f: \underset{\text{triv}}{S^1} \longrightarrow \mathbf{S}^2$ is a model function iff $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a constrained model function.

Proof. let U^2 be a model neighborhood of $\mathbf{S}^2$. $f^{-1}[U^2]$ is open, hence a model neighborhood of $\mathbf{S}^1$.

$\mathbf{S}^1$ is a model neighborhood of $\mathbf{S}^1$. If $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a model function then $f[\mathbf{S}^1]$ is contained in a model neighborhood of $\mathbf{S}^2$ and thus f is constrained.

If f is constrained, let U^1 be a model neighborhood of $\mathbf{S}^1$. Then $f[U^1] \subseteq f[\mathbf{S}^1]$ is contained in a model neighborhood of $\mathbf{S}^2$. $\square$

{lem:modcomp}

Lemma 6.5 (Composition of model functions). Let $\mathbf{S}_i \stackrel{\text{def}}{=} (S_i, \mathcal{S}_i)$, $i \in [1, 3]$, be a model space and $f_i: \mathbf{S}_i \rightarrow \mathbf{S}_{i+1}$, $i = 1, 2$, be a model function.

$f_2 \circ f_1$ is a model function.

Proof. Let $U_i \in \mathcal{S}_i$, $i = 1, 2, 3$.

Since f_2 is a model function, $f_2^{-1}[U_3] \in \mathcal{S}_2$. Since f_1 is a model function, $(f_2 \circ f_1)^{-1}[U_3] = f_1^{-1}[f_2^{-1}[U_3]] \in \mathcal{S}_1$. Since f_1 is a model function, $f_1[U_1]$ is contained in a model neighborhood V_2 . Since f_2 is a model function, $f_2[V_2]$ is contained in a model neighborhood V_3 . Then $(f_2 \circ f_1)[U_1] \subseteq V_3$. $\square$

If each f_i is constrained then $f_2 \circ f_1$ is constrained.

Proof. $f_1[\mathbf{S}^1] \subseteq \mathbf{S}^2$ and $f_2[\mathbf{S}^2] \subseteq \mathbf{S}^3$, hence $f_2 \circ f_1[\mathbf{S}^1] \subseteq \mathbf{S}^3$. $\square$

{def:modelhomeo}

Definition 6.6 (Model homeomorphisms). Let $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, be a model space and $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ be a model function. f is a model homeomorphism iff it is also invertible and its inverse is a model function.

6.2 Model categories

{def:modcat}

Definition 6.7 (Model categories). A category $\mathcal{M}$ is a model category iff

1. the objects of $\mathcal{M}$ are model spaces
2. the morphisms of $\mathcal{M}$ are model functions.
3. composition is functional composition.
4. If $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S}) \in \mathcal{M}$ and $S^i \in \mathcal{S}$, $i = 1, 2$, then $\mathbf{S}^i \stackrel{\text{def}}{=} \text{Mod}(S^i, \mathbf{S}) \in \mathcal{M}$. If further $S^1 \subseteq S^2$ then the inclusion map $\text{Id}_{S^2}^{\mathbf{S}^1}: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{S}$.
5. If $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}$, $U^i \in \mathcal{S}^i$, $i = 1, 2$, $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{M}$ and $f[U^1] \subseteq U^2$ then $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}$.
6. If $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i)$, $i = 1, 2$, is a subspace of $\mathbf{S} \stackrel{\text{def}}{=} (S, \mathcal{S}) \in \mathcal{M}$ and $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{S}$ then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathcal{M}$.

{modcat:inc}

{modcat:rest}

{modcat:morph}

A model category $\mathcal{M}$ satisfies the restricted sheaf condition iff whenever

1. U and V are objects of $\mathcal{S}$.
2. $U_\alpha \overset{\text{mod}}{\subseteq} U$ and $V_\alpha \overset{\text{mod}}{\subseteq} V$, $\alpha < A$, are objects of $\mathcal{M}$.
3. $\mathbf{Set}(U) = \bigcup_{\alpha < A} \mathbf{Set}(U_\alpha)$
4. $\mathbf{Set}(V) = \bigcup_{\alpha < A} \mathbf{Set}(V_\alpha)$
5. $f_\alpha: U_\alpha \longrightarrow V_\alpha$ are morphisms of $\mathcal{M}$.
6. $f: U \longrightarrow V$ is a model function and for every $\alpha < A$, f agrees with f_α on U_α

then f is a morphism of $\mathcal{M}$.

Let $\mathcal{M}^i$, $i = 1, 2$, be a model category. $\mathcal{M}^1$ is a strict subcategory of $\mathcal{M}^2$, abbreviated $\mathcal{M}^1 \overset{\text{strict-cat}}{\subseteq} \mathcal{M}^2$, iff $\mathcal{M}^1 \overset{\text{fullt-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2$ satisfies the restricted sheaf condition.

{lem:modcat}

Lemma 6.8 (Model categories). Let $\mathcal{M}$ be a model category, $\mathbf{S}^i \overset{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}$.

$i = 1, 2$, $U^i \in \mathcal{S}^i$, $\mathbf{U}^i \overset{\text{def}}{=} \text{Mod}(U^i, \mathbf{S}^i)$ and $f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \in \mathcal{M}$.

$f: \mathbf{U}^1 \longrightarrow \mathbf{S}^2 \overset{\text{Ar}}{\in} \mathcal{M}$.

Proof. Since $\text{Id}_{\mathbf{S}^2} \overset{\text{Ar}}{\in} \mathcal{M}$ by item 4 of definition 6.7 above and $f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \overset{\text{Ar}}{\in} \mathcal{M}$

{modcat:inc}

by hypothesis, $f: \mathbf{U}^1 \longrightarrow \mathbf{S}^2 = \text{Id}_{\mathbf{S}^2} \circ f: \mathbf{U}^1 \longrightarrow \mathbf{U}^2 \overset{\text{Ar}}{\in} \mathcal{M}$

□

{def:modtrivcat}

Definition 6.9 (Trivial model categories). Let $\mathbf{S} \overset{\text{def}}{=} (S, \mathcal{S})$ be a model space.

$\mathbf{S}_{\text{submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \mathbf{M} \overset{\text{mod}}{\subseteq} \mathbf{S} \right\}$ is the set of all subspaces of $\mathbf{S}$.

$\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}$, the category of all model functions between elements of $\mathbf{S}_{\text{submod-triv}}$, is the trivial subspace model category of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \mathbf{M} \overset{\text{full-mod}}{\subseteq} \mathbf{S} \right\}$ is the set of all full subspaces of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}$, the category of all full model functions between elements of $\mathbf{S}_{\text{full-submod-triv}}$, is the trivial full subspace model category of $\mathbf{S}$.

Let $\mathbf{S}$ be a set of model spaces.

$\mathbf{S}_{\text{submod-triv}} \overset{\text{def}}{=} \left\{ \mathbf{M} \mid \left(\exists_{\mathbf{M}' \in \mathbf{S}} \right) \mathbf{M} \overset{\text{mod}}{\subseteq} \mathbf{M}' \right\}$ is the set of all model subspaces of model spaces in $\mathbf{S}$.

$\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}$, the category of all model functions between elements of $\mathbf{S}_{\text{submod-triv}}$, is the trivial subspace model category of $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-triv}} \stackrel{\text{def}}{=} \left\{ \mathbf{M} \mid \left(\exists_{\mathbf{M}' \in \mathbf{S}} \mathbf{M} \stackrel{\text{full-mod}}{\subseteq} \mathbf{M}' \right) \right\}$ is the set of all full subspaces in $\mathbf{S}$.

$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}$, the category of all full model functions between elements of $\mathbf{S}_{\text{full-submod-triv}}$, is the trivial full subspace model category of $\mathbf{S}$.

{lem:modtrivcat}

Lemma 6.10 (Trivial model categories). Let $\mathbf{S} \stackrel{\text{def}}{=} (\mathbf{S}, \mathcal{S})$ be a model space.

$$\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}} \stackrel{\text{full-cat}}{\subseteq} \mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}}.$$

Proof. $\left(\bigvee_{\mathbf{M} \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}} \mathbf{M} \stackrel{\text{mod}}{\subseteq} \mathbf{S} \right)$, so $\text{Ob}(\mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}) \subseteq \text{Ob}(\mathbf{S}_{\text{submod-}\mathcal{T}\text{riv}})$.

It remains to show that for a pair of objects in both categories, the Hom sets are the same. Let $\mathbf{M}^i \stackrel{\text{full-mod}}{\subseteq} \mathbf{S}$, $i = 1, 2$.

1. Let $f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}^{\text{Ar}}$. Then f is a model function and thus

$$f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{submod-triv}}^{\text{Ar}}.$$

2. Let $f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{submod-triv}}^{\text{Ar}}$. Then f is a model function and thus

$$f: \mathbf{M}^1 \longrightarrow \mathbf{M}^2 \in \mathbf{S}_{\text{full-submod-}\mathcal{T}\text{riv}}^{\text{Ar}}.$$

□

6.3 (Local) m-morphisms

{def:modlocalMmorph}

Definition 6.11 (Local m-morphisms). Let $\mathcal{M}^i$ be a model category, $i = 1, 2$, and $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. A function $f: U^1 \longrightarrow U^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 iff $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and for every $u \in U^1$ there is a model neighborhood $U_u \subseteq U^1$ for u and a model neighborhood $V_u \subseteq U^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$ be open. A function $f: U^1 \longrightarrow U^2$ is a local $\mathbf{S}^1$ m-morphism of U^1 to U^2 iff it is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

⁵This definition does not require that $\mathcal{S}^i$ be a model neighborhood of $\mathcal{S}^i$.

Let $U^i \in \mathcal{S}^i$, $i = 1, 2$. A function $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of U^1 to U^2 iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and for every $u \in U^1$ there is a model neighborhood $U_u \subseteq U^1$ for u and a model neighborhood $V_u \subseteq U^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: \text{Mod}(U_u, \mathbf{S}^1) \rightarrow \text{Mod}(V_u, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$, be open. A model function $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 iff it is a local $\mathbf{S}^i\text{-}\mathbf{S}^i\text{-}\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is a local $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is a local $\mathcal{M}^i$ m-morphism iff it is a local $\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

The phrases “locally a(n) ... m-morphism” and “a local ... m-morphism” are equivalent.

{def:modMmorph}

Definition 6.12 (M-morphisms). Let $\mathcal{M}^i$ be a model category, $i = 1, 2$, and

$\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 iff $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and f is a morphism of $\mathbf{S}^2$.

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff it is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is an m-morphism of $\mathbf{S}^i$ iff it is an m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be model neighborhoods. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1\text{-}\mathbf{S}^2\text{-}\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of U^1 to U^2 iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$.

Let $U^1 \subseteq \mathbf{S}^1$ and $U^2 \subseteq \mathbf{S}^2$, $i = 1, 2$, be model neighborhoods. A model function $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 iff it is an $\mathbf{S}^i\text{-}\mathbf{S}^i\text{-}\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of U^1 to U^2 .

A model function $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an $\mathcal{M}^1\text{-}\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ iff $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and f is a morphism of $\mathcal{M}^2$.⁵

A model function $f: \mathbf{S}^i \rightarrow \mathbf{S}^i$ is an $\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ iff it is an $\mathcal{M}^i\text{-}\mathcal{M}^i$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^i$.

{lem:modMmorph}

Lemma 6.13 ((Local) m-morphisms). Let $\mathcal{M}^i, \mathcal{M}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^i$, $i = 1, 2$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$ and $\mathbf{S}^i \stackrel{\text{mod}}{\subseteq} \mathbf{S}^i$.

{M-morph:openmorphislocal}

1. Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 .

Proof.

- (a) Since f is an $\mathbf{S}^1\text{-}\mathbf{S}^2$ m-morphism of U^1 to U^2 , $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and f is a morphism of $\mathbf{S}^2$,

(b) U^1 and U^2 are model neighborhoods of $\mathbf{S}^2$. U^1 is a model neighborhood of $\mathbf{S}^1$ by item 5 of ^{def:modsub} definition 2.6 (Model subspaces) on page 18.

{modsub:full}

(c) Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f(u^1) \in U^2$, Then U^1 is a model neighborhood for u^1 and U^2 is a model neighborhood for u^2 . $f[U^1] \subseteq U^2$ by hypothesis. $f: U^1 \rightarrow U^2$ is a morphism of $\mathbf{S}^2$ as shown above.

□

{M-morph:openlocalismorph}

2. Let $U^i \subseteq \mathbf{S}^i$ be a model neighborhood of $\mathbf{S}^i$, $i = 1, 2$. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ local m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .

Proof. $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ by hypothesis. U^1 is a model neighborhood of $\mathbf{S}^1$ by hypothesis, thus a model neighborhood of $\mathbf{S}^2$

For each $u \in U^1$, let U_u be a model subneighborhood of U^1 for u and V_u be a model subneighborhood of U^2 for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \rightarrow V_u$ is a morphism of $\mathbf{S}^2$. Then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a morphism of $\mathbf{S}^2$ by ^{mod:sheaf} item 6 of definition 2.1 (Model spaces) on page 17.

□

{def:model}

{M-morph:opencatmorphisloc}

3. Let $U^i \subseteq \mathbf{S}^i$ be a model neighborhood, $i = 1, 2$. If $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 .

Proof.

(a) Each U^i is a model neighborhood of $\mathbf{S}^i$ by hypothesis.

(b) Since $f: U^1 \rightarrow U^2$ is an $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 ,

$\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $f: \text{Mod}(U^1, \mathbf{S}^1) \rightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^2$ by definition 6.12 (M-morphisms) on page 30.

{def:modMmorph}

(c) Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f(u^1) \in U^2$.

i. Each U^i is a model neighborhood of $\mathbf{S}^i$ for u^i ,

ii. Each $U^i \subseteq U^i$.

iii. $f[U^1] \subseteq U^2$.

□

{M-morph:opencatlocalismorph}

4. Let $\mathcal{M}^2$ satisfy the restricted sheaf condition and each $U^i \subseteq \mathbf{S}^i$ be a model neighborhood, $i = 1, 2$. Every $f: U^1 \rightarrow U^2$ that is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ local m-morphism of U^1 to U^2 is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of U^1 to U^2 .

Proof.

(a) Since each U^i is a model neighborhood of $\mathbf{S}^i$ by hypothesis, each

$\text{Mod}(U^i, \mathbf{S}^i) \in \mathcal{M}^i$ by item 4 of definition 6.7

{modcat:inc}

(b) $f: \text{Mod}(U^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U^2, \mathbf{S}^2)$ is a model function by definition 6.11

{def:modlocalMmorph}

(c) Each $\mathbf{S}^i \in \mathcal{M}^i$ by hypothesis

(d) $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$ by hypothesis

(e) $\mathcal{M}^2$ satisfies the restricted sheaf condition by hypothesis

(f) For every $u^1 \in U^1$ there is a model neighborhood $U_{u^1}^1$ for u^1 and a model neighborhood $U_{u^1}^2$ for $u^2 \stackrel{\text{def}}{=} f(u^1)$ such that $f[U_{u^1}^1] \subseteq U_{u^1}^2$ and $f: \text{Mod}(U_{u^1}^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U_{u^1}^2, \mathbf{S}^2)$ is a morphism of $\mathcal{M}^i$.

i. $\text{Mod}(U_{u^1}^1, \mathbf{S}^1) \subseteq^{\text{mod}} \mathbf{S}^1$.

ii. Each $f: \text{Mod}(U_{u^1}^1, \mathbf{S}^1) \longrightarrow \text{Mod}(U_{u^1}^2, \mathbf{S}^2)$ agrees with f on $U_{u^1}^1$.

iii. Since $\bigcup_{u^1 \in U^1} U_{u^1}^1 = U^1$ and $\bigcup_{u^1 \in U^1} U_{u^1}^2 = U^2$, then $f \in \mathcal{M}^2$ by definition 6.7 (Model categories) on page 27.

{def:modcat}

□

{M-morph:localislocaltosup}

5. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. $f: \mathbf{S}^1 \longrightarrow \mathbf{S}'^2$ is continuous by corollary 2.10 (Model subspaces) on page 20.

{cor:modsub}

Let $u \in \mathbf{S}^1$, U_u be a model neighborhood of $\mathbf{S}^1$ for u and V_u be a model neighborhood of $\mathbf{S}^2$ for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}^2$.

Since $\mathbf{S}^2 \subseteq^{\text{mod}} \mathbf{S}'^2$ by hypothesis, $f: U_u \longrightarrow V_u$ is a morphism of $\mathbf{S}'^2$.

□

{M-morph:morphismorphptosup}

6. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$, then f is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2$ by definition 6.12 (M-morphisms) on page 30 and $\mathbf{S}^2 \subseteq^{\text{mod}} \mathbf{S}'^2$ by hypothesis, so $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}'^2$.

{def:modMmorph}

f is a morphism of $\mathbf{S}^2$ by definition 6.12 (M-morphisms) on page 30, hence a morphism of $\mathbf{S}'^2$.

{def:modMmorph}

□

{M-morph:catlocalislocaltosup}

7. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$, then f is a local $\mathcal{M}'^1$ - $\mathcal{M}'^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}'^2$.

Proof. Let $u \in S^1$, U_u be a model neighborhood of S^1 for u and V_u be a model neighborhood of S^2 for $v \stackrel{\text{def}}{=} f(u)$ such that $f[U_u] \subseteq V_u$ and $f: U_u \rightarrow V_u$ is a morphism of S^2 .

Since $S^2 \stackrel{\text{full-cat}}{\subseteq} S'^2$ by hypothesis, $f: U_u \rightarrow V_u$ is a morphism of S'^2 . $\square$

{M-morph: catmorphismorph}

8. If $f: S^1 \rightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of S^1 to S^2 , then f is an $\mathcal{M}'^1$ - $\mathcal{M}'^2$ m-morphism of S^1 to S'^2 .

Proof. $S^1 \stackrel{\text{full-cat}}{\subseteq} S^2$ by definition 6.12 (M-morphisms) on page 30 and $S^2 \stackrel{\text{full-cat}}{\subseteq} S'^2$ by hypothesis, so $S^1 \stackrel{\text{full-cat}}{\subseteq} S'^2$.

{def: modMmorph}

f is a morphism of S^1 by definition 6.12, hence a morphism of S'^2 . $\square$

{def: modMmorph}
{M-morph: contisstrict}

9. If $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ and $f: S^1 \rightarrow S^2$ is continuous, then f is an m-morphism of S^1_{triv} to S^2_{triv} .

Proof.

f is continuous:

$f: S^1 \rightarrow S^2$ is continuous. $\text{Top}\left(\begin{smallmatrix} S^i \\ \text{triv} \end{smallmatrix}\right) = \text{Top}(S^i) = S^i, i = 1, 2.$

f maps model neighborhoods into model neighborhoods:

S^2 is open in S^2 , hence a model neighborhood of S^2_{triv} . Let U be a model neighborhood of S^1_{triv} . Then $f[U] \subseteq S^2$.

Inverse images of model neighborhoods are model neighborhoods:

Let V be a model neighborhood of S^2_{triv} . V is open in S^2 and f is continuous, so $f^{-1}[V]$ is open in S^1 and hence a model neighborhood of S^1_{triv} .

$\square$

{M-morph: Idislocal}

10. If $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ then $\text{Id}_{S^2}^{S^1}$ is a local m-morphism of S^1 to S^2 . If each S^i is a model neighborhood of S^i then $\text{Id}_{S^2}^{S^1}$ is a morphism of S^2 .

Proof. Let $u \in S^1$ and U_u a model neighborhood of S^1 for u . Since $S^1 \stackrel{\text{mod}}{\subseteq} S^2$, U_u is also a model neighborhood of S^2 for u , and hence $\text{Id}_{U_u}^{\text{Ar}} \in S^2$.

If each S^i is a model neighborhood of S^i , then since $S^1 \stackrel{\text{mod}}{\subseteq} S^2$ by hypothesis, S^1 is a model neighborhood of S^2 . The result follows by item 5 of definition 2.1 (Model spaces) on page 17. $\square$

{def: model}
{mod: inclusion}

{M-morph: Idcat is local}

11. If $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$ and $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2$ then $\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}$ is a morphism of $\mathcal{M}^2$.

Proof. $\mathbf{S}^1 \in^{\text{Ob}} \mathcal{M}^2$ because $\mathcal{M}^1 \subseteq^{\text{cat}} \mathcal{M}^2$. $\text{Id}_{\mathbf{S}^2}^{\text{Ar}} \in \mathcal{M}^2$. Then $\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1} \in \mathcal{M}^2$ by item 4 of definition 6.7 (Model categories) on page 27. $\square$

{def: modcat}
{M-morph: openmorph is trivial}

12. Let $U^i \subseteq \mathbf{S}^i$, $i = 1, 2$, be open. If $f: U^1 \rightarrow U^2$ is a (local) $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 then $f: U^1 \rightarrow U^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of U^1 to U^2 .

Proof. Since $\mathbf{S}^1 \subseteq^{\text{mod}} \mathbf{S}^2$, $\mathbf{S}^1 \subseteq^{\text{subsp}} \mathbf{S}^2$. Since $f: U^1 \rightarrow U^2$ is a model function, it is continuous and hence a morphism of $\mathbf{S}^2$. $\square$

{cor: modMmorph}

Corollary 6.14 ((Local) m-morphisms). Let $\mathcal{M}^i, \mathcal{M}^i \subseteq^{\text{full-cat}} \mathcal{M}^i$, $i = 1, 2$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathbf{S}^i, \mathcal{S}^i) \in^{\text{Ob}} \mathcal{M}^i$, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathbf{S}^i, \mathcal{S}^i) \in^{\text{Ob}} \mathcal{M}^i$ and $\mathbf{S}^i \subseteq^{\text{mod}} \mathbf{S}^i$.

{M-morph: morphis local}

1. If $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{M-morph: sublocal is morph}

2. If U^1 and U^2 are model neighborhoods of $\mathbf{S}^i$ then every function $f: U^1 \rightarrow U^2$ that is a local $\mathbf{S}^i$ m-morphism of U^1 to U^2 is a $\mathbf{S}^i$ m-morphism of U^1 to U^2 .

Proof. Since $f: U^1 \rightarrow U^2$ is a local $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^1 to U^2 then it is an $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^1 to U^2 . $\square$

{M-morph: local is morph}

3. If $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and each $\mathbf{S}^i$ is a model neighborhood of $\mathbf{S}^i$, then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then it is an $\mathbf{S}^1$ - $\mathbf{S}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{M-morph: catmorph is local}

4. If $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \rightarrow \mathbf{S}^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Since f is a $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathbf{S}^1$ - $\mathbf{S}^2$ - $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

5. If $\mathcal{M}^i$ satisfies the restricted sheaf condition and U^j is a model neighborhood of $\mathcal{S}^i$, $j = 1, 2$, then every function $f: U^1 \rightarrow U^2$ that is a local $\mathcal{S}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 is a $\mathcal{S}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 .

{M-morph: subcatlocalismorph}

Proof.

- (a) $f: \text{Mod}(U^1, \mathcal{S}^i) \rightarrow \text{Mod}(U^2, \mathcal{S}^i)$ is a local $\mathcal{S}^i$ - $\mathcal{S}^i$ - $\mathcal{M}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 by definition 6.11 (Local m-morphisms) on page 29
- (b) $f: \text{Mod}(U^1, \mathcal{S}^i) \xrightarrow{\text{lem:modMmorph}} \text{Mod}(U^2, \mathcal{S}^i)$ is a $\mathcal{S}^i$ - $\mathcal{S}^i$ - $\mathcal{M}^i$ - $\mathcal{M}^i$ m-morphism of U^1 to U^2 by item 4 of lemma 6.13.
- (c) The result follows from definition 6.11.

{def:modlocalMmorph}

{M-morph: opencatlocalismorph}

{def:modlocalMmorph}

□

{M-morph: catlocalismorph}

6. Let each S^i be a model neighborhood of $\mathcal{S}^i$, Then every $f: S^1 \rightarrow S^2$ that is a $\mathcal{M}^1$ - $\mathcal{M}^2$ local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ is a $\mathcal{M}^1$ - $\mathcal{M}^2$ m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$.

1. If $f: S^1 \rightarrow S^2$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $\mathcal{S}^2 \stackrel{\text{mod}}{\subseteq} \mathcal{S}'^2$ then f is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}'^2$.

Proof. $f: S^1 \rightarrow S^2$ is continuous by corollary 2.10 (Model subspaces) on page 20. □

{cor:modsub}

2. If $f: S^1 \rightarrow S^2$ is a(n) (local) m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ then f is an m-morphism of S^1 to S^2 .

Proof. f is continuous by definition 6.11 (Local m-morphisms) on page 29. $\text{Top}_{\text{triv}}(S^i) = \text{Top}(\mathcal{S}^i) = S^i, i = 1, 2$. □

{def:modlocalMmorph}

3. If S^i is a model neighborhood of $\mathcal{S}^i$ then Id_{S^i} is a morphism of $\mathcal{S}^i$.

Proof. $\mathcal{S}^i \stackrel{\text{mod}}{\subseteq} \mathcal{S}^i$. □

□

4. If $f: S^1 \rightarrow S^2$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $f^2: S^2 \rightarrow S^3$ is an m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then $f^2 \circ f$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $f^2: S^2 \rightarrow S^3$ is an m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then f^2 is a local m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$. □

□

5. If $f: S^1 \rightarrow S^2$ is an m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^2$ and $f^2: S^2 \rightarrow S^3$ is a local m-morphism of $\mathcal{S}^2$ to $\mathcal{S}^3$ then $f^2 \circ f$ is a local m-morphism of $\mathcal{S}^1$ to $\mathcal{S}^3$.

Proof. Since $f: S^1 \longrightarrow S^2$ is an $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^2$. $\square$

6. If $f: S^1 \longrightarrow S^2$ is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and $f^2: S^2 \longrightarrow S^3$ is an $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then $f^2 \circ f$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $f^2: S^2 \longrightarrow S^3$ is an $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then f^2 is a local $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$. $\square$

7. If $f: S^1 \longrightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ and $f^2: S^2 \longrightarrow S^3$ is a local $\mathcal{M}^2$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^3$ then $f^2 \circ f$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $f: S^1 \longrightarrow S^2$ is an $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then f is a local $\mathcal{M}^1$ - $\mathcal{M}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^2$ to $\mathbf{S}^2$. $\square$

{M-morph:morphistrivmorph}

8. If $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is a (local) $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$ then $f: \mathbf{S}^1 \longrightarrow \mathbf{S}^2$ is an $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$.

Proof. Each S^i is open, so $f: S^1 \longrightarrow S^2$ is a $\mathbf{S}^1$ - $\mathbf{S}^2$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^2$. $\square$

{lem:modlocalMmorphcomp}

Lemma 6.15 (Composition of local $\mathbf{m}$ -morphisms). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (S^i, \mathcal{S}^i) \in \mathcal{M}^i$ and $U^i \subseteq S^i$ be open.

1. If $f^i: U^i \longrightarrow U^{i+1}$, $i = 1, 2$, is a local $\mathbf{S}^i$ - $\mathbf{S}^{i+1}$ $\mathbf{m}$ -morphism of U^i to U^{i+1} then $f^2 \circ f^1$ a local $\mathbf{S}^1$ - $\mathbf{S}^3$ $\mathbf{m}$ -morphism of U^1 to U^{i+1} .

Proof. Since $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$ then $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$, Let $u^1 \in U^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathbf{S}^i$ and $\mathbf{S}^{i+1}$ such that that $f^i[U_{u^i}] \subseteq V_{u^i}$ and $f^i: \text{Mod}(U_{u^i}, \mathbf{S}^i) \longrightarrow \text{Mod}(V_{u^i}, \mathbf{S}^{i+1})$ is a morphism of $\mathbf{S}^{i+1}$.

Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^2 \circ f^1)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \longrightarrow V_{u^3}$ is a morphism of $\mathbf{S}^3$. $\square$

2. If $f^i: S^i \longrightarrow S^{i+1}$, $i = 1, 2$, is a local $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ $\mathbf{m}$ -morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1$ a local $\mathcal{M}^i$ - $\mathcal{M}^{i+2}$ $\mathbf{m}$ -morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Let $u^1 \in \mathbf{S}^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathbf{S}^i$ and $\mathbf{S}^{i+1}$ such that $f^i: U_{u^i} \longrightarrow V_{u^i}$ is a morphism of $\mathcal{M}^{i+1}$. Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^3 \circ f^2)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \longrightarrow V_{u^3}$ is a morphism of $\mathcal{M}^3$ by [item 5 of definition 6.7 \(Model categories\)](#) on page 27. $\square$ {def:modcat}

If each $f^i: \mathbf{S}^i \longrightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is a local m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$, then $f^2 \circ f^1: \mathbf{S}^1 \longrightarrow \mathbf{S}^3$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^3$, $\mathcal{M}^1 \stackrel{\text{mod}}{\subseteq} \mathcal{M}^3$. Let $u \in \mathbf{S}^i$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist a model neighborhood U_u for u , model neighborhoods V_u, V'_u for v and a model neighborhood W_v of w such that $f_1[U_u] \subseteq V'_u$, $f_2[V_v] \subseteq W_v$, $f_1: U_u \longrightarrow V'_u$ is a morphism of $\mathcal{S}^2$ and $f_2: V_v \longrightarrow W_v$ is a morphism of $\mathcal{S}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_v \cap V'_u \neq \emptyset$, $\hat{V}_u$ is a model neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}_u]$ is a model neighborhood for u . $f_1: \hat{U}_u \longrightarrow \hat{V}_u$ and $f_2: \hat{V}_u \longrightarrow W_v$ are morphisms of $\mathcal{S}^3$ by [item 4 of definition 2.1 \(Model spaces\)](#) on page 17 and thus $f_2 \circ f_1: \hat{U}_u \longrightarrow W_v$ is a morphism of $\mathcal{S}^3$. $\square$ {def:model}

If each $f^i: \mathbf{S}^i \longrightarrow \mathbf{S}^{i+1}$ is a local $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1: \mathbf{S}^1 \longrightarrow \mathbf{S}^3$ is a local $\mathcal{M}^1$ - $\mathcal{M}^3$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Let $u \in \mathbf{S}^1$, $v \stackrel{\text{def}}{=} f^1(u)$ and $w \stackrel{\text{def}}{=} f^2(v)$. There exist a model neighborhood U_u for u , model neighborhoods V_v, V'_u for v and a model neighborhood W_v of w such that $f^1[U_u] \subseteq V'_u$, $f^1: U_u \longrightarrow V'_u$ is a morphism of $\mathcal{M}^2$, $f^2[V_v] \subseteq W_v$ and $f^2: V_v \longrightarrow W_v$ is a morphism of $\mathcal{M}^3$. Then $\hat{V}_u \stackrel{\text{def}}{=} V_v \cap V'_u \neq \emptyset$, $\hat{V}_u$ is a model neighborhood of v and $\hat{U}_u \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}_u]$ is a model neighborhood for u . $f^1: \hat{U}_u \longrightarrow \hat{V}_u$ and $f^2: \hat{V}_u \longrightarrow W_v$ are morphisms of $\mathcal{M}^3$ by [item 4 of definition 6.7 \(Model categories\)](#) on page 27 and thus $f^2 \circ f^1: \hat{U}_u \longrightarrow W_v$ is a morphism of $\mathcal{M}^3$. $\square$ {def:modcat}

Corollary 6.16 ([Composition of local m-morphisms]). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathbf{S}^i, \mathcal{S}^i) \in \mathcal{M}^i$.

1. Let $U^j \subseteq \mathbf{S}^i$, $j = 1, 2, 3$, be open. If $f^j: U^j \longrightarrow U^{j+1}$, $j = 1, 2$, is a local $\mathbf{S}^i$ m-morphism of U^j to U^{j+1} then $f^2 \circ f^1$ is a local $\mathbf{S}^i$ m-morphism of U^1 to U^3 .

Proof. Since each $f^j: U^j \longrightarrow U^{j+1}$ is a local $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^j to U^{j+1} then $f^2 \circ f^1$ is a local $\mathbf{S}^i$ - $\mathbf{S}^i$ m-morphism of U^1 to U^3 . $\square$

2. If $f^i: \mathbf{S}^i \longrightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is a local m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1$ is a local m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2$ and $\mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$ then $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$. Let $u^1 \in \mathbf{S}^1$, $u^2 \stackrel{\text{def}}{=} f^1(u^1)$ and $u^3 \stackrel{\text{def}}{=} f^2(u^2)$. Let U_{u^i} and V_{u^i} be model neighborhoods of $\mathbf{S}^i$ and $\mathbf{S}^{i+1}$ such that $f^i: U_{u^i} \rightarrow V_{u^i}$ is a morphism of $\mathbf{S}^{i+1}$. Let $U'_{u^1} \stackrel{\text{def}}{=} U_{u^1} \cap (f^2 \circ f^1)^{-1}[V_{u^3}]$. Then $f^2 \circ f^1: U'_{u^1} \rightarrow V_{u^3}$ is a morphism of $\mathbf{S}^3$. $\square$

Lemma 6.17 (Composition of m-morphisms). Let $\mathcal{M}^i$, $i = 1, 2, 3$, be model categories and $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathcal{S}^i)^{\text{Ob}} \in \mathcal{M}^i$.

{lem:modMmorphcomp}

1. If $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is an m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is an m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since each f^i is a morphism of $\mathbf{S}^{i+1}$ by definition 6.12 (M-morphisms) on page 30 and $\mathbf{S}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^2 \stackrel{\text{mod}}{\subseteq} \mathbf{S}^3$, then each f^i is a morphism of $\mathbf{S}^3$ by definition 6.12 (M-morphisms) on page 30, and thus $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is a morphism of $\mathbf{S}^3$. $\square$

{def:modMmorph}

{def:modMmorph}

2. If $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$, $i = 1, 2$, is an $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1$ is an $\mathcal{M}^1$ - $\mathcal{M}^{i+2}$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$. Since each $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$ is a morphism of $\mathcal{S}^3$, then so is $f^2 \circ f^1$. $\square$

3. If each $f^i: \mathbf{S}^i \rightarrow \mathbf{S}^{i+1}$ is an $\mathcal{M}^i$ - $\mathcal{M}^{i+1}$ m-morphism of $\mathbf{S}^i$ to $\mathbf{S}^{i+1}$ then $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is an $\mathcal{M}^1$ - $\mathcal{M}^3$ m-morphism of $\mathbf{S}^1$ to $\mathbf{S}^3$.

Proof. Since $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ then $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$ and f^i is a morphism of $\mathcal{S}^{i+1}$ by definition 6.12 (M-morphisms) on page 30. Since $\mathcal{M}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^3$, f^1 is a morphism of $\mathcal{S}^3$. and thus $f^2 \circ f^1: \mathbf{S}^1 \rightarrow \mathbf{S}^3$ is a morphism of $\mathcal{S}^3$. $\square$

{def:modMmorph}

7 Spaces and proper functions

{sec:spaces}

Several of the following definitions involve spaces of a restrictive character and specific types of mappings among them. Except where otherwise qualified, the word *space* will have this restricted meaning.

{def:space}

Definition 7.1 (Spaces). A space is a topological space, a model space or either with an additional associated structure.

{def:proper}

Definition 7.2 (Proper functions). Let S^1, S^2 be spaces and $f: S^1 \rightarrow S^2$ a continuous function; f need not preserve any associated algebraic structure⁶. f is a proper⁷ function iff

1. S^1 and S^2 are both Truthspace or both **Truthspace**, and $f(\text{True}) = \text{True}$.
2. S^1, S^2 are topological spaces other than Truthspace.
3. S^1 is a topological space other than Truthspace, S^2 is a model space other than **Truthspace** and the images of open sets are contained in model neighborhoods.
4. S^1 is a model space other than **Truthspace**, S^2 is a topological space other than Truthspace and the inverse images of open sets are model neighborhoods.
5. S^1, S^2 are both model spaces other than **Truthspace** and f is a model function.

{def:singcat}

Definition 7.3 (Singleton categories and sequences of singleton categories). Let S be a topological space. Then the singleton category of S , abbreviated $\underset{Sing}{S}$, is the category whose sole object is S and whose sole morphism is the identity morphism of S to itself.

Let $\mathbf{S}$ be a model space. Then the singleton category of $\mathbf{S}$, abbreviated $\underset{Sing}{\mathbf{S}}$, is the category whose sole object is $\mathbf{S}$ and whose sole morphism is the identity morphism of $\mathbf{S}$ to itself.

Let $\mathbf{S} \stackrel{\text{def}}{=} (S^\alpha, \alpha < A)$ be a sequence of spaces. Then the singleton category sequence of $\mathbf{S}$, abbreviated $\underset{Sing}{\mathbf{S}}$, is $(\underset{Sing}{S^\alpha}, \alpha < A)$.

Let $\mathbf{S}$ be a set of spaces. Then the singleton category of $\mathbf{S}$, abbreviated $\underset{Sing}{\mathbf{S}}$, is

$$\bigcup_{S \in \mathbf{S}} \underset{Sing}{S}.$$

Remark 7.4. By abuse of language the notation $\underset{Sing}{\mathbf{S}}$ will be used to name sequences of categories constructed with this and similar functions.

⁶However, in practice commutation relations will often enforce the preservation of algebraic structures.

⁷In the spirit of [2, footnote, p. 112]

{GenTop}

This nomenclature is an excellent example of the time-honored custom of referring to a problem we cannot handle as abnormal, irregular, improper, degenerate, inadmissible, and otherwise undesirable.

Part V

Signatures

{part:sig}

Given a sequence of spaces, we need a way to characterise a function that takes arguments in those spaces and has a value in them. For this purpose we use a sequence of ordinals where the last ordinal in the sequence identifies the space containing the function's value.

8 Definitions of signatures

Definition 8.1 (Signature). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ be a sequence and $\sigma \stackrel{\text{def}}{=} (\alpha_\beta < A, \beta \leq B)$ be a sequence of ordinals in A . Then σ is a signature over $\mathbf{S}$.

Definition 8.2 ($\mathbf{S}$ -signature of function). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ be a sequence, $\sigma \stackrel{\text{def}}{=} (\alpha_\beta < A, \beta \leq B)$ be a signature over $\mathbf{S}$ and $f: (S_{\alpha_\beta}, \beta < B) \longrightarrow S_{\alpha_B}$ continuous. Then σ is the $\mathbf{S}$ -signature for f and f has $\mathbf{S}$ -signature σ .

Remark 8.3. If $n \in \mathbb{N}$ and $\sigma \stackrel{\text{def}}{=} (\alpha, \beta \leq n)$ then f is an n -ary operation over S_α .

f is proper iff either

1. $S_{\alpha_\beta}, \beta \leq B$ are all topological spaces.
2. $S_{\alpha_\beta}, \beta \leq B$ are all model spaces.

Definition 8.4 ($\mathcal{S}$ -signature of function). Let $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$ be a sequence, $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$ be a sequence of categories with $\left(\bigvee_{\alpha < A}\right) S_\alpha \in \mathcal{S}_\alpha$, $\sigma \stackrel{\text{def}}{=} (\alpha_\beta < A, \beta \leq B)$ be a signature over $\mathcal{S}$ and $f: (S_{\alpha_\beta}, \beta < B) \longrightarrow S_{\alpha_B}$ continuous. Then σ is the $\mathcal{S}$ -signature for f and f has $\mathcal{S}$ -signature σ .

Definition 8.5 ($\mathbf{S}$ -signature of function sequence). Let $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$ be a sequence of signatures over $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A)$, $\mathbf{F} = (F_\gamma, \gamma < \Gamma)$ a sequence of functions where F_γ has $\mathbf{S}$ -signature σ_γ , then Σ is the $\mathbf{S}$ -signature for $\mathbf{F}$ and $\mathbf{F}$ has $\mathbf{S}$ -signature Σ .

Definition 8.6 ($\mathcal{S}$ -signature of function sequence). Let $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$ be a sequence of signatures over $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$, $\mathbf{F} = (F_\gamma, \gamma < \Gamma)$ a sequence of functions where F_γ has $\mathcal{S}$ -signature σ_γ , then Σ is the $\mathcal{S}$ -signature for $\mathbf{F}$ and $\mathbf{F}$ has $\mathcal{S}$ -signature Σ .

9 Functions associated with signatures and function sequences

When characterizing functions with signatures, several auxillary functions are helpful.

{sec:sig}

{def:sigop}

Definition 9.1 (Signature operators). Let $\mathcal{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_\alpha^i, \alpha < A)$, $i = 1, 2$, be sequences of categories with $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A)$ sequences of sets in those categories, $f^i, i = 1, 2$ functions with $\mathcal{S}^i$ -signatures $\sigma = (\alpha_\beta < A, \beta \leq B)$, $\mathbf{g} \stackrel{\text{def}}{=} (g_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha < A)$ a sequence of functions then

$$\langle \sigma | \mathbf{S}^i \rangle \stackrel{\text{def}}{=} \mathbf{S}^i \stackrel{\sigma}{\text{def}} (S_{\alpha_\beta}^i, \beta \leq B) \quad (9.1)$$

selects sets from $\mathbf{S}^i$ according to signature σ .

$$\langle \sigma | \mathbf{g} \rangle \stackrel{\text{def}}{=} \mathbf{g} \stackrel{\sigma}{\text{def}} (g_{\alpha_\beta}, \beta \leq B) \quad (9.2)$$

selects functions from $\mathbf{g}$ according to signature σ .

$$\underline{\mathbf{g}} \stackrel{\sigma}{\text{def}} \bigtimes_{\beta < B} S_{\alpha_\beta}^1 \longrightarrow \bigtimes_{\beta < B} S_{\alpha_\beta}^2 = \langle \sigma | \mathbf{g} \rangle = \bigtimes_{\beta < B} g_{\alpha_\beta} \quad (9.3)$$

is the product of the functions selected by all but the last ordinal in signature σ .

$$\mathbf{g} \stackrel{\sigma}{\cdot} f^1 \stackrel{\text{def}}{=} \text{tail}(\langle \sigma | \mathbf{g} \rangle) \circ f^1 \quad (9.4)$$

composes the last function selected from $\mathbf{g}$ by signature σ with f^1 .

$$f^2 \stackrel{\sigma}{\odot} \mathbf{g} \stackrel{\text{def}}{=} f^2 \circ \underline{\mathbf{g}} \quad (9.5)$$

composes f with all but the last function in $\mathbf{g}$ selected by signature σ .

10 Commutative diagrams with signatures

The conventional usage of the word commutative diagram is for a naturality condition on functions of a single argument. It is convenient to extend the nomenclature to functions of multiple arguments and to sequences of such functions. This is convenient, e.g., as a means of specifying that functions preserve algebraic structures.

{sec:cdsig}

{def:Sigmacomm}

Definition 10.1 (Σ -commutation). Let $\mathcal{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, with $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A) \in \mathcal{S}^i$ a sequence of spaces, $\mathbf{F}^i \stackrel{\text{def}}{=} (F_\gamma^i, \gamma < \Gamma)$ be a sequence of functions with $\mathcal{S}^i$ -signature $\Sigma \stackrel{\text{def}}{=} (\sigma_{\gamma < \Gamma}) \stackrel{\text{def}}{=}$




Figure 3: $f_2 \odot^\sigma \underline{\sigma}$

{fig:fgsigma}




Figure 4: $\underline{\sigma} \cdot f_1$

{fig:gsigmaf}

$$\begin{array}{ccc}
\underline{\sigma_\gamma \mathbf{S}_\gamma^1} & \xrightarrow{F_\gamma^1} & \text{tail} \left(\sigma_\gamma \mathbf{S}_\gamma^1 \right) \\
\downarrow \sigma_\gamma \mathbf{f} & & \downarrow \text{tail} \left(\sigma_\gamma \mathbf{f} \right) \\
\underline{\sigma_\gamma \mathbf{S}_\gamma^2} & \xrightarrow{F_\gamma^2} & \text{tail} \left(\sigma_\gamma \mathbf{S}_\gamma^2 \right)
\end{array}$$

Figure 5: $\mathbf{f}$ Σ -commutes with the function sequences $\mathbf{F}^1, \mathbf{F}^2$

{fig:fF1F2}

$((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$, $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha < A)$ a sequence of functions satisfying the naturality condition $\mathbf{f}^{\sigma_\gamma} F_\gamma^1 = F_\gamma^2 \odot \mathbf{f}$, $\gamma < \Gamma$, then $\mathbf{f}$ Σ -commutes with the function sequences $\mathbf{F}^1, \mathbf{F}^2$.

Remark 10.2. If all of the $\sigma_{\gamma, \alpha_\beta}$ are equal to the same ordinal $\hat{\alpha}$ and each B_γ is finite then $\{(S_\alpha^i, F_\gamma^i) \mid \gamma < \Gamma\}$ is an Ω -algebra and $\mathbf{f}$ is an Ω -homomorphism.

{lem:Sigmacomm}

Lemma 10.3 (Σ -commutation). Let $\mathcal{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A)$, $i \in [1, 3]$, be a sequence of categories, with $\mathbf{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A) \in \mathcal{S}^i$ a sequence of spaces and $\mathbf{F}^i \stackrel{\text{def}}{=} (F_\gamma^i, \gamma < \Gamma)$ be a sequence of functions with $\mathcal{S}^i$ -signature $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$.

Let $\mathbf{f}^i \stackrel{\text{def}}{=} (f_\alpha^i: S_\alpha^i \longrightarrow S_\alpha^{i+1}, \alpha < A)$, $i = 1, 2$, Σ -commute with $\mathbf{F}^i, \mathbf{F}^{i+1}$. Then $\mathbf{f}^2 \circ \mathbf{f}^1$ Σ -commutes with $\mathbf{F}^1, \mathbf{F}^3$.

Proof. The result follows by diagram chasing in fig. 6.

□ {fig:fF1F2F3}

Let $\mathbf{S}^1 \subseteq \mathbf{S}^2$ and $F_\gamma^1 = F_\gamma^2: \bigtimes^{\sigma_\gamma} \mathbf{S}^1 \longrightarrow \text{tail} \left(\sigma_\gamma \mathbf{S}^1 \right)$. Then $\mathbf{Id}_{\mathbf{S}^2}^{S^1}$ Σ -commutes with $\mathbf{F}^1, \mathbf{F}^2$.

Proof. The result follows from definition 9.1 (Signature operators) on page 41 and

{def:sigop}

$$\begin{array}{ccc}
\begin{array}{c} \sigma_\gamma \\ \underline{\mathbf{S}_\gamma^1} \end{array} & \xrightarrow{F_\gamma^1} & \begin{array}{c} \sigma_\gamma \\ \text{tail}(\mathbf{S}_\gamma^1) \end{array} \\
\begin{array}{c} \sigma_\gamma \\ \underline{f^1} \end{array} \downarrow & & \downarrow \begin{array}{c} \sigma_\gamma \\ \text{tail}(f^1) \end{array} \\
\begin{array}{c} \sigma_\gamma \\ \underline{\mathbf{S}_\gamma^2} \end{array} & \xrightarrow{F_\gamma^2} & \begin{array}{c} \sigma_\gamma \\ \text{tail}(\mathbf{S}_\gamma^2) \end{array} \\
\begin{array}{c} \sigma_\gamma \\ \underline{f^2} \end{array} \downarrow & & \downarrow \begin{array}{c} \sigma_\gamma \\ \text{tail}(f^2) \end{array} \\
\begin{array}{c} \sigma_\gamma \\ \underline{\mathbf{S}_\gamma^3} \end{array} & \xrightarrow{F_\gamma^3} & \begin{array}{c} \sigma_\gamma \\ \text{tail}(\mathbf{S}_\gamma^3) \end{array}
\end{array}$$

Figure 6: $f^2 \circ f^1$ Σ -commute with F^1, F^3

{fig:fF1F2F3}

`def:SigmaComm`
definition 10.1 (Σ -commutation) on page 41 :

$$\begin{aligned}
\mathbf{Id}_{\mathbf{S}^2}^{\mathbf{S}^1} \cdot F_\gamma^1 &= \text{Id}_{S_\gamma, \alpha_{B_\gamma}}^{S_\gamma, \alpha_{B_\gamma}} \circ F_\gamma^1 \\
&= F_\gamma^2 \upharpoonright \times_{\underline{\mathbf{S}^1}}^{\sigma_\gamma}, \\
&= F_\gamma^2 \circ \bigtimes_{\beta < B_\gamma} \text{Id}_{S_{\alpha_\beta}^2}^{S_{\alpha_\beta}^1} \\
&= F_\gamma^2 \odot^{\sigma_\gamma} \mathbf{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}
\end{aligned}$$

□

{cor:SigmaComm}

Corollary 10.4 (Σ -commutation). Let $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$, be a sequence of categories, with $\mathbf{S} \stackrel{\text{def}}{=} (S_\alpha, \alpha < A) \in \mathcal{S}$ a sequence of spaces and $\mathbf{F} \stackrel{\text{def}}{=} (F_\gamma, \gamma < \Gamma)$ be a sequence of functions with $\mathcal{S}$ -signature $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$. Then $\mathbf{Id}_{\mathbf{S}}$ Σ -commutes with $\mathbf{F}, \mathbf{F}$.

Proof. $\mathbf{S} \subseteq \mathbf{S}, F_\gamma = F_\gamma: \times_{\underline{\mathbf{S}}}^{\sigma_\gamma} \longrightarrow \text{tail}(\mathbf{S})^{\sigma_\gamma}$ and $\mathbf{Id}_{\mathbf{S}} = \mathbf{Id}_{\mathbf{S}}$. □

Part VI

Prestructures

{part:pre}

Prestructures and prestructure morphisms are generalizations of Ω -algebras and Ω -homomorphisms, and are notational conveniences to simplify imposing commutation relations on multiple unrelated functions of multiple variables.

{def:pre}

Definition 10.5 (Prestructures). Let $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$ be a sequence of categories, $\mathcal{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A)$ a sequence of spaces, $F = (F_\gamma, \gamma < \Gamma)$ a sequence of continuous functions and $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma.\alpha_\beta}, \beta \leq B_\gamma), \gamma < \Gamma)$ a sequence of signatures. Then $P \stackrel{\text{def}}{=} (\mathcal{S}, \mathcal{S}, \Sigma, F)$ is a $\mathcal{S} - \Sigma$ prestructure iff

1. $\mathcal{S}_\alpha$ is either a topological category or a model category.
2. $\mathcal{S} \stackrel{\circ}{\in} \mathcal{S}$.
3. F has $\mathcal{S}$ -signature Σ .

{lem:pre}

Lemma 10.6 (Prestructures). Let

1. $\mathcal{S}^i, i = 1, 2$, be a sequence of categories
2. $\mathcal{S}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{S}^2$,
3. $\mathcal{S} \stackrel{\circ}{\in} \mathcal{S}^1$
4. F a sequence of continuous functions with $\mathcal{S}^1$ -signatures Σ
5. $P^1 \stackrel{\text{def}}{=} (\mathcal{S}^1, \mathcal{S}, \Sigma, F)$ a $\mathcal{S}^1 - \Sigma$ prestructure

Then $P^2 \stackrel{\text{def}}{=} (\mathcal{S}^2, \mathcal{S}, \Sigma, F)$ is a $\mathcal{S}^2 - \Sigma$ prestructure.

Proof. $\mathcal{S} \stackrel{\circ}{\in} \mathcal{S}^2$ by lemma 1.28 (Sequence inclusion) on page 16. The other conditions do not depend on the category sequence. $\square$

{lem:seqin}

Let

1. $\mathcal{S}^i, i = 1, 2$, be a sequence of categories
2. $\mathcal{S}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{S}^2$
3. $\mathcal{S}^i \stackrel{\circ}{\in} \mathcal{S}^i$
4. $\mathcal{S}^1 \stackrel{\circ}{\subseteq} \mathcal{S}^2$

5. $F^2 \stackrel{\text{def}}{=} (F_\gamma^2, \gamma < \Gamma)$ be a sequence of continuous functions with $\mathcal{S}^2$ -signatures
 $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma.\alpha_\beta}, \beta \leq B_\gamma), \gamma < \Gamma)$
 6. $F_\gamma^2 \left[\times \underline{\mathcal{S}^1} \right] \subseteq \text{tail} \left(\begin{smallmatrix} \sigma_\gamma \\ \mathcal{S}^1 \end{smallmatrix} \right)$
 7. $F^1 \stackrel{\text{def}}{=} \left(F_\gamma^1 \stackrel{\text{def}}{=} F_\gamma^2: \times \underline{\mathcal{S}^1} \longrightarrow \text{tail} \left(\begin{smallmatrix} \sigma_\gamma \\ \mathcal{S}^1 \end{smallmatrix} \right), \gamma < \Gamma \right)$
 8. $P^2 \stackrel{\text{def}}{=} (\mathcal{S}^2, S^2, \Sigma, F^2)$ a $\mathcal{S}^2 - \Sigma$ prestructure.
- Then $P^1 \stackrel{\text{def}}{=} (\mathcal{S}^1, S^1, \Sigma, F^1)$ is a $\mathcal{S}^1 - \Sigma$ prestructure.

Proof. $\mathcal{S}^1 \in \mathcal{S}^1$ by hypothesis. F^1 is continuous and has $\mathcal{S}^1$ -signatures Σ by construction. $\square$

{def:premorph}

Definition 10.7 (Morphisms of prestructures). Let $\mathcal{S}^i = (\mathcal{S}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathcal{S}^i \stackrel{\text{def}}{=} (S_\alpha^i, \alpha < A) \in \mathcal{S}^i$ be a sequence of spaces, $F^i = (F_\gamma^i, \gamma < \Gamma)$ be a sequence of continuous functions with $\mathcal{S}^i$ -signatures Σ , $f \stackrel{\text{def}}{=} (f_\alpha: S_\alpha^1 \longrightarrow S_\alpha^2, \alpha < A)$ be a sequence of proper functions and $P^i \stackrel{\text{def}}{=} (\mathcal{S}^i, S^i, \Sigma, F^i)$ be a $\mathcal{S}^i - \Sigma$ prestructure. f is a (sem-strict, strict) prestructure morphism of P^1 to P^2 iff it Σ -commutes with F^1, F^2 , as shown in [fig. 5](#) (**f Σ -commutes with the function sequences F^1, F^2**) on page 43. {fig:fF1F2}

Remark 10.8. If $F^1 = F^2 = \emptyset$ then f is a morphism since it vacuously Σ -commutes with $\emptyset, \emptyset$.

It is a semi-strict prestructure morphism iff $\mathcal{S}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}^2$ and for each α

If $\mathcal{S}_\alpha^i$ is a model category: f_α is a local $\mathcal{S}_\alpha^1 - \mathcal{S}_\alpha^2$ m-morphism of S_α^1 to S_α^2 .

If $\mathcal{S}_\alpha^i$ is a topological category: f_α is a local $\mathcal{S}_\alpha^1 - \mathcal{S}_\alpha^2$ morphism of S_α^1 to S_α^2 .

Otherwise: f_α is a morphism of $\mathcal{S}_\alpha^2$.

It is a strict prestructure morphism iff $\mathcal{S}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{S}^2$ and each function f_α is a morphism of $\mathcal{S}_\alpha^2$.

Remark 10.9. It is not sufficient to require that each f_α be a morphism of $\mathcal{S}_\alpha^1 \cup^{\text{cat}} \mathcal{S}_\alpha^2$ because that would not ensure that the composition of strict prestructure morphisms is strict.

If $\mathcal{S}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{S}^2$ and $F_\gamma^1 = F_\gamma^2: \times \underline{\mathcal{S}^1} \longrightarrow \text{tail} \left(\begin{smallmatrix} \sigma_\gamma \\ \mathcal{S}^1 \end{smallmatrix} \right)$ then $\text{Id}_{P^1} \stackrel{\text{def}}{=} \text{Id}_{\mathcal{S}^1}$.

$\text{Id}_{P^i} \stackrel{\text{def}}{=} \text{Id}_{\mathcal{S}^i} = \text{Id}_{P^i}^{P^i}$.

For some application, certain spaces of a prestructure are of primary interest while the rest serve only in an auxilliary role. For those applications, morphisms that are identical on the primary spaces may be considered equivalent.

{def:premorphKequiv}

Definition 10.10 (**K**-equivalence of prestructure morphisms). Let $\mathcal{S}^i = (\mathcal{S}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathbf{S}^i, \Sigma, \mathbf{F}^i)$ be a $\mathcal{S}^i - \Sigma$ prestructure, $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: \mathcal{S}_\alpha^1 \longrightarrow \mathcal{S}_\alpha^2, \alpha < A)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_\alpha: \mathcal{S}_\alpha^1 \longrightarrow \mathcal{S}_\alpha^2, \alpha < A)$ be prestructure morphisms of $\mathbf{P}^1$ to $\mathbf{P}^2$ and K be a set of ordinals less than A .

$\mathbf{f}$ is **K**-equivalent to $\mathbf{g}$ as a prestructure morphism from $\mathbf{P}^1$ to $\mathbf{P}^2$, abbreviated $\mathbf{f}: \mathbf{P}^1 \longrightarrow \mathbf{P}^2$ is **K**-equivalent to $\mathbf{g}: \mathbf{P}^1 \longrightarrow \mathbf{P}^2$, iff $\left(\bigvee_{\alpha \in K}\right) f_\alpha = g_\alpha$.

By abuse of language we write $\mathbf{f}$ is **K**-equivalent to $\mathbf{g}$ when the prestructures $\mathbf{P}^1$ and $\mathbf{P}^2$ are understood by context.

{lem:premorph}

Lemma 10.11 (Prestructure morphisms). Let $\mathcal{S}^i = (\mathcal{S}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_\alpha^i, \alpha < A) \overset{\circ}{\in} \mathcal{S}^i$ a sequence of spaces, $\mathbf{F}^i = (F_\gamma^i, \gamma < \Gamma)$, be a sequence of continuous functions with $\mathbf{S}^i$ -signatures Σ and $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathbf{S}^i, \Sigma, \mathbf{F}^i)$ be a $\mathcal{S}^i - \Sigma$ prestructure.

If $\mathbf{S}^1 \overset{\circ}{\subseteq} \mathbf{S}^2$ and each $F_\gamma^1 = F_\gamma^2: \bigtimes_{\gamma} \mathbf{S}^1 \longrightarrow \text{tail}\left(\mathbf{S}^1\right)^{\sigma_\gamma}$ then $\text{Id}_{\mathbf{P}^2}^{\mathbf{P}^1}$ is a prestructure morphism from $\mathbf{P}^1$ to $\mathbf{P}^2$.

$\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}$ is a semi-strict prestructure morphism from $\mathbf{P}^1$ to $\mathbf{P}^2$ iff $\mathcal{S}^1 \overset{\text{full-cat}}{\subseteq} \mathcal{S}^2$.

$\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}$ is a strict prestructure morphism from $\mathbf{P}^1$ to $\mathbf{P}^2$ iff each $\text{Id}_{\mathcal{S}_\alpha^2}^{\mathcal{S}_\alpha^1} \overset{\text{Ar}}{\in} \mathcal{S}_\alpha^2$ and $\mathcal{S}^1 \overset{\text{full-cat}}{\subseteq} \mathcal{S}^2$.

Proof. $\text{Id}_{\mathbf{S}^2}^{\mathbf{S}^1}$ Σ -commutes with $\mathbf{F}^1, \mathbf{F}^2$ by lemma 10.3 (**Σ -commutation**) on page 43.

The hypotheses for strictness are precisely those of definition 10.7 (**Morphisms of prestructures**). $\square$

{lem:Sigmacomm}
{def:premorph}

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: \mathcal{S}_\alpha^1 \longrightarrow \mathcal{S}_\alpha^2, \alpha < A)$ be a sequence of proper functions,

If $\mathcal{S}'^i = (\mathcal{S}_\alpha'^i, \alpha < A)$, $i = 1, 2$, is a sequence of categories, $\mathcal{S}^i \overset{\text{cat}}{\subseteq} \mathcal{S}'^i$ and $\mathbf{P}'^i \stackrel{\text{def}}{=} (\mathcal{S}'^i, \mathbf{S}^i, \Sigma, \mathbf{F}^i)$, then $\mathbf{f}$ is a prestructure morphism of $\mathbf{P}'^1$ to $\mathbf{P}'^2$ iff $\mathbf{f}$ is a prestructure morphism of $\mathbf{P}^1$ to $\mathbf{P}^2$.

If $\mathcal{S}^i \overset{\text{full-cat}}{\subseteq} \mathcal{S}'^i$ and $\mathbf{f}$ is a semi-strict (strict) prestructure morphism of $\mathbf{P}^1$ to $\mathbf{P}^2$ then $\mathbf{f}$ is a semi-strict (strict) prestructure morphism of $\mathbf{P}'^1$ to $\mathbf{P}'^2$.

Proof. The commutation relations do not depend on the choice of categories.

If $\mathcal{S}^i \overset{\text{full-cat}}{\subseteq} \mathcal{S}'^i$ and $\mathbf{f}$ is a semi-strict prestructure morphism of $\mathbf{P}^1$ to $\mathbf{P}^2$ then for each α , if

1. Each $\mathcal{S}_\alpha^i$ is a model category: f_α is a local $\mathcal{S}_\alpha^1$ - $\mathcal{S}_\alpha^2$ m-morphism of S_α^1 to S_α^2 and thus a local $\mathcal{S}_\alpha'^1$ - $\mathcal{S}_\alpha'^2$ m-morphism of S_α^1 to S_α^2 .
2. Each $\mathcal{S}_\alpha^i$ is a topological category: f_α is a local $\mathcal{S}_\alpha^1$ - $\mathcal{S}_\alpha^2$ morphism of S_α^1 to S_α^2 and thus a local $\mathcal{S}_\alpha'^1$ - $\mathcal{S}_\alpha'^2$ morphism of S_α^1 to S_α^2 .
3. Otherwise f_α is a morphism of $\mathcal{S}_\alpha^2$ and thus a morphism of $\mathcal{S}_\alpha'^2$.

If $\mathcal{S}^i \xrightarrow{\text{full-cat}} \mathcal{S}'^i$ and $\mathbf{f}$ is a strict prestructure morphism of $\mathbf{P}^1$ to $\mathbf{P}^2$ then each $f_\alpha \xrightarrow{\text{Ar}} \mathcal{S}_\alpha^2 \xrightarrow{\text{full-cat}} \mathcal{S}_\alpha'^2$, $\alpha < A$, and thus $\mathbf{f}$ is a strict prestructure morphism of $\mathbf{P}'^1$ to $\mathbf{P}'^2$. $\square$

$\text{Id}_{\mathbf{P}^i}$ is an identity morphism of $\mathbf{P}^i$.

Proof. $\mathbf{f} \circ \text{Id}_{\mathbf{S}^1} = \mathbf{f}$ and $\text{Id}_{\mathbf{S}^2} \circ \mathbf{f} = \mathbf{f}$ $\square$

{cor:premorph}

Corollary 10.12 (Prestructure morphisms). Let $\mathcal{S} = (\mathcal{S}_\alpha, \alpha < A)$ be a sequence of categories, $\mathbf{S} \stackrel{\text{def}}{=} (\mathcal{S}_\alpha, \alpha < A) \in \mathcal{S}$ be a sequence of spaces, $\mathbf{F} = (F_\gamma, \gamma < \Gamma)$ be a sequence of continuous functions with $\mathcal{S}$ -signatures Σ and $\mathbf{P} \stackrel{\text{def}}{=} (\mathcal{S}, \mathbf{S}, \Sigma, \mathbf{F})$ be a $\mathcal{S} - \Sigma$ prestructure. Then $\text{Id}_{\mathbf{P}}$ is a strict prestructure morphism from $\mathbf{P}$ to $\mathbf{P}$.

Proof. $\mathbf{S} \xrightarrow{\circ} \mathbf{S}$ and $F_\gamma = F_\gamma \upharpoonright \times_{\underline{\mathbf{S}}, \text{tail}}^{\sigma_\gamma} (\mathbf{S})$.

Each $\text{Id}_{\mathcal{S}_\alpha} = \text{Id}_{\mathcal{S}_\alpha} \xrightarrow{\text{Ar}} \mathcal{S}_\alpha$ and $\mathcal{S} \xrightarrow{\text{full-cat}} \mathcal{S}$. $\square$

{lem:premorphcomp}

Lemma 10.13 (Composition of prestructure morphisms). Let

1. $\mathcal{S}^i = (\mathcal{S}_\alpha^i, \alpha < A)$, $i \in [1, 4]$, be a sequence of categories
2. $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{S}_\alpha^i, \alpha < A) \in \mathcal{S}^i$ be a sequence of spaces
3. $\mathbf{F}^i = (F_\gamma^i, \gamma < \Gamma)$, be a sequence of continuous functions with $\mathbf{S}^i$ -signatures Σ
4. $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{S}^i, \mathbf{S}^i, \Sigma, \mathbf{F}^i)$ be a $\mathcal{S}^i - \Sigma$ prestructure
5. $\mathbf{f}^i \stackrel{\text{def}}{=} (f_\alpha^i: \mathcal{S}_\alpha^i \longrightarrow \mathcal{S}_\alpha^{i+1}, \alpha < A)$ and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_\alpha^i: \mathcal{S}_\alpha^i \longrightarrow \mathcal{S}_\alpha^{i+1}, \alpha < A)$ $i = 1, 2, 3$, be (sem-strict, strict) prestructure morphisms of $\mathbf{P}^i$ to $\mathbf{P}^{i+1}$.

$\mathbf{f}^{i+1} \circ \mathbf{f}^i$, $i \in [1, 3]$, is a prestructure morphism and $\mathbf{f}^3 \circ (\mathbf{f}^2 \circ \mathbf{f}^1) = (\mathbf{f}^3 \circ \mathbf{f}^2) \circ \mathbf{f}^1$.

Proof. If $\mathbf{f}^1$ Σ -commutes with $\mathbf{F}^1, \mathbf{F}^2$ and $\mathbf{f}^2$ Σ -commutes with $\mathbf{F}^2, \mathbf{F}^3$ then by lemma 10.3 (Σ -commutation) on page 43 $\mathbf{f}^2 \circ \mathbf{f}^1$ Σ -commutes with $\mathbf{F}^1, \mathbf{F}^3$. Thus $\mathbf{f}^2 \circ \mathbf{f}^1$ is a morphism.

Associativity is just lemma 1.14 (Sequence functions) on page 11. $\square$

{lem:seqfunc}

Let each f^i be a semi-strict prestructure morphism. Then $f^2 \circ f^1$ is a semi-strict prestructure morphism.

Proof. If $\mathcal{S}^1 \xrightarrow{\text{full-cat}} \mathcal{S}^2$ and $\mathcal{S}^2 \xrightarrow{\text{full-cat}} \mathcal{S}^3$ then $\mathcal{S}^1 \xrightarrow{\text{full-cat}} \mathcal{S}^3$. For each $\alpha < A$:

1. Each $\mathcal{S}_\alpha^i$, $i = 1, 2, 3$, is a model category:
Each f_α^i is a local $\mathcal{S}_\alpha^1$ - $\mathcal{S}_\alpha^{i+1}$ m-morphism of S_α^i to S_α^{i+1} . Then $f_\alpha^2 \circ f_\alpha^1$ is a local $\mathcal{S}_\alpha^1$ - $\mathcal{S}_\alpha^{i+2}$ m-morphism of S_α^i to S_α^{i+2} by lemma 6.17 (Composition of m-morphisms) on page 38. {lem:modMmorphcomp}
2. Each $\mathcal{S}_\alpha^i$, $i = 1, 2, 3$, is a topological category:
Each f_α^i is a local $\mathcal{S}_\alpha^i$ - $\mathcal{S}_\alpha^{i+1}$ morphism of S_α^i to S_α^{i+1} . Then $f_\alpha^2 \circ f_\alpha^1$ is a local $\mathcal{S}_\alpha^i$ - $\mathcal{S}_\alpha^{i+2}$ morphism of S_α^i to S_α^{i+2} by lemma 1.10 (Local morphisms) on page 10 {lem:topLocal}
3. Otherwise
Each f_α^i is a morphism of $\mathcal{S}_\alpha^{i+1}$. Then $f_\alpha^2 \circ f_\alpha^1$ is a morphism of $\mathcal{S}_\alpha^{i+2}$.

□

Let each f^i be a strict prestructure morphism. Then $f^2 \circ f^1$ is a strict prestructure morphism.

Proof. If $\mathcal{S}^1 \xrightarrow{\text{full-cat}} \mathcal{S}^2$ and $\mathcal{S}^2 \xrightarrow{\text{full-cat}} \mathcal{S}^3$ then $\mathcal{S}^1 \xrightarrow{\text{full-cat}} \mathcal{S}^3$.

If f_α^1 is a morphism of $\mathcal{S}_\alpha^2 \xrightarrow{\text{full-cat}} \mathcal{S}_\alpha^3$ and f_α^2 is a morphism of $\mathcal{S}_\alpha^3$ then $f_\alpha^2 \circ f_\alpha^1$ is a morphism of $\mathcal{S}_\alpha^3$. □

If each f^i is K -equivalent to g^i then each $f^{i+1} \circ f^i$ is K -equivalent to $g^{i+1} \circ g^i$.

Proof. $\left(\bigvee_{\substack{i \in [1-4] \\ \alpha \in K}} \right) f_\alpha^i = g_\alpha^i$ by definition 10.10 (K -equivalence of prestructure morphisms) on page 47. Then $\left(\bigvee_{\substack{i \in [1-3] \\ \alpha \in K}} \right) f_\alpha^2 \circ f_\alpha^1 = g_\alpha^2 \circ g_\alpha^1$ {def:premorphKequiv} □

Part VII

M-charts and m-atlases

{part:M-charts}

The literature defines fiber bundles and manifolds using the language of charts, atlases and transition functions; it has multiple equivalent definitions. Some authors start with topological spaces and define atlases over them. Some start with abstract sets and define atlases over them, deriving the topology from the atlas. Some start with an indexed set of open patches in a coordinate space and transition functions

among them satisfying a cocycle (compatibility) constraint, and then derive the total space as a quotient space of the disjoint union of the patches. Some authors use maximal atlases while others use equivalence classes of atlases.

This paper uses the first approach, explicitly making the relevant spaces topological spaces, but modifies the definitions of charts in order to make them fit more naturally into the context of local coordinate spaces. It adds a prefix, e.g., C^k , M , in order to avoid confusion with the conventional definitions. An m -atlas based on a topological spaces is closer to the conventional definitions of an atlas for a manifold while an m -atlas based on a model space is suitable for defining both manifolds and fiber bundles. Although simple manifolds and fiber bundles could both be defined directly in terms of maximal m -atlases, this paper has a different perspective, and uses the m -atlases as part of the more general Local Coordinate Space (LCS), presented in part VIII (Local Coordinate Spaces) on page 113.

{part:lcs}

Sections 11 to 12 define m -charts and m -atlases. Maximal m -atlases are essentially the same as manifolds.

Section 13 defines morphisms between m -atlases, referred to as m -atlas morphisms, in a fashion tailored to use in defining local coordinate spaces. It constructs categories of M -atlases and functors, and proves some basic results.

Section 14 defines morphisms between m -atlases in a fashion similar to that conventionally used for C^k maps between differentiable manifolds, in order to provide context. It constructs categories of M -atlases and functors, and proves some basic results.

11 M-charts

sec:M-charts {def:M-chart}

Definition 11.1 (M-charts). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces. An m -chart (U, V, ϕ) of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ consists of

1. A nonvoid model neighborhood $U \in \mathcal{E}$, known as a coordinate patch
2. A model neighborhood $V \in \mathcal{C}$
3. A model homeomorphism $\phi: U \xrightarrow{\cong} V$, known as a coordinate function

Remark 11.2. I consider it clearer to explicate the range, rather than the conventional usage of specifying only the domain and function or the minimalist usage of specifying only the function.

For every $u \in U$, (U, V, ϕ) is an m -chart of $\mathbf{E}$ in the coordinate space $\mathcal{C}$ at u and covers u . $\mathbf{E}$ is the total model space or total space for the chart and $\mathbf{C}$ is the coordinate model space or coordinate space for the chart.

Let E be a topological space and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be a model space. (U, V, ϕ) is an m -chart of E in the coordinate space $\mathcal{C}$ iff it is an m -chart of $\underset{\text{triv}}{E}$ in the coordinate space $\mathcal{C}$.

For every $x \in U$, (U, V, ϕ) is an m-chart of E in the coordinate space $\mathcal{C}$ at x . E is the total space for the chart and $\mathcal{C}$ is the coordinate model space or coordinate space for the chart.

{lem:M-chart}
{M-chart:openC}

Lemma 11.3 (M-charts).

1. Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces, and (U, V, ϕ) be an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ such that every open subset of V is a model neighborhood of $\mathbf{C}$. Then (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. U, V and ϕ satisfy the conditions of the definition:

U is a model neighborhood of $\mathbf{E}$: U is a model neighborhood of E , hence open.

V is a model neighborhood of $\mathbf{C}$: By hypothesis.

ϕ is a model homeomorphism from $\text{Mod}\left(U, \mathbf{E}_{\text{triv}}\right)$ to $\text{Mod}(V, \mathbf{C})$:

- (a) Since ϕ is a model homeomorphism from $\text{Mod}(U, \mathbf{E})$ to $\text{Mod}(V, \mathbf{C})$, it is a homeomorphism.
- (b) If V' is a model subneighborhood of V , then $\phi^{-1}[V']$ is open and hence a model neighborhood of $\mathbf{E}_{\text{triv}}$.
- (c) If U' is a model neighborhood of $\mathbf{E}_{\text{triv}}$ then it is open, hence $\phi[U']$ is an open subset of V and thus by hypothesis a model neighborhood of $\mathbf{C}$.

□

{M-chart:openE}

2. Let (U, V, ϕ) be an m-chart of E in the coordinate space $\mathbf{C}$. (U, V, ϕ) is an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ iff every open subset of U is a model neighborhood of $\mathbf{E}$.

Proof. If (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$ and also an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ then let $U' \subseteq U$ be open. Since (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$, ϕ^{-1} is a model function from $\text{Mod}(V, \mathbf{C})$ to $\mathbf{E}_{\text{triv}}$ and $\phi[U']$ is a model neighborhood of $\mathbf{C}$. Since (U, V, ϕ) is an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$, ϕ is a model function from $\text{Mod}(U, \mathbf{E})$ to $\mathbf{C}$ and thus $\phi^{-1}[\phi[U']]$ is a model neighborhood of $\mathbf{E}$.

If (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$ and every open $U' \subseteq U$ is a model neighborhood of $\mathbf{E}$ then

- (a) Since $U \subseteq U$ is open, U is a model neighborhood of $\mathbf{E}$ by hypothesis.
- (b) V is a model neighborhood of $\mathbf{C}$ by hypothesis.

- (c) If $V' \subseteq V$ is a model neighborhood of $\mathbf{C}$ then $\phi^{-1}[V']$ is open and hence a model neighborhood of $\mathbf{E}$ by hypothesis.

ϕ^{-1} is a model function from $\text{Mod}\left(U, \underset{\text{triv}}{E}\right)$ to $\text{Mod}(V, \mathbf{C})$. if $U' \subseteq U$ is a model neighborhood of $\mathbf{E}$ then U' is open and hence a model neighborhood of $\underset{\text{triv}}{E}$. Then $\phi[U']$ is a model neighborhood of $\mathbf{C}$.

□

{cor:M-chart}

Corollary 11.4 (M-charts). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces, $\mathbf{C}$ be fine grained and (U, V, ϕ) be an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$. Then (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. Since V is open and $\mathbf{C}$ is fine grained, every open subset of V is a model neighborhood of $\mathbf{C}$. The result follows by item 1 of [lemma 11.3 \(M-charts\)](#) on page 51. □

{M-chart:openC}

{def:M-subchart}

Definition 11.5 (Subcharts). Let (U, V, ϕ) be an m-chart of $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and $U' \subseteq U$ a nonvoid model neighborhood of $\mathbf{E}$. Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

Let (U, V, ϕ) be an m-chart of E in the coordinate space $(C, \mathcal{C})$ and $U' \subseteq U$ an open set of E . Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

(U', V', ϕ') is a subchart of (U, V, ϕ) at x iff (U', V', ϕ') is a subchart of (U, V, ϕ) and $x \in U'$.

{lem:M-subchart}

Lemma 11.6 (Subcharts). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces.

Let (U, V, ϕ) be an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ and (U', V', ϕ') a subchart of (U, V, ϕ) . Then (U', V', ϕ') is an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$.

Proof. U' , V' and ϕ' satisfy the conditions of the definition of an m-chart:

1. U' is a model neighborhood of $\mathbf{E}$ by the definition of subchart.
2. Since U' is a model neighborhood and ϕ is a model homeomorphism, $V' = \phi[U']$ is a model neighborhood.
3. Since ϕ and ϕ^{-1} are model functions, so are $\phi' = \phi|_{U', V'}$ and $\phi'^{-1} = \phi^{-1}|_{V', U'}$. Thus ϕ' is a model homeomorphism.

□

Let (U, V, ϕ) be an m-chart of E in the coordinate space $\mathbf{C}$ and (U', V', ϕ') a subchart of (U, V, ϕ) . Then (U', V', ϕ') is an m-chart of E in the coordinate space $\mathbf{C}$.

Proof. U' , V' and ϕ' satisfy the conditions of the definition of an m-chart:

1. U' is open by the definition of subchart.
2. Since ϕ is a homeomorphism and U' is open, $V' = \phi[U']$ is open.
3. Since ϕ is a homeomorphism that maps open sets into model neighborhoods, $\phi|_{U', V'}$ is a homeomorphism that maps open sets into model neighborhoods.

□

{def:M-compat}

Definition 11.7 (M-compatibility). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be model spaces and (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of $\mathbf{E}$ in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) in the coordinate space $\mathbf{C}$ iff either

1. U_1 and U_2 are disjoint
2. The transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$.

(U_1, V_1, ϕ_1) is also properly m-compatible with (U_2, V_2, ϕ_2) in the space $\mathbf{E}$ and the coordinate space $\mathbf{C}$, and properly m-compatible in the space $\mathbf{E}$ and the coordinate space $\mathbf{C}$ with (U_2, V_2, ϕ_2) , iff for every model neighborhoods $U'_i \subseteq U_i$

1. For every morphism $f: U'_1 \longrightarrow U'_2$ of $\mathbf{E}$, the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \longrightarrow \phi_2[U'_2]$ is a morphism of $\mathcal{C}$.
2. For every morphism $g: U'_2 \longrightarrow U'_1$ of $\mathbf{E}$, the composition $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \longrightarrow \phi_1[U'_1]$ is a morphism of $\mathcal{C}$.

Remark 11.8. This is a stronger condition than requiring that for every $f: U_1 \longrightarrow U_2$ and $g: U_2 \longrightarrow U_1$ the compositions be morphisms of $\mathcal{C}$.

Let E be a topological space, $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ be a model space and (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of E in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) iff either

1. U_1 and U_2 are disjoint
2. The transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$

(U_1, V_1, ϕ_1) is also properly m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$, and properly m-compatible in the space E and the coordinate space $\mathbf{C}$ with (U_2, V_2, ϕ_2) , iff for every open $U'_i \subseteq U_i$

1. For every continuous $f: U'_1 \longrightarrow U'_2$, the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \longrightarrow \phi_2[U'_2]$ is a morphism of $\mathcal{C}$.
2. For every continuous $g: U'_2 \longrightarrow U'_1$, the composition $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \longrightarrow \phi_1[U'_1]$ is a morphism of $\mathcal{C}$.

In both cases, the space and coordinate space may be omitted when they are clear from context.

{lem:M-compat}

Lemma 11.9 (M-compatibility).

1. Let (U_i, V_i, ϕ_i) , $i = 1, 2$. be a chart of E in the coordinate space $\mathbf{C}$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ iff (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space $\underset{\text{triv}}{E}$ and the coordinate space $\mathbf{C}$.

Proof. $U_1 \cap U_2$ and the transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is the same for the case of E and the case of $\underset{\text{triv}}{E}$.

Any $U'_i \subseteq U_i$ is a model neighborhood of $\underset{\text{triv}}{E}$ iff it is open.

Any $f: U'_1 \longrightarrow U'_2$ is a morphism of $\underset{\text{triv}}{E}$ iff it is continuous.

Any $g: U'_2 \longrightarrow U'_1$ is a morphism of $\underset{\text{triv}}{E}$ iff it is continuous.

The compositions $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U'_1] \longrightarrow \phi_2[U'_2]$ and $\phi_1 \circ g \circ \phi_2^{-1}: \phi_2[U'_2] \longrightarrow \phi_1[U'_1]$ are the same for the case of E and the case of $\underset{\text{triv}}{E}$. $\square$

{cor:M-compat}

Corollary 11.10. Let (U_i, V_i, ϕ_i) , $i = 1, 2$. be a chart of E in the coordinate space $\mathbf{C}$ and let (U'_i, V'_i, ϕ'_i) be a subchart. If (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. By lemma 11.9 above, (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space E and the coordinate space $\mathbf{C}$ iff (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space $\underset{\text{triv}}{E}$ and the coordinate space $\mathbf{C}$. $\square$

The result follows from lemma 11.9.

{lem:M-compat}
{lem:M-compat-sym}

Lemma 11.11 (Symmetry of m-compatibility). Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(\mathbf{C}, \mathcal{C})$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ iff (U_2, V_2, ϕ_2) is (properly) m-compatible with (U_1, V_1, ϕ_1) .

Proof. It suffices to prove the implication in only one direction.

1. $U_1 \cap U_2 = U_2 \cap U_1$.
2. Since the transition function $t = \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$, so is $t^{-1} = \phi_1 \circ \phi_2^{-1}: \phi_2[U_1 \cap U_2] \longrightarrow \phi_1[U_1 \cap U_2]$.
3. The constraint for proper maximality is intrinsically symmetrical.

$\square$

Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of E in the coordinate space $(C, \mathcal{C})$. Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ iff (U_2, V_2, ϕ_2) is (properly) m-compatible with (U_1, V_1, ϕ_1) .

Proof. The same proof applies as above. $\square$

{lem:M-compatsub}

Lemma 11.12 (M-compatibility of subcharts). Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$, (U_1, V_1, ϕ_1) be (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$ and (U'_i, V'_i, ϕ'_i) be a subchart. Then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. Since subcharts are charts, U'_i and V'_i are model neighborhoods. If $U_1 \cap U_2 = \emptyset$ then $U'_1 \cap U'_2 = \emptyset$ and thus (U'_1, V'_1, ϕ'_1) is m-compatible with (U'_2, V'_2, ϕ'_2) . Otherwise, the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \longrightarrow \phi_2[U_1 \cap U_2]$ is an isomorphism of $\mathcal{C}$ and hence $t_2^1: \phi_1[U'_1 \cap U'_2] \xrightarrow{\cong} \phi_2[U'_1 \cap U'_2]$ is an isomorphism of $\mathcal{C}$.

If (U_1, V_1, ϕ_1) is properly m-compatible with (U_2, V_2, ϕ_2) , $U''_i \subseteq U'_i$, $i = 1, 2$, is a model neighborhood and $f: U''_1 \longrightarrow U''_2$ is a morphism of E , then each $U''_i \subseteq U_i$ and thus by definition 11.7 the composition $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1[U''_1] \longrightarrow \phi_2[U''_2]$ is a morphism of $\mathbf{C}$. $\square$

{def:M-compat}

Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be an m-chart of E in the coordinate space $(C, \mathcal{C})$, (U'_i, V'_i, ϕ'_i) be a subchart and (U_1, V_1, ϕ_1) be (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$. Then (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) .

Proof. By lemma 11.9 (M-compatibility) on page 54, (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) in the space E and the coordinate space $\mathbf{C}$. Then by lemma 11.12 above, (U'_1, V'_1, ϕ'_1) is (properly) m-compatible with (U'_2, V'_2, ϕ'_2) in the space E and the coordinate space $\mathbf{C}$. $\square$

{lem:M-compat}

{lem:M-compatsub}

The result follows from lemma 11.9. $\square$

{lem:M-compat}
{cor:M-compatsub}

Corollary 11.13 (M-compatibility of subcharts). Let (U, V, ϕ) be an m-chart of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and (U', V', ϕ') a subchart. Then (U', V', ϕ') is properly m-compatible with (U, V, ϕ) in the space E and the coordinate space $\mathbf{C}$.

Proof. (U, V, ϕ) is m-compatible with itself and is a subchart of itself, $\square$

Let (U, V, ϕ) be an m-chart of E in the coordinate space $(C, \mathcal{C})$ and (U', V', ϕ') a subchart. Then (U', V', ϕ') is m-compatible with (U, V, ϕ) in the space E and the coordinate space $\mathbf{C}$.

Proof. (U, V, ϕ) is m-compatible with itself and is a subchart of itself, $\square$

Definition 11.14 (Covering by m-charts). Let $\mathbf{A}$ be a set of charts of the topological space E in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. $\mathbf{A}$ covers E iff $\pi_1[\mathbf{A}]$ covers E .

Let $\mathbf{A}$ be a set of charts of the model space $\mathbf{E} = (E, \mathcal{E})$ in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. $\mathbf{A}$ covers $\mathbf{E}$ iff $\pi_1[\mathbf{A}]$ covers E .

12 M-atlases

{sec:M-atlas}

A set of charts can be atlases for different coordinate model spaces even if it is for the same total model space. In order to aggregate atlases into categories, there must be a way to distinguish them. Including the two⁸ spaces in the definitions of the categories serves the purpose.

{def:M-atlas}

Definition 12.1 (M-atlases). Let $\mathbf{A}$ be a set of mutually (properly) m-compatible m-charts of $\mathbf{E} = (E, \mathcal{E})$ in the coordinate space $\mathbf{C} = (C, \mathcal{C})$. Then

1. $\mathbf{A}$ is an (a proper) m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C})$, iff $\mathbf{A}$ covers E .
2. $\mathbf{A}$ is a full (proper) atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C})$, iff $\mathbf{A}$ covers E and $\pi_2[\mathbf{A}]$ covers C .
3. The triple $(\mathbf{A}, \mathbf{E}, \mathbf{C})$ refers to $\mathbf{A}$ considered as an m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$.
4. E is the total model space or total space for the atlas
5. C is the coordinate model space or coordinate space for the atlas.

Let E be a topological space, $\mathbf{C} = (C, \mathcal{C})$ a model space and $\mathbf{A}$ be a set of mutually m-compatible m-charts of E in the coordinate space $\mathbf{C}$. Then

6. $\mathbf{A}$ is an m-atlas of E in the coordinate space $\mathbf{C}$, abbreviated as $\text{isAtl}_{\text{Ob}}(\mathbf{A}, E, \mathbf{C})$, iff $\mathbf{A}$ is an m-atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\mathbf{C}$.
7. $\mathbf{A}$ is a full atlas of E in the coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}(\mathbf{A}, E, \mathbf{C})$, iff $\mathbf{A}$ is a full atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\mathbf{C}$.
8. The triple $(\mathbf{A}, E, \mathbf{C})$ refers to $\mathbf{A}$ considered as an m-atlas of E in the coordinate space $\mathbf{C}$.
9. E is the total space for the atlas.
10. C is the coordinate model space or coordinate space for the atlas.

Let $\mathbf{A}$ be an (a proper) m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$. $\mathbf{A}$ is fine grained iff whenever $(U, V, \phi) \in \mathbf{A}$, every subchart of (U, V, ϕ) is also an atlas of $\mathbf{A}$.

⁸The spaces are redundant, but convenient.

Remark 12.2. $\mathbf{A}$ can be a fine graind (proper) atlas of E in the coordinate space $\mathbf{C}$ even if $\mathbf{C}$ is not fine grained.

Let $\mathbf{A}$ be an (a proper) m-atlas of E in the coordinate space $\mathbf{C}$. $\mathbf{A}$ is fine grained iff whenever $(U, V, \phi) \in \mathbf{A}$, every subchart of (U, V, ϕ) is also an atlas of $\mathbf{A}$.

By abuse of language we write $U \in \mathbf{A}$ for $U \in \pi_1[\mathbf{A}]$.

{lem:M-atlas}

Lemma 12.3 (M-atlases). Let $\mathbf{A}$ be an m-atlas of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. If $\mathbf{C}$ is fine grained then $\mathbf{A}$ is an m-atlas of E in the coordinate space $\mathbf{C}$.

Proof. Let $(U, V, \phi) \in \mathbf{A}$. Then (U, V, ϕ) is an m-chart of E in the coordinate space $\mathbf{C}$ by corollary 11.4 (M-charts) on page 52. $\square$

{cor:M-chart}
{def:M-CompatATL}

Definition 12.4 (M-compatibility with m-atlases). An m-chart (U, V, ϕ) of $(E, \mathcal{E})$ is (properly) m-compatible with an m-atlas $\mathbf{A}$ in the space E and the coordinate space $\mathbf{C}$ iff it is (properly) m-compatible in the space E and the coordinate space $\mathbf{C}$ with every chart in $\mathbf{A}$.

An m-chart (U, V, ϕ) of E is (properly) m-compatible with an m-atlas $\mathbf{A}$ in the space E and the coordinate space $\mathbf{C}$ iff it is (properly) m-compatible in the space E and the coordinate space $\mathbf{C}$ with every chart in $\mathbf{A}$.

{lem:M-CompatATL}

Lemma 12.5 (M-compatibility with m-atlases). Let $\mathbf{A}$ be an m-atlas of $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. If $\mathbf{C}$ is fine grained then $\mathbf{A}$ is an m-atlas of E in the coordinate space $\mathbf{C}$.

Proof.

$\square$

{lem:M-CompatsubATL}

Lemma 12.6 (M-compatibility of subcharts with atlases). Let $\mathbf{A}$ be an (a proper) m-atlas of $(E, \mathcal{E})$ in the coordinate space $(C, \mathcal{C})$ and (U_1, V_1, ϕ_1) an m-chart in $\mathbf{A}$. Then any subchart of (U_1, V_1, ϕ_1) is properly m-compatible with $\mathbf{A}$ in the space E and the coordinate space $\mathbf{C}$.

Proof. Let (U'^1, V'^1, ϕ'^1) be a subchart of (U^1, V^1, ϕ^1) and (U_2, V_2, ϕ_2) be another chart in $\mathbf{A}$. Then (U_1, V_1, ϕ_1) is properly compatible with (U_2, V_2, ϕ_2) by definition 12.1 (M-atlases) on page 56, (U_2, V_2, ϕ_2) is properly m-compatible with itself and (U_2, V_2, ϕ_2) is a subchart of itself. (U'^1, V'^1, ϕ'^1) is properly m-compatible with (U_2, V_2, ϕ_2) by lemma 11.12 (M-compatibility of subcharts) on page 55. Since (U_2, V_2, ϕ_2) is arbitrary, (U'^1, V'^1, ϕ'^1) is properly m-compatible with $\mathbf{A}$ by definition 12.4 (M-compatibility with m-atlases) on page 57. $\square$

{def:M-atlas}

{lem:M-compatsub}
{def:M-CompatATL}

Let $\mathbf{A}$ be an m-atlas of E in the coordinate space $(C, \mathcal{C})$ and (U_1, V_1, ϕ_1) an m-chart in $\mathbf{A}$. Then any subchart of (U_1, V_1, ϕ_1) is properly m-compatible with $\mathbf{A}$ in the space E and the coordinate space $\mathbf{C}$.

Proof. The same proof applies in as above. $\square$

{\lem:M-ATLextensions}

Lemma 12.7 (Extensions of m-atlases). Let $E = (E, \mathcal{E})$ and $C = (C, \mathcal{C})$ be model spaces, A an m-atlas of E in the coordinate space C , and (U_i, V_i, ϕ_i) , $i = 1, 2$, an m-chart of E in the coordinate space C (properly) m-compatible with A in the space E and the coordinate space C . Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) .

Proof. If $U_1 \cap U_2 = \emptyset$ then (U_1, V_1, ϕ_1) is m-compatible with (U_2, V_2, ϕ_2) . Otherwise, ϕ_1 is a model homeomorphism, $U_1 \cap U_2$ is a model neighborhood of E , $\phi_1[U_1 \cap U_2]$ is a model neighborhood of C , $\phi_2[U_1 \cap U_2]$ is a model neighborhood of

C and $\phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2] \xrightarrow{\cong} \phi_2[U_1 \cap U_2]$ is a homeomorphism. It remains to show that $\phi_2 \circ \phi_1^{-1}|_{\phi_1[U_1 \cap U_2]}$ is a model homeomorphism. Let $(U_\alpha, V_\alpha, \phi_\alpha)$, $\alpha < A$,

be charts in A such that $U_1 \cap U_2 \subseteq \bigcup_{\alpha < A} U_\alpha$ and $(\bigcap_{\alpha < A} U_\alpha) \cap U_1 \cap U_2 \neq \emptyset$.

Since the charts are m-compatible with $(U_\alpha, V_\alpha, \phi_\alpha)$, the compositions

$\phi_2 \circ \phi_\alpha^{-1}: \phi_\alpha[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha]$ and $\phi_\alpha \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_\alpha[U_1 \cap U_2 \cap U_\alpha]$ are isomorphisms of C and thus the composition $\phi_2 \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha] =$

$\phi_2 \circ \phi_\alpha^{-1} \circ \phi_\alpha \circ \phi_1^{-1}: \phi_1[U_1 \cap U_2 \cap U_\alpha] \rightarrow \phi_2[U_1 \cap U_2 \cap U_\alpha]$ is an isomorphism of C . Then by [item 6](#), the restricted sheaf property, of [definition 2.1 \(Model spaces\)](#) {\def:model} on page [17](#), $\phi_2 \circ \phi_1^{-1}$ is an isomorphism of C .

If each (U_i, V_i, ϕ_i) is properly m-compatible with A then let $f: U'_1 \rightarrow U'_2$ and $g: U'_2 \rightarrow U'_1$ be morphisms of E and $U'_i \subseteq U_i$, $i = 1, 2$. Let u be an arbitrary point in U_1 and $(U_u, V_u, \phi_u) \in A$ be an arbitrary chart at u . Since (U_1, V_1, ϕ_1) is properly m-compatible with A , it is properly m-compatible with $(U_u, V_u, \phi_u) \in A$, thus the compositions $\phi_2 \circ f \circ \phi_u^{-1}: \phi_u^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2]$ and $\phi_u \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_u^{-1}[U_u]$ are morphisms of C , their composition $\phi_2 \circ f \circ \phi_u^{-1} \circ \phi_u \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2] =$
 $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1^{-1}[U_u] \rightarrow \phi_2^{-1}[U'_2]$ is a morphism of C and by [item 6](#), the restricted sheaf property, of [definition 2.1 \(Model spaces\)](#) on page [17](#), $\phi_2 \circ f \circ \phi_1^{-1}: \phi_1^{-1}[U'_1] \rightarrow \phi_2^{-1}[U'_2]$ is a morphism of C . {\def:model}

The same proof applies in the opposite direction. □

Let E be a topological space, $C = (C, \mathcal{C})$ a model space, A an (a proper) m-atlas of E in the coordinate space C and (U_i, V_i, ϕ_i) , $i = 1, 2$, a chart (properly) m-compatible with A in the space E and the coordinate space C . Then (U_1, V_1, ϕ_1) is (properly) m-compatible with (U_2, V_2, ϕ_2) .

Proof. The same proof applies as above. □

{\def:M-ATLmax}

Definition 12.8 (Maximal m-atlases). Let E and C be model spaces and A an m-atlas of E in the coordinate space C . A is a (proper) maximal m-atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, C)$, iff A cannot be extended by adding an additional (properly) m-compatible chart.

Let E be a topological space, $C = (C, \mathcal{C})$ be a model space and A be an m-atlas of E in the coordinate space C . A is a (properly) maximal m-atlas of E in the

coordinate space $\mathbf{C}$, abbreviated $\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C})$, iff $\mathbf{A}$ is an m-atlas that cannot be extended by adding an additional (properly) m-compatible chart.

Remark 12.9. There is no sheaf condition⁹; the union of a set of coordinate patches in the maximal atlas whose coordinate functions match on the intersections need not be a coordinate patch in the atlas.

$$\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \quad (12.1)$$

$$\text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \quad (12.2)$$

Theorem 12.10 (Existence and uniqueness of maximal m-atlases). Let $\mathbf{A}$ be an (a proper) m-atlas of $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. Then there exists a unique (proper) maximal m-atlas $\text{Atlas}[(\text{proper})](\mathbf{A}, \mathbf{E}, \mathbf{C})$ of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all m-atlases of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ containing $\mathbf{A}$ and (properly) m-compatible in the coordinate space $\mathbf{C}$ with all of the m-charts in $\mathbf{A}$. Let $\mathbf{P}_{\text{max}}$ be a maximal chain of $\mathbf{A}$. Then $\text{Atlas}[(\text{proper})](\mathbf{A}, \mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\text{max}} \mathbf{P}_{\text{max}}$ is a (proper) maximal m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$. Uniqueness follows from lemma 12.7 (Extensions of m-atlases) on page 58. $\square$

Corollary 12.11 (Existence and uniqueness of maximal m-atlases). Let $\mathbf{A}$ be an (a proper) m-atlas of $\mathbf{E}$ in the coordinate space $\mathbf{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$. Then there exists a unique (proper) maximal m-atlas $\text{Atlas}[(\text{proper})](\mathbf{A}, \mathbf{E}, \mathbf{C})$ of $\mathbf{E}$ in the coordinate space $\mathbf{C}$ m-compatible with $\mathbf{A}$.

Proof. $\text{Atlas}[(\text{proper})](\mathbf{A}, \mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \text{Atlas}[(\text{proper})](\mathbf{A}, \mathbf{E}, \mathbf{C})$. $\square$

Definition 12.12 (Sets of m-atlases). Let $\mathbf{E}$ and $\mathbf{C}$ be model spaces.

$$\mathcal{Atl}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, \mathbf{E}, \mathbf{C}) \mid \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \right\} \quad (12.3)$$

Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces.

⁹However, note item 6 (restricted sheaf condition) of definition 2.1 (Model spaces) on page 17.

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ (A, (E, \mathcal{E}), (C, \mathcal{C})) \mid \right. \\ \left. (E, \mathcal{E}) \in \mathbf{E} \wedge (C, \mathcal{C}) \in \mathbf{C} \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, (E, \mathcal{E}), (C, \mathcal{C})) \right\} \quad (12.4)$$

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ (A, E, (C, \mathcal{C})) \mid \right. \\ \left. E \in \mathbf{E} \wedge (C, \mathcal{C}) \in \mathbf{C} \wedge \text{isAtl}_{\text{Ob}}^{(\text{proper})}(\mathbf{A}, E, (C, \mathcal{C})) \right\} \quad (12.5)$$

Lemma 12.13 (Sets of m-atlases). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. {lem:M-ATLsets}

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.6)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.7)$$

Proof. The result follows from definition 12.1 (M-atlases) on page 56, definition 12.8 (Maximal m-atlases) on page 58 and definition 12.12 above. {def:M-ATLmax} {def:M-ATLsets} $\square$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.8)$$

Proof. The result follows from definition 12.8 (Maximal m-atlases) on page 58 and definition 12.12 above. {def:M-ATLmax} {def:M-ATLsets} {cor:M-ATLsets} $\square$

Corollary 12.14 (Sets of m-atlases). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.9)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.10)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.11)$$

$$\mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \subseteq \mathcal{Att}_{\text{Ob}}^{(\text{proper})}(\mathbf{E}, \mathbf{C}) \quad (12.12)$$

Proof. The result follows from definition 12.1 (M-atlases) on page 56, definition 12.8 (Maximal m-atlases) on page 58 and definition 12.12 above. {def:M-ATLmax} {def:M-ATLsets} $\square$

13 M-atlas morphisms and functors

{sec:M-ATLmorph}

This section introduces a taxonomy for morphisms between m-atlases, defines categories of m-atlases, defines functors among them and proves some basic results.

13.1 M-atlas morphisms

{sub:M-ATLmorph}

This subsection introduces the notions of m-atlas morphisms, strict m-atlas morphisms and semi-strict m-atlas morphisms.

13.1.1 Definitions of m-atlas morphisms

{def:M-ATLmorph}

Definition 13.1 (M-atlas morphisms for model spaces). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i and $f \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$ a pair of model functions.

f is an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$ and an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{E}^1, \mathcal{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f_0, f_1)$, iff for any charts $(U^i, V^i, \phi^i: U^i \xrightarrow{\cong} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$ and any $u^1 \in I$, there exists a sub-chart $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\cong} V'^1) \in A^1$ at u^1 , a model neighborhood $U'^2 \subseteq U^2$ of $u^2 \stackrel{\text{def}}{=} f_0(u^1)$ and a chart $(U'^2, V'^2, \phi'^2: U'^2 \xrightarrow{\cong} V'^2) \in A^2$ at u^2 such that $f_0[U'^1] \subseteq U'^2$ and diagram (13.1) below is commutative, as shown in fig. 7 (Completed m-atlas morphism) on page 62.

{fig:AtlM1}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{c} \{u^1\} \xrightarrow{i} U'^1 \xrightarrow[\cong]{\phi^1} V'^1 \xrightarrow{f_1} V'^2, \\ U'^1 \xrightarrow{f_0} U'^2 \xrightarrow[\cong]{\phi'^2} V'^2 \end{array} \right. \quad (13.1) \quad \{\text{eq:AtlM1}\}$$

The triple $(f, (A^1, E^1, C^1), (A^2, E^2, C^2))$ will refer to f considered as an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is also a constrained E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$ and a constrained $\mathcal{E}^1$ - $\mathcal{E}^2$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{E}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f_0, f_1)$, iff f_0 and f_1 are




Figure 7: Completed m-atlas morphism

{fig:AtlM1}

constrained, i.e., $f_0[E^1]$ is contained in a model neighborhood of E^2 and $f_1[C^1]$ is contained in a model neighborhood of C^2 .

Remark 13.2. If $\mathbf{f}$ is constrained then $\text{Top}(E^1)$ is a model neighborhood of E^1 and $\text{Top}(C^1)$ is a model neighborhood of C^1 .

{def:M-ATLmorphTop}

Definition 13.3 (M-atlas morphisms for topological spaces). Let $\mathcal{C}^i$, $i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space and $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$.

A pair of functions $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ is an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^i$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^i$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff it is an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$.

The triple $(\mathbf{f}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2))$ will refer to $\mathbf{f}$ considered as an E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is also a constrained E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, E^1, \mathbf{C}^1, \mathbf{A}^2, E^2, \mathbf{C}^2, f_0, f_1)$, and a constrained E^1 - E^2 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, E^1, \mathcal{C}^1, \mathbf{A}^2, E^2, \mathcal{C}^2, f_0, f_1)$, iff f_1 is constrained, i.e., $f_1[C^1]$ is contained in a model neighborhood of C^2 .

{def:M-ATLident}

Definition 13.4 (M-atlas inclusion and identity morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{C}^i$
3. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
4. $\mathbf{C}^i \in \mathcal{C}^i$

$$5. \mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$$

6. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$

$$7. \mathbf{A}^1 \subseteq \mathbf{A}^2$$

The $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The m-atlas inclusion morphism of $(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$ is $\mathbf{Id}_{(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right), \mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The $\mathbf{E}^i$ m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i, \mathbf{C}^2$ is $\mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The m-atlas identity morphism of $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ is

$\mathbf{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}, (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \right), \mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}$ considered as an $\mathbf{E}^i$ m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

Let

1. $\mathbf{E}^i$ be a topological space, $i = 1, 2$

$$2. \mathbf{E}^1 \stackrel{\text{subsp}}{\subseteq} \mathbf{E}^2$$

3. $\mathcal{C}^i$ be a model category

$$4. \mathbf{C}^i \in \mathcal{C}^i$$

$$5. \mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$$

6. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$

$$7. \mathbf{A}^1 \subseteq \mathbf{A}^2$$

The $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}$.

The m-atlas inclusion morphism of $(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$ is

$\mathbf{Id}_{(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right), \mathbf{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)}$ considered as an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The $\mathbf{E}^i$ m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$ is $\mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}$.

The m-atlas identity morphism of $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ is

$\mathbf{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}, (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \right), \mathbf{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}$ considered as an $\mathbf{E}^i$ m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

{def:M-ATLmorphequiv}

Definition 13.5 (Equivalence of m-atlas morphisms). Let $E^i, C^i, i = 1, 2$, be model spaces, A^i be an m-atlas of E^i in the coordinate space C^i , and

$f \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$ and $g \stackrel{\text{def}}{=} (g_0: E^1 \rightarrow E^2, g_1: C^1 \rightarrow C^2)$ be m-atlas morphisms of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then f is equivalent to g as an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated $f: (A^1, E^1, C^1) \rightarrow (A^2, E^2, C^2)$ is equivalent to $g: (A^1, E^1, C^1) \rightarrow (A^2, E^2, C^2)$ and $(f, (A^1, E^1, C^1), (A^2, E^2, C^2))$ is equivalent to $(g, (A^1, E^1, C^1), (A^2, E^2, C^2))$ iff $f_0 = g_0$.

By abuse of language we write f is equivalent to g when the spaces and atlases are understood by context.

Let $E^i, i = 1, 2$, be a topological space, C^i be a model space, A^i be an m-atlas of E^i in the coordinate space C^i and $f \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$, and $g \stackrel{\text{def}}{=} (g_0: E^1 \rightarrow E^2, g_1: C^1 \rightarrow C^2)$ be m-atlas morphisms of A^1 to A^2 in the coordinate spaces C^1, C^2 . Then f is equivalent to g as an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated $f: (A^1, E^1, C^1) \rightarrow (A^2, E^2, C^2)$ is equivalent to $g: (A^1, E^1, C^1) \rightarrow (A^2, E^2, C^2)$, iff $f_0 = g_0$.

By abuse of language we write f is equivalent to g when the spaces and atlases are understood by context.

13.1.2 Definitions of semi-strict and strict

{def:M-ATLmorphsemi}

Definition 13.6 (Semi-strict m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i and $f \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$ an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a semi-strict E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{semi-strict}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$, iff f_0 is a local m-morphism of E^1 to E^2 , f_1 is a local m-morphism of C^1 to C^2 and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \rightarrow V^2$ is a local C^1 - C^2 m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $E^i \in \mathcal{E}^i$, each $C^i \in \mathcal{C}^i, E^1 \stackrel{\text{strict-mod}}{\subseteq} E^2$ and $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$.

f is a semi-strict E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{semi-strict}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff f_0 is a local m-morphism of E^1 to E^2 , f_1 is a local $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism of C^1 to C^2 and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \rightarrow V^2$ is a local $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $E^1-E^2-\mathcal{E}^1-\mathcal{E}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $E^i \in \mathbf{E}^i$, $E^1 \stackrel{\text{Ob}}{\subseteq} E^2$ and $\mathcal{E}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathcal{E}^2$.

f is a semi-strict $\mathcal{E}^1-\mathcal{E}^2-E^1-E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff f_0 is a local $\mathcal{E}^1-\mathcal{E}^2$ m-morphism of E^1 to E^2 , f_1 is a local m-morphism of C^1 to C^2

and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow U^2$ is a local C^1-C^2 m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathcal{E}^1-\mathcal{E}^2-E^1-E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $C^i \in \mathbf{C}^i$, $\mathcal{E}^1 \stackrel{\text{Ob}}{\subseteq} \mathcal{E}^2$ and $C^1 \stackrel{\text{strict-cat}}{\subseteq} C^2$.

f is a semi-strict $\mathcal{E}^1-\mathcal{E}^2-E^1-E^2-\mathcal{E}^1-\mathcal{E}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2, \mathcal{E}^1, \mathcal{E}^2)$, iff f_0 is a local $\mathcal{E}^1-\mathcal{E}^2$ m-morphism of E^1 to E^2 and f_1 is a local $\mathcal{E}^1-\mathcal{E}^2$ m-morphism of C^1 to C^2 and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local $\mathcal{E}^1-\mathcal{E}^2$ m-morphism of $\phi^1[I]$ to V^2 .

It is also a coherent semi-strict $\mathcal{E}^1-\mathcal{E}^2-E^1-E^2-\mathcal{E}^1-\mathcal{E}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ and $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$.

By abuse of language we will write f is coherent when it is a $(\mathcal{E}^1-\mathcal{E}^2-E^1-E^2-\mathcal{E}^1-\mathcal{E}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 where the model spaces, model categories and atlases are understood by context.

{def:M-ATLmorphstrict}

Definition 13.7 (Strict m-atlas morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathbf{E}^i$
3. $C^i \in \mathbf{C}^i$
4. A^i be an m-atlas of E^i in the coordinate space C^i
5. $f \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ be an m-atlas morphism

f is a strict E^1-E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$, iff f_0 is an m-morphism of E^1 to E^2 , f_1 is an m-morphism of C^1 to C^2 and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$, $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow U^2$ is a morphism of $\text{Mod}(V^2, C^2)$.

It is also a coherent strict E^1-E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff each $E^i \in \mathbf{E}^i$, each $C^i \in \mathbf{C}^i$, $E^1 \stackrel{\text{Ob}}{\subseteq} E^2$ and $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$.

$\mathbf{f}$ is a strict $\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff f_0 is an m-strict morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$, f_1 is an m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of $\mathcal{C}^2$.

It is also a coherent strict $\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff each $E^i \in \mathbf{E}^i, E^1 \stackrel{\text{Ob}}{\subseteq} E^2$ and $\mathcal{C}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathcal{C}^2$.

$\mathbf{f}$ is a strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff f_0 is an m-strict morphism of $\mathcal{E}^1$ to $\mathcal{E}^2$, f_1 is an m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$ and for every

$(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow U^2$ is a morphism of $\text{Mod}(\phi^1[I], \mathbf{C}^2)$.

It is also a coherent strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff $\mathbf{C}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathbf{C}^2, \mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ and $\mathbf{C}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathbf{C}^2$.

$\mathbf{f}$ is a strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f_0 is an m-morphism of $\mathcal{E}^1$ to $\mathcal{E}^2$, f_1 is an m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ and for every $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in \mathbf{A}^i,$

$i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset, \phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of $\mathcal{C}^2$.

It is also a coherent strict $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ and $\mathcal{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$.

By abuse of language we will write $\mathbf{f}$ is coherent when it is a $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2$ $(\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ where the model spaces, model categories and atlases are understood by context.

13.1.3 Abbreviated nomenclature for m-atlas morphisms

{def:M-ATLmorphabbrev}

Definition 13.8 (Abbreviated nomenclature for m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i, \mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ a pair of model functions.

$\mathbf{f}$ is a constrained semi-strict (strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{\text{const}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$, iff it is a constrained $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$ and it is a semi-strict (strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1\text{-}\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in

the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

$\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathcal{E}^1$ - $\mathcal{E}^2$ - $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

strict) $\mathcal{E}^1$ - $\mathcal{E}^2$ - $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{C}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

$\mathbf{E}^1$ - $\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ - (semi-strict, strict)

$\mathbf{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ - $\mathcal{E}^1$ - $\mathcal{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{C}^2, f_0, f_1, \mathcal{E}^1, \mathcal{E}^2)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

strict) $\mathbf{E}^1$ - $\mathbf{E}^1$ - $\mathcal{E}^1$ - $\mathcal{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ - $\mathcal{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate space $\mathbf{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, f_0, f_1, \mathcal{E}^1)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

$\mathbf{E}^1$ - $\mathbf{E}^1$ - $\mathcal{E}^1$ - $\mathcal{E}^1$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^1$.

Let $\mathcal{C}^i, i = 1, 2$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, $\mathbf{E}^i$ be a topological space, $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$, $f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2$ be continuous, $f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2$

be a model function and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$.

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1)$, and a (constrained) (semi-strict, strict) (semi-strict, strict)

$\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(\mathbf{A}^1, \mathbf{E}^1, \mathcal{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathcal{C}^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) (semi-strict, strict)

$\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model spaces $\mathbf{C}^1, \mathbf{C}^2$.

triv triv

$\mathbf{f}$ is a (constrained) (semi-strict, strict) $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas (constrained) morphism

of A^1 to A^2 in the coordinate space C^1 , abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(A^1, E^1, C^1, A^2, E^2, f_0, f_1)$, iff it is a (constrained) (semi-strict, strict) $E^1 - E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^1 .

f is a (constrained) (semi-strict, strict) $E^1 - E^2 - \mathcal{C}^1 - \mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2)$, iff it is a (constrained) (semi-strict, strict) $E^1 - E^2 - \mathcal{C}^1 - \mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate model spaces C^1, C^2 .

f is a (constrained) (semi-strict, strict) $E^1 - E^2 - \mathcal{C}^1$ m-atlas morphism of A^1 to A^2 in the coordinate model category $\mathcal{C}^1$, abbreviated as

$\text{isAtl}_{\text{Ar}}^{(\text{const})}(A^1, E^1, C^1, A^2, E^2, f_0, f_1, \mathcal{C}^1)$, iff it is a (constrained) (semi-strict, strict) $E^1 - E^2 - \mathcal{C}^1 - \mathcal{C}^1$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^1$.

Similar definitions apply with restrictions on the admissible atlases, i.e., f is a (constrained) (semi-strict, strict) full (maximal, full maximal) $\mathcal{C}^1 - \mathcal{C}^2 - E^1 - E^2 - \mathcal{C}^1 - \mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1 , abbreviated as

$$\begin{aligned} & \text{isAtl}_{\text{Ar}}^{(\text{const})}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2, \mathcal{C}^1, \mathcal{C}^2) \quad \Longleftrightarrow \\ & \quad \text{full (max,max-full)} \\ & \quad \text{(semi-strict,strict)} \\ & \text{isAtl}_{\text{Ar}}^{(\text{const})}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{C}^1, \mathcal{C}^2, \mathcal{C}^1, \mathcal{C}^2) \quad \wedge \\ & \quad \text{semi-strict (strict)} \\ & \text{isAtl}_{\text{Ob}}(A^1, E^1, C^1) \quad \wedge \\ & \quad \text{full (max,max-full)} \\ & \text{isAtl}_{\text{Ob}}(A^2, E^2, C^2) \\ & \quad \text{full (max,max-full)} \end{aligned}$$

13.1.4 Proclamations on m-atlas morphisms

{lem:M-ATLmorph}

Lemma 13.9 (M-atlas morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}^i, \mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathcal{C}^i \xrightarrow{\text{full-cat}} \mathcal{C}^i, \mathcal{C}^i \xrightarrow{\text{full-cat}} \mathcal{C}^i$
3. $E^i \in \mathcal{C}^i, C^i \in \mathcal{C}^i, C^i \in \mathcal{C}^i, C^i \xrightarrow{\text{mod}} C^i$
4. $f'_1: C'^1 \rightarrow C'^2$ be a model function such that $f'_1[C^1] \subseteq C^2$ and $f_1 \stackrel{\text{def}}{=} f'_1 \upharpoonright_{C^1, C^2}$ is a model function

5. A^i be an M-atlas of E^i in the coordinate space C^i
6. $\hat{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ be a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2$ $(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2
7. $\hat{f}' \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f'_1: C^1 \longrightarrow C'^2)$,
 $\hat{f}'' \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f'_1: C'^1 \longrightarrow C'^2)$ be pairs of model functions

Then

{M-ATLmorph: fineistopmorph}

1. Let $E^i \stackrel{\text{def}}{=} \text{Top}(E^i)$, $i = 1, 2$. If each C^i is fine grained then
 $\hat{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ is a (semi-strict, strict) $E^1\text{-}E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof.

- (a) A^i is an m-atlas of E^i in the coordinate space C^i by lemma 12.3 (M-atlases) on page 57. {lem:M-atlas}
- (b) $f_0: E^1 \longrightarrow E^2$ is a model function, hence continuous, hence by lemma 6.4 (Model functions and trivial model spaces) on page 26, $f_0: \begin{smallmatrix} E^1 \\ \text{triv} \end{smallmatrix} \longrightarrow \begin{smallmatrix} E^2 \\ \text{triv} \end{smallmatrix}$ is a model function. {lem:modfunctriv}
- (c) $f_1: C^1 \longrightarrow C^2$ is a model function by hypothesis.
- (d) Any model neighborhood of E^i is open, hence a model neighborhood of $\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}$.
- (e) Let $(U^i, V^i, \phi^i) \in A^i$, $i = 1, 2$. All of the model neighborhoods of E^i mentioned in definition 13.1 (M-atlas morphisms for model spaces) on page 61 are also model neighborhoods of $\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}$. {def:M-ATLmorph}

□

{M-ATLmorph: istosup}

2. $\hat{f}'$ is an $E^1\text{-}E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C'^2 and $\hat{f}''$ is an $E^1\text{-}E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C'^1, C'^2 .

Proof. $f_0: E^1 \longrightarrow E^2$ is a model function by definition 13.1 (M-atlas morphisms for model spaces) on page 61. {def:M-ATLmorph}

$f'_1: C'^1 \longrightarrow C'^2$ and $f'_1: C^1 \longrightarrow C'^2$ are model functions by hypothesis.

$C^i \stackrel{\text{mod}}{\subseteq} C'^i$ by hypothesis,

C^i is an open subset of C'^i by lemma 2.9 (Model subspaces) on page 19. {lem:modsub}

The expanded model space does not change the atlases or the commutation.

□

- {M-ATLmorph:strictissemi}
3. If f is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 then f is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2
- Proof.* There are four overlapping cases:
- E^i without category:** f_0 is an m-morphism of E^1 to E^2 by definition 13.7 (Strict m-atlas morphisms) on page 65. f_0 is a local m-morphism of E^1 to E^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphstrict}
{lem:modMmorph}
- C^i without category:** f_1 is an m-morphism of C^1 to C^2 by definition 13.7. f_1 is a local m-morphism of C^1 to C^2 by lemma 6.13. {def:M-ATLmorphstrict}
{lem:modMmorph}
- E^i with category:** f_0 is an $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.7. f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by lemma 6.13. {def:M-ATLmorphstrict}
{lem:modMmorph}
- C^i with category:** f_1 is a $\mathcal{E}^1\text{-}\mathcal{C}^2$ m-morphism of C^1 to C^2 by definition 13.7. f_1 is a local $\mathcal{E}^1\text{-}\mathcal{C}^2$ m-morphism of C^1 to C^2 by lemma 6.13. {def:M-ATLmorphstrict}
{lem:modMmorph}
-
- {M-ATLmorph:strictSstrictis}
4. If f is a coherent semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , then f is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .
- Proof.* There are four overlapping cases.
- E^i without category:** f_0 is a local m-morphism of E^1 to E^2 by definition 13.6 {def:M-ATLmorphsemi}
(Semi-strict m-atlas morphisms) on page 64. $E^1 \stackrel{\text{strict-mod}}{\subseteq} E^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\text{Set}(E^2)$ is a model neighborhood of E^2 by definition 2.6 (Model subspaces) on page 18. The result follows by item 6 (Restricted sheaf condition) of definition 2.6. {def:M-ATLmorphsemi}
{def:modsub}
{mod:sheaf}
- E^i with category:** f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ by definition 13.6. $\mathcal{E}^2$ satisfies the restricted sheaf condition by definition 6.7 (Model categories) on page 27 and the result follows. {def:M-ATLmorphsemi}
{def:ModCatMorphsemi}
- C^i without category:** f_0 is a local m-morphism of C^1 to C^2 by definition 13.6 {def:M-ATLmorphsemi}
(Semi-strict m-atlas morphisms) on page 64. $C^1 \stackrel{\text{strict-mod}}{\subseteq} C^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\text{Set}(C^2)$ is a model neighborhood of C^2 by definition 2.6 (Model subspaces) on page 18. The result follows by item 6 (Restricted sheaf condition) of definition 2.6. {def:M-ATLmorphsemi}
{def:modsub}
{mod:sheaf}
- C^i with category:** f_0 is a local $\mathcal{E}^1\text{-}\mathcal{C}^2$ m-morphism of C^1 to C^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{E}^2$ by definition 13.6. $\mathcal{E}^2$ satisfies the restricted sheaf condition by definition 6.7 (Model categories) on page 27 and the result follows. {def:M-ATLmorphsemi}
{def:ModCatMorphsemi}

□

{M-ATLmorph:SstrictisSstri

5. If f is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 then f' is a semi-strict $(\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-})E^1\text{-}E^2(-\mathcal{E}'^1\text{-}\mathcal{E}'^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C'^2 .

Proof. There are four overlapping cases.

- E^i without category:** f_0 is a (strict) local m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. {def:M-ATLmorphsemi}
- E^i with category:** f_0 is a (strict) local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^i \overset{\text{full-cat}}{\subseteq} \mathcal{E}'^i$ by hypothesis. f_0 is a (strict) local $\mathcal{E}'^1\text{-}\mathcal{E}'^2$ m-morphism of E^1 to E^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphsemi}
- C^i without category:** f_1 is a (strict) local m-morphism of C^1 to C^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $C^2 \overset{\text{mod}}{\subseteq} C'^2$ by hypothesis. f_1 is a (strict) local m-morphism of C^1 to C'^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphsemi}
- C^i with category:** f_0 is a local $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of C^1 to C^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^i \overset{\text{full-cat}}{\subseteq} \mathcal{E}'^i$ by hypothesis. f_0 is a (strict) local $\mathcal{E}'^1\text{-}\mathcal{E}'^2$ m-morphism of C^1 to C^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphsemi}

□

{M-ATLmorph:strictisstrict

6. If f is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(-\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 then f' is a strict $(\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-})E^1\text{-}E^2(-\mathcal{E}'^1\text{-}\mathcal{E}'^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C'^2 .

Proof. There are four overlapping cases.

- E^i without category:** f_0 is an m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. {def:M-ATLmorphsemi}
- E^i with category:** f_0 is an $\mathcal{E}^1\text{-}\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^i \overset{\text{full-cat}}{\subseteq} \mathcal{E}'^i$ by hypothesis. f_0 is an $\mathcal{E}'^1\text{-}\mathcal{E}'^2$ m-morphism of E^1 to E^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphsemi}
- C^i without category:** f_1 is an m-morphism of C^1 to C^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $C^2 \overset{\text{mod}}{\subseteq} C'^2$ by hypothesis. f_1 is an m-morphism of C^1 to C'^2 by lemma 6.13 ((Local) m-morphisms) on page 30. {def:M-ATLmorphsemi}

$\mathcal{C}^i$ with category: f_0 is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is a $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by lemma 6.13 ((Local) m-morphisms) on page 30.

□

{M-ATLmorph:SstrictlocalisS

7. If $\mathbf{f}$ is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$ and f'_1 is a local m-morphism of $\mathcal{C}'^1$ to $\mathcal{C}'^2$ (a local $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism) of $\mathcal{C}'^1$ to $\mathcal{C}'^2$) then $\mathbf{f}''$ is a semi-strict $(\mathcal{E}^{i1}\text{-}\mathcal{E}^{i2}\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^{i1}\text{-}\mathcal{E}^{i2})$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}'^1$, $\mathcal{C}'^2$.

Proof. There are four overlapping cases:

E^i without categories: f_0 is a local m-morphism of E^1 to E^2 by definition 13.6 (Semi-strict m-atlas morphisms) on page 64.

{def:M-ATLmorphsemi}

E^i with categories: f_0 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.6. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is a local $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism of E^1 to E^2 by lemma 6.13 ((Local) m-morphisms) on page 30.

{def:M-ATLmorphsemi}

{lem:modMmorph}

$\mathcal{C}^i$ without categories: f'_1 is a local m-morphism of $\mathcal{C}'^1$ to $\mathcal{C}'^2$ by hypothesis.

$\mathcal{C}^i$ with categories: f'_1 is a local $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by hypothesis.

□

{M-ATLmorph:strictmorphisS

8. If $\mathbf{f}$ is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}^1$, $\mathcal{C}^2$ and f'_1 is an m-morphism of $\mathcal{C}'^1$ to $\mathcal{C}'^2$ (a $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism) of $\mathcal{C}'^1$ to $\mathcal{C}'^2$) then $\mathbf{f}''$ is a strict $(\mathcal{E}^{i1}\text{-}\mathcal{E}^{i2}\text{-})E^1\text{-}E^2(\text{-}\mathcal{E}^{i1}\text{-}\mathcal{E}^{i2})$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathcal{C}'^1$, $\mathcal{C}'^2$.

Proof. There are four overlapping cases:

E^i without categories: f_0 is an m-morphism of E^1 to E^2 by definition 13.1 (M-atlas morphisms for model spaces) on page 61.

{def:M-ATLmorph}

E^i with categories: f_0 is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.1. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^i$ by hypothesis. f_0 is an $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism of E^1 to E^2 by lemma 6.13 ((Local) m-morphisms) on page 30.

{def:M-ATLmorph}

{lem:modMmorph}

$\mathcal{C}^i$ without categories: f'_1 is an m-morphism of $\mathcal{C}'^1$ to $\mathcal{C}'^2$ by hypothesis.

$\mathcal{C}^i$ with categories: f'_1 is a $\mathcal{E}^{i1}$ - $\mathcal{E}^{i2}$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ by hypothesis.

□

Corollary 13.10 (M-atlas morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}'^i, i = 1, 2$, be model categories
2. $\mathcal{C}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}'^i$
3. $\mathbf{C}^i \in \mathcal{C}^i$
4. $\mathbf{C}'^i \in \mathcal{C}'^i$
5. $\mathbf{C}^i \stackrel{\text{mod}}{\subseteq} \mathbf{C}'^i$
6. $f'_1: \mathbf{C}'^1 \rightarrow \mathbf{C}'^2$ be a model function such that $f'_1[\mathbf{C}^1] \subseteq \mathbf{C}^2$ and $f_1 \stackrel{\text{def}}{=} f'_1 \upharpoonright_{\mathbf{C}^1, \mathbf{C}^2}$ is a model function
7. E^i be a topological space
8. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$
9. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \rightarrow E^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ be a (semi-strict, strict) $(\mathcal{C}^1\text{-}\mathcal{C}^2\text{-})E^1\text{-}E^2$ $(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$

If $\mathbf{f}$ is a semi-strict (strict) $E^1\text{-}E^2\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ then $\mathbf{f}$ is a semi-strict (strict) $E^1\text{-}E^2\text{-}\mathcal{C}'^1\text{-}\mathcal{C}'^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. The result follows from definition 13.8 (Abbreviated nomenclature for m-atlas morphisms) on page 66. $\square$

{def:M-ATLmorphabbrev}

{lem:M-ATLmorphequiv}

Lemma 13.11 (Equivalence of m-atlas morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}'^i, i = 1, 2$, be model categories
2. $\mathbf{E}^i \in \mathcal{C}^i, \mathbf{C}^i \in \mathcal{C}^i$,
3. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$,
4. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_0: \mathbf{E}^1 \rightarrow \mathbf{E}^2, g_1: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ be equivalent m-atlas morphisms of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

If $\mathbf{f}$ is a (coherent) semi-strict $(\mathcal{C}^1\text{-}\mathcal{C}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, then $\mathbf{g}$ is a (coherent) semi-strict $(\mathcal{C}^1\text{-}\mathcal{C}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.




Figure 8: Uncompleted equivalent m-atlas morphisms

{fig:AtlE1}

Proof. Let $v^1 \in C^1$ be an arbitrary point. Let $(U^1, V^1, \phi^1) \in A^1$ be an arbitrary chart such that $v^1 \in V^1$, $u^1 \stackrel{\text{def}}{=} \phi^{1-1}(v^1)$, $(U^2, V^2, \phi^2) \in A^2$ be an arbitrary chart at $u^2 \stackrel{\text{def}}{=} f_0(u^1)$, $v^2 \stackrel{\text{def}}{=} \phi^2(u^2)$ and $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2]$, as shown in fig. 8 (Uncompleted equivalent m-atlas morphisms).

{fig:AtlE1}

Let $(fU'^1, fV'^1, \phi^1: fU'^1 \xrightarrow{\cong} fV'^1) \in A^1$ be a subchart of (U^1, V^1, ϕ^1) at u^1 , $fU'^2 \subseteq U^2$ a model neighborhood of E^2 for u^2 and

$(fU'^2, f\hat{V}'^2, f\phi'^2: fU'^2 \xrightarrow{\cong} f\hat{V}'^2) \in A^2$ be a chart at u^2 as in definition 13.1 (M-atlas morphisms for model spaces) on page 61, i.e.,

{def:M-ATLmorph}

$$f_0[U'^1] \subseteq fU'^2 \quad (13.2) \quad \{\text{eq:AtlM1}\}$$

$$f\phi'^2 \circ f_0: fU'^1 \longrightarrow f\hat{V}'^2 = f_1 \circ \phi^1: fU'^1 \longrightarrow f\hat{V}'^2 \quad (13.3)$$

$f\hat{V}'^2$ is a model neighborhood of C^2 by definition 11.1 (M-charts) on page 50. $f_1: fV'^1 \longrightarrow f\hat{V}'^2 = f\phi'^2 \circ f_0 \circ \phi^{1-1}$

{def:M-chart}

Let $(gU'^1, gV'^1, g\phi^1: gU'^1 \xrightarrow{\cong} gV'^1) \in A^1$ be a subchart of (U^1, V^1, ϕ^1) at u^1 , $gU'^2 \subseteq U^2$ a model neighborhood of E^2 for u^2 and

$(gU'^2, g\hat{V}'^2, g\phi'^2: gU'^2 \xrightarrow{\cong} g\hat{V}'^2) \in A^2$ be a chart at u^2 as in definition 13.1 (M-atlas morphisms for model spaces) on page 61, i.e.,

{def:M-ATLmorph}

$$g_0[U'^1] \subseteq {}^g U'^2 \quad (13.4)$$

$$\begin{aligned} {}^g \phi'^2 \circ f_0: {}^g U'^1 &\longrightarrow {}^g \hat{V}'^2 = {}^g \phi'^2 \circ g_0: {}^g U'^1 \longrightarrow {}^g \hat{V}'^2 \\ &= g_1 \circ \phi^1: {}^g U'^1 \longrightarrow {}^g \hat{V}'^2 \end{aligned} \quad (13.5)$$

${}^g \hat{V}'^2$ is a model neighborhood of $\mathbf{C}^2$ by definition 11.1.

{def:M-chart}

$U'^1 \stackrel{\text{def}}{=} f U'^1 \cap {}^g U'^1$ is a model neighborhood of $\mathbf{E}^1$, $V'^1 \stackrel{\text{def}}{=} f U'^1 \cap {}^g U'^1$ is a model neighborhood of $\mathbf{C}^1$, $U'^2 \stackrel{\text{def}}{=} f U'^2 \cap {}^g U'^2$ is a model neighborhood of $\mathbf{E}^2$ and $V'^2 \stackrel{\text{def}}{=} f \hat{V}'^2 \cap {}^g \hat{V}'^2$ is a model neighborhood of $\mathbf{C}^2$ by ^{mod:closed} item 2 of definition 2.1 (Model spaces) on page 17. $f_1[V'^1] \subseteq V'^2$ and $g_1[V'^1] \subseteq V'^2$.

{def:model}

$f_0 \upharpoonright U'^1, U'^2 = g_0 \upharpoonright U'^1, U'^2$ is a morphism of $\mathbf{E}^2$ by lemma 6.13 ((Local) m-morphisms) on page 30. $f_1 \upharpoonright V'^1, V'^2 =$ is a morphism of $\mathbf{C}^2$ by lemma 6.13.

{lem:modMmorph}

{lem:modMmorph}

${}^g \phi'^2 \circ f \phi'^2: V'^2 \longrightarrow V'^2$ is an isomorphism of $\mathbf{C}^2$ by definition 11.7 (M-compatibility).

{def:M-compat}

$$\begin{aligned} g_1 &= {}^g \phi'^2 \circ f_0 \circ \phi^{1-1} \\ &= {}^g \phi'^2 \circ f_0 \circ \phi^{1-1} \\ &= {}^g \phi'^2 \circ f \phi'^{2-1} \circ f \phi'^2 \circ f_0 \circ \phi^{1-1} \\ &= {}^g \phi'^2 \circ f \phi'^{2-1} \circ f_1 \end{aligned} \quad (13.6)$$

There are four overlapping cases.

E^i without category: $f_0 = g_0$ by definition 13.5 (Equivalence of m-atlas morphisms) on page 64. f_0 is a local m-morphism of $\mathbf{E}^1$ to $\mathbf{E}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64.

{def:M-ATLmorphequiv}

{def:M-ATLmorphsemi}

If f is coherent then $\mathbf{E}^1 \stackrel{\text{strict-mod}}{\subseteq} \mathbf{E}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64.

{def:M-ATLmorphsemi}

C^i without category: f_1 is a local m-morphism of $\mathbf{C}^1$ to $\mathbf{C}^2$.

$\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a local m-morphism of $\phi^1[I]$ to $\mathbf{C}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64.

{def:M-ATLmorphsemi}

Let $\hat{V}'^2 \subseteq f \hat{V}'^2 \cap {}^g \hat{V}'^2$ be a model neighborhood of $\mathbf{C}^2$ at v^2 such that

$f_1: V'^1 \stackrel{\text{def}}{=} f_1^{-1}[\hat{V}'^2] \longrightarrow \hat{V}'^2$ is a morphism of $\mathbf{C}^2$.

The following, shown in ^{fig:AtLE2} figs. 9 to 10, are model neighborhoods of $\mathbf{C}^2$ by definition 6.1 (Model functions) on page 25:

{def:modfunc}

$$\begin{aligned} V'^2 &\stackrel{\text{def}}{=} \hat{f}^{-1}[\hat{V}'^2] \\ U'^2 &\stackrel{\text{def}}{=} \phi^{2-1}[V'^2] \\ U'^1 &\stackrel{\text{def}}{=} f_0^{-1}[U'^2] \end{aligned}$$




Figure 9: Completed equivalent m-atlas morphisms

{fig:At1E2}

Then

$$\begin{aligned}
 g_1: V'^1 \longrightarrow \hat{V}'^2 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad \hat{f}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_1: V'^1 \longrightarrow \hat{V}'^2
 \end{aligned}$$




Figure 10: Completed equivalent m-atlas morphisms

{fig:AtlE3}

is a composition of morphisms of $\mathcal{C}^2$, hence a morphism.

If $\mathbf{f}$ is coherent then $\mathcal{C}^1 \xrightarrow{\text{strict-mod}} \mathcal{C}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. {def:M-ATLmorphsemi}

E^i with category: $f_0 = g_0$ by definition 13.5. f_0 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 by definition 13.6. {def:M-ATLmorphequiv}
{def:M-ATLmorphsemi}

If $\mathbf{f}$ is coherent then $\mathcal{E}^1 \xrightarrow{\text{strict-cat}} \mathcal{E}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. {def:M-ATLmorphsemi}

$\mathcal{C}^i$ with category: f_1 is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^2$ and $\phi^2 \circ f_0 \circ \phi^{1-1}: \phi^1[I] \rightarrow V^2$ is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of $\text{Mod}(\phi^1[I], \mathcal{C}^2)$ to $\text{Mod}(V^2, \mathcal{C}^2)$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64. {def:M-ATLmorphsemi}

Let $U'^2 \subseteq {}^f U'^2 \cap {}^g U'^2$ be a model neighborhood of $\mathcal{C}^2$ at u^2 such that

$f_0: U'^1 \stackrel{\text{def}}{=} f_0^{-1}[U'^2] \rightarrow U'^2$ is a morphism of $\mathcal{E}^2$.

Let $V'^2 \stackrel{\text{def}}{=} \phi^2[U'^2]$, $f\hat{V}'^2 \stackrel{\text{def}}{=} \hat{f}[fV'^2]$, $g\hat{V}'^2 \stackrel{\text{def}}{=} \hat{g}[gV'^2]$, $fV'^1 \stackrel{\text{def}}{=} f_1^{-1}[f\hat{V}'^2]$ and $gV'^1 \stackrel{\text{def}}{=} g_1^{-1}[g\hat{V}'^2]$. All of these are model neighborhoods by definition 6.1 (Model functions) on page 25 and the corresponding relative model subspaces of V^i are objects of $\mathcal{V}^i$ by item 5 of definition 6.7 (Model categories) on page 27. The restrictions of morphisms to them are morphisms by item 5 of definition 6.7.

Then

$$\begin{aligned}
 g_1: V'^1 \longrightarrow \hat{V}'^2 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad \hat{f}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \phi^2: U'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_0: U'^1 \longrightarrow U'^2 \quad \circ \\
 &\quad \phi^{1-1}: V'^1 \longrightarrow U'^1 \quad \circ \\
 &= \hat{g}: V'^2 \longrightarrow \hat{V}'^2 \quad \circ \\
 &\quad \hat{f}^{-1}: \hat{V}'^2 \longrightarrow V'^2 \quad \circ \\
 &\quad f_1: V'^1 \longrightarrow \hat{V}'^2 \quad \circ
 \end{aligned}$$

is a composition of morphisms of $\mathcal{C}^2$, hence a morphism.

If $\mathbf{f}$ is coherent then $\mathcal{C}^1 \stackrel{\text{strict-cat}}{\subseteq} \mathcal{C}^2$ by definition 13.6 (Semi-strict m-atlas morphisms) on page 64.

□ {cor:M-ATLmorphequiv}

Corollary 13.12 (Equivalence of m-atlas morphisms). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathbf{E}^i \in \mathcal{C}^i$
3. $\mathbf{C}^i \in \mathcal{C}^i$
4. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$
5. $\mathbf{A}^1$ be full
6. $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be a coherent semi-strict $(\mathcal{C}^1\text{-}\mathcal{C}^2)\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2$ $(\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.
7. $\mathbf{g} \stackrel{\text{def}}{=} (g_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, g_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be an m-atlas morphisms of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ equivalent to $\mathbf{f}$

Then $\mathbf{g}$ is a strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Proof. $\mathbf{g}$ is a coherent semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})\mathbf{E}^1\text{-}\mathbf{E}^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ by lemma 13.11 (Equivalence of m-atlas morphisms) on page 73. The result follows from item 4 of lemma 13.9 (M-atlas morphisms) on page 68. $\square$

{lem:M-ATLmorphequiv}
{M-ATLmorph:strictSstrictis
{lem:M-ATLmorphcomp}

Lemma 13.13 (Composition of m-atlas morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2, 3$, be a model category, $\mathbf{E}^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{C}^i$,
2. $\mathbf{A}^i$ an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$
3. $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_0^i: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}, g_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ $i = 1, 2$, be equivalent (semi-strict, strict) $(\mathcal{E}^i\text{-}\mathcal{E}^{i+1}\text{-})\mathbf{E}^i\text{-}\mathbf{E}^{i+1}(\text{-}\mathcal{C}^i\text{-}\mathcal{C}^{i+1})$ m-atlas morphisms of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$.

Then:

{M-morphcomp:morphcompmorph}

1. $\mathbf{f}^2 \circ \mathbf{f}^1$ is an $\mathbf{E}^1\text{-}\mathbf{E}^3$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. $f_0^2 \circ f_0^1$ and $f_1^2 \circ f_1^1$ are model functions by lemma 6.5 (Composition of model functions) on page 27.

{lem:modcomp}

Let $(U^i, V^i, \phi^i) \in \mathbf{A}^i, i = 1, 3$, be a chart, $I \stackrel{\text{def}}{=} U^1 \cap (f_0^2 \circ f_0^1)^{-1}[U^3] \neq \emptyset$.

Let $u^1 \in I, u^2 \stackrel{\text{def}}{=} f_0^1(u^1), u^3 \stackrel{\text{def}}{=} f_0^2(u^2)$ and $(U^2, V^2, \phi^2) \in \mathbf{A}^2$ be a chart at u^2 .

Since $\mathbf{f}^1$ is a morphism by hypothesis, there exists a subchart

$(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , a model neighborhood $U'^2 \subseteq U^2$ of

u^2 and a chart $(U'^2, \hat{V}'^2, \phi'^2: U'^2 \xrightarrow{\sim} \hat{V}'^2) \in \mathbf{A}^2$ at u^2 such that $f_0^1[U'^1] \subseteq U'^2$

and diagram (13.1) in definition 13.1 (M-atlas morphisms for model spaces) on page 61 is commutative, as shown in fig. 7 (Completed m-atlas morphism) on page 62.

{def:M-ATLmorph}
{fig:AtlM1}

Since $\mathbf{f}^2$ is a morphism by hypothesis, there exists a subchart

$(U''^2, V''^2, \phi^2: U''^2 \xrightarrow{\sim} V''^2) \in \mathbf{A}^2$ at u^2 , a model neighborhood $U'^3 \subseteq U^3$

of u^3 and a chart $(U'^3, \hat{V}'^3, \phi'^3: U'^3 \xrightarrow{\sim} \hat{V}'^3) \in \mathbf{A}^3$ at u^3 such that $f_0^2[U'^2] \subseteq$




Figure 11: Uncompleted composition of m-atlas morphisms

{fig:morphcompM1}

U'^3 and diagram (13.7) below is commutative, as shown in figs. 11 and 12.

{fig:morphcompM1,fig:morphcompM2}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^2\} \rightarrow U''^2, \phi^2: U''^2 \xrightarrow{\cong} V'^2, f_1: V''^2 \rightarrow \hat{V}'^3, \\ f_0: U'^2 \rightarrow U'^3, \phi'^3: U'^3 \xrightarrow{\cong} \hat{V}'^3 \end{array} \right. \quad (13.7) \quad \{\text{eq:AtlM2}\}$$

Define $U'''^1 \stackrel{\text{def}}{=} f_0^{1-1}[U''^2]$, $V'''^1 \stackrel{\text{def}}{=} f_1^{1-1}[V''^2] = \phi^1[U'''^1]$, Figure 13 is commutative. $\square$

{M-morphcomp: semicompsemi}

2. If each f^i , $i = 1, 2$, is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2)\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism then $\mathbf{f}^2 \circ \mathbf{f}^1$ is a semi-strict $(\mathcal{E}^1\text{-}\mathcal{E}^2)\text{-}\mathbf{E}^1\text{-}\mathbf{E}^2(\text{-}\mathcal{E}^1\text{-}\mathcal{E}^2)$ m-atlas morphism.

Proof. There are four overlapping cases.




Figure 12: Partially completed composition of m-atlas morphisms

{fig:morphcompM2}

E^i without category: If each $E^i \subseteq E^{i+1}$ and each f_0^i is a local m-morphism of E^i to E^{i+1} , then by lemma 6.17 (Composition of m-morphisms) on page 38 $f_0^2 \circ f_0^1$ is a local m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

C^i without category: If each $C^i \subseteq C^{i+1}$ and each f_1^i is a local m-morphism of C^i to C^{i+1} , then by lemma 6.17 $f_1^2 \circ f_1^1$ is a local morphism of C^1 to C^3 . Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 3$, with $I^1 \stackrel{\text{def}}{=} U^1 \cap (f_0^2 \circ f_0^1)^{-1}[U^3] \neq \emptyset$. For every $u^1 \in I^1$ let $u^2 \stackrel{\text{def}}{=} f_0^1(u^1)$, $u^3 \stackrel{\text{def}}{=} f_0^2(u^2)$ and $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in A^2$ be a chart at u^2 , $I^2 \stackrel{\text{def}}{=} U^1 \cap f_0^{1-1}[U^2] \neq \emptyset$ and $I^2 \stackrel{\text{def}}{=} U^2 \cap f_0^{2-1}[U^3] \neq \emptyset$. If $\phi^2 \circ f_0^1 \circ \phi^{1-1}: \phi^1[I^1] \rightarrow V^2$ is a local m-morphism of $\text{Mod}(\phi^1[I^1], C^1)$ to $\text{Mod}(V^2, C^2)$ and $\phi^3 \circ f_0^2 \circ \phi^{2-1}: \phi^2[I^2] \rightarrow V^3$ is a local m-morphism of $\text{Mod}(\phi^1[I^2], C^1)$ to $\text{Mod}(V^2, C^2)$ then by definition 6.11 there exist model neighborhoods $V'^i \subseteq V^i, i \in [1, 3]$, such that $f^i[V'^i] \subseteq V'^{i+1}$ and {def:modlocalMmorph}




Figure 13: Completed composition of M-atlas morphisms

{fig:morphcompM3}

$\phi^{i+1} \circ f_0^i \circ \phi^{i-1}: V'^i \longrightarrow V'^{i+1}$ is an m-morphism of $\mathcal{C}^{i+1}$, $i = 1, 2$. Then

$$\begin{aligned} \phi^3 \circ f_0^2 \circ f_0^1 \circ \phi^{1-1}: V'^1 \longrightarrow V'^3 &= \phi^3 \circ f_0^2 \circ \phi^{2-1}: V'^2 \longrightarrow V'^3 \circ \\ &\phi^2 \circ f_0^1 \circ \phi^{1-1}: V'^1 \longrightarrow V'^2 \end{aligned}$$

is an m-morphism of $\mathcal{C}^3$. Since $u^1 \in I$ is arbitray, $\phi^3 \circ f_0^2 \circ f_0^1 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3$ is a local m-morphism of $\text{Mod}(\phi^1[I], \mathcal{C}^3)$ to $\text{Mod}(V^3, \mathcal{C}^3)$.

E^i with category: If each $\mathcal{E}^i \overset{\text{full-cat}}{\subseteq} \mathcal{E}^{i+1}$ and each f_0^i is a local $\mathcal{E}^i$ - $\mathcal{E}^{i+1}$ m-morphism of E^i to E^{i+1} , then by lemma 6.17 (Composition of m-morphisms) on page 38 $f_0^2 \circ f_0^1$ is a local $\mathcal{E}^1$ - $\mathcal{E}^3$ m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

$\mathcal{C}^i$ with category: If each $\mathcal{C}^i \overset{\text{full-cat}}{\subseteq} \mathcal{C}^{i+1}$ and each f_1^i is a local $\mathcal{C}^i$ - $\mathcal{C}^{i+1}$ m-morphism of $\mathcal{C}^i$ to $\mathcal{C}^{i+1}$, then by lemma 6.17 (Composition of m-morphisms) on page 38 $f_1^2 \circ f_1^1$ is a local $\mathcal{C}^1$ - $\mathcal{C}^3$ m-morphism of $\mathcal{C}^1$ to $\mathcal{C}^3$. {lem:modMmorphcomp}

□

{M-morphcomp:strictcompstr}

3. If each $f^i, i = 1, 2$, is strict then $f^2 \circ f^1$ is strict.

Proof. There are four overlapping cases.

E^i without category: If each $E^i \subseteq^{mod} E^{i+1}$ and each f_0^i is an m-morphism of E^i to E^{i+1} , then by lemma 6.17 (Composition of m-morphisms) on page 38 $f_0^2 \circ f_0^1$ is an m-morphism of E^1 to E^3 . {lem:modMmorphcomp}

C^i without category: If each $C^i \subseteq^{mod} C^{i+1}$ and each f_1^i is an m-morphism of C^i to C^{i+1} , then by lemma 6.17 (Composition of m-morphisms) on page 38 $f_1^2 \circ f_1^1$ is an m-morphism of C^1 to C^3 . {lem:modMmorphcomp}

E^i with category: If each $\mathcal{E}^i \subseteq^{full-cat} \mathcal{E}^{i+1}$ and each f_0^i is a morphism of $\mathcal{E}^{i+1}$, then $f_0^2 \circ f_0^1$ is a morphism of $\mathcal{E}^3$.

C^i with category: If each $\mathcal{C}^i \subseteq^{full-cat} \mathcal{C}^{i+1}$ and each f_1^i is a morphism of $\mathcal{C}^i$, then $f_1^2 \circ f_1^1$ is a morphism of $\mathcal{C}^3$.

□

{M-morphcomp:constcompconst}

4. If each f^i is constrained then $f^2 \circ f^1$ is constrained.

Proof. This is a corollary of lemma 6.5 (Composition of model functions) on page 27. {lem:modcomp}

□

{M-morphcomp:compequivcomp}

5. $f^2 \circ f^1$ is equivalent to $g^2 \circ g^1$.

Proof. $f_0^1 = g_0^1$ and $f_0^2 = g_0^2$, hence $f_0^2 \circ f_0^1 = g_0^2 \circ g_0^1$. □

{cor:M-ATLmorphcomp}

Corollary 13.14 (Composition of m-atlas morphisms). Let $\mathcal{E}^1$ be a model category, $E^1 \in^{Ob} \mathcal{E}^1$, $\mathcal{E}^i, i = 1, 2, 3$, be a model category, $C^i \in^{Ob} \mathcal{E}^i$, A^i be an m-atlas of E^1 in the coordinate space C^i and $f^i \stackrel{def}{=} (f_0^i: E^1 \rightarrow E^1, f_1^i: C^i \rightarrow C^{i+1}), i = 1, 2$, be a (semi-strict, strict) $(\mathcal{E}^1-)E^1(-\mathcal{E}^i-\mathcal{E}^{i+1})$ m-atlas morphism of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} .

Then

1. $f^2 \circ f^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $(\mathcal{E}^1-)E^1(-\mathcal{E}^1-\mathcal{E}^3)$ m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof. Since each $\mathbf{f}^i$ is a (semi-strict, strict) $(\mathcal{E}^1 - \mathcal{E}^1 -)E^1 - E^1(-\mathcal{E}^i - \mathcal{E}^{i+1})$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$ by definition 13.8

{def:M-ATLmorphabbrev}

(Abbreviated nomenclature for m-atlas morphisms) on page 66, $\mathbf{f}^2 \circ \mathbf{f}^1$ is a (semi-strict, strict) $(\mathcal{E}^1 - \mathcal{E}^1 -)E^1 - E^1(-\mathcal{E}^1 - \mathcal{E}^3)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$ by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79. Apply definition 13.8. $\square$

{M-morphcomp:morphcompmorph}
{def:M-ATLmorphabbrev}

Let $\mathcal{E}^i, i = 1, 2, 3$, be a model category, $E^i \in \mathcal{E}^i$, $\mathcal{E}^1$ be a model category, $\mathbf{C}^1 \in \mathcal{E}^1$, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^1$ and

$\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^1 \longrightarrow \mathbf{C}^1), i = 1, 2$, be a (semi-strict, strict) $(\mathcal{E}^i - \mathcal{E}^{i+1} -)E^i - E^{i+1}(-\mathcal{E}^1)$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate space $\mathbf{C}^1$.

2. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $(\mathcal{E}^1 - \mathcal{E}^3 -)E^1 - E^3(-\mathcal{E}^1)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\mathbf{C}^1$.

Proof. Apply definition 13.8 as above. $\square$

{def:M-ATLmorphabbrev}

Let $\mathcal{E}^1, \mathcal{E}^1$ be model categories, $E^1 \in \mathcal{E}^1, \mathbf{C}^1 \in \mathcal{E}^1$, $\mathbf{A}^i, i = 1, 2, 3$, be an m-atlas of E^1 in the coordinate space $\mathbf{C}^1$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^1 \longrightarrow E^1, f_1^i: \mathbf{C}^1 \longrightarrow \mathbf{C}^1), i = 1, 2$, be a (semi-strict, strict) E^1 m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate space $\mathbf{C}^1$.

3. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate space $\mathbf{C}^1$.

Proof. Apply definition 13.8 as above. $\square$

{def:M-ATLmorphabbrev}

Let $\mathcal{E}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{E}^i, E^i$ be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), i = 1, 2$, be a (semi-strict, strict) $E^i - E^{i+1}(-\mathcal{E}^i - \mathcal{E}^{i+1})$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

4. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) $E^1 - E^3 E^1 - E^3(-\mathcal{E}^1 - \mathcal{E}^3)$ m-atlas morphisms of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 13.8 as above. $\square$

{def:M-ATLmorphabbrev}

Let $\mathcal{E}^i, \mathcal{E}^i, i \in [1, 3]$, be model categories, $E^i \in \mathcal{E}^i, \mathbf{C}^i \in \mathcal{E}^i$ $\mathbf{A}^i$ an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ a (semi-strict, strict) $E^i - E^{i+1}$ m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

5. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 - E^3 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 13.8 as above. □ {def:M-ATLmorphabbrev}

Let $\mathcal{C}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ be a (semi-strict, strict) E^i - E^{i+1} m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

6. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 - E^3 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 13.8 as above. □ {def:M-ATLmorphabbrev}

Let $\mathcal{C}^i, i = 1, 2, 3$, be a model category, $\mathbf{C}^i \in \mathcal{C}^i$, E^i be a topological space, $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$, $i = 1, 2$, be a (semi-strict, strict) E^i - E^{i+1} m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then

7. $\mathbf{f}^2 \circ \mathbf{f}^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a (semi-strict, strict) E^1 - E^3 m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. Apply definition 13.8 as above. □ {def:M-ATLmorphabbrev}

{lem:M-ATLid}

Lemma 13.15 (M-atlas identity). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$
3. $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$
4. $E^i \in \mathcal{C}^i$
5. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
6. $\mathbf{C}^i \in \mathcal{C}^i$
7. $\mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2$
8. $\mathbf{A}^i$ be a maximal m-atlas of E^i in the coordinate space $\mathbf{C}^i$

9. $A^1 \subseteq A^2$

The inclusion morphism $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a semi-strict $(\mathcal{C}^1 - \mathcal{C}^2 -)E^1 - E^2 - (\mathcal{C}^1 - \mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. $\mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is an m-atlas morphism: Let $(U^i, V^i, \phi^i) \in A^i, i = 1, 2$, be charts

such that $I \stackrel{\text{def}}{=} U^1 \cap U^2 \neq \emptyset$. Since each U^i is a model neighborhood of E^i and

$E^1 \stackrel{\text{mod}}{\subseteq} E^2$ by hypothesis, $U^2 \cap \text{Top}(E^1)$ is a model neighborhood of E^1 by item 4 of definition 2.6 (Model subspaces) on page 18 and {modsub: rest}

$I = U^1 \cap U^2 \cap \text{Top}(E^1)$ is a model neighborhood of $E^1 \stackrel{\text{mod}}{\subseteq} E^2$.

Define

1. $f \stackrel{\text{def}}{=} \mathbf{Id}_{(E^2, C^2)}^{(E^1, C^1)}$
2. $U'^1 \stackrel{\text{def}}{=} I$
3. $V'^1 \stackrel{\text{def}}{=} \phi^1[I]$
4. $U'^2 \stackrel{\text{def}}{=} U'^1 = I,$
5. $V'^2 \stackrel{\text{def}}{=} \phi^2[U'^2]$
6. $\hat{V}'^2 \stackrel{\text{def}}{=} V'^1$
7. $\phi'^2 \stackrel{\text{def}}{=} \phi^1: U'^2 = U'^1 \xrightarrow{\cong} \hat{V}'^2 = V'^1.$

Then

1. $U'^1 = I$ is a model neighborhood of E^1 .
2. Since ϕ^1 is a model isomorphism, V'^1 is a model neighborhood of C^1 .
3. $\phi^1 \upharpoonright_{U'^1, V'^1}$ is a model isomorphism of C^1 .
4. Since A^i is maximal, $(U'^1, V'^1, \phi^1 \upharpoonright_{U'^1, V'^1})$ is an m-chart of E^1 in the coordinate space C^1 .
5. Since A^1 is maximal, $(U'^1, V'^1, \phi^1: U'^1 \xrightarrow{\cong} V'^1) \in A^1$.
6. Since $E^1 \stackrel{\text{mod}}{\subseteq} E^2$, $U'^2 = U'^1$ is a model neighborhood of E^2 .
7. Since ϕ^2 is a model isomorphism, V'^2 is a model neighborhood of C^2 .
8. Since $C^1 \stackrel{\text{mod}}{\subseteq} C^2$, $\hat{V}'^2 = V'^1$ is a model neighborhood of C^2 .
9. $\phi^1 \upharpoonright_{U'^2, \hat{V}'^2}$ is a model isomorphism of C^2 .
10. $(U'^2, \hat{V}'^2, \phi^1 \upharpoonright_{U'^2, \hat{V}'^2})$ is an m-chart of E^2 in the coordinate space C^2 .
11. $f_0[U'^1] = \text{Id}_{E^2}^{E^1}[U'^1] = U'^1 \subseteq U'^2$

12. Diagram (13.1) in definition is commutative, as shown in fig. 7 (Completed m-atlas morphism) on page 62. {fig:At1M1}

$\text{Id}_{(E^1, C^1)}^{(E^1, C^1)}$ is semi-strict: Let $u^1 \in E^1$. Since the model neighborhoods of E^1 form an open cover, there is a model neighborhood U of E^1 at u^1 . Id_U is a morphism of E^1 and of $\mathcal{E}^1$; since $E^1 \stackrel{\text{mod}}{\subseteq} E^2$ and $\mathcal{E}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}^2$, Id_U is also a morphism of E^2 and of $\mathcal{E}^2$. Thus $\text{Id}_{E^2}^{E^1}$ is a local m-morphism of E^1 to E^2 and a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism E^1 to E^2 .

Let $v^1 \in C^1$. Since the model neighborhoods of C^1 form an open cover, there is a model neighborhood V of C^1 at v^1 . Id_V is a morphism of C^1 and of $\mathcal{C}^1$; since $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ and $\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$, Id_V is also a morphism of C^2 and of $\mathcal{C}^2$. Thus $\text{Id}_{C^2}^{C^1}$ is a local m-morphism of C^1 to C^2 and a local $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism C^1 to C^2 . □

If each $\text{Top}(E^i) \in E^i$ ($E^i \in \mathcal{E}^i$) and each $\text{Top}(C^i) \in C^i$ ($C^i \in \mathcal{C}^i$) then $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a strict $(\mathcal{E}^1$ - $\mathcal{E}^2$)- E^1 - E^2 -($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. There are four overlapping cases

Each $\text{Top} E^i \in E^i$: $\text{Id}_{E^2}^{E^1}$ is a morphism of E^2 by item 5 of definition 2.1 (Model spaces) on page 17, hence an m-morphism of E^1 to E^2 . {mod:inclusion}

Each $E^i \in \mathcal{E}^2$: $\text{Id}_{E^2}^{E^1}$ is a morphism of $\mathcal{E}^2$ by item 6 of definition 6.7 (Model categories) on page 27, hence an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 . {modcat:morph}

Each $\text{Top} C^i \in C^i$: $\text{Id}_{C^2}^{C^1}$ is a morphism of C^2 by item 5 of definition 2.1, hence an m-morphism of C^1 to C^2 . {mod:inclusion}

Each $C^i \in \mathcal{C}^2$: $\text{Id}_{C^2}^{C^1}$ is a morphism of $\mathcal{C}^2$ by item 6 of definition 6.7, hence a $\mathcal{C}^1$ - $\mathcal{C}^2$ m-morphism of C^1 to C^2 . {modcat:morph}

□

13.2 Categories of m-atlases and functors

This subsection defines categories of m-atlases and functors among them, and proves some basic results. {sub:M-cat}

13.2.1 M-atlas categories

In the following, ... will refer to one of

$$\begin{aligned} &A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1 \\ &A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1 \end{aligned}$$

depending on context.

Definition 13.16 (Sets of m-atlas morphisms). Let E^i and C^i . $i = 1, 2$, be model spaces. Then the set of (constrained) (semi-strict, strict) (full, maximal, full maximal) E^1 - E^2 m-atlas morphisms in the coordinate spaces C^1, C^2 is

{def:M-ATLmorphsets}

$$\begin{aligned} \mathcal{Atl}_{Ar}^{(const)}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \\ \begin{aligned} & \begin{aligned} & \text{(full,max,max-full)} \\ & \text{(semi-strict,strict)} \end{aligned} \\ & \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ & \left. \text{isAtl}^{(const)}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \end{aligned} \end{aligned} \quad (13.8)$$

Let E^i , $i = 1, 2$, be a topological space and $C^1 = (C^i, \mathcal{C}^i)$, $i = 1, 2$, be a model space. Then the set of (constrained) (semi-strict, strict) (full, maximal, full maximal) E^1 - E^2 m-atlas morphisms in the coordinate spaces C^1, C^2 is

$$\begin{aligned} \mathcal{Atl}_{Ar}^{(const)}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \\ \begin{aligned} & \begin{aligned} & \text{(full,max,max-full)} \\ & \text{(semi-strict,strict)} \end{aligned} \\ & \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ & \left. \text{isAtl}^{(const)}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \end{aligned} \end{aligned} \quad (13.9)$$

Let

1. E and C be sets of model spaces
2. $P \stackrel{\text{def}}{=} E \times C$

$$\begin{aligned}
& \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \\
& \quad \begin{array}{l} (\text{full, max, max-full}) \\ (\text{semi-strict, strict}) \end{array} \\
& \quad \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\
& \quad \left. (E^i, C^i) \in P \wedge \text{isAtl}^{(\text{const})}_{(\text{full, max, max-full}, \text{semi-strict, strict})}(\dots) \right\} \quad (13.10)
\end{aligned}$$

Let

1. $\mathcal{E}$ be a model category
2. $\mathbf{C}$ be a set of model spaces
3. $P \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \mathbf{C}$

$$\begin{aligned}
& \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(\mathbf{E}, \mathcal{E}) \stackrel{\text{def}}{=} \\
& \quad \begin{array}{l} (\text{full, max, max-full}) \\ (\text{semi-strict, strict}) \end{array} \\
& \quad \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\
& \quad \left. (E^i, C^i) \in P \wedge \text{isAtl}^{(\text{const})}_{(\text{full, max, max-full}, \text{semi-strict, strict})}(\dots, \mathcal{E}, \mathcal{E}) \right\} \quad (13.11)
\end{aligned}$$

Let

1. $\mathbf{E}$ be a set of model spaces
2. $\mathbf{C}$ be a model category
3. $P \stackrel{\text{def}}{=} \mathbf{E} \times \text{Ob}(\mathcal{E})$

$$\begin{aligned}
& \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(\mathcal{E}, \mathcal{E}) \stackrel{\text{def}}{=} \\
& \quad \begin{array}{l} (\text{full, max, max-full}) \\ (\text{semi-strict, strict}) \end{array} \\
& \quad \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\
& \quad \left. (E^i, C^i) \in P \wedge \text{isAtl}^{(\text{const})}_{(\text{full, max, max-full}, \text{semi-strict, strict})}(\dots, \mathcal{E}, \mathcal{E}) \right\} \quad (13.12)
\end{aligned}$$

Let

1. $\mathcal{E}$ and $\mathcal{C}$ be model categories

2. $P \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \text{Ob}(\mathcal{C})$

$$\begin{aligned} \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(\mathcal{E}, \mathcal{C}) &\stackrel{\text{def}}{=} \\ &\quad \begin{matrix} (\text{full}, \text{max}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{matrix} \\ &\quad \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ &\quad \left. (E^i, C^i) \in P \wedge \text{isAtl}_{\begin{matrix} (\text{full}, \text{max}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{matrix}}^{(\text{const})}(..., \mathcal{E}, \mathcal{E}, \mathcal{C}, \mathcal{C}) \right\} \end{aligned} \quad (13.13)$$

Lemma 13.17 (Sets of m-atlas morphisms). Let E and C be sets of model spaces.

{lem:M-ATLmorphsets}

$$\begin{aligned} \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) &\subseteq \mathcal{A}tl_{\text{Ar}}(E, C) \\ &\quad \begin{matrix} (\text{full}, \text{max}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{matrix} \\ \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) &\subseteq \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) \\ &\quad \begin{matrix} \text{max-full} & \text{full} \\ (\text{semi-strict}, \text{strict}) & (\text{semi-strict}, \text{strict}) \end{matrix} \end{aligned}$$

Proof. The result follows from $\stackrel{\text{def:M-ATLmorphabbrev}}{\text{definition 13.8}}$ (Abbreviated nomenclature for m-atlas morphisms) on page 66 and definition 13.16 above. $\square$

{def:M-ATLmorphsets}
{cor:M-ATLmorphsets}

Corollary 13.18 (Sets of m-atlas morphisms). Let E and C be sets of topological spaces.

$$\begin{aligned} \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) &\subseteq \mathcal{A}tl_{\text{Ar}}(E, C) \\ &\quad \begin{matrix} (\text{full}, \text{max}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{matrix} \\ \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) &\subseteq \mathcal{A}tl_{\text{Ar}}^{(\text{const})}(E, C) \\ &\quad \begin{matrix} \text{max-full} & \text{full} \\ (\text{semi-strict}, \text{strict}) & (\text{semi-strict}, \text{strict}) \end{matrix} \end{aligned}$$

Proof. The result follows from $\stackrel{\text{def:M-ATLmorphabbrev}}{\text{definition 13.8}}$ (Abbreviated nomenclature for m-atlas morphisms) on page 66 and definition 13.16 above. $\square$

{def:M-ATLmorphsets}
{def:M-cat}

Definition 13.19 (Categories $\mathcal{A}tl(E, C)$). Let E and C be sets of model spaces. Let $P \stackrel{\text{def}}{=} E \times C$.

$$\mathcal{Atl}^{(\text{const})}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \begin{matrix} (\text{full}, \text{max}, \text{max-full}) \\ (\text{semi-strict}, \text{strict}) \end{matrix} \left(\mathcal{Atl}_{\text{Ob}}(\mathbf{E}, \mathbf{C}), \mathcal{Atl}_{\text{Ar}}^{(\text{const})}(\mathbf{E}, \mathbf{C}), \circ^A \right) \quad (13.14)$$

Let $\mathbf{E}^i$ and $\mathbf{C}^i$, $i = 1, 2$, be model spaces, $\mathbf{A}^i$ be an atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2)$ be a (strict, semi-strict) $\mathbf{E}^1$ - $\mathbf{E}^2$ M-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The identity functor $\mathcal{F}_{\mathbf{M}, \text{Id}}$ is

$$\mathcal{F}_{\mathbf{M}, \text{Id}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \quad (13.15)$$

This nomenclature will be justified below.

$$\mathcal{F}_{\mathbf{M}, \text{Id}} \left((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right) \stackrel{\text{def}}{=} ((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \quad (13.16)$$

These are fine grained iff $\mathbf{C}$ is fine grained.

Lemma 13.20 ($\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ is a category). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Then

`{lem:M-ATLiscat}`
`{Mcat:morph}`

1. $\mathcal{Atl}^{(\text{const})}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ of $\mathcal{Atl}_{\text{Ob}}^{(\text{const})}(\mathbf{E}, \mathbf{C})$ is $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)}$.

Proof. Let $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ and let

$m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^{i+1}, \mathbf{E}^{i+1}, \mathbf{C}^{i+1}))$ be a morphism of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$. Then

Composition:

$m^2 \circ m^1 = ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^3, \mathbf{E}^3, \mathbf{C}^3))$ is a morphism of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79.

`{M-morphcomp:morphcompmorph}`

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

`{lem:labcomp}`

Identity:

$\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{cor:M-ATLiscat}

Corollary 13.21 (Subcategories of $\mathcal{Att}(E, C)$). Let E and C be sets of model spaces.

Then $\mathcal{Att}_{(full, max, max-full)}^{(const)}(E, C)$ is a subcategory of $\mathcal{Att}_{(semi-strict, strict)}^{(const)}(E, C)$.

Remark 13.22. They are not, in general, full subcategories.

The identity functor $\mathcal{F}_{M, Id}$ is a functor from each of the subcategories to each of the containing categories.

{def:M-cat-cat}

Definition 13.23 (Categories $\mathcal{Att}(\mathcal{E}, \mathcal{C})$). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Let $P \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \text{Ob}(\mathcal{C})$. Then

$$\mathcal{Att}_{\text{Ob}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E \in \mathcal{E} \\ C \in \mathcal{C}}} \mathcal{Att}_{\text{Ob}}^{(full, max, max-full)}(E, C) \quad (13.17)$$

$$\mathcal{Att}_{\text{Ar}}^{(const)}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left\{ \left((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2) \right) \mid \right. \\ \left. (E^i, C^i) \in P \wedge \text{isAtl}_{\text{Ar}}^{(const)}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1, \mathcal{E}, \mathcal{C}) \right\} \quad (13.18)$$

$$\mathcal{Att}^{(const)}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} (\mathcal{Att}_{\text{Ob}}(\mathcal{E}, \mathcal{C}), \mathcal{Att}_{\text{Ar}}^{(const)}(\mathcal{E}, \mathcal{C}), \overset{A}{\circ}) \quad (13.19)$$

{lem:M-cat-ATLiscat}

Lemma 13.24 ($\mathcal{Att}(\mathcal{E}, \mathcal{C})$ is a category). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Then

$\mathcal{Att}_{(full, max, max-full)}^{(const)}(\mathcal{E}, \mathcal{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i)

of $\mathcal{Att}_{(full, max, max-full)}^{(const)}(\mathcal{E}, \mathcal{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{A}tl(\mathcal{E}, \mathcal{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$ be a morphism of $\mathcal{A}tl(\mathcal{E}, \mathcal{C})$. Then

Composition:

$f_0^2 \circ f_0^1$ is a morphism of $\mathcal{E}$, $f_1^2 \circ f_1^1$ is a morphism of $\mathcal{C}$ and $f^2 \circ f^1 = (f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^2 by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79.

{M-morphcomp: morphcompmorph}

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{def:T-cat}

Definition 13.25 (Categories $\mathcal{A}tl(E, C)$ of topological spaces). Let E be a set of topological spaces and C a set of model spaces. Let $P \stackrel{\text{def}}{=} E \times C$. Then

$$\mathcal{A}tl_{\text{Ob}}(E, C) \stackrel{\text{def}}{=} \bigcup_{\substack{E^1 \in E \\ C^1 \in C}} \mathcal{A}tl_{\text{Ob}}(E^1, C^1) \quad (13.20)$$

$$\mathcal{A}tl_{\text{Ar}}(E, C) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^1, C^1) \in P \\ (E^2, C^2) \in P}} \mathcal{A}tl_{\text{Ar}}(E^1, C^1, E^2, C^2) \quad (13.21)$$

$$\mathcal{A}tl(E, C) \stackrel{\text{def}}{=} (\mathcal{A}tl_{\text{Ob}}(E, C), \mathcal{A}tl_{\text{Ar}}(E, C), \circ^A) \quad (13.22)$$

Similar definitions apply with restrictions on the admissible atlases, the admissible morphisms, or both:

1. $\mathcal{A}tl_{\text{full}}$
2. $\mathcal{A}tl_{\text{S-max}}$
3. $\mathcal{A}tl_{\text{max}}$
4. $\mathcal{A}tl_{\text{S-max-full}}$
5. $\mathcal{A}tl_{\text{max-full}}$
6. $\mathcal{A}tl_{\text{full semi-strict}}$

7. $\mathcal{Att}_{\text{S-max semi-strict}}$
8. $\mathcal{Att}_{\text{max semi-strict}}$
9. $\mathcal{Att}_{\text{S-max-full semi-strict}}$
10. $\mathcal{Att}_{\text{max-full semi-strict}}$
11. $\mathcal{Att}_{\text{full strict}}$
12. $\mathcal{Att}_{\text{S-max strict}}$
13. $\mathcal{Att}_{\text{max strict}}$
14. $\mathcal{Att}_{\text{S-max-full strict}}$
15. $\mathcal{Att}_{\text{max-full strict}}$

e.g.,

$$\mathcal{Att}_{\text{S-max-full Ob}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{(E^1, \mathbf{C}^1) \in \mathbf{P}} \mathcal{Att}_{\text{S-max-full Ob}}(E^1, \mathbf{C}^1) \quad (13.23)$$

$$\mathcal{Att}_{\text{S-max-full Ar semi-strict}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^1, \mathbf{C}^1) \in \mathbf{P} \\ (E^2, \mathbf{C}^2) \in \mathbf{P}}} \mathcal{Att}_{\text{S-max-full Ar semi-strict}}(E^1, \mathbf{C}^1, E^2, \mathbf{C}^2) \quad (13.24)$$

$$\mathcal{Att}_{\text{S-max-full semi-strict}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} (\mathcal{Att}_{\text{S-max-full Ob}}(\mathbf{E}, \mathbf{C}), \mathcal{Att}_{\text{S-max-full Ar semi-strict}}(\mathbf{E}, \mathbf{C}), \overset{A}{\circ}) \quad (13.25)$$

Let $(\mathbf{A}^1, E^1, \mathbf{C}^1) \in \mathcal{Att}_{\text{Ob}}(\mathbf{E}, \mathbf{C})$.

$$\text{Id}_{(\mathbf{A}^1, E^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} ((\text{Id}_{E^1}, \text{Id}_{\mathbf{C}^1}), (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^1, E^1, \mathbf{C}^1)) \quad (13.26)$$

Lemma 13.26 ($\mathcal{Att}(\mathbf{E}, \mathbf{C})$ of topological spaces is a category). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces. Then $\mathcal{Att}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object $(\mathbf{A}^1, E^1, \mathbf{C}^1)$ is $\text{Id}_{(\mathbf{A}^1, E^1, \mathbf{C}^1)}$.

{\lem:T-ATLiscat}

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{At}\ell(E, C)$. and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$ be a morphism of $\mathcal{At}\ell(E, C)$. Then

Composition:

$((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3))$ is a morphism of $\mathcal{At}\ell(E, C)$ by corollary 13.14 (Composition of m-atlas morphisms) on page 83. {cor:M-ATLmorphcomp}

Associativity:

Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19. {lem:labcomp}

□

Similar results follow with restrictions on the admissible atlases, the admissible morphisms, or both:

1. $\mathcal{At}\ell_{\text{full}}(\mathcal{E}, \mathcal{C})$
2. $\mathcal{At}\ell_{\text{S-max}}(\mathcal{E}, \mathcal{C})$
3. $\mathcal{At}\ell_{\text{max}}(\mathcal{E}, \mathcal{C})$
4. $\mathcal{At}\ell_{\text{S-max-full}}(\mathcal{E}, \mathcal{C})$
5. $\mathcal{At}\ell_{\text{max-full}}(\mathcal{E}, \mathcal{C})$
6. $\mathcal{At}\ell_{\text{full semi-strict}}(\mathcal{E}, \mathcal{C})$
7. $\mathcal{At}\ell_{\text{S-max semi-strict}}(\mathcal{E}, \mathcal{C})$
8. $\mathcal{At}\ell_{\text{max semi-strict}}(\mathcal{E}, \mathcal{C})$
9. $\mathcal{At}\ell_{\text{S-max-full semi-strict}}(\mathcal{E}, \mathcal{C})$
10. $\mathcal{At}\ell_{\text{max-full semi-strict}}(\mathcal{E}, \mathcal{C})$
11. $\mathcal{At}\ell_{\text{full strict}}(\mathcal{E}, \mathcal{C})$
12. $\mathcal{At}\ell_{\text{S-max strict}}(\mathcal{E}, \mathcal{C})$

13. $\mathcal{Att}_{\max}^{\text{strict}}(\mathcal{E}, \mathcal{C})$
14. $\mathcal{Att}_{\text{S-max-full}}^{\text{strict}}(\mathcal{E}, \mathcal{C})$
15. $\mathcal{Att}_{\max\text{-full}}^{\text{strict}}(\mathcal{E}, \mathcal{C})$

Definition 13.27 ($\mathcal{E}_{\text{triv}}$ and $\mathcal{Att}_{\text{triv}}(E, \mathcal{C})$). Let E be a set of topological spaces and $\mathcal{C}$ a set of model spaces. Then

{def:M-triv}

$$\mathcal{E}_{\text{triv}} \stackrel{\text{def}}{=} \left\{ E^\mu \mid E^\mu \in E \right\} \quad (13.27)$$

$$\mathcal{Att}_{\text{triv}}(E, \mathcal{C}) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{triv}}\left(\mathcal{E}_{\text{triv}}, \mathcal{C}\right) \quad (13.28)$$

13.2.2 M-atlas functors

{def:Func-M-Top}

Definition 13.28 ($\mathcal{F}_{\text{M,Top}}$). Let $E^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i)$, $i = 1, 2$, and $\mathcal{C}^i \stackrel{\text{def}}{=} (\mathcal{C}^i, \mathcal{C}^i)$ be model spaces and A^i be an m-atlas of E^i in the coordinate space $\mathcal{C}^i$. Then

$$\mathcal{F}_{\text{M,Top}}(A^i, E^i, \mathcal{C}^i) \stackrel{\text{def}}{=} (A^i, E^i, \mathcal{C}^i) \quad (13.29)$$

Let $f_0: E^1 \longrightarrow E^2$ and $f_1: \mathcal{C}^1 \longrightarrow \mathcal{C}^2$ be model functions. Then

$$\mathcal{F}_{\text{M,Top}}((f_0: E^1 \longrightarrow E^2, f_1: \mathcal{C}^1 \longrightarrow \mathcal{C}^2), (A^1, E^1, \mathcal{C}^1), (A^2, E^2, \mathcal{C}^2)) \stackrel{\text{def}}{=} ((f_0: E^1 \longrightarrow E^2, f_1: \mathcal{C}^1 \longrightarrow \mathcal{C}^2), (A^1, E^1, \mathcal{C}^1), (A^2, E^2, \mathcal{C}^2)) \quad (13.30)$$

Lemma 13.29 ($\mathcal{F}_{\text{M,Top}}$ maps M-atlases to M-atlases if the coordinate model space is fine grained.). Let $E \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathcal{C}$ be model spaces, $\mathcal{C}$ be fine grained and A be an M-atlas of E in the model space $\mathcal{C}$. Then A is an M-atlas of E in the model space $\mathcal{C}$.

{lem:Func-M-Top}

Proof. Let (U, V, ϕ) be an M-chart in A . Every open subset of V is a model neighborhood of $\mathcal{C}$ since it is fine grained by hypothesis. Then (U, V, ϕ) is an M-chart of E in the model space $\mathcal{C}$ by lemma 11.3 (M-charts) on page 51.

{lem:M-chart}

The definition of mutually compatible charts neighborhood of $\mathcal{C}$. (U, V, ϕ) is also an M-chart of E in the coordinate space $\mathcal{C}$:

Since U is a model neighborhood of E , it is open and hence a model neighborhood of $\mathcal{E}_{\text{triv}}$.

V is a model neighborhood of $\mathcal{C}$ by hypothesis.

Since U' is a model neighborhood of $\mathcal{E}_{\text{triv}}$ and hence open, and ϕ is a homeomorphism, $\phi[U']$ is open. Since V is a model neighborhood of $\mathcal{C}$, $\mathcal{C}$ is fine grained and $\phi[U'] \subseteq V$ is open, $\phi[U']$ is a model neighborhood of $\mathcal{C}$,

Since V' is a model neighborhood of $\mathbf{C}$, it is open, $\phi^{-1}[V']$ is open and hence a model neighborhood of $\mathbf{E}$. $\square$

Theorem 13.30 ($\mathcal{F}_{\mathbf{M}, \text{Top}}$ is a functor if $\mathbf{C}$ is fine grained.). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces and $\mathbf{C}$ be fine grained. Then $\mathcal{F}_{\mathbf{M}, \text{Top}}$ is a functor from $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ to $\mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})$

{the:Func-M-Top}

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i), \mathbf{C}^i), i \in [1, 3]$, be an object of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$, $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{\mathbf{M}, \text{Top}} o^i = (\mathbf{A}^i, E^i, \mathbf{C}^i)$ the corresponding object of $\mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})$, $m^i \stackrel{\text{def}}{=} ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o^i, o^{i+1}), i = 1, 2$, be a morphism of $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ and $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{\mathbf{M}, \text{Top}} m^i = ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o'^i, o'^{i+1})$ the corresponding morphism of $\mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})$.

$\mathcal{F}_{\mathbf{M}, \text{Top}}$ satisfies these criteria:

Preservation of objects:

$\mathbf{A}^i$ is an M-atlas of E^i in the coordinate space $\mathbf{C}^i$ by lemma 13.29 ($\mathcal{F}_{\mathbf{M}, \text{Top}}$ maps M-atlases to M-atlases if the coordinate model space is fine grained) above. {lem:Func-M-Top}

Preservation of morphisms:

$f_0^i: E^i \longrightarrow E^{i+1}$ is continuous and $f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}$ is a model function.

Preservation of endpoints:

$$\mathcal{F}_{\mathbf{M}, \text{Top}} o^i = o'^i \in \mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})^{\text{Ob}}$$

$$\mathcal{F}_{\mathbf{M}, \text{Top}} m^i = ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o'^i, o'^{i+1}) \in \mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})^{\text{Ar}}$$

is a morphism from o'^i to o'^{i+1} .

Identity:

Let $((\text{Id}_{E^i}, \text{Id}_{\mathbf{C}^i}), o^i, o^i)$ be an identity morphism of o^i in $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$. Then $((\text{Id}_{E^i}: \text{Top}(E^i) \longrightarrow \text{Top}(E^i), \text{Id}_{\mathbf{C}^i}), o'^i, o'^i)$ is an identity morphism of o'^i in $\mathcal{Atl}(\text{Top}(\mathbf{E}), \mathbf{C})$.

Composition:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}, \text{Top}}(m^2 \stackrel{A}{\circ} m^1) &= \mathcal{F}_{\mathbf{M}, \text{Top}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o'^1, o'^3) \\ &= ((f_0^2, f_1^2), o'^1, o'^2) \stackrel{A}{\circ} ((f_0^1, f_1^1), o'^2, o'^3) \\ &= \mathcal{F}_{\mathbf{M}, \text{Top}}((f_0^2, f_1^2), o'^2, o'^3) \stackrel{A}{\circ} \mathcal{F}_{\mathbf{M}, \text{Top}}((f_0^1, f_1^1), o'^1, o'^2) \\ &= \mathcal{F}_{\mathbf{M}, \text{Top}} m^2 \stackrel{A}{\circ} \mathcal{F}_{\mathbf{M}, \text{Top}} m^1 \end{aligned}$$

□

{def:Func-Top-M}

Definition 13.31 ($\mathcal{F}_{\text{Top,M}}$). Let E^i , $i = 1, 2$, be a topological space, $\mathbf{C}^i$ a model space and $\mathbf{A}^i$ an m-atlas of E^i and $\mathbf{A}$ an m-atlas of E in the coordinate space $\mathbf{C}^i$. Then

$$\mathcal{F}_{\text{Top,M}}(\mathbf{A}^i, E^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}, \underset{\text{triv}}{E^i}, \mathbf{C}^i) \quad (13.31)$$

Let $f_0: E^1 \longrightarrow E^2$ be continuous and $f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2$ be a model function. Then

$$\begin{aligned} \mathcal{F}_{\text{Top,M}}((f_0: E^1 \longrightarrow E^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2), (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2)) &\stackrel{\text{def}}{=} \\ ((f_0: \underset{\text{triv}}{E^1} \longrightarrow \underset{\text{triv}}{E^2}, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2), (\mathbf{A}^1, \underset{\text{triv}}{E^1}, \mathbf{C}^1), (\mathbf{A}^2, \underset{\text{triv}}{E^2}, \mathbf{C}^2)) &\quad (13.32) \end{aligned}$$

{lem:Func-Top-M}

Lemma 13.32 ($\mathcal{F}_{\text{Top,M}}$ maps M-atlases to M-atlases). Let E be a topological space, $\mathbf{C}$ be a model space and $\mathbf{A}$ be an M-atlas of E in the model space $\mathbf{C}$. Then $\mathbf{A}$ is an M-atlas of $\underset{\text{triv}}{E}$ in the model space $\mathbf{C}$.

Proof. Let (U, V, ϕ) be a M-chart in $\mathbf{A}$, $U' \subseteq U$ a model neighborhood of $\underset{\text{triv}}{E}$ and $V' \subseteq V$ a model neighborhood of $\mathbf{C}$. (U, V, ϕ) is also an M-chart of E in the coordinate space $\mathbf{C}$. □

{the:Func-Top-M}

Theorem 13.33 ($\mathcal{F}_{\text{Top,M}}$ is a functor). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of model spaces, Then $\mathcal{F}_{\text{Top,M}}$ is a functor from $\mathcal{At}(\mathbf{E}, \mathbf{C})$ to $\mathcal{At}(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$.

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, E^i, \mathbf{C}^i)$, $i \in [1, 3]$, be an object of $\mathcal{At}(\mathbf{E}, \mathbf{C})$, $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \underset{\text{triv}}{E^i}, \mathbf{C}^i)$

the corresponding object of $\mathcal{At}(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$, $m^i \stackrel{\text{def}}{=} ((f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}), o^i, o^{i+1})$,

$i = 1, 2$, be a morphism of $\mathcal{At}(\mathbf{E}, \mathbf{C})$ and $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$ the corresponding morphism of $\mathcal{At}(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$.

$\mathcal{F}_{\text{Top,M}}$ satisfies these criteria:

Preservation of endpoints: $\mathbf{A}^i$ is an M-atlas of $\underset{\text{triv}}{E^i}$ in the coordinate space $\mathbf{C}^i$ by

lemma 13.29 ($\mathcal{F}_{\text{M,Top}}$ maps M-atlases to M-atlases if the coordinate model space is fine grained) above. $f_0^i: E^i \longrightarrow E^{i+1}$ is continuous and $f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}$ is a model function.

{lem:Func-M-Top}

$$\mathcal{F}_{\text{Top,M}} o^i = o^i \stackrel{\text{Ob}}{\in} \mathcal{At}(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$$

$$\mathcal{F}_{\text{Top,M}} m^i = m^i \stackrel{\text{Ar}}{\in} \mathcal{At}(\underset{\text{triv}}{\mathbf{E}}, \mathbf{C})$$

is a morphism from o^i to o^{i+1} .

Identity: $\text{Id}_{o^i} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{C^i}), o^i, o^i)$. Then $\mathcal{F}_{\text{Top}, M} \text{Id}_{o^i} = ((\text{Id}_{E^i}, \text{Id}_{C^i}), o^i, o^i)$ is an identity morphism of o^i .

Composition:

$$\begin{aligned}
\mathcal{F}_{\text{Top}, M}(m^2 \overset{A}{\circ} m^1) &= \mathcal{F}_{\text{Top}, M}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\
&= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\
&= ((f_0^2, f_1^2), o^1, o^2) \overset{A}{\circ} ((f_0^1, f_1^1), o^2, o^3) \\
&= \mathcal{F}_{\text{Top}, M}((f_0^2, f_1^2), o^2, o^3) \overset{A}{\circ} \mathcal{F}_{\text{Top}, M}((f_0^1, f_1^1), o^1, o^2) \\
&= \mathcal{F}_{\text{Top}, M} m^2 \overset{A}{\circ} \mathcal{F}_{\text{Top}, M} m^1
\end{aligned}$$

□

$\mathcal{F}_{M, \text{Top}} \circ \mathcal{F}_{\text{Top}, M}$ is the identity functor on $\mathcal{ATL}(E, C)$.

Proof.

$$\begin{aligned}
\mathcal{F}_{M, \text{Top}} \circ \mathcal{F}_{\text{Top}, M} o^i &= \mathcal{F}_{M, \text{Top}} o^i \\
&= o^i
\end{aligned}$$

$$\begin{aligned}
\mathcal{F}_{M, \text{Top}} \circ \mathcal{F}_{\text{Top}, M} m^i &= \mathcal{F}_{M, \text{Top}} m^i \\
&= m^i
\end{aligned}$$

□

14 Classic m-atlas morphisms and functors

{sec:M-ATLcmorph}

This subsection introduces an alternate definition of and taxonomy for morphisms between m-atlases, defines categories of m-atlases, defines functors among them and proves some basic results. It introduces the notion of classic m-atlas morphisms, which is very similar to the conventional definitions for a manifold, although it does not require the space to be locally Euclidean. The reader can safely skip it, but it does help to establish a context for other notions that are used in the exposition of local coordinate spaces.

14.1 Classic m-atlas morphisms

{sub:M-ATLcmorph}

This subsection introduces the notion of classic m-atlas morphisms, and proves some basic results.

14.1.1 Definitions of classic m-atlas morphisms

{def:M-ATLcmorph}

Definition 14.1 (Classic m-atlas morphisms for model spaces). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

$f: E^1 \longrightarrow E^2$ is a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$. and a classic $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f)$, iff $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ ($\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2$) and for any $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset, \phi^2 \circ f \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of C^2 ($\mathcal{C}^2$).

f is also a constrained classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{constrained,classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1)$, and a constrained classic $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{constrained,classic}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f_0, f_1)$, iff f is constrained, i.e., $f[E]$ is contained in a model neighborhood of E^2 .

Let f be a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The triple $((f), (A^1, E^1, C^1), (A^2, E^2, C^2))$ will refer to f considered as a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ and $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{def:M-ATLcmorphTop}

Definition 14.2 (Classic m-atlas morphisms for topological spaces). Let $\mathcal{E}^i, i = 1, 2$, be a model category, $\mathcal{E}^i$ be a topological category, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

$f: E^1 \longrightarrow E^2$ is a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$. and a classic $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate model categories $\mathcal{C}^1, \mathcal{C}^2$, abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, \mathcal{E}^1, \mathcal{C}^1, A^2, \mathcal{E}^2, \mathcal{C}^2, f)$, iff f is a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

The triple $((f), (A^1, E^1, C^1), (A^2, E^2, C^2))$ will refer to f considered as a classic $E^1-E^2(-\mathcal{C}^1-\mathcal{C}^2)$ and $\mathcal{E}^1-\mathcal{E}^2(-\mathcal{C}^1-\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

{def:M-ATLcid}

Definition 14.3 (Classic m-atlas inclusion and identity morphisms). Let

1. $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $E^i \in \mathcal{E}^i$
3. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
4. $C^i \in \mathcal{C}^i$

$$5. \mathbf{C}^1 \stackrel{\text{mod}}{\subseteq} \mathbf{C}^2 (\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2)$$

6. $\mathbf{A}^i$ be an m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$

$$7. \mathbf{A}^1 \subseteq \mathbf{A}^2$$

The classic $\mathbf{E}^1\text{-}\mathbf{E}^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\text{Id}_{(\mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{E}^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} \text{Id}_{\mathbf{E}^2}^{\mathbf{E}^1}$.

The classic m-atlas inclusion morphism of $(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)$ is $\text{Id}_{(\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)}^{(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(\mathbf{E}^2)}^{(\mathbf{E}^1)}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right), \text{Id}_{\mathbf{E}^2}^{\mathbf{E}^1}$ considered as a classic $\mathbf{E}^1\text{-}\mathbf{E}^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The classic $\mathbf{E}^i(-\mathcal{C}^i)$ m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$ is $\text{Id}_{(\mathbf{E}^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} \text{Id}_{\mathbf{E}^i}$.

The classic m-atlas identity morphism of $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ is $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} (\mathbf{Id}_{(\mathbf{E}^i)}, (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)), \text{Id}_{\mathbf{E}^i}$ considered as a classic $\mathbf{E}^i(-\mathcal{C}^i)$ m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

Let

1. E^i be a topological space, $i = 1, 2$,

$$2. E^1 \stackrel{\text{subsp}}{\subseteq} E^2$$

3. $\mathcal{C}^i$ be a model category

$$4. \mathbf{C}^i \in \mathcal{C}^i$$

$$5. \mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2$$

6. $\mathbf{A}^i$ be an m-atlas of E^i in the coordinate space $\mathbf{C}^i$

$$7. \mathbf{A}^1 \subseteq \mathbf{A}^2$$

The classic $E^1\text{-}E^2(-\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas inclusion morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ is $\text{Id}_{\mathbf{E}^2}^{\mathbf{E}^1}$.

The classic m-atlas inclusion morphism of $(\mathbf{A}^1, E^1, \mathbf{C}^1)$ to $(\mathbf{A}^2, E^2, \mathbf{C}^2)$ is $\text{Id}_{(\mathbf{A}^2, E^2, \mathbf{C}^2)}^{(\mathbf{A}^1, E^1, \mathbf{C}^1)} \stackrel{\text{def}}{=} \left(\mathbf{Id}_{(E^2)}^{(E^1)}, (\mathbf{A}^1, E^1, \mathbf{C}^1), (\mathbf{A}^2, E^2, \mathbf{C}^2) \right), \text{Id}_{E^2}^{\mathbf{E}^1}$ considered as a classic $E^1\text{-}E^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The classic E^i m-atlas identity morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$ is $\text{Id}_{(\mathbf{E}^i, \mathbf{C}^i)}$.

The classic m-atlas identity morphism of $(\mathbf{A}^i, E^i, \mathbf{C}^i)$ is $\text{Id}_{(\mathbf{A}^i, E^i, \mathbf{C}^i)} \stackrel{\text{def}}{=} (\mathbf{Id}_{(E^i)}, (\mathbf{A}^i, E^i, \mathbf{C}^i), (\mathbf{A}^i, E^i, \mathbf{C}^i)), \text{Id}_{E^i}$ considered as an E^i m-atlas morphism of $\mathbf{A}^i$ in the coordinate space $\mathbf{C}^i$.

14.1.2 Definitions of semi-strict and strict

{def:M-ATLcmorphsemi}

Definition 14.4 (Semi-strict classic m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a semi-strict classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff f is a local m-morphism of E^1 to E^2 .

Remark 14.5. Definitions of semi-strict classic m-atlas morphisms for topological spaces would be pointless, as any semi-strict classic m-atlas morphism would be strict.

It is a semi-strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2)$, iff f is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a semi-strict classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{C}^1, \mathcal{C}^2)$, iff f is a local m-morphism of E^1 to E^2 .

It is a semi-strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f is a local $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

{def:M-ATLcmorphstrict}

Definition 14.6 (Strict classic m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i$ and A^i be an m-atlas of E^i in the coordinate space C^i .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a strict classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff f is an m-morphism of E^1 to E^2 .

It is a strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as

$\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2)$, iff f is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

Let $f: E^1 \longrightarrow E^2$ be a classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a strict classic E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{C}^1, \mathcal{C}^2)$, iff f is an m-morphism of E^1 to E^2 .

It is a strict classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 - $\mathcal{C}^1$ - $\mathcal{C}^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}^1, \mathcal{E}^2, \mathcal{C}^1, \mathcal{C}^2)$, iff f is an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 .

14.1.3 Abbreviated nomenclature for classic m-atlas morphisms

{def:M-ATLcmorphabbrev}

Definition 14.7 (Abbreviated nomenclature for classic m-atlas morphisms). Let $\mathcal{E}^i$, $\mathcal{C}^i$, $i = 1, 2$, be model categories, $E^i \in \mathcal{E}^i$, $C^i \in \mathcal{C}^i$, A^i be an m-atlas of E^i in the coordinate space C^i and $f: E^1 \rightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate space C^1 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{constrained}), \text{classic}}(A^1, E^1, C^1, A^2, E^2, f)$, iff it is a (full) (maximal) (constrained) (full, max, max-full) (semi-strict, strict)

(semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^1 .

f is a (full) (maximal) (constrained) (semi-strict, strict) classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate space C^1 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{constrained}), \text{classic}}(A^1, E^1, C^1, A^2, E^2, f, \mathcal{E}^1, \mathcal{E}^2)$, iff it is a (full) (maximal) (constrained) (full, max, max-full) (semi-strict, strict)

(semi-strict, strict) classic $\mathcal{E}^1$ - $\mathcal{E}^2$ - E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^1 .

Let $\mathcal{E}^i$, $i = 1, 2$, be a model category, $C^i \in \mathcal{E}^i$, E^i be a topological space, A^i be an m-atlas of E^i in the coordinate space C^i and $f: E^1 \rightarrow E^2$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{constrained}), \text{classic}}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff it is a (full) (maximal) (constrained) (full, max, max-full) (semi-strict, strict)

(semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

f is a (full) (maximal) (constrained) (semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate space C^1 , abbreviated as $\text{isAtl}_{\text{Ar}}^{(\text{constrained}), \text{classic}}(A^1, E^1, C^1, A^2, E^2, f)$, iff it is a (full) (maximal) (constrained) (full, max, max-full) (semi-strict, strict)

(semi-strict, strict) classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^1 .

14.1.4 Proclamations on classic m-atlas morphisms

{lem:M-ATLcmorph}

Lemma 14.8 (Classic m-atlas morphisms). Let E be a set of topological spaces, $E^i \in E, i = 1, 2, C^i$ be a model space, A^i be an m-atlas of E^i in the coordinate space C^i .

1. Let $f: E^1_{\text{triv}} \longrightarrow E^2_{\text{triv}}$ be a classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and $C^1 \overset{\text{mod}}{\subseteq} C^2$. Then f is a strict constrained classic $E_{\mathcal{T}riv} - E_{\mathcal{T}riv} - E^1 - E^2$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof.

f is constrained: $f[E^1] \subseteq E^2$ by hypothesis. E^2 is a model neighborhood of E^2_{triv} by definition 3.1 (Trivial model spaces) on page 21.

{def:trivmod}

f is strict: $E_{\mathcal{T}riv} \overset{\text{full-cat}}{\subseteq} E_{\mathcal{T}riv}$. f is continuous by definition 6.1 (Model functions) on page 25. $E^i \in E$ by hypothesis, $E^i \overset{\text{Ob}}{\in} E_{\mathcal{T}riv}$ and $f \overset{\text{Ar}}{\in} E_{\mathcal{T}riv}$ by definition 3.1. $C^1 \overset{\text{mod}}{\subseteq} C^2$ by definition 14.1 (Classic m-atlas morphisms for model spaces) on page 100.

{def:modfunc}

{def:trivmod}

{def:M-ATLcmorph}

□

2. If E^1 is an open subspace of E^2 then $f: E^1_{\text{triv}} \longrightarrow E^2_{\text{triv}}$ is also a strict constrained classic E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof. Each $E^i \in E$ by hypothesis and each $E^i \overset{\text{Ob}}{\in} E_{\mathcal{T}riv}$ by definition 3.1.

{def:trivmod}

$E^1 \overset{\text{open-subsp}}{\subseteq} E^2$ by hypothesis.

$E^1_{\text{triv}} \overset{\text{mod}}{\subseteq} E^2_{\text{triv}}$ by lemma 3.2 (Trivial model spaces) on page 21.

{lem:trivmod}

$f \overset{\text{Ar}}{\in} E_{\mathcal{T}riv}$ by definition 3.1.

□

{def:trivmod}

Let

1. $\mathcal{C}^i, \mathcal{C}^i, \mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories

2. $\mathcal{E}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{E}'^i$
3. $\mathcal{C}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}'^i$
4. $\mathcal{C}'^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}'^2$
5. $E^i \in \mathcal{E}^i$
6. $C^i \in \mathcal{C}^i$
7. $C'^i \in \mathcal{C}'^i$
8. $C^i \stackrel{\text{mod}}{\subseteq} C'^i$
9. A^i be an m-atlas of E^i in the coordinate space C^i
10. $f: E^1 \longrightarrow E^2$ be a (semi-strict, strict) $(\mathcal{E}^1\text{-}\mathcal{E}^2\text{-})E^1\text{-}E^2(\text{-}\mathcal{C}^1\text{-}\mathcal{C}^2)$ m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 (model function?)

If f is a (semi-strict, strict) $\mathcal{E}^1\text{-}\mathcal{E}^2\text{-}E^1\text{-}E^2(E^1\text{-}E^2)$ classic m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , then f is a (semi-strict, strict) $\mathcal{E}'^1\text{-}\mathcal{E}'^2\text{-}E^1\text{-}E^2(E^1\text{-}E^2)$ classic m-atlas morphism of A^1 to A^2 in the coordinate spaces C'^1, C'^2 .

Proof.

f remains a classic m-atlas morphism with the expanded categories. $\mathcal{C}'^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}'^2$ by hypothesis.

Let $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset$.

$\phi^2 \circ f \circ \phi^{-1}$ is a morphism of $\mathcal{C}^2 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}'^2$ by definition 14.1.

{def:M-ATLcmorph}

f remains semi-strict (strict). If f is a local morphism of $\mathcal{E}^2$ then f is a local morphism of $\mathcal{E}'^2$. If f is a local morphism of E^2 then f is a local morphism of E'^2 . If $f \stackrel{\text{Ar}}{\in} \mathcal{E}^2$ then $f \stackrel{\text{Ar}}{\in} \mathcal{E}'^2$. If f is a morphism of E^2 then f is a morphism of E'^2 .

□

{lem:M-ATLcmorphcomp}

Lemma 14.9 (Composition of classic m-atlas morphisms). Let $\mathcal{E}^i, \mathcal{C}^i, i \in [1, 3]$, be a model category, $E^i \in \mathcal{E}^i, C^i \in \mathcal{C}^i, A^i$ be an m-atlas of E^i in the coordinate space C^i and $f^i: E^i \longrightarrow E^{i+1}, i = 1, 2$, be a (semi-strict, strict) classic $E^i\text{-}E^{i+1}$ m-atlas morphism of A^i to A^{i+1} in the coordinate spaces C^i, C^{i+1} .

$f^2 \circ f^1: E^1 \longrightarrow E^3$ is a (semi-strict, strict) classic $E^1\text{-}E^3$ m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 .

Proof.

$f^2 \circ f^1$ is a **classic E^1 - E^3 m-atlas morphism**. If $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^2$ and $\mathcal{C}^2 \subseteq^{\text{cat}} \mathcal{C}^3$ then $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^3$. Let $(U^1, V^1, \phi^1: U^1 \xrightarrow{\sim} V^1) \in \mathbf{A}^1$, $(U^3, V^3, \phi^3: U^3 \xrightarrow{\sim} V^3) \in \mathbf{A}^3$ with $I \stackrel{\text{def}}{=} U^1 \cap (f^2 \circ f^1)^{-1}[U^3] \neq \emptyset$, $(f^2 \circ f^1)^{-1}[U^3]$ is a model neighborhood by definition 6.1 (Model functions) on page 25 and I is a model neighborhood by item 2 of definition 2.1 (Model spaces) on page 17 . {def:modfunc}
{mod:closed}

For any $u^1 \in I$ and any chart $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in \mathbf{A}^2$ at $f^1(u^1)$, define $I^{1,2} \stackrel{\text{def}}{=} U^1 \cap f^{1-1}[U^2]$, $I^{2,3} \stackrel{\text{def}}{=} U^2 \cap f^{2-1}[U^3]$ and $I_{u^1}^{1,3} \stackrel{\text{def}}{=} U^1 \cap f^{1-1}[I^{2,3}] \subseteq I$. $I_{u^1}^{1,3}$ is a model neighborhood by definition 6.1 and item 2 of definition 2.1 . {def:modfunc
{mod:closed}

$\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I^{1,2}] \longrightarrow V^2$ is a morphism of $\mathcal{C}^2 \subseteq^{\text{cat}} \mathcal{C}^3$ by hypothesis and $\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^2$ is a morphism of $\mathcal{C}^2 \subseteq^{\text{cat}} \mathcal{C}^3$ by item 4. {mod:restriction}
of definition 2.1. $\phi^3 \circ f^2 \circ \phi^{2-1}: \phi^2[I^{2,3}] \longrightarrow V^3$ is a morphism of $\mathcal{C}^3$ by hypothesis and thus $(\phi^3 \circ f^2 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^3) = (\phi^3 \circ f^2 \circ \phi^{2-1}: \phi^2[I^{2,3}] \longrightarrow V^3) \circ (\phi^2 \circ f^1 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^2)$ is a morphism of $\mathcal{C}^3$. {def:model}
 $(\phi^3 \circ f^2 \circ \phi^{1-1}: \phi^1[I_{u^1}^{1,3}] \longrightarrow V^3)$ agrees with $(\phi^3 \circ f^2 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3)$ on $\phi^1[I_{u^1}^{1,3}]$ and $\phi^1[I] = \bigcup_{u^1 \in I} \phi^1[I_{u^1}^{1,3}]$, thus $(\phi^3 \circ f^2 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^3)$ is a morphism of $\mathcal{C}^3$ by item 6 of definition 2.1. {def:model}

$f^2 \circ f^1$ is **(semi-strict, strict)**. If $E^1 \subseteq^{\text{mod}} E^2$ and $E^2 \subseteq^{\text{mod}} E^3$ then $E^1 \subseteq^{\text{mod}} E^3$. If $\mathcal{C}^1 \subseteq^{\text{mod}} \mathcal{C}^2$ and $\mathcal{C}^2 \subseteq^{\text{mod}} \mathcal{C}^3$ then $\mathcal{C}^1 \subseteq^{\text{mod}} \mathcal{C}^3$. If $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^2$ and $\mathcal{C}^2 \subseteq^{\text{cat}} \mathcal{C}^3$ then $\mathcal{C}^1 \subseteq^{\text{cat}} \mathcal{C}^3$.

If f^1 is a local morphism of E^2 and f^2 is a local morphism of E^3 then $f^2 \circ f^1$ is a local morphism of E^3 by lemma 6.15 (Composition of local m-morphisms) on page 36. {lem:modlocalMmorphcomp}

If f^1 is a local $\mathcal{C}^1$ - $\mathcal{C}^2$ morphism of E^2 and f^2 is a local $\mathcal{C}^2$ - $\mathcal{C}^3$ morphism of E^3 then $f^2 \circ f^1$ is a local $\mathcal{C}^1$ - $\mathcal{C}^3$ morphism of E^3 by lemma 6.15. {lem:modlocalMmorphcomp}

If f^1 is a morphism of $E^2 E^3$ and f^2 is a morphism of E^3 then $f^2 \circ f^1$ is a morphism of E^3 .

□

Similar results apply with restrictions on the admissible atlases.

{lem:M-ATLcid}

Lemma 14.10 (Classic m-atlas inclusion and identity). Let

1. $\mathcal{C}^i, \mathcal{C}^i, i = 1, 2$, be model categories
2. $\mathcal{C}^1 \subseteq^{\text{full-cat}} \mathcal{C}^2$

3. $E^i \in \mathcal{E}^i$
4. $E^1 \stackrel{\text{mod}}{\subseteq} E^2$
5. $C^i \in \mathcal{C}^i$
6. $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ ($\mathcal{C}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{C}^2$)
7. A^i be a maximal m-atlas of E^i in the coordinate space C^i
8. $A^1 \subseteq A^2$

The classic m-atlas inclusion morphism $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)} = \mathbf{Id}_{(E^2)}^{(E^1)}$ is a semi-strict classic E^1 - E^2 ($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and a classic $\mathcal{E}^1$ - $\mathcal{E}^2$ ($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof.

$\text{Id}_{E^2}^{E^1}$ is a classic m-atlas morphism:

Let $(U^i, V^i, \phi^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap U^2 \neq \emptyset$.

Since $A^1 \subseteq A^2$, (U^1, V^1, ϕ^1) is compatible with (U^2, V^2, ϕ^2) ,

$\phi^2 \circ \text{Id}_{E^2}^{E^1} \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2 = \phi^2 \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a morphism of C^2 and a morphism of $\mathcal{C}^2$.

$\text{Id}_{E^2}^{E^1}$ is semi-strict:

□

If each $\text{Top}(E^i) \in E^i$ ($E^i \in \mathcal{E}^i$) and $C^1 \stackrel{\text{mod}}{\subseteq} C^2$ ($\mathcal{C}^1 \stackrel{\text{cat}}{\subseteq} \mathcal{C}^2$) then the classic m-atlas inclusion morphism $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a strict classic ($\mathcal{E}^1$ - $\mathcal{E}^2$)- E^1 - E^2 ($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 .

Proof.

$\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a classic m-atlas morphism $\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is a classic E^1 - E^2 ($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 and a classic $\mathcal{E}^1$ - $\mathcal{E}^2$ ($\mathcal{C}^1$ - $\mathcal{C}^2$) m-atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 . by the above proof.

$\text{Id}_{(E^2, C^2)}^{(E^1, C^1)}$ is strict: If each $\text{Top}(E^i) \in E^i$ then $\text{Id}_{E^2}^{E^1}$ is a morphism of E^2 by item 5 of definition 2.1 (Model spaces) on page 17, hence an m-morphism of E^1 to E^2 . {mod:inclusion}

If each $E^i \in \mathcal{E}^i$ $\text{Id}_{E^2}^{E^1}$ is a morphism of $\mathcal{E}^2$ by item 6 of definition 6.7 (Model categories) on page 27, hence an $\mathcal{E}^1$ - $\mathcal{E}^2$ m-morphism of E^1 to E^2 . {modcat:morph}

□

{cor:M-ATLcid}

Corollary 14.11 (Classic m-atlas identity). Let $\mathcal{E}, \mathcal{C}$, be model categories, $\mathbf{E} \in \mathcal{E}$, $\mathbf{C} \in \mathcal{C}$, $\mathbf{A}$ an m-atlas of $\mathbf{E}$ in the coordinate spaces $\mathcal{C}$.

The identity morphism $\mathbf{f} \stackrel{\text{def}}{=} \text{Id}_{(\mathbf{A}, \mathbf{E}, \mathbf{C})}$ is a strict m-atlas morphism of $\mathbf{A}$ to $\mathbf{A}$ in the coordinate spaces $\mathcal{C}, \mathcal{C}$.

14.2 Categories of classic m-atlases and functors

{sub:C-cat}

This subsection defines categories of m-atlases with classic m-atlas morphisms and functors among them, and proves some basic results.

14.2.1 Classic m-atlas categories

Definition 14.12 (Sets of classic m-atlas morphisms). Let $\mathbf{E}^i$ and $\mathbf{C}^i$. $i = 1, 2$, be model spaces. Then

$$\mathcal{Atl}_{\text{Ar}}^{\text{classic}}(\mathbf{E}^1, \mathbf{C}^1, \mathbf{E}^2, \mathbf{C}^2) \stackrel{\text{def}}{=} \left\{ \left((f), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2) \right) \mid \text{isAtl}_{\text{Ar}}^{\text{classic}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f) \right\} \quad (14.1)$$

(full,max,max-full)
(semi-strict,strict)

{def:C-cat}

Definition 14.13 (Categories $\mathcal{Atl}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces.

Let $P \stackrel{\text{def}}{=} \mathbf{E} \times \mathbf{C}$. Then

$$\mathcal{Atl}_{\text{Ob}}^{\text{classic}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{(\mathbf{E}^\mu, \mathbf{C}^\mu) \in P} \mathcal{Atl}_{\text{Ob}}^{\text{classic}}(\mathbf{E}^\mu, \mathbf{C}^\mu) \quad (14.2)$$

(full,max,max-full)
(semi-strict,strict)

$$\mathcal{Atl}_{\text{Ar}}^{\text{classic}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{(\mathbf{E}^\mu, \mathbf{C}^\mu) \in P \\ (\mathbf{E}^\nu, \mathbf{C}^\nu) \in P}} \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(\mathbf{E}^\mu, \mathbf{C}^\mu, \mathbf{E}^\nu, \mathbf{C}^\nu) \quad (14.3)$$

(full,max,max-full)
(semi-strict,strict)

$$\mathcal{Atl}^{\text{classic}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left(\mathcal{Atl}_{\text{Ob}}^{\text{classic}}(\mathbf{E}, \mathbf{C}), \mathcal{Atl}_{\text{Ar}}^{\text{classic}}(\mathbf{E}, \mathbf{C}), \circ^A \right) \quad (14.4)$$

(full,max,max-full) (full,max,max-full) (full,max,max-full)

Let $\mathbf{E}^i$ and $\mathbf{C}^i$, $i = 1, 2$, be model spaces, $\mathbf{A}^i$ be an atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and $f: \mathbf{E}^1 \rightarrow \mathbf{E}^2$ be a (strict, semi-strict) $\mathbf{E}^1$ - $\mathbf{E}^2$ classic M-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

The identity functor $\mathcal{F}_{\text{Classic}, \text{Id}}$ is

$$\mathcal{F}_{\text{Classic}, \text{Id}}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \quad (14.5)$$

$$\mathcal{F}_{\text{Classic}, \text{Id}}((f: \mathbf{E}^1 \longrightarrow \mathbf{E}), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} ((f: \mathbf{E}^1 \longrightarrow \mathbf{E}^2), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \quad (14.6)$$

This nomenclature will be justified below.

{lem:M-ATLCiscat}

Lemma 14.14 ($\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$ is a category). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Then $\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$ is a category and the identity morphism for an object $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$ of $\mathcal{A}t\ell_{\text{Ob}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$ is $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)}$.

Proof. Let $(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, $i \in [1, 3]$, be an object of $\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$ and let $m^i \stackrel{\text{def}}{=} ((f^i), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^{i+1}, \mathbf{E}^{i+1}, \mathbf{C}^{i+1}))$ be a morphism of $\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$. Then

Composition: $f^2 \circ f^1$ is a classic $\mathbf{E}^1\text{-}\mathbf{E}^3$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$ by lemma 14.9 (Composition of classic m-atlas morphisms) on page 105 and $(f^2) \circ^0 (f^1) = (f^2 \circ f^1)$.

{lem:M-ATLcmorphcomp}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{cor:ATLCiscat}

Corollary 14.15 (Subcategories of $\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces. Then all of the following are subcategories of $\mathcal{A}t\ell^{\text{Classic}}(\mathbf{E}, \mathbf{C})$.

1. $\mathcal{A}t\ell_{\text{full}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
2. $\mathcal{A}t\ell_{\text{S-max}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
3. $\mathcal{A}t\ell_{\text{max}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
4. $\mathcal{A}t\ell_{\text{S-max-full}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
5. $\mathcal{A}t\ell_{\text{max-full}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
6. $\mathcal{A}t\ell_{\text{semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
7. $\mathcal{A}t\ell_{\text{full semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$

8. $\mathcal{Att}_{\text{S-max semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
9. $\mathcal{Att}_{\text{max semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
10. $\mathcal{Att}_{\text{S-max-full semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
11. $\mathcal{Att}_{\text{max-full semi-strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
12. $\mathcal{Att}_{\text{strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
13. $\mathcal{Att}_{\text{full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
14. $\mathcal{Att}_{\text{S-max strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
15. $\mathcal{Att}_{\text{max strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
16. $\mathcal{Att}_{\text{S-max-full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$
17. $\mathcal{Att}_{\text{max-full strict}}^{\text{Classic}}(\mathbf{E}, \mathbf{C})$

Remark 14.16. They are not, in general, full subcategories.

The identity functor $\mathcal{F}_{\text{Classic}, \text{Id}}$ is a functor from each of the subcategories to each of the containing categories.

{def:C-cat-cat}

Definition 14.17 (Categories $\mathcal{Att}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories.

Let $\mathbf{P} \stackrel{\text{def}}{=} \text{Ob}(\mathcal{E}) \times \text{Ob}(\mathcal{C})$. Then

$$\mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{\mathbf{E} \in \mathcal{E} \\ \mathbf{C} \in \mathcal{C}}} \mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathbf{E}, \mathbf{C}) \quad (14.7)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left\{ ((f), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. (E^i, C^i) \in \mathbf{P} \wedge \text{isAtI}_{\text{Ar}}(A^1, E^1, C^1, A^2, E^2, C^2, f, \mathcal{E}, \mathcal{C}) \right\} \quad (14.8)$$

$$\mathcal{Att}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} (\mathcal{Att}_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}), \mathcal{Att}_{\text{Ar}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}), \overset{A}{\circ}) \quad (14.9)$$

Similar definitions apply with restrictions on the admissible atlases, the admissible morphisms, or both:

1. $\mathcal{Att}_{\text{full}}^{\text{Classic}}$
2. $\mathcal{Att}_{\text{S-max}}^{\text{Classic}}$
3. $\mathcal{Att}_{\text{max}}^{\text{Classic}}$
4. $\mathcal{Att}_{\text{S-max-full}}^{\text{Classic}}$
5. $\mathcal{Att}_{\text{max-full}}^{\text{Classic}}$
6. $\mathcal{Att}_{\text{full semi-strict}}^{\text{Classic}}$
7. $\mathcal{Att}_{\text{S-max semi-strict}}^{\text{Classic}}$
8. $\mathcal{Att}_{\text{max semi-strict}}^{\text{Classic}}$
9. $\mathcal{Att}_{\text{S-max-full semi-strict}}^{\text{Classic}}$
10. $\mathcal{Att}_{\text{max-full semi-strict}}^{\text{Classic}}$
11. $\mathcal{Att}_{\text{full strict}}^{\text{Classic}}$
12. $\mathcal{Att}_{\text{S-max strict}}^{\text{Classic}}$
13. $\mathcal{Att}_{\text{max strict}}^{\text{Classic}}$
14. $\mathcal{Att}_{\text{S-max-full strict}}^{\text{Classic}}$
15. $\mathcal{Att}_{\text{max-full strict}}^{\text{Classic}}$

e.g.,

$$\mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{(E^\mu, C^\mu) \in \mathbf{P}} \mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(E^\mu, C^\mu) \quad (14.10)$$

$$\mathcal{A}tl_{\text{Ar}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^\mu, C^\mu) \in \mathbf{P} \\ (E^\nu, C^\nu) \in \mathbf{P}}} \mathcal{A}tl_{\text{Ar}}^{\text{Classic}}(E^\mu, C^\mu, E^\nu, C^\nu) \quad (14.11)$$

$$\mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}) \stackrel{\text{def}}{=} \left(\mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C}), \mathcal{A}tl_{\text{Ar}}^{\text{Classic}}(E, C), \circ^A \right) \quad (14.12)$$

Let $(A^1, E^1, C^1) \in \mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$.

$$\text{Id}_{(A^1, E^1, C^1)} \stackrel{\text{def}}{=} ((\text{Id}_{E^1}, \text{Id}_{C^1}), (A^1, E^1, C^1), (A^1, E^1, C^1)) \quad (14.13)$$

{lem:C-cat-ATLCiscat}

Lemma 14.18 ($\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is a category). Let $\mathcal{E}$ and $\mathcal{C}$ be model categories. Then $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is a category and the identity morphism for an object (A^i, E^i, C^i) of $\mathcal{A}tl_{\text{Ob}}^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ is $\text{Id}_{(A^i, E^i, C^i)}$.

Proof. Let (A^i, E^i, C^i) , $i \in [1, 3]$, be an object of $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$ and let $m^i \stackrel{\text{def}}{=} ((f^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$, $i = 1, 2$, be a morphism of $\mathcal{A}tl^{\text{Classic}}(\mathcal{E}, \mathcal{C})$. Then

Composition: $f^2 \circ f^1$ is a morphism of $\mathcal{E}$ and is an E^1 - E^3 m-atlas morphism of A^1 to A^3 in the coordinate spaces C^1, C^3 by lemma 14.9 (Composition of classic m-atlas morphisms) on page 105. {lem:M-ATLcmorphcomp}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

Similar results follow with restrictions on the admissible atlases, the admissible morphisms, or both.

14.2.2 Classic m-atlas functors

{def:Func-M-C}

Definition 14.19 ($\mathcal{F}_{\text{M,Classic}}$). Let E^i , $i = 1, 2$, C^i be model spaces and A^i an m-atlas of E^i in the coordinate model space C^i . Then

$$\mathcal{F}_{\text{M,Classic}}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A^i, E^i, C^i) \quad (14.14)$$

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,)$ be an M-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$. Then

$$\begin{aligned} \mathcal{F}_{\text{M,Classic}}((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2, f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2,), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) &\stackrel{\text{def}}{=} \\ ((f_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) &\quad (14.15) \end{aligned}$$

Theorem 14.20 ($\mathcal{F}_{\text{M,Classic}}$ is a functor). Let $\mathbf{E}$ and $\mathbf{C}$ be sets of model spaces, Then $\mathcal{F}_{\text{M,Classic}}$ is a functor from $\mathcal{At}\ell(\mathbf{E}, \mathbf{C})$ to $\mathcal{At}\ell^{\text{classic}}(\mathbf{E}, \mathbf{C})$.

{the:Func-M-C}

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i)$, $i \in [1, 3]$, be an object of $\mathcal{At}\ell(\mathbf{E}, \mathbf{C})$ and

$m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1})$, $i = 1, 2$, be a morphism of $\mathcal{At}\ell(\mathbf{E}, \mathbf{C})$.

$\mathcal{F}_{\text{M,Classic}}$ satisfies these criteria:

Preservation of endpoints:

$$\begin{aligned} \mathcal{F}_{\text{M,Classic}} o^i &= o^i \stackrel{\text{Ob}}{\in} \mathcal{At}\ell^{\text{classic}}(\mathbf{E}, \mathbf{C}) \\ \mathcal{F}_{\text{Top,M}} m^i &= ((f_0: \mathbf{E}^i \longrightarrow \mathbf{E}^{i+1}), (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i), (\mathbf{A}^{i+1}, \mathbf{E}^{i+1}, \mathbf{C}^{i+1})) \\ &\text{is a morphism from } o^i \text{ to } o^{i+1}. \end{aligned}$$

Identity: Let $(\text{Id}_{\mathbf{E}^i}, \text{Id}_{\mathbf{C}^i}), o^i, o^i$ be an identity morphism of o^i in $\mathcal{At}\ell(\mathbf{E}, \mathbf{C})$. Then $(\text{Id}_{\mathbf{E}^i}: \mathbf{E}^i \longrightarrow \mathbf{E}^i), o^i, o^i$ is an identity morphism of o^i in $\mathcal{At}\ell^{\text{classic}}(\mathbf{E}, \mathbf{C})$.

Composition:

$$\begin{aligned} \mathcal{F}_{\text{M,Classic}}(m^2 \stackrel{A}{\circ} m^1) &= \mathcal{F}_{\text{M,Classic}}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ &= ((f_0^2 \circ f_0^1), o^1, o^3) \\ &= ((f_0^2), o^1, o^2) \stackrel{A}{\circ} ((f_0^1), o^2, o^3) \\ &= \mathcal{F}_{\text{M,Classic}}((f_0^2, f_1^2), o^2, o^3) \stackrel{A}{\circ} \\ &\quad \mathcal{F}_{\text{M,Classic}}((f_0^1, f_1^1), o^1, o^2) \\ &= \mathcal{F}_{\text{M,Classic}} m^2 \stackrel{A}{\circ} \mathcal{F}_{\text{M,Classic}} m^1 \end{aligned}$$

□

Part VIII

Local Coordinate Spaces

This part of the paper defines local coordinate spaces, morphisms among them and categories of them. The definitions given here are based on m-atlas morphisms, but they could as well have been based on classic m-atlas morphisms.

{part:lcs}

15 Local $\mathbf{M}$ -Sigma and $\mathcal{M}$ -Sigma coordinate spaces

Definition 15.1 (Local $\mathbf{M} - \Sigma$ and $\mathcal{M} - \Sigma$ coordinate spaces). Let $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{M}_\alpha, \alpha < A)$ be a sequence of categories, $\mathbf{M} \stackrel{\text{def}}{=} (M_\alpha, \alpha < A)$ be a sequence of spaces and $F \stackrel{\text{def}}{=} (F_\gamma, \gamma < \Gamma)$ a sequence of functions. $\mathbf{L} \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}, \mathbf{A}, F, \Sigma)$ is a (maximal) local $\mathbf{M} - \Sigma$ coordinate space, abbreviated $\max[*]isLCS_{\text{Ob}}\mathbf{L}$, a (maximal) local $\mathcal{M} - \Sigma$ coordinate space, abbreviated $\max[*]isLCS_{\text{Ob}}\mathbf{L}$, and a (maximal) local $A - \Sigma$ coordinate space, abbreviated $\max[*]isLCS_{\text{Ob}}(\mathbf{L}, A)$, iff

1. $\mathcal{M}_1$ is a model category
2. $(\mathcal{M}, \mathbf{M}, \Sigma, F)$ is a $\mathcal{M}$ - Σ prestructure
3. $\mathbf{A}$ is a (maximal) m-atlas of M_0 in M_1 .
4. $\forall \gamma < \Gamma$, if $F_\gamma: (M_{\sigma_\gamma, \alpha_\beta}, \beta < B_\gamma) \longrightarrow \mathbb{T}$ is a constraint function then $F_\gamma[\bigwedge_{\beta < B_\gamma} M_{\sigma_\gamma, \alpha_\beta}] = \{\text{True}\}$, i.e.,
$$\left(\bigvee_{(s_\beta \in M_{\sigma_\gamma, \alpha_\beta}, \beta < B_\gamma)} \right) F_\gamma(s_\beta, \beta < B_\gamma) = \text{True}.$$

Define the total space $(E, \mathcal{E})$ to be M_0 , the Coordinate space $(C, \mathcal{C})$ to be M_1 , the adjunct spaces, if any, to be $M_\alpha, \alpha > 1$, and the adjunct functions if any, to be $F_\alpha, \alpha < A$.

Remark 15.2. There are alternative approaches that are beyond the scope of this paper, e.g.,

1. Requiring the atlas to be full would eliminate certain pathologies.
2. A more complex definition would more easily accommodate structures with more than one atlas, e.g.,
 - (a) A fiber bundle with a differential structure
 - (b) Multiple fiber bundles on the same base space.

Lemma 15.3 (Local $* - \Sigma$ coordinate spaces). let $\mathcal{M}^i, i = 1, 2$, be a sequence of categories, $\mathcal{M}_j^2, j = 0, 1$, model categories, $\mathcal{M}^1 \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$ and $\mathbf{L}^1 \stackrel{\text{def}}{=} (\mathcal{M}^1, \mathbf{M}, \mathbf{A}, F, \Sigma)$ a local $\mathcal{M}^1 - \Sigma$ coordinate space, Then $\mathbf{L}^2 \stackrel{\text{def}}{=} (\mathcal{M}^2, \mathbf{M}, \mathbf{A}, F, \Sigma)$ is a local $\mathcal{M}^2 - \Sigma$ coordinate space,

Proof. $\mathbf{L}^2$ satisfies the conditions of definition 15.1 (Local $\mathbf{M} - \Sigma$ and $\mathcal{M} - \Sigma$ coordinate spaces):

1. $\mathcal{M}_0^2$ and $\mathcal{M}_1^2$ are model categories by hypothesis.
2. $(\mathcal{M}^2, \mathbf{M}, \Sigma, F)$ is a $\mathcal{M}^2$ - Σ prestructure by lemma 10.6 (Prestructures) on page 45.

3. A is a maximal m-atlas of M_0 in M_1 by hypothesis.
4. $\forall \gamma < \Gamma$, if $F_\gamma: (M_{\sigma_\gamma, \alpha_\beta}, \beta < B_\gamma) \longrightarrow \mathbb{T}$ is a constraint function then

$$F_\gamma[\bigwedge_{\beta < B_\gamma} M_{\sigma_\gamma, \alpha_\beta}] = \{\text{True}\}, \text{ i.e.,}$$

$$\left(\bigvee_{(s_\beta \in M_{\sigma_\gamma, \alpha_\beta}, \beta < B_\gamma)} \right) F_\gamma(s_\beta, \beta < B_\gamma) = \text{True}.$$

□

Definition 15.4 (LCS_{Ob}).

$$\text{LCS}_{\text{Ob}}(\mathcal{M}, \Sigma) \stackrel{\text{def}}{=} \left\{ L \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}, \mathbf{A}, \mathbf{F}, \Sigma) \mid \mathbf{M} \stackrel{()}{\in} \mathcal{M} \wedge \text{isLCS}_{\text{Ob}} L \right\} \quad (15.1)$$

A variety of conditions can be imposed on the transition functions $t_\beta^\alpha = \phi_\beta \circ \phi_\alpha^{-1}$ by appropriate choice of category.

Remark 15.5. While some of cases in section 17 (Examples) on page 120 require constraint functions, the specific examples worked out in parts IX and X below do not.

{Examples}

{part:man,part:bun}

16 Morphisms of local coordinate spaces

{sec:lcsmorph}

Informally, a morphism between two local coordinate spaces is a sequence of functions that is compatible with the functions and m-atlases of the two local coordinate spaces. The definition implicitly uses commutation relations, which allows a variety of properties to fall out automatically, e.g., preservation of fibers, being a homomorphism.

{def:LCSmorph}

Definition 16.1 (Morphisms of local coordinate spaces). Let

$\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i$
a sequence of spaces, $L^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma)$ a local $\mathbf{M}^i$ - Σ coordinate space and
 $\mathbf{f} \stackrel{\text{def}}{=} (f_\alpha: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^2, \alpha < A)$.

$\mathbf{f}$ is a morphism from L^1 to L^2 , abbreviated
 $\text{isLCS}_{\text{Ar}}(\mathcal{M}^1, \mathbf{M}^1, \mathbf{A}^1, \mathbf{F}^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, \mathbf{A}^2, \mathbf{F}^2, \mathbf{f})$, iff

1. $\mathbf{f}$ is a morphism from the prestructure $\mathbf{P}^1 \stackrel{\text{def}}{=} (\mathcal{M}^1, \mathbf{M}^1, \Sigma, \mathbf{F}^1)$ to the prestructure $\mathbf{P}^2 \stackrel{\text{def}}{=} (\mathcal{M}^2, \mathbf{M}^2, \Sigma, \mathbf{F}^2)$.
2. (f_0, f_1) is a $\mathcal{M}_0^1$ - $\mathcal{M}_0^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{M}_1^1$ - $\mathcal{M}_1^2$.

$\mathbf{f}: L^1 \longrightarrow L^2$ refers to $\mathbf{f}$ regarded as a morphism from L^1 to L^2 .

$\mathbf{f}$ is a semi-strict morphism from L^1 to L^2 , abbreviated
 $\text{isLCS}_{\text{Ar}}(\mathcal{M}^1, \mathbf{M}^1, \mathbf{A}^1, \mathbf{F}^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, \mathbf{A}^2, \mathbf{F}^2, \mathbf{f})$, iff
semi-strict

1. f is a semi-strict morphism from the prestructure $P^1 \stackrel{\text{def}}{=} (\mathcal{M}^1, \mathbf{M}^1, \Sigma, F^1)$ to the prestructure $P^2 \stackrel{\text{def}}{=} (\mathcal{M}^2, \mathbf{M}^2, \Sigma, F^2)$.
2. (f_0, f_1) is a semi-strict $\mathcal{M}_0^1\text{-}\mathcal{M}_0^2\text{-}\mathbf{M}_0^1\text{-}\mathbf{M}_0^2\text{-}\mathcal{M}_1^1\text{-}\mathcal{M}_1^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model categories $\mathcal{M}_1^1\text{-}\mathcal{M}_1^2$.

f is a strict morphism from $(\mathcal{M}^1, \mathbf{M}^1, \mathbf{A}^1, F^1, \Sigma)$ to $(\mathcal{M}^2, \mathbf{M}^2, \mathbf{A}^2, F^2, \Sigma)$, abbreviated $\text{isLCS}_{\text{Ar}}(\mathcal{M}^1, \mathbf{M}^1, \mathbf{A}^1, F^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, \mathbf{A}^2, F^2, f)$, iff

1. f is a strict morphism from the prestructure $P^1 \stackrel{\text{def}}{=} (\mathcal{M}^1, \mathbf{M}^1, \Sigma, F^1)$ to the prestructure $P^2 \stackrel{\text{def}}{=} (\mathcal{M}^2, \mathbf{M}^2, \Sigma, F^2)$.
2. (f_0, f_1) is a strict $\mathcal{M}_0^1\text{-}\mathcal{M}_0^2\text{-}\mathbf{M}_0^1\text{-}\mathbf{M}_0^2\text{-}\mathcal{M}_1^1\text{-}\mathcal{M}_1^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{M}_1^1, \mathbf{M}_1^2$.

{def:LCSmorphKequiv}

Definition 16.2 (K -equivalence of local coordinate space morphisms). Let

$\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i$ a sequence of spaces, $P^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, F^i, \Sigma)$, $L^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, F^i, \Sigma)$ a local $\mathbf{M}^i\text{-}\Sigma$ coordinate space $f \stackrel{\text{def}}{=} (f_\alpha: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^2, \alpha < A)$ and $g \stackrel{\text{def}}{=} (g_\alpha: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^2, \alpha < A)$ morphisms from L^1 to L^2 and K be a set of ordinals less than A .

f is K -equivalent to g iff f is K -equivalent to g as a prestructure morphism, i.e., $\left(\bigvee_{\alpha \in K} f_\alpha = g_\alpha\right)$.

{lem:LCSmorph}

Lemma 16.3 (morphisms of local coordinate spaces). Let $\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A)$, $i = 1, 2$,

be a sequence of categories, $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i$ a sequence of spaces, $P^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, F^i, \Sigma)$, $L^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, F^i, \Sigma)$ a local $\mathbf{M}^i\text{-}\Sigma$ coordinate space and $f \stackrel{\text{def}}{=} (f_\alpha: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^2, \alpha < A)$, $i = 1, 2, 3$, a (semi-strict, strict) morphism from L^2 to L^2 .

If $\mathcal{M}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}^2$, $\mathbf{S}^i \stackrel{\circ}{\subseteq} \mathbf{S}^2$, $\mathbf{A}^i = \mathbf{A}^2$ and $F_\gamma^i = F_\gamma^2$: $\bigtimes_{\gamma}^{\sigma_\gamma} \mathbf{S}^i \longrightarrow \text{tail} \left(\mathbf{S}^i \right)$ then $\text{Id}_{\mathbf{M}^2}^{\mathbf{M}^1}$ is a strict morphism from L^1 to L^2 .

Proof. $\text{Id}_{P^2}^{P^1}$ is a strict morphism from P^1 to P^2 by lemma 10.11 (Prestructure morphisms) on page 47

{lem:premorph}

$\text{Id}_{(\mathbf{M}_0^2, \mathbf{M}_1^2)}^{(\mathbf{M}_0^1, \mathbf{M}_1^1)}$ is a strict $\mathcal{M}_0^1\text{-}\mathcal{M}_0^2\text{-}\mathbf{M}_0^1\text{-}\mathbf{M}_0^2\text{-}\mathcal{M}_1^1\text{-}\mathcal{M}_1^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{M}_1^1, \mathbf{M}_1^2$. by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79. $\square$

{M-morphcomp:morphcompmorph}

Let $\mathcal{M}'^i \stackrel{\text{def}}{=} (\mathcal{M}'^i_\alpha, \alpha < A)$, $i = 1, 2$, be a sequence of categories and $L'^i \stackrel{\text{def}}{=} (\mathcal{M}'^i, \mathbf{M}^i, \mathbf{A}^i, F^i, \Sigma)$ be a local coordinate space. Then $\mathbf{f}$ is a morphism from L'^1 to L'^2

Proof. The commutation relations do not depend on the categories. $\square$

Let $\mathcal{M}'^i \stackrel{\text{def}}{=} (\mathcal{M}'^i_\alpha, \alpha < A)$, $i = 1, 2$, be a sequence of categories, $\mathcal{M}_0'^i$ and $\mathcal{M}_1'^i$ be model categories, $\mathcal{M}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}'^i$, $\mathbf{P}'^i \stackrel{\text{def}}{=} (\mathcal{M}'^i, \mathbf{M}^i, F^i, \Sigma)$ and $L'^i \stackrel{\text{def}}{=} (\mathcal{M}'^i, \mathbf{M}^i, \mathbf{A}^i, F^i, \Sigma)$. Then

1. L'^i is a local $\mathbf{M}^i$ - Σ coordinate space and $\mathbf{f}$ is a (semi-strict, strict) morphism from L'^1 to L'^2 .
2. If $\mathbf{f}$ is a semi-strict morphism from L^1 to L^2 then $\mathbf{f}$ is a semi-strict morphism from L'^1 to L'^2 .
3. If $\mathbf{f}$ is a strict morphism from L^1 to L^2 then $\mathbf{f}$ is a strict morphism from L'^1 to L'^2 .

Proof. Each L'^i is a local $\mathbf{M}^i$ - Σ coordinate space:

1. $\mathcal{M}_0'^i$ and $\mathcal{M}_1'^i$ are model categories by hypothesis.
2. $\mathbf{P}'^i$ is a $\mathcal{M}'^i$ - Σ prestructure by lemma 10.6 (Prestructures) on page 45. {lem:pre}
3. $\mathbf{A}^i$ is a maximal m-atlas of $\mathbf{M}_0^i$ in $\mathbf{M}_1^i$ by hypothesis.
4. All constraint functions evaluate to True by hypothesis.

Each $\mathbf{f}$ is a (semi-strict, strict) morphism from L'^1 to L'^2 :

1. $\mathbf{f}$ is a morphism from the prestructure $\mathbf{P}'^1$ to the prestructure $\mathbf{P}'^2$ by lemma 10.11 (Prestructure morphisms) on page 47. {lem:premorph}
 If $\mathbf{f}$ is a semi-strict (strict) morphism from the prestructure $\mathbf{P}^1$ to the prestructure $\mathbf{P}^2$ then $\mathbf{f}$ is a semi-strict (strict) morphism from the prestructure $\mathbf{P}'^1$ to the prestructure $\mathbf{P}'^2$ by lemma 10.11. {lem:premorph}
2. (f_0, f_1) is a $\mathbf{M}_0^1$ - $\mathbf{M}_0^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model spaces $\mathbf{M}_i^1, \mathbf{M}_i^2$ by hypothesis.
 If (f_0, f_1) is a semi-strict (strict) $\mathcal{M}_0^1$ - $\mathcal{M}_0^2$ - $\mathbf{M}_0^1$ - $\mathbf{M}_0^2$ - $\mathcal{M}_1^1$ - $\mathcal{M}_1^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model spaces $\mathbf{M}_i^1, \mathbf{M}_i^2$ then (f_0, f_1) is a semi-strict (strict) $\mathcal{M}_0'^1$ - $\mathcal{M}_0'^2$ - $\mathbf{M}_0'^1$ - $\mathbf{M}_0'^2$ - $\mathcal{M}_1'^1$ - $\mathcal{M}_1'^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate model spaces $\mathbf{M}_i^1, \mathbf{M}_i^2$ by lemma 13.9 (M-atlas morphisms) on page 68. {lem:M-ATLmorph}

$\square$

{lem:LCSmorphcomp}

Lemma 16.4 (Composition of local coordinate space morphisms). Let

$\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A), i \in [1, 4]$, be a sequence of categories, $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i$ a sequence of spaces, $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{F}^i, \Sigma)$, $\mathbf{L}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma)$ a local $\mathbf{M}^i$ - Σ coordinate space and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_\alpha^i: \mathbf{M}_\alpha^i \longrightarrow \mathbf{M}_\alpha^{i+1}, \alpha < A), i = 1, 2, 3$, a (semi-strict, strict) morphism from $\mathbf{L}^i$ to $\mathbf{L}^{i+1}$.

$\mathbf{f}^{i+1} \circ \mathbf{f}^i$ is a (semi-strict, strict) morphism from $\mathbf{L}^i$ to $\mathbf{L}^{i+2}$:

Proof. $\mathbf{f}^{i+1} \circ \mathbf{f}^i$ satisfies the criteria of definition 16.1 (Morphisms of local coordinate spaces) on page 115. {def:LCSmorph}

1. If each $\mathbf{f}^j$ is a (semi-strict, strict) morphism from the prestructure $\mathbf{P}^j$ to the prestructure $\mathbf{P}^{j+1}$ then $\mathbf{f}^{i+1} \circ \mathbf{f}^i$ is a (semi-strict, strict) morphism from the prestructure $\mathbf{P}^i$ to the prestructure $\mathbf{P}^{i+2}$ by lemma 10.11 (Prestructure morphisms) on page 47. {lem:premorph}

If each $\mathbf{f}^j$ is a semi-strict (strict) morphism from the prestructure $\mathbf{P}^j$ to the prestructure $\mathbf{P}^{j+1}$ then $\mathbf{f}^{i+1} \circ \mathbf{f}^i$ is a semi-strict (strict) morphism from the prestructure $\mathbf{P}^i$ to the prestructure $\mathbf{P}^{i+2}$ by lemma 10.11. {lem:premorph}

2. $(f_0^{i+1} \circ f_0^i, f_1^{i+1} \circ f_1^i)$ is a semi-strict (strict) $\mathcal{M}_0^i - \mathcal{M}_0^{i+1} - \mathbf{M}_0^i - \mathbf{M}_0^{i+1} - \mathcal{M}_1^i - \mathcal{M}_1^{i+1}$ m-atlas morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+2}$ in the coordinate model spaces $\mathbf{M}_i^i, \mathbf{M}_i^{i+2}$ by lemma 13.9 (M-atlas morphisms) on page 68. If $(f_0^{i+1} \circ f_0^i, f_1^{i+1} \circ f_1^i)$ is a semi-strict (strict) $\mathcal{M}_0^i - \mathcal{M}_0^{i+1} - \mathbf{M}_0^i - \mathbf{M}_0^{i+1} - \mathcal{M}_1^i - \mathcal{M}_1^{i+1}$ m-atlas morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+2}$ in the coordinate model spaces $\mathbf{M}_i^i, \mathbf{M}_i^{i+2}$ then $(f_0^{i+1} \circ f_0^i, f_1^{i+1} \circ f_1^i)$ is a semi-strict (strict) $\mathcal{M}_0^i - \mathcal{M}_0^{i+1} - \mathbf{M}_0^i - \mathbf{M}_0^{i+1} - \mathcal{M}_1^i - \mathcal{M}_1^{i+1}$ m-atlas morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+2}$ in the coordinate model spaces $\mathbf{M}_i^i, \mathbf{M}_i^{i+2}$ by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79. {lem:M-ATLmorphcomp} {M-morphcomp:morphcompmorph}

□

{lem:LCSmorphcompKequiv}

Lemma 16.5 (Composition of $\mathbf{K}$ -equivalent local coordinate space morphisms). Let

1. $\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A), i = 1, 2, 3$, be a sequence of categories,
2. $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i$ be a sequence of spaces,
3. $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{F}^i, \Sigma)$ be a prestructure
4. $\mathbf{L}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma)$ be a local $\mathbf{M}^i$ - Σ coordinate space
5. $\mathbf{f}^1 \stackrel{\text{def}}{=} (f_\alpha^1: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^{i+1}, \alpha < A)$ and $\mathbf{g}^1 \stackrel{\text{def}}{=} (g_\alpha^1: \mathbf{M}_\alpha^1 \longrightarrow \mathbf{M}_\alpha^{i+1}, \alpha < A), i = 1, 2, 3$, be morphisms from $\mathbf{L}^i$ to $\mathbf{L}^{i+1}$
6. $\mathbf{K}$ be a set of ordinals less than A

If each $\mathbf{f}^i$ is $\mathbf{K}$ -equivalent to $\mathbf{g}^i$ then $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1$ is $\mathbf{K}$ -equivalent to $\mathbf{f}^2 \overset{\circ}{\circ} \mathbf{f}^1$.

Proof. The result follows from lemma 10.13 (Composition of prestructure morphisms) on page 48. □

{lem:premorphcomp}

Definition 16.6 (LCS_{Ar}). Let $\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_\alpha^i, \alpha < A), i = 1, 2$, be sequences of categories and $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \alpha_\beta < A, \beta \leq B_\gamma), \gamma < \Gamma)$ a signature over $\mathcal{M}^i$.

{def:LCSmorphAr}

$$\text{LCS}_{\text{Ar}}(\mathcal{M}^1, \Sigma, \mathcal{M}^2) \stackrel{\text{def}}{=} \left\{ (f, (\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma), (\mathcal{M}^2, \mathbf{M}^2, A^2, F^2, \Sigma)) \mid \right. \\ \left. \mathbf{M}^i \in \mathcal{M}^i \wedge \text{isLCS}_{\text{Ar}}(\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, A^2, F^2, f) \right\} \quad (16.1)$$

$$\text{LCS}(\mathcal{M}^i, \Sigma) \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}(\mathcal{M}^i, \Sigma), \text{LCS}_{\text{Ar}}(\mathcal{M}^i, \Sigma, \mathcal{M}^i), \overset{A}{\circ}) \quad (16.2)$$

If $\mathcal{M}^1 \overset{\text{full-cat}}{\subseteq} \mathcal{M}^2$ then

$$\text{LCS}_{\text{Ar}}(\mathcal{M}^1, \Sigma, \mathcal{M}^2) \stackrel{\text{def}}{=} \left\{ (f, (\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma), (\mathcal{M}^2, \mathbf{M}^2, A^2, F^2, \Sigma)) \mid \right. \\ \left. \mathbf{M}^i \in \mathcal{M}^i \wedge \text{isLCS}_{\text{Ar}}(\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, A^2, F^2, f) \right\} \quad (16.3)$$

$$\text{LCS}_{\text{Ar}}^{\text{strict}}(\mathcal{M}^1, \Sigma, \mathcal{M}^2) \stackrel{\text{def}}{=} \left\{ (f, (\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma), (\mathcal{M}^2, \mathbf{M}^2, A^2, F^2, \Sigma)) \mid \right. \\ \left. \mathbf{M}^i \in \mathcal{M}^i \wedge \text{isLCS}_{\text{Ar}}^{\text{strict}}(\mathcal{M}^1, \mathbf{M}^1, A^1, F^1, \Sigma, \mathcal{M}^2, \mathbf{M}^2, A^2, F^2, f) \right\} \quad (16.4)$$

$$\text{LCS}_{\text{semi-strict}}(\mathcal{M}^1, \Sigma) \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}(\mathcal{M}^1, \Sigma), \text{LCS}_{\text{Ar}}^{\text{semi-strict}}(\mathcal{M}^1, \Sigma, \mathcal{M}^1), \overset{A}{\circ}) \quad (16.5)$$

$$\text{LCS}_{\text{strict}}(\mathcal{M}^1, \Sigma) \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}(\mathcal{M}^1, \Sigma), \text{LCS}_{\text{Ar}}^{\text{strict}}(\mathcal{M}^1, \Sigma, \mathcal{M}^1), \overset{A}{\circ}) \quad (16.6)$$

Let $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathbf{M}_\alpha^i, \alpha < A) \in \mathcal{M}^i, i = 1, 2$, be a sequence of spaces, $\mathbf{P}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, F^i, \Sigma)$ and $\mathbf{L}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, A^i, F^i, \Sigma)$ be a local $\mathbf{M}^i$ - Σ coordinate space.

If $\mathcal{M}^1 \overset{\text{full-cat}}{\subseteq} \mathcal{M}^2$ then $\mathbf{S}^1 \subseteq \mathbf{S}^2, A^1 = A^2$ and each $F_\gamma^1 = F_\gamma^2: \times_{\mathbf{S}^1}^{\sigma_\gamma} \rightarrow \text{tail}(\mathbf{S}^1)$

then

$$\text{Id}_{L^2}^{L^1} \stackrel{\text{def}}{=} (\text{Id}_{M^2}^{M^1}, L^1, L^2) \quad (16.7)$$

$$\text{Id}_{L^i} \stackrel{\text{def}}{=} \text{Id}_{L^i}^{L^i} \quad (16.8)$$

This nomenclature is justified below.

{the:LCSiscat}

Theorem 16.7 ($\text{LCS}(\mathcal{M}, \Sigma)$ is a category). Let $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{M}_\alpha, \alpha < A)$ be a sequence of categories and $\Sigma \stackrel{\text{def}}{=} (\sigma_\gamma, \gamma < \Gamma) \stackrel{\text{def}}{=} ((\sigma_{\gamma, \alpha_\beta}, \beta \leq B_\gamma), \gamma < \Gamma)$ a signature over $\mathcal{M}$. Then $\text{LCS}(\mathcal{M}, \Sigma)$, $\text{LCS}_{\text{semi-strict}}(\mathcal{M}, \Sigma)$ and $\text{LCS}_{\text{strict}}(\mathcal{M}, \Sigma)$ are categories.

Let $L^i \stackrel{\text{Ob}}{\in} \text{LCS}(\mathcal{M}, \Sigma)$. Then Id_{L^i} is the identity morphism of L^i :

Proof. Let $L^i \stackrel{\text{def}}{=} (M^i, A^i, F^i, \Sigma)$ be an object of $\text{LCS}(\mathcal{M}, \Sigma)$, let $P^i \stackrel{\text{def}}{=} (\mathcal{M}, M^i, \Sigma, F^i)$ and let $m^i \stackrel{\text{def}}{=} (f^i \stackrel{\text{def}}{=} (f_\alpha^i: M_\alpha^i \longrightarrow M_\alpha^{i+1}, \alpha < A), L^i, L^{i+1})$ be a morphism.

Composition: $f^{i+1} \circ f^i$ is a prestructure morphism from M^i to M^{i+2} by lemma 10.11 {lem:premorph} (Prestructure morphisms) on page 47 and $(f_0^{i+1} \circ f_0^i, f_1^{i+1} \circ f_1^i)$ is an m-atlas morphism from A^i to A^{i+1} by item 1 of lemma 13.13 {M-morphcomp: morphcompmorph} (Composition of m-atlas morphisms) on page 79.

If each f^i is a strict prestructure morphism from P^i to P^{i+1} then $f^{i+1} \circ f^i$ is a strict prestructure morphism from M^i to M^{i+2} by lemma 10.11. {lem:premorph}

If each (f_0^i, f_1^i) is a strict (semi-strict) $\mathcal{M}_0\text{-}\mathcal{M}_0\text{-}M_0^i\text{-}M_0^{i+1}\text{-}\mathcal{M}_1\text{-}\mathcal{M}_1$ m-atlas morphism from A^i to A^{i+1} in the coordinate spaces M_1^i, M_1^{i+1} then $(f_0^{i+1} \circ f_0^i, f_1^{i+1} \circ f_1^i)$ is a strict (semi-strict) $\mathcal{M}_0\text{-}\mathcal{M}_0\text{-}M_0^i\text{-}M_0^{i+1}\text{-}\mathcal{M}_1\text{-}\mathcal{M}_1$ m-atlas morphism from A^i to A^{i+1} in the coordinate spaces M_1^i, M_1^{i+1} by item 1 of lemma 13.13 {M-morphcomp: morphcompmorph} (Composition of m-atlas morphisms) on page 79.

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: Id_{L^1} is the identity morphism of L^1 by lemma 1.19. {lem:labcomp}

□

17 Examples

sec:ex {Examples}

Functors can be constructed among special cases of local coordinate spaces and many mathematical structures, although this paper only gives details for two of them.

Example 17.1 (Manifolds). Choosing the coordinate category as open subsets of a Banach space or more generally a Fréchet space, yields a manifold; the definitions and results of [part IX \(Equivalence of manifolds\)](#) on page 122, also cover manifolds with boundaries and manifolds with tails. {part:man}

Example 17.2 (Fiber bundles). Choosing the coordinate category as the space of products of open sets in a topological space (base) with a fixed fiber and the morphisms as fiber-preserving maps yields a fiber bundle; [part X \(Equivalence of fiber bundles\)](#) on page 146 covers the more general case of a restricted group of transition functions on fibers. {part:bun}

Example 17.3 (Lie groups). Let G be a topological group with group operation $\star$ and identity $\mathbf{1}_G$, C a linear space¹⁰ and A a maximal C^k -atlas¹¹ of G in the coordinate space C . Define $\hat{\star}: G \times G \longrightarrow G$, $\text{isCk}_{\star,A}: G \times G \longrightarrow \text{Truthspace}$ and L by:

$$\hat{\star}(g_1, g_2) \stackrel{\text{def}}{=} g_1 \star g_2^{-1} \quad (17.1)$$

$$\text{isCk}_{\star, \mathbf{1}_G, A}(g_1, g_2) \stackrel{\text{def}}{=} \begin{cases} \text{True} & \hat{\star} \text{ is } C^k \text{ at } (g_1, g_2) \\ \text{False} & \hat{\star} \text{ is not } C^k \text{ at } (g_1, g_2) \end{cases} \quad (17.2)$$

$$\mathbf{M} \stackrel{\text{def}}{=} \left(G, \underset{\text{triv}}{C}, \underset{C^k\text{-triv}}{C}, \text{Truthspace} \right) \quad (17.3)$$

$$\mathcal{M} \stackrel{\text{def}}{=} \underset{\text{sing}}{\mathbf{M}} \quad (17.4)$$

$$\mathbf{F} \stackrel{\text{def}}{=} (\star, \text{isCk}_{\star, A}) \quad (17.5)$$

$$\Sigma \stackrel{\text{def}}{=} ((0, 0, 0), (0, 0, 2)) \quad (17.6)$$

$$\mathbf{L} \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}, A, \mathbf{F}, \Sigma) \quad (17.7)$$

Then L is a local coordinate space iff $\hat{\star}$ is C^k ; the constrain function $\text{isCk}_{\star, A}$ expresses this condition. For $k = \infty$, that LCS is equivalent to a Lie group.

Example 17.4 (Fiber bundle with global sections). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho, A)$ be a fiber bundle¹², $s: X \longrightarrow E$ and

$$\text{isSection}_{E, X, \pi, s}(x) \stackrel{\text{def}}{=} \begin{cases} \text{True} & \pi \circ s(x) = x \\ \text{False} & \pi \circ s(x) \neq x \end{cases} \quad (17.8)$$

a constraint function. Then a simple modification of [definition 29.4 \(Functor from fiber bundles to Local Coordinate Spaces\)](#) on page 177 gives a local coordinate space equivalent to $\mathbf{B}$ with the global section s . {def:BunLCS}

¹⁰See [definition 18.1 \(Linear spaces\)](#) on page 122 and [definition 18.5 \(Trivial \$C^k\$ linear model spaces\)](#) on page 123. {def:trivck}

¹¹See [definition 20.6 \(Maximal \$C^k\$ -atlases\)](#) on page 129. {def:Ck-ATLmax}

¹²See [definition 28.1 \(fiber bundles\)](#) on page 174. {def:Bun}

Example 17.5 (Minkowsky manifolds with foliations). Similarly to the other examples, a local coordinate space can represent a manifold, a global section of the tensor bundle, a constraint function for the signature, a global time coordinate, constraint functions enforcing diferentiability and a constraint function enforcing a foliation.

Part IX

Equivalence of manifolds

{part:man}

For a manifold¹³ the coordinate category is open subsets of a Banach space or more generally a Fréchet space, with an appropriate choice of morphisms. Choosing a separating hyperplane and half space, with open sets in the chosen half space, allows manifolds with boundary. Similarly, choosing a ball with tails allows a manifold with tails.

For differentiable manifolds the coordinate category is similar, but the morphisms are limited to those sufficiently differentiable, in order to impose a diferentiability constraint on the transition functions $t_\beta^\alpha = \phi_\beta \circ \phi_\alpha^{-1}$.

This part of the paper defines C^k -atlases, C^k -manifolds, local coordinate spaces equivalent to C^k -manifolds, categories of them and functors, and gives some basic results.

18 Linear spaces and linear model spaces

sec:lin

This subsection defines spaces and related model spaces suitable for use as the coordinate spaces of generalized C^k manifolds.

def:lin

Definition 18.1 (Linear spaces). Let C be a locally arcwise connected topological subspace of a real (complex) Banach or Fréchet space. Then C is a real (complex) linear space. If C has a non-void interior, i.e., contains a ball, then it is non-degenerate.

Remark 18.2. This paper uses the term *ball* to refer to balls of the underlying space but uses the terms *open set* and *neighborhoods* to refer to the relative topology.

Let $\mathcal{C}$ be a small category whose objects are real (complex) linear spaces and whose morphisms are C^k functions. Then $\mathcal{C}$ is a C^k linear category. If each object of $\mathcal{C}$ is non-degenerate then $\mathcal{C}$ is non-degenerate.

Let C be a real (complex) linear space. Then C_{C^k-Triv} , the category of all C^k functions between nonvoid open subspaces of C , is the trivial C^k linear category of C .

Lemma 18.3 (Open subsets of linear spaces are locally arcwise connected). Let U be an open subset of the real (complex) linear space S . Then U is locally arcwise connected.

¹³The literature has several different definitions of a manifold. This paper uses one chosen for ease of exposition.

Proof. An open subset of a locally arcwise connected space is locally arcwise connected. $\square$

Definition 18.4 (Linear model spaces). Let S be a real (complex) linear space and $\mathcal{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ a model space for S . Then $\mathcal{S}$ is a real (complex) linear model space.

Let $\mathcal{S} \stackrel{\text{def}}{=} (S, \mathcal{S})$ be a real (complex) linear model space such that every morphism of $\mathcal{S}$ is a C^k function. Then $\mathcal{S}$ is a real (complex) C^k linear model space.

Let $\mathcal{S}$ be a small category whose objects are real (complex) C^k linear model spaces and whose morphisms are C^k model functions. Then $\mathcal{S}$ is a C^k linear model category.

{def:trivck}

Definition 18.5 (Trivial C^k linear model spaces). Let C be a real (complex) linear space and $\mathcal{C}$ the category of all C^k functions between open sets of C . Then

$\mathcal{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ is the trivial C^k linear model space of C and $\mathcal{C} \stackrel{\text{def}}{=} (C, \mathcal{C})$ is a real (complex) trivial C^k linear model space.

Let $\mathcal{C}$ be a set of real (complex) linear spaces.

$\mathcal{C} \stackrel{\text{def}}{=} \left\{ \mathcal{C}' \mid \mathcal{C}' \in \mathcal{C} \right\}$ is the set of trivial C^k linear model spaces of $\mathcal{C}$.

The category of trivial C^k linear model spaces of $\mathcal{C}$, abbreviated $\mathcal{C} \stackrel{\text{def}}{=} \mathcal{C} \stackrel{\text{def}}{=} \mathcal{C}$, is the category whose objects are $\mathcal{C} \stackrel{\text{def}}{=} \mathcal{C}$ and whose morphisms are all the C^k functions among them.

$\mathcal{C} \stackrel{\text{def}}{=} \mathcal{C} \stackrel{\text{def}}{=} \mathcal{C}$, the category of open trivial C^k model spaces in $\mathcal{C}$, is the category

whose objects are $\left\{ \mathcal{U} \mid \mathcal{U} \in \mathcal{C} \right\}$, the trivial C^k linear model spaces of non-null open sets of spaces in $\mathcal{C}$, and whose morphisms are all the C^k functions among them.

{lem:trivck}

Lemma 18.6 (Trivial C^k linear model spaces). Let C^i , $i = 1, 2$, be a real (complex) linear space and $\mathcal{C}^i \stackrel{\text{def}}{=} \text{Cat} \left(\mathcal{C}^i \right)$.

1. $\mathcal{C}^i$ is a C^k linear model space.

Proof.

(a) $\text{Ob}(\mathcal{C}^i)$ is an open cover for C^i .

(b) $\text{Ob}(\mathcal{C}^i)$ is closed under finite intersections.

(c) The morphisms of $\mathcal{C}^i$ are C^k , hence continuous.

(d) If $f: A \rightarrow B$ is a morphism, $A' \in \mathcal{C}^i \subseteq A \in \mathcal{C}^i$, $B' \in \mathcal{C}^i \subseteq B \in \mathcal{C}^i$ and $f[A'] \subseteq B'$ then $f|_{A'}: A' \rightarrow B'$ is C^k and thus a morphism.

(e) If $A' \in \mathcal{C}^i \subseteq A \in \mathcal{C}^i$ then the inclusion map $i: A' \hookrightarrow A$ is C^k and thus a morphism.

(f) Restricted sheaf condition: Whenever

- i. U_α and $V_\alpha, \alpha < A$, are objects of $\mathcal{C}^i$.
 - ii. $f_\alpha: U_\alpha \longrightarrow V_\alpha$ are morphisms of $\mathcal{C}^i$.
 - iii. $U \stackrel{\text{def}}{=} \bigcup_{\alpha < A} U_\alpha \stackrel{\text{Ob}}{\in} \mathcal{C}^i$,
 - iv. $V \stackrel{\text{def}}{=} \bigcup_{\alpha < A} V_\alpha \stackrel{\text{Ob}}{\in} \mathcal{C}^i$
 - v. $f: U \longrightarrow V$ is a continuous function and for every $\alpha < A$, f agrees with f_α on U_α
- then f is C^k and thus a morphism of $\mathcal{C}^i$.

□

2. Let $f: C^1 \longrightarrow C^2$ be C^k . Then f is a model function from $\underset{C^k\text{-triv}}{C^1}$ to $\underset{C^k\text{-triv}}{C^2}$.

Proof. Let U^i be a model neighborhood of $\underset{C^k\text{-triv}}{C^i}, i = 1, 2$.

(a) Since U^2 is a model neighborhood of $\underset{C^k\text{-triv}}{C^2}$, U^2 is open, $f^{-1}[U^2]$ is open and thus a model neighborhood of $\underset{C^k\text{-triv}}{C^1}$.

(b) $f[U^1] \subseteq C^2$. Since C^2 is open, it is a model neighborhood of $\underset{C^k\text{-triv}}{C^2}$.

□

3. Let C^1 be an open subspace of C^2 . Then $\underset{C^k\text{-triv}}{C^1} \stackrel{\text{mod}}{\subseteq} \underset{C^k\text{-triv}}{C^2}$.

Proof. By definition 18.5 above, $\underset{C^k\text{-triv}}{C^i} = (C^i, \mathcal{C}^i)$, where $\mathcal{C}^i$ is the category of all C^k functions between open sets of C^i . {def:trivck}

$$\underset{C^k\text{-triv}}{C^1} \stackrel{\text{mod}}{\subseteq} \underset{C^k\text{-triv}}{C^2} :$$

- (a) C^1 is an open subspace of C^2 by hypothesis.
- (b) Every object of $\mathcal{C}^1$ is open in C^1 , thus open in C^2 and thus an object of $\mathcal{C}^2$. If $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^1$ then U and V are open in C^1 and f is C^k in C^1 , thus U and V are open in C^2 and f is C^k in C^2 and $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^2$. If $f: U \xrightarrow{\text{Ob}} V$ is a morphism of $\mathcal{C}^2$, $U \stackrel{\text{Ob}}{\in} \mathcal{C}^1$ and $V \stackrel{\text{Ob}}{\in} \mathcal{C}^1$, then U and V are open in C^1 and f is C^k in C^2 , thus C^k in C^1 , and $f: U \xrightarrow{\text{Ar}} V \in \mathcal{C}^1$.

- (c) If $U \in \mathcal{C}^2$ then U is open in C^2 , hence $U \cap C^1$ is open in C^2 , hence $U \cap C^1$ is open in C^1 and $U \in \mathcal{C}^1$.
- (d) If $U \in \mathcal{C}^2$ and $U \subseteq C^1$ then U is open in C^2 , hence open in C^1 , hence $U \in \mathcal{C}^1$.

□

{def:singCk}

Definition 18.7 (C^k singleton categories). Let $\mathbf{C}$ be a C^k linear model space. Then the C^k singleton category of $\mathbf{C}$, abbreviated $\mathbf{C}^{C^k-Sing}$, is the category whose sole object is $\mathbf{C}$ and whose morphisms are all of the C^k model functions from $\mathbf{C}$ to itself.

19 C^k charts

sec:ckcharts

{def:Ck-chart}

Definition 19.1 (C^k charts). Let C be a linear space and E a topological space. A C^{k14} chart (U, V, ϕ) of E in the coordinate space C consists of

1. A nonvoid open subset $U \subseteq E$, known as a coordinate patch
2. An open subset $V \subseteq C$
3. A homeomorphism $\phi: U \xrightarrow{\cong} V$, known as a coordinate function

Remark 19.2. I consider it clearer to explicate the range, rather than the conventional usage of specifying only the domain and function or the minimalist usage of specifying only the function.

A chart (U, V, ϕ) is non-degenerate iff V contains a ball of the underlying Banach or Fréchet space.

{lem:Ck-chart}

Lemma 19.3 (C^k charts). Let C be a linear space and E a topological space. The triple (U, V, ϕ) is a C^k chart of E in the coordinate space C iff it is an m -chart of E in the coordinate space $\mathbf{C}^{C^k-triv}$.

Proof. U is a model neighborhood of E iff it is a nonvoid open set of E . V is a model neighborhood of $\mathbf{C}^{C^k-triv}$ iff it is a nonvoid open set of C . ϕ is required to be a homeomorphism from U to V in either case. □

{def:Ck-subchart}

Definition 19.4 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and U' be a nonvoid open subset of U . Then $(U', V', \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U', V'})$ is a subchart of (U, V, ϕ) .

By abuse of language we will write (U', V', ϕ) for (U', V', ϕ') .

¹⁴With $k \in \mathbb{N} \cup \{\infty, \omega\}$.

Lemma 19.5 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and (U', V', ϕ') be a triple. Then (U', V', ϕ') is a subchart of (U, V, ϕ) iff it is a subchart of (U, V, ϕ) considered as a chart of E in the coordinate space C . {lem:Ck-subchart}

Proof. (U', V', ϕ') satisfies the conditions of definition 19.1; {def:Ck-chart}

1. U is open by definition 19.1 {def:Ck-chart}

2. U' is required to be a nonvoid open subset of U in either case, and thus a model neighborhood of E .

3. ϕ is a homeomorphism, so $V' = \phi[U']$ is open, and thus a model neighborhood of C .

4. U' is a model neighborhood of E iff it is an open set of E .

5. ϕ is a homeomorphism, so $\phi|_{U'}: U' \xrightarrow{\cong} \phi[U']$ is also.

□ {cor:Ck-subchart}

Corollary 19.6 (C^k subcharts). Let (U, V, ϕ) be a C^k chart of E in the coordinate space C and (U', V', ϕ') be a subchart of (U, V, ϕ) . Then (U', V', ϕ') is a C^k chart of E in the coordinate space C .

Proof. The result follows from lemma 19.3 (C^k charts) above, lemma 11.3 (M-charts) on page 51 and definition 11.5 (Subcharts) on page 52. {lem:Ck-chart} {def:M-subchart} {Ck-compat}

Definition 19.7 (C^k compatibility). Let (U, V, ϕ) and (U', V', ϕ') be C^k charts of E in the coordinate space C . Then (U, V, ϕ) is C^k compatible with (U', V', ϕ') iff either

1. U and U' are disjoint

2. The transition function $t = \phi' \circ \phi^{-1}|_{\phi[U \cap U']}$ is a C^k diffeomorphism.

{lem:Ck-compatSYM}

Lemma 19.8 (Symmetry of C^k compatibility). Let (U, V, ϕ) and (U', V', ϕ') be C^k charts of E in the coordinate space C . Then (U, V, ϕ) is C^k compatible with (U', V', ϕ') iff (U', V', ϕ') is C^k compatible with (U, V, ϕ) .

Proof. It suffices to prove the implication in only one direction.

1. $U \cap U' = U' \cap U$.

2. Since the transition function $t = \phi' \circ \phi^{-1}|_{\phi[U \cap U']}$ is a C^k diffeomorphism of C , so is $t^{-1} = \phi \circ \phi'^{-1}|_{\phi'[U' \cap U']}$.

□

Lemma 19.9 (C^k compatibility of subcharts). Let (U_i, V_i, ϕ_i) , $i = 1, 2$, be C^k charts of E in the coordinate space C , (U'_i, V'_i, ϕ'_i) be subcharts and (U_1, V_1, ϕ_1) be C^k compatible with (U_2, V_2, ϕ_2) . Then (U'_1, V'_1, ϕ'_1) is C^k compatible with (U'_2, V'_2, ϕ'_2) .

Proof. If $U_1 \cap U_2 = \emptyset$ then $U'_1 \cap U'_2 = \emptyset$. If $U'_1 \cap U'_2 = \emptyset$ then (U'_1, V'_1, ϕ'_1) is C^k compatible with (U'_2, V'_2, ϕ'_2) . Otherwise, the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism and hence $t_2^1 \upharpoonright_{\phi_1[U'_1 \cap U'_2]}: \phi_1[U'_1 \cap U'_2] \xrightarrow{\cong} \phi_2[U'_1 \cap U'_2]$ is a C^k diffeomorphism. $\square$

Corollary 19.10 (C^k compatibility with subcharts). Let (U, V, ϕ) be a C^k charts of E in the coordinate space C and (U', V', ϕ') a subchart. Then (U', V', ϕ') is C^k compatible with (U, V, ϕ) .

Proof. (U, V, ϕ) is C^k compatible with itself and is a subchart of itself. $\square$

Definition 19.11 (Covering by C^k charts). Let $\mathbf{A}$ be a set of charts of E in the coordinate space C . $\mathbf{A}$ covers E iff $\pi_1[\mathbf{A}]$ covers E , i.e., $E = \bigcup \pi_1[\mathbf{A}]$.

20 C^k -atlases

{sec:ckatlases}

A set of charts can be atlases for different coordinate spaces even if it is for the same total space. In order to aggregate them into categories, there must be a way to distinguish them. Including the two¹⁵ spaces in the definitions of the categories serves the purpose.

{def:Ck-atlas}

Definition 20.1 (C^k -atlases). Let $\mathbf{A}$ be a set of mutually C^k compatible charts of E in the coordinate space C . $\mathbf{A}$ is a C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C)$, iff $\mathbf{A}$ covers E .

$\mathbf{A}$ is a full C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{\text{full}, C^k}(\mathbf{A}, E, C)$, iff

1. $\pi_1[\mathbf{A}]$ covers E
2. $\pi_2[\mathbf{A}]$ covers C .

$\mathbf{A}$ is non-degenerate iff it contains a non-degenerate chart.

By abuse of language we write $U \in \mathbf{A}$ for $U \in \pi_1[\mathbf{A}]$.

Let E be a topological space and C a linear space. Then

$$\mathcal{Atl}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \right\} \quad (20.1)$$

$$\mathcal{Atl}_{\text{Ob}}^{\text{full}, C^k}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{\text{full}, C^k}(\mathbf{A}, E, C) \right\} \quad (20.2)$$

¹⁵The total space is redundant, but convenient.

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then

$$\mathcal{Att}_{\text{Ob}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E_\mu \in \mathbf{E} \\ C_\mu \in \mathbf{C}}} \mathcal{Att}_{\text{Ob}}(E_\mu, C_\mu) \quad (20.3)$$

$$\mathcal{Att}_{\text{Ob}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \begin{array}{c} \text{isAtl}^{\text{C}^k}_{\text{Ob}}(\mathbf{A}, E, C) \\ \text{full} \end{array} \right\} \quad (20.4)$$

{lem:Ck-atlas}

Lemma 20.2 (C^k -atlases). Let E be a topological space, C be a linear space and $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C .

If $\mathbf{A}$ is non-degenerate then C is non-degenerate.

Proof. Let (U, V, ϕ) be a non-degenerate chart of $\mathbf{A}$. V contains a ball and is contained in C . $\square$

If C is non-degenerate and $\mathbf{A}$ is full then $\mathbf{A}$ is non-degenerate.

Proof. Let B be a ball in C with center v . Since $\mathbf{A}$ is full, it contains a chart (U, V, ϕ) with $v \in V$. Then V contains a ball with center v . $\square$

{def:CkCompat}

Definition 20.3 (Compatibility of charts with C^k -atlases). A chart (U, V, ϕ) of E in the coordinate space C is C^k compatible with a C^k -atlas $\mathbf{A}$ iff it is C^k compatible with every chart in the atlas.

{lem:CkCompat}

Lemma 20.4 (Compatibility of subcharts with C^k -atlases). Let $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C and $\mathbf{C}_1 = (U_1, V_1, \phi_1)$ a C^k chart in $\mathbf{A}$. Then any subchart of $\mathbf{C}_1$ is C^k compatible with $\mathbf{A}$.

Proof. Let $\mathbf{C}' = (U', V', \phi')$ be a subchart of $\mathbf{C}_1$ and $\mathbf{C}_2 = (U_2, V_2, \phi_2)$ another chart in $\mathbf{A}$.

1. If $U_1 \cap U_2 = \emptyset$, then $U' \cap U_2 = \emptyset$.
2. If $U' \cap U_2 = \emptyset$ then $\mathbf{C}'$ is C^k compatible with $\mathbf{C}_2$.
3. Otherwise the transition function $t_2^1 \stackrel{\text{def}}{=} \phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism and thus $t_2^1 \upharpoonright_{\phi_1[U' \cap U_2]}$ is a C^k diffeomorphism.

$\square$

{Ck-atl:extensions}

Lemma 20.5 (Extensions of C^k -atlases). Let $\mathbf{A}$ be a C^k atlas of E in the coordinate space C and (U_i, V_i, ϕ_i) , $i = 1, 2$, be a C^k chart of E in the coordinate space C C^k compatible with $\mathbf{A}$ in the coordinate space C . Then (U_1, V_1, ϕ_1) is C^k compatible with (U_2, V_2, ϕ_2) in the coordinate space C .

Proof. If $U_1 \cap U_2 = \emptyset$ then (U_1, V_1, ϕ_1) is C^k compatible with (U_2, V_2, ϕ_2) . Otherwise, $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}: \phi_1[U_1 \cap U_2] \xrightarrow{\cong} \phi_2[U_1 \cap U_2]$ is a homeomorphism. It remains to show that $\phi_2 \circ \phi_1^{-1} \upharpoonright_{\phi_1[U_1 \cap U_2]}$ is a C^k diffeomorphism. Let $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\alpha < A$, be charts in $\mathbf{A}$ such that $U_1 \cap U_2 \subseteq \bigcup_{\alpha < A} U'_\alpha$ and $U_1 \cap U_2 \cap U'_\alpha \neq \emptyset$, $\alpha < A$. Since the charts are C^k compatible with $(U'_\alpha, V'_\alpha, \phi'_\alpha)$, $\phi_2 \circ \phi'_\alpha^{-1} \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ and $\phi'_\alpha \circ \phi_1^{-1} \upharpoonright_{U_1 \cap U_2 \cap U'_\alpha}$ are C^k diffeomorphisms and thus $\phi_2 \circ \phi_1^{-1} = \phi_2 \circ \phi'_\alpha^{-1} \circ \phi'_\alpha \circ \phi_1^{-1}$ is a C^k diffeomorphism. $\square$

{def:Ck-ATLmax}

Definition 20.6 (Maximal C^k -atlases). Let E be a topological space, C be a linear space and $\mathbf{A}$ be a non-degenerate C^k -atlas of E in the coordinate space C .

$\mathbf{A}$ is a maximal C^k -atlas of E in the coordinate space C , abbreviated $\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C)$, iff $\mathbf{A}$ is a C^k -atlas that cannot be extended by adding an additional C^k compatible chart.

$$\text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \stackrel{\text{def}}{=} \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \wedge \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \quad (20.5)$$

Theorem 20.7 (Existence and uniqueness of maximal C^k -atlases). Let $\mathbf{A}$ be a C^k -atlas of E in the coordinate space C . Then there exists a unique maximal C^k -atlas $\text{Atlas}_{\text{max}}^{C^k}(\mathbf{A}, E, C)$ of E in the coordinate space C compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all C^k -atlases of E in the coordinate space C containing $\mathbf{A}$ and C^k compatible in the coordinate space C with all of the C^k charts in $\mathbf{A}$. Let $\mathbf{P}$ be a maximal chain of $\mathbf{A}$. Then $\mathbf{A}' = \bigcup_{\max} \mathbf{P}$ is a maximal C^k atlas of E in the coordinate space C C^k compatible with $\mathbf{A}$. Uniqueness follows from lemma 20.5 (Extensions of C^k -atlases) on page 128. $\square$

{Ck-atl:extensions}

{def:Ck-ATLsets}

Definition 20.8 (Sets of C^k -atlases). Let E be a topological space and C be a linear space.

$$\text{Att}_{\text{Ob}}^{C^k}(E, C) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, E, C) \mid \text{isAtl}_{\text{Ob}}^{C^k}(\mathbf{A}, E, C) \right\} \quad (20.6)$$

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then

$$\text{Att}_{\text{Ob}}^{C^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{E \in \mathbf{E} \\ C \in \mathbf{C}(\text{full, max, max-full})}} \text{Att}_{\text{Ob}}^{C^k}(E, C) \quad (20.7)$$

21 C^k -atlas morphisms and functors

{sec:ckmorph}

This section introduces a taxonomy for morphisms between C^k -atlases, defines categories of C^k -atlases, defines functors among them, defines functors between them and categories of m-atlases, defines inverse functors and proves some basic results.

21.1 C^k -atlas morphisms

This subsection introduces the notions of C^k -atlas morphisms, and proves some basic results.

21.1.1 Definitions of C^k -atlas morphisms

{def:Ck-ATLmorph}

Definition 21.1 (C^k -atlas morphisms). Let E^i , $i = 1, 2$, be a topological space, C^i a linear space, $\mathbf{A}^i$ a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ a pair¹⁶ of functions.

$\mathbf{f}$ is an E^1 - E^2 C^k -morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1)$, iff

1. $f_0: E^1 \longrightarrow E^2$ is a continuous function.
2. $f_1: C^1 \longrightarrow C^2$ is a C^k function.
3. for any $u^1 \in E^1$, any chart $(U^1, V^1, \phi^1: U^1 \xrightarrow{\sim} V^1) \in \mathbf{A}^1$ at u^1 and any chart $(U^2, V^2, \phi^2: U^2 \xrightarrow{\sim} V^2) \in \mathbf{A}^2$ at $f_0(u^1)$ there exists a subchart $(U'^1, V'^1, \phi'^1: U'^1 \xrightarrow{\sim} V'^1) \in \mathbf{A}^1$ at u^1 , an open set $U'^2 \subseteq U^2$ and a chart $(U'^2, V'^2, \phi'^2: U'^2 \xrightarrow{\sim} V'^2) \in \mathbf{A}^2$ at $f_0(u^1)$ such that $f_0[U'^1] \subseteq U'^2$ and diagram (13.1) in definition 13.1 (M-atlas morphisms for model spaces) on page 61 is commutative, as shown in fig. 7 (Completed m-atlas morphism) on page 62.

{def:M-ATLmorph}
{fig:AtlM1}

The identity morphism of $(\mathbf{A}^i, E^i, C^i)$ is

$$\text{Id}_{(\mathbf{A}^i, E^i, C^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{C^i}), (\mathbf{A}^i, E^i, C^i), (\mathbf{A}^i, E^i, C^i)) \quad (21.1)$$

This nomenclature will be justified later.

Let E^i , $i = 1, 2$, be topological spaces and C^i be linear spaces. Then

$$\text{Atl}_{\text{Ar}}^{C^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid \text{isAtl}_{\text{Ar}}^{C^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (21.2)$$

¹⁶The conventional definition uses only the first of the two functions and a slightly different compatibility condition.

$$\mathcal{Atl}_{\text{Ar}}^{\text{C}^k}_{\text{full}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (21.3)$$

$$\mathcal{Atl}_{\text{Ar}}^{\text{C}^k}_{\text{max}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}_{\text{max}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (21.4)$$

$$\mathcal{Atl}_{\text{Ar}}^{\text{C}^k}_{\text{S-max}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}_{\text{S-max}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (21.5)$$

$$\mathcal{Atl}_{\text{Ar}}^{\text{C}^k}_{\text{max-full}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}_{\text{max-full}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (21.6)$$

$$\mathcal{Atl}_{\text{Ar}}^{\text{C}^k}_{\text{S-max-full}}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \left\{ ((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \mid \right. \\ \left. \text{isAtl}_{\text{Ar}}^{\text{C}^k}_{\text{S-max-full}}(A^1, E^1, C^1, A^2, E^2, C^2, f_0, f_1) \right\} \quad (21.7)$$

{def:Ck-ATLmorphequiv}

Definition 21.2 (Equivalence of C^k -atlas morphisms). Let E^i , $i = 1, 2$, be a topological space, C^i a linear space, A^i a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ and $\mathbf{g} \stackrel{\text{def}}{=} (g_0, g_1)$ E^1 - E^2 C^k morphisms of A^1 to A^2 in the coordinate spaces C^1, C^2 .

$\mathbf{f}$ is C^k -equivalent to $\mathbf{g}$ iff $f_0 = g_0$.

21.1.2 Proclamations on C^k -atlas morphisms

{lem:Ck-ATLmorph}

Lemma 21.3 (C^k -atlas morphisms). Let $\mathbf{E}$ be a set of topological spaces, $\mathbf{C}$ a set of C^k linear spaces, $E^i \in \mathbf{E}$, $i = 1, 2$, $C^i \in \mathbf{C}$, A^i a C^k -atlas of E^i in the coordinate space C^i and $\mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2)$ a pair of functions.

$\mathbf{f}$ is an E^1 - E^2 C^k morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff $\mathbf{f}$ is a strict $\mathbf{E} - \mathbf{E} - E^1 - E^2 - \mathbf{C} - \mathbf{C}$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1_{\text{C}^k\text{-triv}}, \mathbf{C}^2_{\text{C}^k\text{-triv}}$.

Proof. The model neighborhoods of $\mathbf{C}^i_{\text{Ck-triv}}$ are the open sets of $\mathbf{C}^i$, $\mathbf{E}_{\mathcal{T}riv}^{\text{full-cat}} \subseteq \mathbf{E}_{\mathcal{T}riv}$, the morphisms of $\mathbf{E}_{\mathcal{T}riv}$ are the continuous functions between spaces in $\mathbf{E}$, $\mathbf{C}_{\text{Ck-triv}}^{\text{full-cat}} \subseteq \mathbf{C}_{\text{Ck-triv}}$, and the morphisms of $\mathbf{C}_{\text{Ck-triv}}$ are the C^k functions between spaces in $\mathbf{C}$. $\square$

$\mathbf{f}$ is an E^1 - E^2 C^k -morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ iff $\mathbf{f}$ is a strict E^1 - E^2 - $\mathbf{C}_{\text{Ck-triv}}^{\text{full-cat}}$ - $\mathbf{C}_{\text{Ck-triv}}^{\text{full-cat}}$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1_{\text{Ck-triv}}, \mathbf{C}^2_{\text{Ck-triv}}$.

Proof. The model neighborhoods of $\mathbf{C}^i_{\text{Ck-triv}}$ are the open sets of $\mathbf{C}^i$, $\mathbf{E}_{\mathcal{T}riv}^{\text{full-cat}} \subseteq \mathbf{E}_{\mathcal{T}riv}$, $\mathbf{C}_{\text{Ck-triv}}^{\text{full-cat}} \subseteq \mathbf{C}_{\text{Ck-triv}}$ and the morphisms of $\mathbf{C}_{\text{Ck-triv}}$ are the C^k functions between spaces in $\mathbf{C}$. $\square$

{cor:Ck-ATLmorph}

Corollary 21.4 (C^k -atlas morphisms). Let $E^i, i = 1, 2, 3$, be a topological space, $\mathbf{C}^i$ a linear space, $\mathbf{A}^i$ a C^k -atlas of E^i in the coordinate space $\mathbf{C}^i$ and (f_0^i, f_1^i) an E^i - E^{i+1} C^k -morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$. Then $(f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is a E^1 - E^3 C^k -morphism of $\mathbf{A}^1$ to $\mathbf{A}^3$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^3$.

Proof. The result follows from item 1 of $\text{lemma 13.13}^{\text{lem:M-ATLmorphcomp}}$ (**Composition of m-atlas morphisms**) on page 79. $\square$

{M-morphcomp:morphcompmorph}

{lem:Ck-ATLmorphequiv}

Lemma 21.5 (Composition of equivalent C^k -atlas morphisms). Let $E^i, i = 1, 2, 3$, be a topological space, $\mathbf{C}^i$ be a C^k linear space, $\mathbf{A}^i$ be a C^k -atlas of E^i in the coordinate space $\mathbf{C}^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_0^i: E^i \longrightarrow E^{i+1}, f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_0^i: E^i \longrightarrow E^{i+1}, g_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1})$ be C^k -equivalent E^i - E^{i+1} m-atlas morphisms of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$.

Then $\mathbf{f}^2 \circ \mathbf{f}^1$ is C^k -equivalent to $\mathbf{g}^2 \circ \mathbf{g}^1$.

Proof. $f_0^1 = g_0^1$ and $f_0^2 = g_0^2$, hence $f_0^2 \circ f_0^1 = g_0^2 \circ g_0^1$. $\square$

21.2 Categories of C^k atlases and functors

21.2.1 Categories of C^k atlases

{def:Ck-ATLcat}

Definition 21.6 (Categories of C^k atlases). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Let $\mathbf{P} \stackrel{\text{def}}{=} \mathbf{E} \times \mathbf{C}$. Then

$$\mathcal{Att}_{\text{Ar}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \bigcup_{\substack{(E^\mu, C^\mu) \in \mathbf{P} \\ (E^\nu, C^\nu) \in \mathbf{P}}} \mathcal{Att}_{\text{Ar}}^{\text{C}^k}(E^\mu, C^\mu, E^\nu, C^\nu) \quad (21.8)$$

$$\mathcal{Att}_{(\max)}^{C^k}(E, \mathbf{C}) \stackrel{\text{def}}{=} \left(\mathcal{Att}_{(\max)}^{C^k}(\mathbf{E}, \mathbf{C}), \mathcal{Att}_{(\max)}^{C^k}(\mathbf{E}, \mathbf{C}), \circ^A \right) \quad (21.9)$$

{lem:Ck-ATLiscat}

Lemma 21.7 ($\mathcal{Att}_{(\max)}^{C^k}(\mathbf{E}, \mathbf{C})$ is a category). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then each of $\mathcal{Att}_{(\max)}^{C^k}(\mathbf{E}, \mathbf{C})$ is a category.

Let $(A^i, E^i, C^i) \in \mathcal{Att}_{(\max)}^{C^k}(\mathbf{E}, \mathbf{C})$. Then $\text{Id}_{(A^i, E^i, C^i)}$ is the identity morphism for (A^i, E^i, C^i) .

Proof. Let $(A^i, E^i, C^i), i \in [1, 3]$, be an object of $\mathcal{Att}^{C^k}(\mathbf{E}, \mathbf{C})$ and let $m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), (A^i, E^i, C^i), (A^{i+1}, E^{i+1}, C^{i+1}))$, $i = 1, 2$, be a morphism of $\mathcal{Att}^{C^k}(\mathbf{E}, \mathbf{C})$. Then

Composition: $((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3))$ is a morphism of $\mathcal{Att}^{C^k}(\mathbf{E}, \mathbf{C})$ by corollary 21.4 (**C^k-atlas morphisms**) on page 132.

{cor:Ck-ATLmorph}

Associativity: Composition is associative by lemma 1.19 (**Tuple composition for labeled morphisms**) on page 13.

{lem:labcomp}

Identity: $\text{Id}_{(A^i, E^i, C^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

21.2.2 C^k atlas functors

{def:Func-Ck-M}

Definition 21.8 (Functors from C^k atlases to m-atlases). Let $E^i, i = 1, 2$, be a topological space, C^i a linear space, A^i C^k-atlases of E^i in the coordinate space C^i , $f_0: E^1 \rightarrow E^2$ continuous and $f_1: C^1 \rightarrow C^2$ C^k. Then

$$\mathcal{F}_{C^k, M}^{C^k}(A^i, E^i, C^i) \stackrel{\text{def}}{=} (A^i, E^i, C^i)_{\text{triv } C^k\text{-triv}} \quad (21.10)$$

$$\mathcal{F}_{C^k, M}^{C^k}((f_0, f_1), (A^1, E^1, C^1), (A^2, E^2, C^2)) \stackrel{\text{def}}{=} \left((f_0, f_1), (A^1, E^1, C^1)_{\text{triv } C^k\text{-triv}}, (A^2, E^2, C^2)_{\text{triv } C^k\text{-triv}} \right) \quad (21.11)$$

{the:Func-Ck-M}

Theorem 21.9 (Functors from C^k atlases to m-atlases). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then $\mathcal{F}_{C^k, M}^{C^k}$ is a functor from $\mathcal{Att}_{\text{triv } C^k\text{-triv}}^{C^k}(\mathbf{E}, \mathbf{C})$ to $\mathcal{Att}_{\text{triv } C^k\text{-triv}}^{C^k}(\mathbf{E}, \mathbf{C})$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, E^i, C^i), i \in [1, 3]$, be an object of $\mathcal{Att}^{C^k}(\mathbf{E}, \mathbf{C})$, $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{C^k, M}^{C^k} o^i = (A^i, E^i, C^i)_{\text{triv } C^k\text{-triv}}, m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i), o^i, o^{i+1}), i = 1, 2$, be a morphism from o^i to o^{i+1} and $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{C^k, M}^{C^k} m^i = ((f_0^i, f_1^i).o'^i, o'^{i+1})$, be the corresponding morphism in $\mathcal{Att}_{\text{triv } C^k\text{-triv}}^{C^k}(\mathbf{E}, \mathbf{C})$.

Preservation of endpoints:

$$\begin{aligned}\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^i &= (\mathbf{A}^i, \mathbf{E}_{\text{triv}}^i, \mathbf{C}_{\mathbf{C}^k - \text{triv}}^i) \\ &= o'^i \\ \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} m^i &= m'^i \\ &= ((f_0^i, f_1^i), o'^i, o'^{i+1})\end{aligned}$$

$\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} m^i$ is a morphism from $\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^i$ to $\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^{i+1}$:

Identity:

$$\begin{aligned}\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} \text{Id}_{o^i} &= \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} ((\text{Id}_{E^1}, \text{Id}_{C^1}), o^i, o^i) \\ &= ((\text{Id}_{E_{\text{triv}}^i}, \text{Id}_{C^i}), o'^i, o'^i) \\ &= ((\text{Id}_{E_{\text{triv}}^i}, \text{Id}_{C^i}), \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^i, \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^i) \\ &= \text{Id}_{\mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} o^i}\end{aligned}$$

Composition:

$$\begin{aligned}m^2 \circ^A m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3) \\ \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} (m^2 \circ^A m^1) &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o'^1, o'^3) \\ &= ((f_0^2, f_1^2), o'^2, o'^3) \circ^A ((f_0^1, f_1^1), o'^1, o'^2) \\ &= \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} ((f_0^2, f_1^2), o^2, o^3) \circ^A \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} ((f_0^1, f_1^1), o^1, o^2) \\ &= \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} m^2 \circ^A \mathcal{F}_{\mathbf{C}^k, \mathbf{M}}^{\mathbf{C}^k} m^1\end{aligned}$$

□

{def:Func-M-Ck}

Definition 21.10 (Functors from m-atlases to $\mathbf{C}^k$ atlases). Let $\mathbf{E}^i$, $i = 1, 2$, be model spaces, $\mathbf{C}^i$ be linear model spaces, $\mathbf{A}^i$ a maximal m-atlas of $\mathbf{E}^i$ in the coordinate space $\mathbf{C}^i$ and (f_0, f_1) an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$, Then

$$\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{\mathbf{C}^k}(\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (\mathbf{A}^i, \pi_1(\mathbf{E}^i), \pi_1(\mathbf{C}^i)) \quad (21.12)$$

$$\begin{aligned}\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{\mathbf{C}^k}((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) &\stackrel{\text{def}}{=} \\ ((f_0, f_1), (\mathbf{A}^1, \pi_1(\mathbf{E}^1), \pi_1(\mathbf{C}^1)), (\mathbf{A}^2, \pi_1(\mathbf{E}^2), \pi_1(\mathbf{C}^2))) &\quad (21.13)\end{aligned}$$

Theorem 21.11 (Functors from m-atlases to C^k atlases). Let $\mathbf{E}$ be a set of model spaces and $\mathbf{C}$ a set of C^k linear model spaces. Then $\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}$ is a functor from $\mathcal{Atl}(\mathbf{E}, \mathbf{C})$ to $\mathcal{Atl}^{C^k}(\pi_1[\mathbf{E}], \pi_1[\mathbf{C}])$.

Proof. Let $o^i \stackrel{\text{def}}{=} (A^i, (E^i, \mathcal{E}^i), (C^i, \mathcal{C}^i))$, $i \in [1, 3]$, be an object of $\mathcal{Atl}^{C^k}(\mathbf{E}, \mathbf{C})$ and

$m^i \stackrel{\text{def}}{=} ((f_0^i, f_1^i).o^i, o^{i+1})$, $i = 1, 2$, be a morphism from o^i to o^{i+1} .

$\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(m^1)$ is a morphism from $\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} o^1$ to $\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} o^2$:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(m^1) &= \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}((f_0^1, f_1^1), (A^1, E^1, C^1), (A^2, E^2, C^2)) &= \\ ((f_0^1, f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^2, \pi_1(E^2), \pi_1(C^2))) & \end{aligned}$$

$\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}$ maps identity functions to identity functions:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} \text{Id}_{(A^i, E^i, C^i)} &= \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}((\text{Id}_{E^i}, \text{Id}_{C^i}), (A^i, E^i, C^i), (A^i, E^i, C^i)) &= \\ ((\text{Id}_{\pi_1(E^i)}, \text{Id}_{\pi_1(C^i)}), (A^i, \pi_1(E^i), \pi_1(C^i)), (A^i, \pi_1(E^i), \pi_1(C^i))) &= \\ ((\text{Id}_{E^i}, \text{Id}_{C^i}), \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(A^i, E^i, C^i), \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(A^i, E^i, C^i)) &= \\ \text{Id}_{\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(A^i, E^i, C^i)} & \end{aligned}$$

$$\mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} m^2 \circ^A \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} m^1 = \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(m^2 \circ^A m^1):$$

$$\begin{aligned} m^2 \circ^A m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, E^1, C^1), (A^3, E^3, C^3)) \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}((A^i, E^i, C^i)) &= (A^i, \pi_1(E^i), \pi_1(C^i)) \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(m^i) &= ((f_0^i, f_1^i), (A^i, \pi_1(E^i), \pi_1(C^i)), (A^{i+1}, \pi_1(E^{i+1}), \pi_1(C^{i+1}))) \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} m^2 \circ^A \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k} m^1 &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^3, \pi_1(E^3), \pi_1(C^3))) \\ \mathcal{F}_{\mathbf{M}, \mathbf{C}^k}^{C^k}(m^2 \circ m^1) &= ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), (A^1, \pi_1(E^1), \pi_1(C^1)), (A^3, \pi_1(E^3), \pi_1(C^3))) \end{aligned}$$

□

22 Classic C^k -atlas morphisms and functors

{sec:Ck-ATLcmorph}

This section introduces an alternate definition of and taxonomy for morphisms between C^k -atlases, defines categories of C^k -atlases, defines functors among them and proves some basic results. It introduces the notion of classic C^k -atlas morphisms, which is equivalent to the conventional definitions for a manifold. The reader can safely skip it, but it does help to establish a context for other notions that are used in the exposition of local coordinate spaces.

22.1 Classic C^k -atlas morphisms

{sub:Ck-ATLcmorph}

This subsection introduces the notion of classic C^k -atlas morphisms, and proves some basic results.

22.1.1 Definitions of classic C^k -atlas morphisms

{def:Ck-ATLcmorph}

Definition 22.1 (Classic C^k -atlas morphisms). Let $E^i, i = 1, 2$, be a topological space, C^i be a linear space, A^i be a C^k -atlases of E^i in the coordinate space C^i and f be a continuous function.

f is a E^1 - E^2 classic C^k morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 , abbreviated as $\text{isAtl}_{\text{Ar}}^{\text{classic}, C^k}(A^1, E^1, C^1, A^2, E^2, C^2, f)$, iff for any $(U^i, V^i, \phi^i: U^i \xrightarrow{\sim} V^i) \in A^i, i = 1, 2$, with $I \stackrel{\text{def}}{=} U^1 \cap f^{-1}[U^2] \neq \emptyset, \phi^2 \circ f \circ \phi^{1-1}: \phi^1[I] \longrightarrow V^2$ is a C^k function.

Remark 22.2. Definitions of constrained, semistrict and strict classic C^k -atlas morphisms would be pointless, as any classic C^k -atlas morphism would be constrained and strict.

22.1.2 Proclamations on classic C^k -atlas morphisms

{lem:Ck-ATLcmorph}

Lemma 22.3 (Classic C^k -atlas morphisms). Let E be a set of topological spaces, C a set of C^k linear spaces, $E^i \in E, i = 1, 2, C^i \in C, A^i$ a C^k -atlas of E^i in the coordinate space C^i and $f: E^1 \longrightarrow E^2$ a function.

Let E^1 be an open subspace of E^2 . Then f is an E^1 - E^2 classic C^k -atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff f is a strict E^1 - E^2 classic m-atlas morphism of A^1 to A^2 in the coordinate spaces $\underset{C^k\text{-triv}}{C^1}, \underset{C^k\text{-triv}}{C^2}$.

f is an E^1 - E^2 classic C^k -atlas morphism of A^1 to A^2 in the coordinate spaces C^1, C^2 iff f is a strict E^1 - E^2 - $\underset{C^k\text{-op-}\mathcal{T}riv}{C} - \underset{C^k\text{-op-}\mathcal{T}riv}{C}$ classic m-atlas morphism of A^1 to A^2 in the coordinate spaces $\underset{C^k\text{-triv}}{C^1}, \underset{C^k\text{-triv}}{C^2}$.

23 C^k manifolds

{sec:ckman}

Conventionally a manifold is different from its atlases, but $\mathcal{Atl}_{\text{max}}^{C^k}(E, C)$ in definition 21.6 (Categories of C^k atlases) on page 132 encourages treating them on an

{def:Ck-ATLcat}

equal footing. All of the results for maximal atlases carry directly over to results for manifolds.

Remark 23.1. The variations of the LCS^{C^k} constructors in definition 23.6 (Functors from C^k manifolds to Local 2- $\emptyset$ coordinate spaces) on page 137 are very similar, as are the variations of the $\mathcal{F}_{\text{Man,LCS}}^{\text{C}^k}$ functors; likewise the variations of the $\mathcal{F}_{\text{LCS,Man}}^{\text{C}^k}$ functors in definition 23.9 (Functor from Local 2- $\emptyset$ coordinate spaces to C^k manifolds) on page 145 are very similar. {def:CktoLCS}
{def:LCS to Ck}

The corresponding results and proofs in theorem 23.8 (Functors from C^k manifolds to local 2- $\emptyset$ coordinate spaces) on page 141 are likewise very similar, as are the corresponding results and proofs in theorem 23.10 (Functor from Local 2- $\emptyset$ coordinate spaces to C^k manifolds) on page 145. {the:CktoLCS}
{the:LCS to Ck}
{def:man}

Definition 23.2 (C^k manifolds). Let E be a topological space, C a linear space and $\mathbf{A}$ a maximal¹⁷ C^k -atlas of E in the coordinate space C . Then $(E, C, \mathbf{A})$ is a C^k manifold.

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ be a set of linear spaces. Then

$$\text{Man}_{\text{Ob}}^{\text{C}^k}(E, C) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{max}}^{\text{C}^k}(E, C) \quad (23.1)$$

Remark 23.3. The manifold $(E, C, \mathbf{A})$ corresponds to the object $(\mathbf{A}, E, C)$. {def:manmorph}

Definition 23.4 (C^k manifold morphisms). Let $S^i \stackrel{\text{def}}{=} (E^i, C^i, \mathbf{A}^i)$, $i = 1, 2$, be C^k manifolds and $(f_0: E^1 \rightarrow E^2, f_1: C^1 \rightarrow C^2)$ be an E^1 - E^2 C^k -morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces C^1, C^2 . Then (f_0, f_1) is a C^k morphism of S^1 to S^2 .

Let E^i , $i = 1, 2$, be a topological space and C^i be a linear space. Then

$$\text{Man}_{\text{Ar}}^{\text{C}^k}(E^1, C^1, E^2, C^2) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{Ar}}^{\text{C}^k}(E^1, C^1, E^2, C^2) \quad (23.2)$$

Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then

$$\text{Man}_{\text{Ar}}^{\text{C}^k}(E, C) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{Ar}}^{\text{C}^k}(E, C) \quad (23.3)$$

$$\text{Man}^{\text{C}^k}(E, C) \stackrel{\text{def}}{=} \mathcal{Att}_{\text{max}}^{\text{C}^k}(E, C) \quad (23.4)$$

Theorem 23.5 (Categories of C^k manifolds). Let $\mathbf{E}$ be a set of topological spaces and $\mathbf{C}$ a set of linear spaces. Then $\text{Man}^{\text{C}^k}(E, C)$ is a category and the identity morphism of $(\mathbf{A}, E, C)$ is an identity morphism.

Proof. The result follows directly from definitions 23.2 and 23.4 above and lemma 21.7. {lem:Ck-ATLiscat}
($\mathcal{Att}^{\text{C}^k}(E, C)$ is a category) on page 133. □

Definition 23.6 (Functors from C^k manifolds to Local 2- $\emptyset$ coordinate spaces). Let $\mathbf{E}$ be a set of topological spaces, $\mathbf{C}$ be a set of C^k linear spaces, $\mathcal{E}$ and $\mathcal{E}$ model {def:CktoLCS}

¹⁷Requiring that the atlas be full would eliminate some pathologies.

categories, $\mathcal{C}$ and $\hat{\mathcal{C}}$ $\mathbb{C}^k$ linear model categories, $\mathcal{M}^{\text{def}} = (\mathcal{C}, \mathcal{C})$, $\hat{\mathcal{M}}^{\text{def}} = (\hat{\mathcal{C}}, \hat{\mathcal{C}})$,
 $\mathcal{M}_{\text{triv}}^{\text{def}} = \left(\begin{smallmatrix} E \\ \mathcal{T}riv \end{smallmatrix}, \begin{smallmatrix} C \\ \mathbb{C}^k\text{-}\mathcal{T}riv \end{smallmatrix} \right)$. $\mathcal{M}_{\text{op-}\mathcal{T}riv}^{\text{def}} = \left(\begin{smallmatrix} E \\ \mathcal{T}riv \end{smallmatrix}, \begin{smallmatrix} C \\ \mathbb{C}^k\text{-op-}\mathcal{T}riv \end{smallmatrix} \right)$. $\mathcal{M}_{\text{triv}}^{\text{full-cat}} \subseteq \mathcal{M}^{\text{full-cat}} \subseteq \hat{\mathcal{M}}$,
 $\mathcal{M}_{\text{op-}\mathcal{T}riv}^{\text{cat}} \subseteq \hat{\mathcal{M}}$.

Let $E^i \in E$, $i = 1, 2$, $C^i \in C$, $\mathbf{M}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}, \begin{smallmatrix} C^i \\ \mathbb{C}^k\text{-triv} \end{smallmatrix} \right)$, $\mathcal{M}_{\text{Sing}}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathbf{M}_0^i & \mathbf{M}_1^i \\ \text{Sing} & \mathbb{C}^k\text{-Sing} \end{smallmatrix} \right)$, A^i a

maximal $\mathbb{C}^k$ -atlas of E^i in C^i , $\mathcal{F}_1^{\text{min}}(A^i, E^i, C^i) \in \hat{\mathcal{C}}$,

$\mathcal{F}_2^{\text{min}}(A^i, E^i, C^i) \in \hat{\mathcal{C}}$, (E^i, C^i, A^i) a $\mathbb{C}^k$ manifold, $\mathbf{S}^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $\mathbf{M}_{\text{min}}^i \stackrel{\text{def}}{=}$

$\left(\mathcal{F}_1^{\text{min}}(A^i, E^i, C^i), \mathcal{F}_2^{\text{min}}(A^i, E^i, C^i) \right)$ and $\mathcal{M}_{\text{triv Sing}}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathbf{M}_0^i & \mathbf{M}_1^i \\ \text{triv} & \text{triv} \\ \text{Sing} & \mathbb{C}^k\text{-Sing} \end{smallmatrix} \right)$, Then

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k} \mathbf{S}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathcal{M}^i & \mathbf{M}^i & A^i & \emptyset & \emptyset \\ \text{Sing} \end{smallmatrix} \right) \quad (23.5)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}} \mathbf{S}^i \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}^i, A^i, \emptyset, \emptyset) \quad (23.6)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}_{\text{min}}} \mathbf{S}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathcal{M}^i & \mathbf{M}_{\text{min}}^i & A^i & \emptyset & \emptyset \\ \text{triv} & \text{min} \\ \text{Sing} \end{smallmatrix} \right) \quad (23.7)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \hat{\mathcal{M}}} \mathbf{S}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \hat{\mathcal{M}} & \mathbf{M}_{\text{min}}^i & A^i & \emptyset & \emptyset \end{smallmatrix} \right) \quad (23.8)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k}((f_0, f_1), \mathbf{S}^1, \mathbf{S}^2) \stackrel{\text{def}}{=} ((f_0, f_1), \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k} \mathbf{S}^1, \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k} \mathbf{S}^2) \quad (23.9)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}}((f_0, f_1), \mathbf{S}^1, \mathbf{S}^2) \stackrel{\text{def}}{=} ((f_0, f_1), \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}} \mathbf{S}^1, \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}} \mathbf{S}^2) \quad (23.10)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}_{\text{min}}}((f_0, f_1), \mathbf{S}^1, \mathbf{S}^2) \stackrel{\text{def}}{=} ((f_0, f_1), \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}_{\text{min}}} \mathbf{S}^1, \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \mathcal{M}_{\text{min}}} \mathbf{S}^2) \quad (23.11)$$

$$\mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \hat{\mathcal{M}}}((f_0, f_1), \mathbf{S}^1, \mathbf{S}^2) \stackrel{\text{def}}{=} ((f_0, f_1), \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \hat{\mathcal{M}}} \mathbf{S}^1, \mathcal{F}_{\text{Man,LCS}}^{\mathbb{C}^k, \hat{\mathcal{M}}} \mathbf{S}^2) \quad (23.12)$$

$$\text{LCS}_{\text{Ob}}^{\text{Ck}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \mid \right. \\ \left. E \in \mathbf{E} \wedge C \in \mathbf{C} \wedge \text{isAtI}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \right\}_{\max} \quad (23.13)$$

$$\text{LCS}_{\text{Ar}}^{\text{Ck}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}}((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\ \left. E^i \in \mathbf{E} \wedge C^i \in \mathbf{C} \wedge \text{isAtI}_{\text{Ar}}^{\text{Ck}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1) \right\} \quad (23.14)$$

$$\text{LCS}^{\text{Ck}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}^{\text{Ck}}(\mathbf{E}, \mathbf{C}), \text{LCS}_{\text{Ar}}^{\text{Ck}}(\mathbf{E}, \mathbf{C}), \circ^{\mathbf{A}}) \quad (23.15)$$

$$\text{LCS}_{\text{Ob}}^{\text{Ck}, \mathcal{M}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \mid \right. \\ \left. E \in \mathbf{E} \wedge C \in \mathbf{C} \wedge \text{isAtI}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \right\}_{\max} \quad (23.16)$$

$$\text{LCS}_{\text{Ar}}^{\text{Ck}, \mathcal{M}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}}((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\ \left. E^i \in \mathbf{E} \wedge C^i \in \mathbf{C} \wedge \text{isAtI}_{\text{Ar}}^{\text{Ck}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1) \right\} \quad (23.17)$$

$$\text{LCS}^{\text{Ck}, \mathcal{M}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}^{\text{Ck}, \mathcal{M}}(\mathbf{E}, \mathbf{C}), \text{LCS}_{\text{Ar}}^{\text{Ck}, \mathcal{M}}(\mathbf{E}, \mathbf{C}), \circ^{\mathbf{A}}) \quad (23.18)$$

$$\text{LCS}_{\text{Ob}}^{\text{Ck}, \min}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \min} \mathbf{S} \mid \right. \\ \left. \mathbf{S} \stackrel{\text{def}}{=} (\mathbf{A}, \mathbf{E}, \mathbf{C}) \wedge E \in \mathbf{E} \wedge C \in \mathbf{C} \wedge \text{isAtI}_{\text{Ob}}^{\text{Ck}}(\mathbf{A}, \mathbf{E}, \mathbf{C}) \right\}_{\max} \quad (23.19)$$

$$\text{LCS}_{\text{Ar}}^{\text{Ck}, \min}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \min}((f_0, f_1), (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \mid \right. \\ \left. E^i \in \mathbf{E} \wedge C^i \in \mathbf{C} \wedge \text{isAtI}_{\text{Ar}}^{\text{Ck}}(\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1, \mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2, f_0, f_1) \right\}_{\max} \quad (23.20)$$

$$\text{LCS}_{\min}^{\text{C}^k}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left(\text{LCS}_{\text{Ob}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}), \text{LCS}_{\text{Ar}}^{\text{C}^k}(\mathbf{E}, \mathbf{C}), \circ^A \right) \quad (23.21)$$

$$\text{LCS}_{\text{Ob}}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man, LCS}}^{\text{C}^k, \hat{\mathcal{M}}} \mathbf{S} \mid \mathbf{S} \stackrel{\text{def}}{=} (\mathbf{A}, E, C) \wedge E \in \mathbf{E} \wedge C \in \mathbf{C} \wedge \text{isAtl}_{\text{Ob}}^{\text{C}^k}(\mathbf{A}, E, C) \right\} \quad (23.22)$$

$$\text{LCS}_{\text{Ar}}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Man, LCS}}^{\text{C}^k, \hat{\mathcal{M}}}((f_0, f_1), (\mathbf{A}^1, E^1, C^1), (\mathbf{A}^2, E^2, C^2)) \mid E^i \in \mathbf{E} \wedge C^i \in \mathbf{C} \wedge \text{isAtl}_{\text{Ar}}^{\text{C}^k}(\mathbf{A}^1, E^1, C^1, \mathbf{A}^2, E^2, C^2, f_0, f_1) \right\} \quad (23.23)$$

$$\text{LCS}_{\min}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C}) \stackrel{\text{def}}{=} \left(\text{LCS}_{\text{Ob}}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C}), \text{LCS}_{\text{Ar}}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C}), \circ^A \right) \quad (23.24)$$

{lem:CktoLCSequiv}

Lemma 23.7 ((0)-equivalence of local 2- $\emptyset$ coordinate spaces). Let

1. $\mathbf{E}$ be a set of topological spaces, $\mathbf{C}$ be a set of C^k linear spaces
2. $\mathcal{E}$ and $\hat{\mathcal{E}}$ be model categories, $\mathcal{C}$ and $\hat{\mathcal{C}}$ be C^k linear model categories
3. $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{E}, \mathcal{C})$, $\hat{\mathcal{M}} \stackrel{\text{def}}{=} (\hat{\mathcal{E}}, \hat{\mathcal{C}})$
4. $\mathcal{M}_{\text{triv}} \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathbf{E} \\ \mathcal{T}riv \end{smallmatrix}, \begin{smallmatrix} \mathbf{C} \\ \text{C}^k\text{-}\mathcal{T}riv \end{smallmatrix} \right)$, $\mathcal{M}_{\text{op-}\mathcal{T}riv} \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathbf{E} \\ \mathcal{T}riv \end{smallmatrix}, \begin{smallmatrix} \mathbf{C} \\ \text{C}^k\text{-op-}\mathcal{T}riv \end{smallmatrix} \right)$
5. $\mathcal{M}_{\text{triv}}^{\text{full-cat}} \subseteq \mathcal{M}^{\text{full-cat}} \subseteq \hat{\mathcal{M}}, \mathcal{M}_{\text{op-}\mathcal{T}riv}^{\text{cat}} \subseteq \hat{\mathcal{M}}$
6. $E^i \in \mathbf{E}, i = 1, 2, C^i \in \mathbf{C}$
7. $\mathbf{M}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} E^i \\ \text{triv} \end{smallmatrix}, \begin{smallmatrix} C^i \\ \text{C}^k\text{-triv} \end{smallmatrix} \right)$, $\mathcal{M}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} \mathbf{M}_0^i \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} \mathbf{M}_1^i \\ \text{C}^k\text{-Sing} \end{smallmatrix} \right)$
8. $\mathbf{A}^i$ be a maximal C^k -atlas of E^i in C^i
9. $\mathcal{F}_1^{\text{C}^k}(\mathbf{A}^i, E^i, C^i) \in \hat{\mathcal{E}}, \mathcal{F}_2^{\text{C}^k}(\mathbf{A}^i, E^i, C^i) \in \hat{\mathcal{C}}$
10. $(E^i, C^i, \mathbf{A}^i)$ be a C^k manifold
11. $\mathbf{S}^i \stackrel{\text{def}}{=} (\mathbf{A}^i, E^i, C^i)$

$$12. \mathbf{M}_{\min}^i \stackrel{\text{def}}{=} (\mathcal{F}_1^{\text{C}^k}(\mathbf{A}^i, E^i, C^i), \mathcal{F}_2^{\text{C}^k}(\mathbf{A}^i, E^i, C^i)), \mathcal{M}_{\text{triv Sing}}^i \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{M}_0^i & \mathbf{M}_1^i \\ \text{triv Sing} & \text{triv C}^k\text{-Sing} \end{pmatrix}$$

$$13. \mathbf{f} \stackrel{\text{def}}{=} (f_0: E^1 \longrightarrow E^2, f_1: C^1 \longrightarrow C^2) \text{ and } \mathbf{g} \stackrel{\text{def}}{=} (g_0: E^1 \longrightarrow E^2, g_1: C^1 \longrightarrow C^2) \text{ be } E^1\text{-}E^2 \text{ C}^k \text{ morphisms of } A^1 \text{ to } A^2 \text{ in the coordinate spaces } C^1, C^2$$

$$\mathbf{f}: \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k} \mathbf{S}^1 \longrightarrow \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k} \mathbf{S}^2 \text{ is (0)-equivalent to } \mathbf{g}: \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k} \mathbf{S}^1 \longrightarrow \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k} \mathbf{S}^2 \text{ iff } \mathbf{f} \text{ is C}^k\text{-equivalent to } \mathbf{g}.$$

Proof. Both equivalences are defined as $f_0 = g_0$. □

Theorem 23.8 (Functors from C^k manifolds to local 2- $\emptyset$ coordinate spaces). Let $\mathbf{E}$ be a set of topological spaces, $\mathbf{C}$ be a set of C^k linear spaces, $\mathcal{E}$ and $\hat{\mathcal{E}}$ model categories, $\mathcal{C}$ and $\hat{\mathcal{C}}$ C^k linear model categories, $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{E}, \mathcal{C})$, $\hat{\mathcal{M}} \stackrel{\text{def}}{=} (\hat{\mathcal{E}}, \hat{\mathcal{C}})$, $\mathcal{M}_{\text{triv}} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{E}_{\text{triv}} & \mathbf{C}_{\text{C}^k\text{-}\mathcal{F}_{\text{triv}}} \end{pmatrix}$.

$$\mathcal{M}_{\text{op-}\mathcal{F}_{\text{triv}}} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{E}_{\text{op-}\mathcal{F}_{\text{triv}}} & \mathbf{C}_{\text{C}^k\text{-op-}\mathcal{F}_{\text{triv}}} \end{pmatrix}. \mathcal{M}_{\text{triv}}^{\text{cat}} \subseteq \mathcal{M}^{\text{cat}} \subseteq \hat{\mathcal{M}}_{\text{op-}\mathcal{F}_{\text{triv}}}^{\text{cat}}, \mathcal{M}_{\text{op-}\mathcal{F}_{\text{triv}}}^{\text{cat}} \subseteq \hat{\mathcal{M}}.$$

Then $\text{LCS}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$, $\text{LCS}^{\text{C}^k, \mathcal{M}}(\mathbf{E}, \mathbf{C})$, $\text{LCS}_{\min}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ and $\text{LCS}_{\min}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C})$ are categories and the identity morphism of $\mathbf{L}^\alpha \stackrel{\text{def}}{=} ((\mathcal{E}, \mathcal{C}), (E^\alpha, C^\alpha), A^\alpha, \emptyset, \emptyset)$ is that given in definition 16.6 (LCS_{Ar}) on page 119: $\text{Id}_{\mathbf{L}^\alpha} \stackrel{\text{def}}{=} ((\text{Id}_{E^\alpha}, \text{Id}_{C^\alpha}), \mathbf{L}^\alpha, \mathbf{L}^\alpha)$. {def:LCSmorphAr}

Proof. $\text{LCS}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$, $\text{LCS}^{\text{C}^k, \mathcal{M}}(\mathbf{E}, \mathbf{C})$, $\text{LCS}_{\min}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ and $\text{LCS}_{\min}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C})$ satisfy the definition of a category:

Composition: Let $m^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k}((f_0^i, f_1^i), \mathbf{S}^i, \mathbf{S}^{i+1})$ be a morphism of $\text{LCS}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$.

$m^2 \circ^A m^1 = \mathcal{F}_{\text{Man,LCS}}^{\text{C}^k}((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), \mathbf{S}^1, \mathbf{S}^3)$. By item 1 of lemma 13.13 (**Composition of m-atlas morphisms**) on page 79 $(f_0^2 \circ f_0^1, f_1^2 \circ f_1^1)$ is an m-atlas morphism. {lem:M-ATLmorphcomp} {M-morphcomp:morphcompmorph}

Associativity: Composition is associative by lemma 1.19 (**Tuple composition for labeled morphisms**) on page 13. {lem:labcomp}

Unit: $\text{Id}_{\mathbf{L}^\alpha}$ is an identity morphism by lemma 1.19. {lem:labcomp}

The same proof applies to $\text{LCS}^{\text{C}^k, \mathcal{M}}(\mathbf{E}, \mathbf{C})$, $\text{LCS}_{\min}^{\text{C}^k}(\mathbf{E}, \mathbf{C})$ and $\text{LCS}_{\min}^{\text{C}^k, \hat{\mathcal{M}}}(\mathbf{E}, \mathbf{C})$. □

Let $E^i \in \mathbf{E}$, $i \in [1, 3]$, $C^i \in \mathbf{C}$, $\mathbf{M}^i \stackrel{\text{def}}{=} \begin{pmatrix} E^i & C^i \\ \text{triv C}^k\text{-triv} & \end{pmatrix}$, $\mathcal{M}_{\text{Sing}}^i \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{M}_0^i & \mathbf{M}_1^i \\ \text{Sing} & \text{C}^k\text{-Sing} \end{pmatrix}$, A^i a

maximal C^k -atlas of E^i in C^i , $\mathcal{F}_1^{\text{C}^k}(\mathbf{A}^i, E^i, C^i) \in \hat{\mathcal{E}}$, $\mathcal{F}_2^{\text{C}^k}(\mathbf{A}^i, E^i, C^i) \in \hat{\mathcal{C}}$, (E^i, C^i, A^i)

a C^k manifold, $\mathbf{S}^i \stackrel{\text{def}}{=} (A^i, E^i, C^i)$, $\mathbf{M}^i \stackrel{\text{def}}{=} (\mathcal{F}_{\min}^{C^k}(A^i, E^i, C^i), \mathcal{F}_2^{\min}(A^i, E^i, C^i))$,
 $\mathcal{M}_{\text{triv}}^i \stackrel{\text{def}}{=} (\mathcal{M}_{\text{triv}}^i, \mathcal{M}_{\text{triv}}^i)$, $L^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Man,LCS}}^{C^k} \mathbf{S}^i = ((\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), (\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), A^i, \emptyset, \emptyset)$,
 $L^i, \mathcal{M} \stackrel{\text{def}}{=} \mathcal{F}_{\text{Man,LCS}}^{C^k, \mathcal{M}} \mathbf{S}^i = (\mathcal{M}, (\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), A^i, \emptyset, \emptyset)$, $L^i \stackrel{\text{def}}{=} (\mathcal{M}_{\text{triv}}^i, (\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), A^i, \emptyset, \emptyset)$,
 $L^i, \hat{\mathcal{M}} \stackrel{\text{def}}{=} (\hat{\mathcal{M}}, (\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), A^i, \emptyset, \emptyset)$, $f^i \stackrel{\text{def}}{=} (f_0^i, f_1^i)$, $f_0^i: E^i \longrightarrow E^{i+1}$ continuous
and $f_1^i: C^i \longrightarrow C^{i+1}$ C^k .
 $\mathcal{F}_{\text{Man,LCS}}^{C^k} \mathbf{S}^i$ is a local $\mathcal{M}^i$ - $\emptyset$ coordinate space and $\mathcal{F}_{\text{Man,LCS}}^{C^k, \mathcal{M}} \mathbf{S}^i$ is a local $\mathcal{M}$ - $\emptyset$ coordinate space.

Proof. $\mathcal{F}_{\text{Man,LCS}}^{C^k, \mathcal{M}} \mathbf{S}^i$ satisfies the conditions of definition 15.1 (Local $\mathbf{M}-\Sigma$ and $\mathcal{M}-\Sigma$ coordinate spaces) on page 114: {def:LCS}

1. Model categories: $\mathcal{E}_{\text{triv}}^i$ and $\mathcal{C}_{\text{triv}}^i$ are model spaces
2. $(\mathcal{M}, (\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), \emptyset, \emptyset)$ satisfies the conditions of definition 10.5 (Prestructures) on page 45 {def:pre}
3. Maximal m-atlas: A^i is a maximal C^k atlas by hypothesis and a maximal atlas of $\mathcal{E}_{\text{triv}}^1$ in the coordinate space $\mathcal{C}_{\text{triv}}^i$ by construction.
4. Constraint functions: There are no adjunct functions so there are no constraint functions.

$\mathcal{M}_{\text{sing}}^i \stackrel{\text{cat}}{\subseteq} \mathcal{M}$, so $\mathcal{F}_{\text{Man,LCS}}^{C^k, \mathcal{M}} \mathbf{S}^i$ is a local $\mathcal{M}$ - $\emptyset$ coordinate space by lemma 15.3 (Local $\ast$ - Σ coordinate spaces) on page 114. {lem:LCS} □

$\mathcal{F}_{\text{Man,LCS}}^{C^k} \mathbf{S}^i$ is a local $\mathcal{M}_{\text{triv}}^i$ - $\emptyset$ coordinate space and $\mathcal{F}_{\text{Man,LCS}}^{C^k, \hat{\mathcal{M}}} \mathbf{S}^i$ is a local $\hat{\mathcal{M}}$ - $\emptyset$ coordinate space.

Proof. $\mathcal{F}_{\text{Man,LCS}}^{C^k} \mathbf{S}^i$ satisfies the conditions of definition 15.1 (Local $\mathbf{M}-\Sigma$ and $\mathcal{M}-\Sigma$ coordinate spaces) on page 114: {def:LCS}

1. Model categories: $\mathcal{E}_{\text{triv}}^1$ and $\mathcal{C}_{\text{triv}}^i$ are model spaces
2. $((\mathcal{E}_{\text{triv}}^i, \mathcal{C}_{\text{triv}}^i), \emptyset, \emptyset)$ satisfies the conditions of definition 10.5 (Prestructures) on page 45 {def:pre}

3. Maximal m-atlas: A^i is a maximal C^k atlas by hypothesis and a maximal atlas of E^1 in the coordinate space $\underset{\text{triv}}{C^k\text{-op-}\mathcal{T}riv}$ by construction.
4. Constraint functions: There are no adjunct functions so there are no constraint functions.

$\mathcal{M}_{\text{triv}}^i \xrightarrow{\text{cat}} \hat{\mathcal{M}}$, so $\mathcal{F}_{\text{Man,LCS}}^{C^k, \hat{\mathcal{M}}} S^i$ is a local $\hat{\mathcal{M}}$ - $\emptyset$ coordinate space by lemma 15.3. $\square$

{lem:LCS}

(f_0^1, f_1^1) is a morphism from L^1 to L^2 and a strict morphism from $L^{1, \mathcal{M}}$ to $L^{2, \mathcal{M}}$.

Proof.

Prestructure morphism: f^1 Σ -commutes with $\emptyset, \emptyset$. f_0^1 is continuous, hence a morphism of $\mathcal{C}$. f_1^1 is C^k , hence a morphism of $\mathcal{C}$.

m-atlas morphism: (f_0^1, f_1^1) is a $\underset{\text{triv}}{E^1 - E^2}$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\underset{C^k\text{-triv}}{C^1}, \underset{C^k\text{-triv}}{C^i}$.

(f_0^1, f_1^1) is a strict $\underset{\text{Sing}}{E^1 - E^2} - \underset{\text{Sing}}{E^1 - E^2} - \underset{C^k\text{-triv}}{C^1} - \underset{C^k\text{-triv}}{C^1}$ m-atlas morphism of A^1

to A^2 in the coordinate spaces $\underset{C^k\text{-triv}}{C^1}, \underset{C^k\text{-triv}}{C^i}$ by lemma 21.3 (C^k -atlas morphisms) on page 131. $\square$

{lem:Ck-ATLmorph}

$\mathcal{F}_{\text{Man,LCS}}^{C^k}((f_0^1, f_1^1), S^1, S^2)$ is a morphism from $\underset{\text{triv}}{L^1}$ to $\underset{\text{triv}}{L^{2, \mathcal{M}}}$ and $\mathcal{F}_{\text{Man,LCS}}^{C^k}((f_0^1, f_1^1), S^1, S^2)$ is a strict morphism from $\underset{\text{triv}}{L^{1, \mathcal{M}}}$ to $\underset{\text{triv}}{L^{2, \mathcal{M}}}$.

Proof.

Prestructure morphism: (f_0^1, f_1^1) Σ -commutes with $\emptyset, \emptyset$.

m-atlas morphism: (f_0^1, f_1^1) is an m-atlas morphism from $(\underset{\text{triv}}{E^1}, \underset{C^k\text{-op-}\mathcal{T}riv}{C^1})$ to $(\underset{\text{triv}}{E^2}, \underset{C^k\text{-op-}\mathcal{T}riv}{C^2})$.

$\square$

$\mathcal{F}_{\text{Man,LCS}}^{C^k, \mathcal{M}}$ is a functor from $\text{Man}^{C^k}(E, C)$ to $\text{LCS}^{C^k, \mathcal{M}}(E, C)$.

Proof.

Preservation of endpoints:

$$\begin{aligned}\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2) &= ((f_0^1, f_1^1), L^i, L^i) \\ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \mathbf{S}^i &= \left(\begin{pmatrix} E^i & C^i \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, A^i, \emptyset, \emptyset \right) \\ &= L^i\end{aligned}$$

Composition: $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \left(((f_0^2, f_1^2), \mathbf{S}^2, \mathbf{S}^3) \overset{A}{\circ} ((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2) \right) =$
 $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \left(((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), \mathbf{S}^1, \mathbf{S}^3) \right) =$
 $((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), L^1, L^3) =$
 $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^2, f_1^2), \mathbf{S}^2, \mathbf{S}^3) \overset{A}{\circ} \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2)$

Identity: $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}(\text{Id}_{\mathbf{S}^i}) = \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}(\text{Id}_{E^i}, \text{Id}_{C^i}, \mathbf{S}^i, \mathbf{S}^i) =$
 $\left(\begin{pmatrix} \text{Id}_{E^i} & \text{Id}_{C^i} \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, L^1, L^1 \right) =$

□

$\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}$ is a functor from $\text{Man}^{\text{Ck}}(E, C)$ to $\text{LCS}_{\text{min}}^{\text{Ck}}(E, C)$.

Proof.

Preservation of endpoints:

$$\begin{aligned}\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2) &= ((f_0^1, f_1^1), L^i, L^i) \\ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \mathbf{S}^i &= \left(\begin{pmatrix} E^i & C^i \\ \text{triv} & \text{Ck-op-triv} \end{pmatrix}, A^i, \emptyset, \emptyset \right) \\ &= L^i\end{aligned}$$

Composition: $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \left(((f_0^2, f_1^2), \mathbf{S}^2, \mathbf{S}^3) \overset{A}{\circ} ((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2) \right) =$
 $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}} \left(((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), \mathbf{S}^1, \mathbf{S}^3) \right) =$
 $((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), L^1, L^3) =$
 $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^2, f_1^2), \mathbf{S}^2, \mathbf{S}^3) \overset{A}{\circ} \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2)$

Identity: $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}(\text{Id}_{\mathbf{S}^i}) = \mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}(\text{Id}_{E^i}, \text{Id}_{C^i}, \mathbf{S}^i, \mathbf{S}^i) =$
 $\left(\begin{pmatrix} \text{Id}_{E^i} & \text{Id}_{C^i} \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, \begin{pmatrix} E^i & C^i \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, A^i, \emptyset, \emptyset, \begin{pmatrix} E^i & C^i \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, A^i, \emptyset, \emptyset \right) =$
 $\text{Id}_{\left(\begin{pmatrix} E^i & C^i \\ \text{triv} & \text{Ck-triv} \end{pmatrix}, A^i, \emptyset, \emptyset \right)} \stackrel{\text{def}}{=} \text{Id}_{\mathcal{F}_{\text{Man,LCS}}^{\text{Ck},\mathcal{M}}(\mathbf{S}^i)}$

□

{def:LCS to Ck}

Definition 23.9 (Functor from Local 2- $\emptyset$ coordinate spaces to C^k manifolds). Let $\mathbf{C}^i \stackrel{\text{def}}{=} (C^i, \mathcal{C}^i)$, $i = 1, 2$, be a C^k linear model space, E^i a model space, $\mathcal{M}_0^i$ a model category, $\mathcal{M}_1^i$ a C^k linear model category, $\mathcal{M}^i \stackrel{\text{def}}{=} (\mathcal{M}_0^i, \mathcal{M}_1^i)$, $\mathbf{M}^i \stackrel{\text{def}}{=} (E^i, \mathbf{C}^i)$, $\mathbf{M}^i \in \mathcal{M}^i$, $\mathbf{A}^i$ maximal m-atlases of E^i in $\mathbf{C}^i$, $f_0: E^1 \longrightarrow E^2$, $f_1: \mathbf{C}^1 \longrightarrow \mathbf{C}^2$, $f \stackrel{\text{def}}{=} (f_0, f_1)$ and $\mathbf{L}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \emptyset, \emptyset)$ a local 2- $\emptyset$ coordinate spaces. Then

$$\mathcal{F}_{\text{LCS,Man}}^{C^k} \mathbf{L}^i \stackrel{\text{def}}{=} (E^i, \mathbf{C}^i, \mathbf{A}^i) \quad (23.25)$$

$$\mathcal{F}_{\text{LCS,Man}}^{C^k} (f, \mathbf{L}^1, \mathbf{L}^2) \stackrel{\text{def}}{=} (f, (E^1, \mathbf{C}^1, \mathbf{A}^1), (E^2, \mathbf{C}^2, \mathbf{A}^2)) \quad (23.26)$$

{the:LCS to Ck}

Theorem 23.10 (Functor from Local 2- $\emptyset$ coordinate spaces to C^k manifolds). Let $\mathbf{C}^i \stackrel{\text{def}}{=} (C^i, \mathcal{C}^i)$, $i = 1, 2$, be C^k linear model spaces, $E^i \stackrel{\text{def}}{=} (E^i, \mathcal{C}^i)$ model spaces, $\mathbf{A}^i$ maximal C^k -atlases of E^i in $\mathbf{C}^i$, $\mathbf{M}^i \stackrel{\text{def}}{=} (E^i, \mathbf{C}^i)$, $\mathcal{M}^i \stackrel{\text{def}}{=} \mathbf{M}^i$, $f_0^i: E^i \longrightarrow E^{i+1}$, $f_1^i: \mathbf{C}^i \longrightarrow \mathbf{C}^{i+1}$ and $\mathbf{L}^i \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \emptyset, \emptyset)$ a local 2- $\emptyset$ coordinate space. Then

1. $\mathcal{F}_{\text{LCS,Man}}^{C^k} \mathbf{L}^i \stackrel{\text{def}}{=} (E^i, \mathbf{C}^i, \mathbf{A}^i)$ is a C^k manifold.

Proof. $(E^i, \mathbf{C}^i, \mathbf{A}^i)$ satisfies the conditions of definition 23.2 (**C^k manifolds**) on page 137: (E^i) is a topological space, $\mathbf{C}^i$ is a C^k linear space and $\mathbf{A}^i$ is a maximal C^k -atlas E^i in $\mathbf{C}^i$. □

2. $\mathcal{F}_{\text{LCS,Man}}^{C^k}$ is a functor from $\text{LCS}^{C^k, \mathcal{M}}(E, \mathbf{C})$ to $\text{Man}^{C^k}(E, \mathbf{C})$.

Proof. Let $\mathbf{S}^i = (\mathbf{A}^i, E^i, \mathbf{C}^i)$.

Preservation of endpoints:

$$\begin{aligned} \mathcal{F}_{\text{LCS,Man}}^{C^k} (f_0^i, f_1^i), \mathbf{L}^i, \mathbf{L}^{i+1} &= (f_0^i, f_1^i), \mathbf{S}^i, \mathbf{S}^{i+1} \\ \mathcal{F}_{\text{LCS,Man}}^{C^k} (\mathbf{L}^i) &= \mathbf{S}^i \end{aligned}$$

$$\begin{aligned} \textbf{Composition: } \mathcal{F}_{\text{LCS,Man}}^{C^k} \left(((f_0^2, f_1^2), \mathbf{L}^2, \mathbf{L}^3) \stackrel{A}{\circ} ((f_0^1, f_1^1), \mathbf{L}^1, \mathbf{L}^2) \right) &= \\ \mathcal{F}_{\text{LCS,Man}}^{C^k} \left(((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), \mathbf{L}^1, \mathbf{L}^3) \right) &= \\ ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), \mathbf{S}^1, \mathbf{S}^3) &= \\ ((f_0^2, f_1^2), \mathbf{S}^2, \mathbf{S}^3) \stackrel{A}{\circ} ((f_0^1, f_1^1), \mathbf{S}^1, \mathbf{S}^2) &= \\ \mathcal{F}_{\text{LCS,Man}}^{C^k} \left(((f_0^2, f_1^2), \mathbf{L}^2, \mathbf{L}^3) \right) \stackrel{A}{\circ} \mathcal{F}_{\text{LCS,Man}}^{C^k} \left(((f_0^1, f_1^1), \mathbf{L}^1, \mathbf{L}^2) \right) & \end{aligned}$$

$$\textbf{Identity: } \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}(\text{Id}_{L^i}) = \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}(\text{Id}_{E^i}, \text{Id}_{C^i}), L^i, L^i) = (\text{Id}_{E^i}, \text{Id}_{C^i}), S^i, S^i) = \text{Id}_{\mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}}(L^i)$$

□

3. $\mathcal{F}_{\text{LCS,Man}}^{\text{Ck}} \circ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}}$ is the identity functor on $\mathcal{Att}^{\text{Ck}}(E, C)$.

$$\textit{Proof. } \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}} \circ \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}}(S^i) = \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}(L^i) = S^i$$

□

4. $\mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}} \circ \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}$ is the identity functor on $\text{LCS}^{\text{Ck}}(E, C)$.

$$\textit{Proof. } \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}} \circ \mathcal{F}_{\text{LCS,Man}}^{\text{Ck}}(L^i) = \mathcal{F}_{\text{Man,LCS}}^{\text{Ck}, \mathcal{M}}(S^i) = L^i$$

□

Part X

Equivalence of fiber bundles

{part:bun}

For fiber bundles¹⁸, the adjunct spaces are the base space $X = (X, \mathcal{X})$, the fiber $Y = (Y, \mathcal{Y})$ and the group G ; the category of the coordinate space is the category of Cartesian products $\{U \times Y \mid U \in \mathcal{X}\}$ of model neighborhoods in the base space with the entire fiber, with morphisms $t: U \times Y \longrightarrow U \times Y$ that preserve the fibers, i.e., $\pi_1 \circ t = \pi_1$, and are generated by the group action on the fiber (Equation (25.1)).

The sole adjunct functions are the projection $\pi: E \rightarrow X$, the group operation and the group action on the fiber.

Remark 23.11. While this presentation makes the projection π explicit, there is a less traditional approach that derives the projection and is in some ways simpler.

This part of the paper defines bundle atlases, fiber bundles, local coordinate spaces equivalent to fiber bundles, categories of them and functors, and gives basic results.

Definition 23.12 (Trivial group category of groups). Let G be a set of topological groups. The trivial group category of G , abbreviated $\mathbf{G}_{\text{group-}\mathcal{T}riv}$, is the category of all continuous homomorphisms between groups of G . By abuse of language it will be shortened to $\mathbf{G}_{\mathcal{T}riv}$ when the meaning is clear from context.

Definition 23.13 (Group actions). Let Y be a topological space, G a topological group, ρ an effective group action of G on Y , $y \in Y$ and $g \in G$. Then $y \star g \stackrel{\text{def}}{=} \rho(y, g)$.

¹⁸The literature has several definitions of fiber bundle. This paper uses one chosen for clarity of exposition. It differs from [6, p. 8] in that, e.g., it uses the machinery of maximal atlases rather than equivalence classes of coordinate bundles, the nomenclature differs in several minor regards.

{TopFib}

Let X be a topological space and $x \in X$. Then $(x, y) \star g \stackrel{\text{def}}{=} (x, y \star g)$.

A $\star$ with a subscript, superset, underset or overset refers to the group action ρ with the same subscript, superset, underset or overset.

Remark 23.14. This notation is only used when it is clear from context what the group action is.

{def:ProtoBun}

Definition 23.15 (Protobundles). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$, where E, X and Y are topological spaces, G a topological group, $\pi: E \rightarrow X$ a continuous surjection and ρ an effective group action of G on Y . Then $\mathbf{B}$ is a protobundle.

Remark 23.16. While this definition does not itself require E to have a local product structure nor require π to have the Covering Homotopy Property, only those protobundles having an atlas are of interest, and for them [definition 26.1 \(Bundle atlases\)](#) on page 161 imposes additional constraints.

{def:Bun-Atl}

{def:Xpimod}

Definition 23.17 (X - π -fine grained model spaces). Let $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ be a model space, X be a topological space and the projection $\pi: E \rightarrow X$ be a continuous function. $\mathbf{E}$ is a X - π -fine grained model space iff every $U \in \mathbf{E}^{\text{Ob}}$ is of the form $\pi^{-1}[V]$ for $V \subseteq X$ open and for every open sets $V' \subseteq V \subseteq X$, if $\pi^{-1}[V] \in \mathbf{E}^{\text{Ob}}$ then $\pi^{-1}[V'] \in \mathbf{E}^{\text{Ob}}$.

Remark 23.18. The spaces used for fiber bundles have a local product structure with a surjective projection, although neither is required by this definition.

{def:Xpitriv}

Definition 23.19 (X - π -trivial model spaces). Let E and X be topological spaces and the projection $\pi: E \rightarrow X$ be a continuous function.

Let $\mathcal{E}$ be the category whose objects are all $\pi^{-1}[V]$ for $V \subseteq X$ open and whose morphisms are all continuous $f: U^1 \rightarrow U^2$ that preserve fibers, i.e.,

$$\left(\bigvee_{x,y \in U^1} \right) \pi(x) = \pi(y) \vdash \pi(f(x)) = \pi(f(y))$$

Then the X - π -trivial model space of E is $\mathbf{E} \stackrel{\text{def}}{=} (E, \mathcal{E})$ and $\mathbf{E}_{X-\pi-\text{triv}}$ is a X - π -trivial model space.

{lem:Xpitriv}

Lemma 23.20 (X - π -trivial model spaces). Let E and X be topological spaces and the projection $\pi: E \rightarrow X$ be a continuous function. Then $\mathbf{E}_{X-\pi-\text{triv}}$ is a X - π -fine grained model space.

Proof. 1. Every $U \in \mathbf{E}^{\text{Ob}}$ is of the form $\pi^{-1}[V]$ for $V \subseteq X$ open and for every open sets by [definition 23.19](#) above.

{def:Xpitriv}

2. For every open $V' \subset X$, $\pi^{-1}[V'] \in \mathbf{E}_{X-\pi-\text{triv}}^{\text{Ob}}$.

□

24 G -rho model spaces

`def:Grhomod {sec:grhomod}`

Definition 24.1 (G - ρ -model spaces). Let $\mathbf{XY} \stackrel{\text{def}}{=} (X \times Y, \mathcal{XY})$ be a model space, G a topological group and ρ an effective group action of G on Y such that the objects of $\mathcal{XY}$ are products of open sets with Y and the morphisms are fiber-preserving automorphisms generated by the group action, i.e.,

$$\left(\bigvee_{U \in \mathcal{XY}}^{\text{Ob}} \right) \left(\exists_{V \in \mathcal{X}_{\text{op}}} \right) U = V \times Y \quad (24.1)$$

$$\left(\bigvee_{t \in \mathcal{XY}}^{\text{Ar}} : V \times Y \xrightarrow{\sim} V \times Y \right) \left(\exists_{g \in G^V} \right) \left(\bigvee_{(x,y) \in V \times Y} t(x,y) = (x, y \star g(x)) \right) \quad (24.2) \quad \{\text{eq:fGrho}\}$$

Then $\mathbf{XY}$ is a G - ρ model space of $X \times Y$, abbreviated $\text{isG}\rho(\mathbf{XY}, Y, G, \rho)$,

$G_{\mathbf{XY},t} \stackrel{\text{def}}{=} \text{range}(g)$ is the set of group elements associated with t and

$G_{\mathbf{XY}} \stackrel{\text{def}}{=} \bigcup_{t \in \mathcal{XY}}^{\text{Ar}} G_{\mathbf{XY},t}$ is the set of group elements associated with $\mathbf{XY}$ morphisms.

$\mathbf{XY}$ is also a fine grained G - ρ model space of $X \times Y$ iff for every $U \times Y \in \mathcal{XY}^{\text{Ob}}$ and every open $U' \subseteq U$, the product $U' \times Y$ is an object of $\mathcal{XY}$.

`{lem:Grhomod}`

Lemma 24.2 (G - ρ -model spaces). Let $E^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i)$, $i = 1, 2$, be a model space, $\mathbf{XY}^i$ be a G^i - ρ^i model space of $X^i \times Y^i$ and $A^i \stackrel{\text{def}}{=} \{(U^\alpha, V^\alpha \times Y^i, \phi^\alpha) | \alpha < A^i\}$ be an m-atlas of E^i in the coordinate space $\mathbf{XY}^i$.

1. Let $f \in \mathcal{XY}^i : V \times Y \xrightarrow{\sim} V \times Y$. The function g in eq. (24.2) is unique. `{eq:fGrho}`

Proof. The group action is effective. □

$G_{\mathbf{XY}^i}^i$ is unique.

Proof. The function g is unique. □

2. Let $(U^{i,j}, V^{i,j} \times Y^i, \phi^{i,j}) \in A^i$, $j = 1, 2$, with $I \stackrel{\text{def}}{=} U^{i,1} \cap U^{i,2} \neq \emptyset$. Then $\pi_1 \circ \phi^{i,1} = \pi_1 \circ \phi^{i,2}$ on I .

Proof. By definition 24.1 above, the transition function is of the form $t(x, y) = (x, y \star^i g(x))$. Then $\pi_1 \circ t(x, y) = \pi_1 \circ (x, y \star^i g(x)) = x$. □

`{def:Grhomod}`

3. Let $\pi^{i,\alpha} \stackrel{\text{def}}{=} \pi_1 \circ \phi^{i,\alpha}$, $\alpha < A^i$, and let $\pi^i : E^i \rightarrow X$ be the function agreeing with each $\pi^{i,\alpha}$ on $U^{i,\alpha}$. Then π^i is continuous and $\pi^{i,\alpha-1}[V^{i,\alpha} \times Y^i] = U^{i,\alpha}$.

Proof. π^i is well defined by the above result. Let $W \subseteq X^i \times Y^i$ be open. Each $W \cap V^{i,\alpha} \times Y^i$ is open, so each $\pi^{i,\alpha^{-1}}[W \cap V^{i,\alpha} \times Y^i]$ is open, Then $\pi^{i-1}[W \cap V^{i,\alpha} \times Y^i] = \bigcup_{\alpha < A} \pi^{\alpha-1}[W \cap V^\alpha \times Y^i]$ is open. $\square$

4. $\mathbf{A}^i$ is an m-atlas of $\frac{E^i}{X^i - \pi^i - \text{triv}}$ in the coordinate space $\mathbf{XY}^i$.

Proof. Let $(U^{i,j}, V^{i,j} \times Y^i, \phi^{i,j}) \in \mathbf{A}^i$, $j = 1, 2$. Then $(U^{i,j} = \pi^{i-1}[V^{i,j}]$ by the above, and hence is a neighborhood of $\frac{E^i}{X^i - \pi^i - \text{triv}}$.

The compatibility constraints are the same in both cases.

If $V^{i,j} \subseteq V^{i,j}$ is open then $\pi^{i,j-1}[V^{i,j}]$ is a neighborhood of $\frac{E^i}{X^i - \pi^i - \text{triv}}$ by definition 23.19 (X - π -trivial model spaces) on page 147. $\square$

5. If $\mathbf{XY}^i$ is a fine grained G^i - ρ^i model space of $X^i \times Y^i$ then $\mathbf{E}^i$ is an X^i - π_1 -fine grained model space.

Proof. $\mathbf{E}^i$ satisfies the conditions of definition 23.17 (X - π -fine grained model spaces) on page 147: $\{\text{def: } X\text{pimod}\}$

(a) For $V \subseteq X^i$, $\pi_1^{i-1}[V] = V \times Y^i$.

(b) If $\pi_1^{i-1}[V] = V \times Y^i$ is a model neighborhood, V' is open and $V' \subseteq V$ then $\pi_1^{i-1}[V'] = V' \times Y^i$ is a model neighborhood, $\square$

6. Let $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ be a $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{XY}^1, \mathbf{XY}^2$. Then $\mathbf{f}$ is a $\frac{E^1}{X^1 - \pi^1 - \text{triv}} - \frac{E^2}{X^2 - \pi^2 - \text{triv}}$ m-atlas morphism of $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces $\mathbf{XY}^1, \mathbf{XY}^2$.

Proof. $\mathbf{f}_1$ is a model function by hypothesis, hence continuous.

$\mathbf{f}_0: \mathbf{E}^1 \longrightarrow \mathbf{E}^2$ is a model function by hypothesis. Let U^2 be a model neighborhood of $\frac{E^2}{X^2 - \pi^2 - \text{triv}}$ and $U^1 \stackrel{\text{def}}{=} f_0^{-1}[U^2]$. Then U^2 is of the form $\pi^{2-1}[V^2]$

with V^2 open. Let $V^1 \stackrel{\text{def}}{=} f_1^{-1}[V^2]$. Since $\mathbf{f}_1$ is continuous, V^1 is open and thus $\pi^{1-1}[V^1] = U^1$ is open and a model neighborhood of $\frac{E^1}{X^1 - \pi^1 - \text{triv}}$.

Let $(U^i, V^i \times Y^i, \phi^i) \in \mathbf{A}^i$, $i = 1, 2$, and $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$. Each V^i is open, thus $U^i = \pi^{i-1}[V^i]$ is a model neighborhood of $\frac{E^i}{X^i - \pi^i - \text{triv}}$.

$$\begin{array}{ccc}
((x^1, y^1), g^1) & \xrightarrow{f_C \times f_G} & ((x^2, y^2), g^2) \\
\downarrow \star^1 & & \downarrow \star^2 \\
(x^1, y^1 \star g^1) & \xrightarrow{f_C} & (x^2, y^2 \star g^2)
\end{array}$$

Figure 14: Preserving group action

{fig:Grho}

Let $u^1 \in I$, $(U'^1, V'^1 \times Y^i, \phi^1)$, $U'^2 \subseteq U^2$, $(U'^2, \hat{V}'^2 \times Y^2, \phi'^2)$ be as in definition 13.1 (M-atlas morphisms for model spaces) on page 61. By the above, $(U'^1, V'^1 \times Y^i, \phi^1)$ and $U'^2 \subseteq U^2$, $(U'^2, \hat{V}'^2 \times Y^2, \phi'^2)$ are M-charts of $\begin{smallmatrix} E^1 \\ X^1 - \pi^1 - \text{triv} \end{smallmatrix}$ $\begin{smallmatrix} E^2 \\ X^2 - \pi^2 - \text{triv} \end{smallmatrix}$ in the coordinate space $\mathbf{XY}^1$ ($\mathbf{XY}^2$). The commutation relations are not changed by the change of base space. $\square$

{def:M-ATLmorph}

Definition 24.3 (Morphisms of G - ρ -model spaces). Let X^i, Y^i , $i = 1, 2$, be topological spaces, G^i a topological group, ρ^i an effective group action on Y^i and

{def:Grhomorph}

$\mathbf{XY}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{XY}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$. Then a model function $f_C: \mathbf{XY}^1 \rightarrow \mathbf{XY}^2$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$, abbreviated is $G\rho\text{morph}(\mathbf{XY}^1, G^1, \rho^1, \mathbf{XY}^2, G^2, \rho^2, f_C)$ iff it preserves the group action, i.e., there is a continuous homomorphism $f_G: G^1 \rightarrow G^2$ such that fig. 14 (Preserving group action) is commutative, i.e., eq. (24.3) below holds.

{fig:Grho}

$$\left(\exists_{f_G: G^1 \rightarrow G^2} \right) \left(\bigvee_{\substack{(x,y) \in X^1 \times Y^1 \\ g \in G^1}} \right) f_C((x, y) \star^1 g) = f_C((x, y)) \star^2 f_G(g) \quad \text{eq. (24.3)}$$

{lem:Grhomorph}

Lemma 24.4 (Morphisms of G - ρ -model spaces). Let X^i, Y^i , $i = 1, 2, 3$, be topological spaces, G^i a topological group, ρ^i an effective group action on Y^i , $\mathbf{XY}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{XY}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$ and $f_C^i: X^i \times Y^i \rightarrow X^{i+1} \times Y^{i+1}$ a G^i - G^{i+1} - ρ^i - ρ^{i+1} morphism of $X^i \times Y^i$ to $X^{i+1} \times Y^{i+1}$.

{grhomorph:fgunique}

1. The function f_G in eq. (24.3) is unique.

{eq:Grho}

Proof. The group action is effective. $\square$

{grhomorph:fiber}

2. f_C^i preserves fibers, i.e., $\pi_1(f_C^i(x, y)) = \pi_1(f_C^i(x, y'))$ for x in X^i and y, y' in Y^i ,

Proof. Since ρ^i is effective, there is a $g \in G^i$ such that $y' = y \star^i g$ and thus by definition 24.3

{def:Grhomorph}

$$\begin{aligned}\pi_1(f_C^i(x, y')) &= \\ \pi_1(f_C^i(x, y \star^i g)) &= \\ \pi_1(f_C^i(x, y) \star^{i+1} f_G^i(g)) &= \\ \pi_1(f_C^i(x, y)) &\end{aligned} \quad (24.4)$$

□

{grhomorph:fxunique}

3. There exists a unique function $f_X^i: X^i \rightarrow X^{i+1}$ such that $f_X^i \circ \pi_1 = \pi_1 \circ f_C^i$.

Proof. Define $f_X^i(x) = \pi_1(f_C^i(x, y))$, where y is an arbitrary point of Y^i . It does not depend on the choice of y because f_C^i preserves fibers. □

f_X^i is continuous.

Proof. Let $V^{i+1} \subseteq X$ be open. Then $(\pi_1 \circ f_C^i)^{-1}[V^{i+1}]$ is open and is of the form $V^i \times Y$ with $V^i \subseteq X^i$ open. $f_X^i[V^{i+1}] = V^i$ and is thus open. □

Remark 24.5. f_C^i may be a twisted product: there need not exist $f_Y^i: Y^i \rightarrow Y^{i+1}$ such that $f_C^i = f_X^i \times f_Y^i$.

{grhomorph:comp}

4. $f_C^2 \circ f_C^1$ is a G^1 - G^3 - ρ^1 - ρ^3 morphism of $X^1 \times Y^1$ to $X^3 \times Y^3$.

Proof. Let $f_G^i: G^i \rightarrow G^{i+1}$ be a continuous homomorphism such that

$$\left(\bigvee_{\substack{(x^i, y^i) \in X^i \times Y^i \\ g^i \in G^i}} \right) f_C^i((x^i, y^i) \star g^i) = f_C^i((x^i, y^i)) \star f_G^i(g^i) \quad (24.5) \quad \{\text{eq:Grho2}\}$$

Let $(x^1, y^1) \in X^1 \times Y^1$ and $g^1 \in G^1$. Then fig. 15 (Preserving group actions) is commutative and {fig:Grho2}

$$\begin{aligned}f_C^2 \circ f_C^1((x^1, y^1) \star g^1) &= f_C^2(f_C^1((x^1, y^1)) \star f_G^1(g^1)) \\ &= f_C^2(f_C^1((x^1, y^1))) \star f_G^2 \circ f_G^1(g^1) \\ &= f_C^2 \circ f_C^1((x^1, y^1)) \star f_G^2 \circ f_G^1(g^1)\end{aligned}$$

□

{def:Grhomodcat}

Definition 24.6 (Categories of G - ρ -model spaces). Let $X^\alpha, Y^\alpha, \alpha < A$, be topological spaces, G^α a topological group, ρ^α an effective group action on Y^α ,




Figure 15: Preserving group actions

{fig:Grho2}

$\mathbf{XY}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{XY}^\alpha)$ a G^α - ρ^α model space of $X^\alpha \times Y^\alpha$,

$\mathcal{XY}\mathcal{E}\rho_{\text{Ob}} \stackrel{\text{def}}{=} \{\mathbf{XY}^\alpha \mid \alpha < A\}$,

$\mathcal{XY}\mathcal{E}\rho_{\text{Ar}} \stackrel{\text{def}}{=} \{f_C: X^\alpha \times Y^\alpha \longrightarrow X^\beta \times Y^\beta \mid$

$\text{isG}\rho\text{morph}(\mathbf{XY}^\alpha, G^\alpha, \rho^\alpha, \mathbf{XY}^\beta, G^\beta, \rho^\beta, f_C) \wedge \alpha < A \wedge \beta < A\}$

and $\mathcal{XY}\mathcal{E}\rho \stackrel{\text{def}}{=} (\mathcal{XY}\mathcal{E}\rho_{\text{Ob}}, \mathcal{XY}\mathcal{E}\rho_{\text{Ar}})$. Then any subcategory of $\mathcal{XY}\mathcal{E}\rho$ is a G - ρ model category.

{def:Grhomodtriv}

Definition 24.7 (Trivial G - ρ -model spaces). Let X and Y be topological spaces, G a topological group, ρ an effective group action of G on Y and $\mathcal{XY}$ the category of all products of open subsets of X with Y and all homeomorphisms induced by the group action, i.e.,

$$\text{Ob}(\mathcal{XY}) \stackrel{\text{def}}{=} \left\{ V \times Y \mid V \in X_{\text{op}} \right\} \quad (24.6) \quad \{\text{eq:Grho2}\}$$

$$\text{Ar}(\mathcal{XY}) \stackrel{\text{def}}{=} \left\{ t: V \times Y \xrightarrow{\sim} V \times Y \mid \right. \\ \left. V \in X_{\text{op}} \wedge \pi_1 \circ t = \pi_1 \wedge \left(\exists_{g \in G^V} \right) \left(\bigvee_{\substack{x \in V \\ y \in Y}} f(x, y) = (x, y \star g(x)) \right) \right\} \quad (24.7)$$

Then the trivial G - ρ model space of X, Y , abbreviated $X, Y_{G-\rho-\text{triv}}$, is $(X \times Y, \mathcal{XY})$ and

$X, Y_{G-\rho-\text{triv}}$ is a trivial G - ρ model space of X, Y .

Remark 24.8. Let G' be a topological group and ρ' an effective group action of G' on Y such that $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix} = \begin{smallmatrix} X, Y \\ G'-\rho'-\text{triv} \end{smallmatrix}$. Although G' must be isomorphic to G , it need not have the same topology.

The identity morphism of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ is $\text{Id}_{\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}} \stackrel{\text{def}}{=} \text{Id}_{X \times Y}$.

Lemma 24.9 (Trivial G - ρ -model spaces). Let X and Y be topological spaces, G a topological group and ρ an effective group action on Y .

$\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ is a G - ρ model space of $X \times Y$.

Proof. $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ satisfies the conditions of definition 24.1 (G - ρ -model spaces) on page 148:

1. Product with fiber:

$$\left(\bigvee_{U \in \pi_2 \left(\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix} \right)}^{\text{Ob}} \right) \left(\bigvee_{V \in X_{\text{op}}} U = V \times Y \right)$$

by definition 24.7.

2. Generated by ρ : Let $f \in \pi_2 \left(\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix} \right)$: $V \times Y \xrightarrow{\sim} V \times Y$. Then

$$\left(\bigvee_{g \in G^V} \right) \left(\bigvee_{(x,y) \in V \times Y} \right) f(x, y) = (x, y \star g(x))$$

by definition 24.7.

□

$\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ is a fine grained G - ρ model space of $X \times Y$.

Proof. Let $V \times Y$ be a model neighborhood and V' an open subset. $V' \times Y$ is open by definition 24.7.

□

Let $E^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i)$, $i = 1, 2$, be a model space, $\mathbf{X}\mathbf{Y}^i$ be a fine grained G^i - ρ^i model space of $X^i \times Y^i$, and $A^i \stackrel{\text{def}}{=} \{(U^\alpha, V^\alpha \times Y^i, \phi^\alpha) | \alpha < A\}$ be an m-atlas of E in the coordinate space $\mathbf{X}\mathbf{Y}^i$.

A^i is an m-atlas of $\begin{smallmatrix} E^i \\ X^i-\pi^i-\text{triv} \end{smallmatrix}$ in the coordinate space $\begin{smallmatrix} X^i, Y^i \\ G^i-\rho^i-\text{triv} \end{smallmatrix}$.

Proof. $\mathbf{X}\mathbf{Y}^i$ is a fine grained G^i - ρ^i model space of $X^i \times Y^i$ by hypothesis.

A^i is an m-atlas of E in the coordinate space $\mathbf{X}\mathbf{Y}^i$ by hypothesis.

A^i is an m-atlas of $\frac{E^i}{X^i-\pi^i-\text{triv}}$ in the coordinate space $\mathbf{XY}^i$ by lemma 24.2 (*G- ρ -model spaces*) on page 148. {lem:Grhomod}

Let $(U^i, V^i \times Y^i, \phi^i)A^i, i = 1, 2$. Then $(U^i, V^i \times Y^i, \phi^i)$ is an m-chart of $\frac{E^i}{X^i-\pi^i-\text{triv}}$ in the coordinate space $\mathbf{XY}^i$:

1. V^i is open by definition 24.1 (*G- ρ -model spaces*) on page 148 and $\pi^{i-1}[V^i \times Y^i] = U^i$ by lemma 24.2, hence U^i is a model neighborhood of $\frac{E^i}{X^i-\pi^i-\text{triv}}$ by definition 24.7 (*Trivial G- ρ -model spaces*) on page 152. {def:Grhomod}
{def:Grhomodtriv}
2. $[V^i \times Y^i]$ is a model neighborhood of $\frac{E^i}{X^i-\pi^i-\text{triv}}$ by the above.
3. The transition function $t_j^i = \phi^j \circ \phi^{i-1}$ is of the form $t_j^i(x^i, y^i) = (X^i, y^i \star g_j^i(x^i))$

□

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_0, f_1)$ be a E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathbf{XY}^1, \mathbf{XY}^2$. Then $\mathbf{f}$ is a $\frac{E^1}{X^1-\pi^1-\text{triv}} - \frac{E^2}{X^2-\pi^2-\text{triv}}$ m-atlas morphism of A^1 to A^2 in the coordinate spaces $\frac{X^1, Y^1}{G^1-\rho^1-\text{triv}}, \frac{X^2, Y^2}{G^2-\rho^2-\text{triv}}$.

Proof. $\mathbf{f}_1$ is a model function by hypothesis, hence continuous.

$\mathbf{f}_0: \frac{E^1}{X^1-\pi^1-\text{triv}} \longrightarrow \frac{E^2}{X^2-\pi^2-\text{triv}}$ is a model function by hypothesis. Let U^2 be a model neighborhood of $\frac{E^2}{X^2-\pi^2-\text{triv}}$ and $U^1 \stackrel{\text{def}}{=} f_0^{-1}[U^2]$. Then U^2 is of the form $\pi^{2-1}[V^2]$ with V^2 open. Let $V^1 \stackrel{\text{def}}{=} f_1^{-1}[V^2]$. Since $\mathbf{f}_1$ is continuous, V^1 is open and thus $\pi^{1-1}[V^1] = U^1$ is open and a model neighborhood of $\frac{E^1}{X^1-\pi^1-\text{triv}}$. Let $(U^i, V^i \times Y^i, \phi^i) \in A^i, i = 1, 2$, and $I \stackrel{\text{def}}{=} U^1 \cap f_0^{-1}[U^2] \neq \emptyset$. Each V^i is open, thus $U^i = \pi^{i-1}[V^i]$ is a model neighborhood of $\frac{E^i}{X^i-\pi^i-\text{triv}}$ and each $V^i \times Y^i$ is a model neighborhood of $\frac{X^i, Y^i}{G^i-\rho^i-\text{triv}}$.

Let $u^1 \in I, (U'^1, V'^1 \times Y^1, \phi^1), U'^2 \subseteq U^2, (U'^2, \hat{V}'^2 \times Y^2, \phi'^2)$ be as in definition 13.1 (*M-atlas morphisms for model spaces*) on page 61. By the above, $(U'^1, V'^1 \times Y^1, \phi^1)$ and $U'^2 \subseteq U^2, (U'^2, \hat{V}'^2 \times Y^2, \phi'^2)$ are M-charts of $\frac{E^1}{X^1-\pi^1-\text{triv}} (\frac{E^2}{X^2-\pi^2-\text{triv}})$ in the coordinate space $\frac{X^1, Y^1}{G^1-\rho^1-\text{triv}} (\frac{X^2, Y^2}{G^2-\rho^2-\text{triv}})$. The commutation relations are not changed by the change of spaces. {def:M-ATLmorph}

{cor:Grhomodtriv}

Corollary 24.10. Let X and Y be topological spaces, G a topological group and ρ an effective group action on Y . Then $\frac{X, Y}{G-\rho-\text{triv}}$ is a X - π_1 -fine grained model space.

Proof. The proof follows from lemma 24.2 (*G- ρ -model spaces*) on page 148 and lemma 24.9 above. {lem:Grhomodtriv}
{lem:Grhomod}

□

{\lem:Grhomodtrivmorph}

Lemma 24.11 (Morphisms of trivial G - ρ -model spaces). Let X^i and Y^i , $i = 1, 2$, be topological spaces, G^i a topological group, ρ^i an effective group action of G^i on Y^i and $(f_X: X^1 \longrightarrow X^2, f_Y: Y^1 \longrightarrow Y^2)$ a pair of continuous functions such that $f_C \stackrel{\text{def}}{=} f_X \times f_Y: \underset{G^1-\rho^1-\text{triv}}{X^1, Y^1} \longrightarrow \underset{G^2-\rho^2-\text{triv}}{X^2, Y^2}$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$. Then f_C is a model function from $\underset{G^1-\rho^1-\text{triv}}{X^1, Y^1}$ to $\underset{G^2-\rho^2-\text{triv}}{X^2, Y^2}$.

Proof.

1. f_C is a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$ by hypothesis, hence continuous.
2. Let $m^2 \stackrel{\text{def}}{=} U^2 \times Y^2$ be a model neighborhood of $\underset{G^2-\rho^2-\text{triv}}{X^2, Y^2}$. Then $U^1 \stackrel{\text{def}}{=} f_X^{-1}(U^2)$ is open and $f_C^{-1}(m^2) = U^1 \times Y^1$ is a model function.
3. Let $m^1 \stackrel{\text{def}}{=} U^1 \times Y^1$ be a model neighborhood of $\underset{G^1-\rho^1-\text{triv}}{X^1, Y^1}$. $X^2 \times Y^2$ is a model neighborhood of $\underset{G^2-\rho^2-\text{triv}}{X^2, Y^2}$ and $f_C(m^1) \subseteq X^2 \times Y^2$.

□

Remark 24.12. The hypothesis precludes the possibility that f_C is twisted.

{\def:Grhomodtrivcat}

Definition 24.13 (Categories of trivial G - ρ -model spaces). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles.

Remark 24.14. The component E^α plays no role in this definition.

The trivial coordinate model category of $\mathbf{B}$, $\underset{\text{Bun-}\mathcal{T}\text{riv}}{\mathbf{B}}$, is the category with objects all trivial G_α - ρ_α model spaces of X_α, Y_α , $\alpha < A$ and morphisms all continuous functions compatible with the group action:

$$\underset{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ob}}}{\mathbf{B}} \stackrel{\text{def}}{=} \left\{ \underset{G-\rho-\text{triv}}{X, Y} \mid (E, X, Y, \pi, G, \rho) \in \mathbf{B} \right\} \quad (24.8)$$

$$\underset{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ar}}}{\mathbf{B}} \stackrel{\text{def}}{=} \left\{ f_C: \underset{G^1-\rho^1-\text{triv}}{X^1 \times Y^1} \longrightarrow \underset{G^2-\rho^2-\text{triv}}{X^2 \times Y^2} \mid \left(\exists \begin{matrix} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i) \in \mathbf{B} \\ f_G: G^1 \longrightarrow G^2 \end{matrix} \right) \text{isG}\rho\text{morph} \left(\underset{G-\rho-\text{triv}}{X^1, Y^1}, G^1, \rho^1, \underset{G-\rho-\text{triv}}{X^2, Y^2}, G^2, \rho^2, f^C \right) \right\} \quad (24.9)$$

$$\underset{\text{Bun-}\mathcal{T}\text{riv}}{\mathbf{B}} \stackrel{\text{def}}{=} \left(\underset{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ob}}}{\mathbf{B}}, \underset{\text{Bun-}\mathcal{T}\text{riv}_{\text{Ar}}}{\mathbf{B}} \right) \quad (24.10)$$

Remark 24.15. The morphisms f_C may be twisted products: there need not exist $f_Y: Y^1 \longrightarrow Y^2$ such that $f_C = f_X \times f_Y$.

The trivial product coordinate model category of $\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$, is the category with objects all trivial G^α - ρ^α model spaces of $X^\alpha, Y^\alpha, \alpha < A$ and morphisms all products of continuous functions compatible with the group action:

$$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}} \stackrel{\text{def}}{=} \mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}} \quad (24.11)$$

$$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}} \stackrel{\text{def}}{=} \left\{ f_C \stackrel{\text{def}}{=} f_X \times f_Y: X^1 \times Y^1 \longrightarrow X^2 \times Y^2 \mid \left(\exists_{\substack{(E^i, X^i, Y^i, \pi^i, G^i, \rho^i) \in \mathbf{B} \\ f_G: G^1 \longrightarrow G^2}} \text{is} G\text{-}\rho \text{ morphism} \left(\begin{smallmatrix} X^1, Y^1 \\ G-\rho\text{-triv} \end{smallmatrix}, \begin{smallmatrix} X^2, Y^2 \\ G-\rho\text{-triv} \end{smallmatrix}, G^2, \rho^2, f_C \right) \right) \right\} \quad (24.12)$$

$$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}} \stackrel{\text{def}}{=} \left(\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}, \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}} \right) \quad (24.13)$$

Any subcategory of $\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}$ is a trivial coordinate model category and any subcategory of $\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$ is a trivial product coordinate model category.

{lem:Grhomodtrivcat}

Lemma 24.16 (The trivial coordinate model category of $\mathbf{B}$ is a category). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha), \alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles.

Then $\mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}$ is a category and the identity morphism for object $\mathbf{B}^\alpha$ is $\text{Id}_{X^\alpha \times Y^\alpha}$.

Proof.

Composition: The composition of G - ρ morphisms is a G - ρ morphism by lemma 24.4 (Morphisms of G - ρ -model spaces) on page 150. {lem:Grhomorph}

Associativity: Morphisms are simply functions and composition of morphisms is simply composition of functions.

Unit: The identity morphisms are simply identity functions and composition of morphisms is simply composition of functions.

□

$\mathbf{B}_{\text{Bun-prod-}\mathcal{T}\text{riv}}$ is a category and the identity morphism for object $\mathbf{B}^\alpha$ is $\text{Id}_{X^\alpha \times Y^\alpha}$.

Proof.

Composition: The composition of G - ρ morphisms is a G - ρ morphism by lemma 24.4_{lem:Grhomorph}

(Morphisms of G - ρ -model spaces) on page 150. Let $X^i, Y^i \in^{Ob}_{G^i-\rho^i-\mathbf{triv}} \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\mathbf{riv}}$,

$i = 1, 2, 3$, $f_C^i = f_X^i \times f_Y^i: X^i \times Y^i \longrightarrow X^{i+1} \times Y^{i+1} \in^{Ar}_{\text{Bun-prod-}\mathcal{T}\mathbf{riv}} \mathbf{B}_{\text{Bun-prod-}\mathcal{T}\mathbf{riv}}$. Then $f_C^{i+1} \circ f_C^i = (f_X^{i+1} \circ f_X^i) \times (f_Y^{i+1} \circ f_Y^i)$.

Associativity: Morphisms are simply functions and composition of morphisms is simply composition of functions.

Unit: The identity morphisms are simply identity functions and composition of morphisms is simply composition of functions.

□

25 Bundle charts

Definition 25.1 (Y - π -bundle charts). Let E, X, Y be topological spaces and $\pi: E \longrightarrow X$. sec:buncharts {def:Ypichart}

Then $(U, V \times Y, \phi: U \xrightarrow{\cong} V \times Y)$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$ iff




Figure 16: ϕ preserves fibers

{fig:Ypichart}

1. U , known as a coordinate patch, is a novoid open subset of E ,
2. $V \times Y$ is a novoid open subset of $X \times Y$
3. ϕ , known as a coordinate function, is a homeomorphism that preserves fibers.
i.e., $\pi_1 \circ \phi = \pi$.

$(U, V \times Y, \phi)$ covers u iff $u \in U$.

{lem:Ypichart}

Lemma 25.2 (Y - π -bundle charts of m -atlases in G - ρ model spaces). Let

1. E, X, Y be topological spaces

2. G a topological group
3. ρ an effective group action of G on Y
4. $\mathbf{XY} \stackrel{\text{def}}{=} (X \times Y, \mathcal{XY})$ a G - ρ model space of $X \times Y$
5. $\mathbf{A}$ an m-atlas of E in the coordinate space $\mathbf{XY}$

There is a unique $\pi: E \rightarrow X$ such that every chart $(U, V \times Y, \phi) \in \mathbf{A}$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$. Further, π is continuous.

Proof. Let e be an arbitrary point of E and let $(U, V \times Y, \phi) \in \mathbf{A}$ be an arbitrary m-chart in $\mathbf{A}$ that covers e . Define $\pi(e) \stackrel{\text{def}}{=} \pi_1 \circ \phi(e)$.

To show that $\pi(e)$ is well defined let $(U', V' \times Y, \phi')$ be another m-chart in $\mathbf{A}$ that covers e . By definition 24.7 (Trivial G - ρ -model spaces) on page 152, the transition function is of the form $t(x, y) = (x, y \star g(x))$. Then $\pi_1 \circ t = \pi_1$ and $\pi_1 \circ \phi'(e) = \pi_1 \circ \phi(e)$. {def:Grhomodtriv}

Let $V \subseteq X$ be open and let $e \in \pi_1^{-1}[V]$. Let $(U', V' \times Y, \phi')$ be an m-chart of $\mathbf{A}$ that covers e . $V \cap V'$ is open, π_1 is continuous and ϕ' is continuous, so $\pi^{-1}[V'] = \phi'^{-1}[\pi_1^{-1}[V']]$ is open. $\pi^{-1}(V)$ is thus a union of open sets, hence open. {lem:GrhomodtrivYpichart}

Lemma 25.3 (Y - π -bundle charts in $\mathbf{X}, Y$). Let E, X and Y be topological spaces, G a topological group, ρ an effective group action of G on Y and $\pi: E \rightarrow X$. $(U, V \times Y, \phi)$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$ iff $(U, V \times Y, \phi)$ is an m-chart of E in the coordinate space $\mathbf{X}, Y$ and ϕ preserves fibers. {lem:GrhomodtrivYpichart}

Proof. If $(U, V \times Y, \phi)$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$ then by definition 25.1 (Y - π -bundle charts) on page 157 {def:Ypichart}

1. U is a nonvoid open subset of E .
2. $V \times Y$ is an open subset of $X \times Y$, hence a model neighborhood of $\mathbf{X}, Y$ by definition 24.7 (Trivial G - ρ -model spaces) on page 152. {def:Grhomodtriv}
3. ϕ is a homeomorphism
4. $(U, V \times Y, \phi)$ is an m-chart of E in the coordinate space $\mathbf{X}, Y$ by definition 24.7. {def:Grhomodtriv}
5. ϕ is a homeomorphism that preserves fibers. i.e., $\pi_1 \circ \phi = \pi$.

If $(U, V \times Y, \phi)$ is an m-chart of E in the coordinate space $\mathbf{X}, Y$ and ϕ preserves fibers then by definition 24.7 {def:Grhomodtriv}

1. U is a nonvoid open subset of E .

2. $V \times Y$ is an m -chart of E in the coordinate space X, Y , hence an open subset of $X \times Y$. $G-\rho-\text{triv}$
3. ϕ is a homeomorphism
4. ϕ preserves fibers by hypothesis.

□

{def:piepi}

Lemma 25.4 (Properties of projection). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $v \in V$. Then

$\pi|_U$ is a surjection.

Proof. Let $v \in V$, $y \in Y$ and $u \stackrel{\text{def}}{=} \phi^{-1}((v, y)) \in U$. Then $\pi(u) = v$. □

$\pi^{-1}[\{v\}]$ is homeomorphic to Y .

Proof. ϕ and ϕ^{-1} are homeomorphisms, so their restrictions are homeomorphisms and thus $\phi^{-1}[\{v\}]$ is homeomorphic to $\{v\} \times Y$, which is homeomorphic to Y . □

{def:Ypisub}

Definition 25.5 (Y - π subcharts). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and U' be a nonvoid open subset of U . Then

$(U', V' \times Y, \phi') \stackrel{\text{def}}{=} (U', \phi[U'], \phi|_{U'})$ is a subchart of $(U, V \times Y, \phi)$.

Lemma 25.6 (Y - π subcharts). Let $(U, V \times Y, \phi)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $(U', V' \times Y, \phi')$ a subchart of $(U, V \times Y, \phi)$. Then $(U', V' \times Y, \phi')$ is a Y - π -bundle chart of E in the coordinate space $X \times Y$.

Proof. $(U', V' \times Y, \phi')$ satisfies the conditions of definition 25.1: {def:Ypicart}

1. $U' \subseteq E$ is open.
2. $\phi[U']$ is open since ϕ is a homeomorphism.
3. $\phi|_{U'}: U' \rightarrow V' \times Y$ is the restriction of a homeomorphism and thus a homeomorphism. $\phi|_{U'}$ preserves fibers because ϕ does.

□

{def:Grho-compat}

Definition 25.7 (G - ρ -compatibility). Let E, X, Y be topological spaces, G a topological group, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y , $\pi: E \rightarrow X$ surjective and $(U^\mu, V^\mu \times Y, \phi^\mu), (U^\nu, V^\nu \times Y, \phi^\nu)$ Y - π -bundle charts. Then $(U^\mu, V^\mu \times Y, \phi^\mu)$ and $(U^\nu, V^\nu \times Y, \phi^\nu)$ are G - ρ -compatible iff either

1. U^μ and U^ν are disjoint
2. The transition function $t_\nu^\mu \stackrel{\text{def}}{=} \phi^\mu \circ \phi^{\nu-1}|_{\phi^\nu[U^\mu \cap U^\nu]}$ is generated by the group action, i.e., there is a continuous function $g_\nu^\mu: \pi_1[\phi^\nu[U^\mu \cap U^\nu]] \rightarrow G$ such that

$$\left(\bigvee_{(x,y) \in \phi^\nu[U^\mu \cap U^\nu]} \right) t_\nu^\mu(x, y) = (x, y \star g_\nu^\mu(x)) \quad \text{transition group} \quad (25.1)$$

{lem:Grho-compat}

Lemma 25.8 (G - ρ -compatibility). Let E, X, Y be topological spaces, G a topological group, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y , $\pi: E \rightarrow X$ surjective and $(U^i, V^i \times Y, \phi^i)$, $i = 1, 2$, Y - π -bundle charts. Then $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are G - ρ -compatible iff $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are compatible m-charts of $\begin{smallmatrix} E \\ \text{triv} \end{smallmatrix}$ in the coordinate space $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$.

Proof.

Each U^i is a model neighborhood of $\begin{smallmatrix} E \\ \text{triv} \end{smallmatrix}$ and each $V^i \times Y$ is a model neighborhood of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$.

2. If $U^1 \cap U^2 = \emptyset$ then the lemma is trivially true.
3. If the charts are G - ρ -compatible then the transition function $t_2^1 \stackrel{\text{def}}{=} \phi^1 \circ \phi^{2^{-1}} \upharpoonright_{\phi^2[U^1 \cap U^2]}$ is generated by the group action, i.e., there is a continuous function $g_2^1: \pi_1[\phi^2[U^1 \cap U^2]] \rightarrow G$ such that $\left(\bigvee_{(x,y) \in \phi^2[U^1 \cap U^2]} \right) t_2^1(x, y) = (x, y \star g_2^1(x))$. Then $\pi_1 \circ t_2^1 = \pi_1$, t_2^1 is a morphism of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ and the charts are compatible m-charts of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$.
4. If the charts are compatible m-charts of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$ then the transition function is an isomorphism of $\begin{smallmatrix} X, Y \\ G-\rho-\text{triv} \end{smallmatrix}$, hence generated by the group action, and the charts are G - ρ -compatible.

□

{cor:Grho-compatsym}

Corollary 25.9 (Symmetry of G - ρ compatibility). Let $(U^\mu, V^\mu \times Y, \phi^\mu)$ and $(U^\nu, V^\nu \times Y, \phi^\nu)$ be Y - π -bundle charts of E in the coordinate space $X \times Y$. Then $(U^\mu, V^\mu \times Y, \phi^\mu)$ is G - ρ -compatible with $(U^\nu, V^\nu \times Y, \phi^\nu)$ iff $(U^\nu, V^\nu \times Y, \phi^\nu)$ is G - ρ -compatible with $(U^\mu, V^\mu \times Y, \phi^\mu)$.

Proof. The result follows from lemma 11.11 (Symmetry of m-compatibility) on page 54. □

{lem:M-compatsym}

{lem:Grho-compatsub}

Lemma 25.10 (G - ρ -compatibility of subcharts). Let $(U^i, V^i \times Y, \phi^i)$ be a Y - π -bundle chart of E in the coordinate space $X \times Y$, $(U'^i, V'^i \times Y, \phi'^i)$ a subchart and $(U^1, V^1 \times Y, \phi^1)$ be G - ρ -compatible with $(U^2, V^2 \times Y, \phi^2)$. Then $(U'^1, V'^1 \times Y, \phi'^1)$ is G - ρ -compatible with $(U'^2, V'^2 \times Y, \phi'^2)$.

Proof. If $U^1 \cap U^2 = \emptyset$ then $U'^1 \cap U'^2 = \emptyset$. If $U'^1 \cap U'^2 \neq \emptyset$ then $(U'^1, V'^1 \times Y, \phi'^1)$ is G - ρ -compatible with $(U'^2, V'^2 \times Y, \phi'^2)$. Otherwise, the transition function

$t_2^1 \stackrel{\text{def}}{=} \phi^1 \circ \phi^{2^{-1}} \upharpoonright_{\phi^2[U^1 \cap U^2]}$ is generated by the group action and hence $t_2^1 \upharpoonright_{\phi^2[U'^1 \cap U'^2]}: \phi^2[U'^1 \cap U'^2] \xrightarrow{\cong} \phi^1[U'^1 \cap U'^2]$ is generated by the group action. □

Corollary 25.11 (G - ρ -compatibility with subcharts). Let $(U, V \times Y, \phi)$, be a Y - π -bundle chart of E in the coordinate space $X \times Y$ and $(U', V' \times Y, \phi')$ a subchart. Then $(U', V' \times Y, \phi')$ is G - ρ -compatible with $(U, V \times Y, \phi)$.

Proof. $(U, V \times Y, \phi)$ is G - ρ -compatible with itself and is a subchart of itself, $\square$

Definition 25.12 (Covering by Y - π -bundle charts). Let $\mathbf{A}$ be a set of Y - π -bundle charts of E in the coordinate space $X \times Y$. $\mathbf{A}$ covers E iff $E = \bigcup \pi_1[\mathbf{A}]$.

Lemma 25.13. Let $\mathbf{A}$ be a set of Y - π -bundle charts of E in the coordinate space $X \times Y$ that covers E and $x \in X$. Then $\pi^{-1}[\{x\}]$ is homeomorphic to Y .

Proof. Since $\mathbf{A}$ covers E , there is a chart (U, V, ϕ) in $\mathbf{A}$ containing x . Then $\pi^{-1}[\{x\}]$ is homeomorphic to Y by lemma 25.4 (Properties of projection) on page 159. $\square$ {def:piepi}

26 Bundle atlases

{sec:BunAtlases}

A set of charts can be atlases for different fiber bundles even if it is for the same total model space, base space and fiber. In order to aggregate atlases into categories, there must be a way to distinguish them. Including the spaces¹⁹, group and group action in the definitions of the categories serves the purpose.

{def:Bun-Atl}

Definition 26.1 (Bundle atlases). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$, be a protobundle. Then $(\mathbf{A}, \mathbf{B})$ is a bundle atlas, $\mathbf{A}$ is a bundle atlas of $\mathbf{B}$, abbreviated $\text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ and $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$, abbreviated $\text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E, X, Y, \pi, G, \rho)$, iff $\mathbf{A}$ consists of a set of mutually G - ρ -compatible Y - π -bundle charts of E in the coordinate space $X \times Y$ that covers E ²⁰.

By abuse of language we write $U \in \mathbf{A}$ for $U \in \pi_1[\mathbf{A}]$.

Remark 26.2. The definition of a π - G - ρ -bundle atlas is by design similar to the definition of a coordinate bundle in [6, p. 7], but there are significant differences. This paper will use the term bundle atlas to avoid confusion. {TopFib}

Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\mathcal{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}) \stackrel{\text{def}}{=} \{(\mathbf{A}, \mathbf{B}^\alpha) | \text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)\} \quad (26.1)$$

$$\mathcal{Atl}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\alpha < A} \mathcal{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \quad (26.2)$$

¹⁹The spaces are redundant, but convenient.

²⁰There is no need to introduce the concept of a full π - G - ρ -bundle atlas because a π - G - ρ -bundle atlas is automatically full.

Lemma 26.3 (Bundle atlases). Let E, X, Y be topological spaces, G a topological group, $\pi: E \rightarrow X$ surjective and $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y . {lem:Bun-Atl}

$\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ iff it is an m-atlas of $\underset{X-\pi-\text{triv}}{E}$ in the coordinate space $\underset{G-\rho-\text{triv}}{X, Y}$.

Proof. Let $(U^i, V^i \times Y, \phi^i) \in \mathbf{A}, i = 1, 2$, be Y - π -bundle charts. Then by lemma 25.8 (G- ρ -compatibility) on page 160, $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are G - ρ -compatible iff $(U^1, V^1 \times Y, \phi^1)$ and $(U^2, V^2 \times Y, \phi^2)$ are compatible m-charts of $\underset{G-\rho-\text{triv}}{E}$ in the coordinate space $\underset{G-\rho-\text{triv}}{X, Y}$. {lem:Grho-compat} $\square$

If $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ then every chart of $\mathbf{A}$ is of the form $(U, V \times Y, \phi)$.

Proof. The result follows from definition 24.7 (Trivial G - ρ -model spaces) on page 152. {def:Grhomodtriv} $\square$ {cor:Bun-Atl}

Corollary 26.4 (Bundle atlases). Let E, X, Y be topological spaces, G a topological group, $\pi: E \rightarrow X$ surjective, $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y and $\mathbf{A}$ an m-atlas of $\underset{\text{triv}}{E}$ in the coordinate space $X \times Y$ such that for every $e \in E$ there is a Y - π -bundle chart of E in the coordinate space $X \times Y$ in $\mathbf{A}$ that covers e .

Then $\mathbf{A}$ is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$.

Proof. The result follows from lemma 25.2 (Y - π -bundle charts of m-atlases in G - ρ model spaces) on page 157. {lem:Ypichart} $\square$

Definition 26.5 (Compatibility of charts with bundle atlases). A Y - π -bundle chart $(U, V \times Y, \phi)$ of E in the coordinate space $X \times Y$ is G - ρ -compatible with a π - G - ρ -bundle atlas $\mathbf{A}$ iff it is G - ρ -compatible with every chart in the atlas. {def:piGrho-atl:extensions}

Lemma 26.6 (Compatibility of subcharts with bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ and $(U, V \times Y, \phi)$ a Y - π -bundle chart in $\mathbf{A}$. Then any subchart of $(U, V \times Y, \phi)$ is G - ρ -compatible with $\mathbf{A}$. {lem:BunCompat}

Proof. The result follows from lemma 12.6 (M-compatibility of subcharts with atlases) on page 57 and lemma 25.8 (G - ρ -compatibility) on page 160. {lem:M-CompatsubATL} $\square$

Lemma 26.7 (Extensions of bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -atlas of E in the coordinate space $X \times Y$ and $(U_i, V_i, \phi_i), i = 1, 2$, be a π - G - ρ chart of E in the coordinate space $X \times Y$ G - ρ -compatible with $\mathbf{A}$ in the coordinate space $X \times Y$. Then (U_1, V_1, ϕ_1) is G - ρ -compatible with (U_2, V_2, ϕ_2) in the coordinate space $X \times Y$. {lem:Grho-compat} {lem:piGrho-atl:extensions}

Proof. The result follows from lemma 12.7 (Extensions of m-atlases) on page 58 and lemma 25.8 (G - ρ -compatibility) on page 160. {lem:M-ATLextensions} $\square$ {lem:Grho-compat}

{def:Bun-ATLmax}

Definition 26.8 (Maximal bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. $\mathbf{A}$ is a maximal π - G - ρ -bundle atlas, abbreviated $\text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E, X, Y, G, \pi, \rho)$, iff it cannot be extended by adding an additional G - ρ -compatible Y - π -bundle chart.

Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\text{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \left\{ (\mathbf{A}, \mathbf{B}^\alpha) \mid \text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \right\} \quad (26.3)$$

$$\text{Atl}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\alpha < A} \text{Atl}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha) \quad (26.4)$$

{lem:Bun-ATLmax}

Lemma 26.9 (Maximal bundle atlases). Let $\mathbf{A}$ be a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. $\mathbf{A}$ is a maximal π - G - ρ -bundle atlas iff it is a maximal m-atlas of E in the coordinate space X, Y .
 $\text{triv} \quad G-\rho-\text{triv}$

Proof. The result follows from lemma 25.8 (G - ρ -compatibility) on page 160 and lemma 26.7 above. $\square$

{lem:Grho-compat}

Theorem 26.10 (Existence and uniqueness of maximal π - G - ρ -bundle atlases). Let $\mathbf{B} \stackrel{\text{def}}{=} (E, X, Y, \pi, G, \rho)$ be a protobundle and $\mathbf{A}$ a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$. Then there exists a unique maximal π - G - ρ -bundle atlas $\text{Atlas}_{\text{max}}^{\text{Bun}}(\mathbf{A}, \mathbf{B})$ of E in the coordinate space $X \times Y$ G - ρ -compatible with $\mathbf{A}$.

Proof. Let $\mathbf{P}$ be the set of all π - G - ρ -bundle atlases E in the coordinate space $X \times Y$ containing $\mathbf{A}$ and G - ρ compatible in the coordinate space $X \times Y$ with $\mathbf{A}$. Let $\mathbf{P}_{\text{max}}$ be a maximal chain of $\mathbf{P}$. Then $\mathbf{A}' = \bigcup_{\text{max}} \mathbf{P}$ is a maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$ G - ρ compatible with $\mathbf{A}$. Uniqueness follows from definition 26.5 (Compatibility of charts with bundle atlases) on page 162. $\square$

{def:piGrho-atl:extensions}

{lem:mGrho}

Lemma 26.11 (Existence and uniqueness of projection for atlases in G - ρ -model spaces). Let E, X and Y be topological spaces, G a topological group, ρ an effective action of G on Y , $\mathbf{C} \stackrel{\text{def}}{=} (X \times Y, \mathcal{X}\mathcal{Y})$ a G - ρ model space of $X \times Y$ and $\mathbf{A}$ an m-atlas of E in the coordinate model space $\mathbf{C}$. Then there exists a unique function $\pi: E \rightarrow X$ such that for any chart (U, V, ϕ) in $\mathbf{A}$, $\pi|_U = \pi_1 \circ \phi$. If $\mathbf{A}$ is full then π is surjective.

Proof. Let (U, V, ϕ) be an arbitrary chart in $\mathbf{A}$ and define $\pi(e \in U) = \pi_1 \circ \phi(e)$. $\pi(e)$ does not depend on the choice of chart because the morphisms of a G - ρ model

space preserve fibers. π is continuous because it is continuous on each coordinate patch.

Let $x \in X$. If $\mathbf{A}$ is full then there exists a chart (U, V, ϕ) in $\mathbf{A}$ such that $x \in \pi_1[V]$. Let u be an arbitrary point in $\phi^{-1}[\{x\} \times Y]$. Then $\pi(u) = \pi_1(\phi(u)) = x$. $\square$

27 Bundle-atlas morphisms and functors

{sec:bunatlmorph}

This section introduces a taxonomy for morphisms between bundle atlases, defines categories of bundle atlases and constructs functors from them to categories of $\mathbf{B}$ $\text{Bun-prod-}\mathcal{T}riv$ atlases. It only constructs reverse functors for subcategories of

27.1 Bundle-atlas morphisms

27.1.1 Definitions of bundle-atlas morphisms

{def:Bun-ATLmorph}

Definition 27.1 (Bundle-atlas morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle and let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$.

Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\text{isAtl}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$, iff

1. all four functions are continuous
2. f_G is a homomorphism
3. $\mathbf{f}$ commutes with π^i and ρ^i , i.e.,

$$(a) \quad \pi^2 \circ f_E = f_X \circ \pi^1$$

$$(b) \quad \left(\bigvee_{\substack{y \in Y^1 \\ g \in G^1}} \right) f_Y(y) \star^2 f_G(g) = f_Y(y \star^1 g)$$

4. For any charts $(U^i, V^i \times Y, \phi^i: U^i \xrightarrow{\cong} V^i \times Y) \in \mathbf{A}^i$, $i = 1, 2$, and any $u^1 \in I \stackrel{\text{def}}{=} U^1 \cap f_E^{-1}[U^2]$, there exists a subchart $(U'^1, V'^1 \times Y, \phi^1: U'^1 \xrightarrow{\cong} V'^1 \times Y) \in \mathbf{A}^1$ at u^1 , an open set $U'^2 \subseteq U^2$ and a chart $(U'^2, V'^2 \times Y, \phi'^2) \in \mathbf{A}^2$ at $f_E(u^1)$ such that $f_E[U'^1] \subseteq U'^2$ and diagram (27.1) below is commutative, as shown in fig. 17.

{fig:Bun-At1M1}

$$D \stackrel{\text{def}}{=} \left\{ \begin{array}{l} i: \{u^1\} \rightarrow U'^1, \phi^1: U'^1 \xrightarrow{\cong} V'^1 \times Y, f_X \times f_Y: V'^1 \times Y \rightarrow \hat{V}'^2 \times Y, \\ f_E: U'^1 \rightarrow U'^2, \phi'^2: U'^2 \xrightarrow{\cong} \hat{V}'^2 \times Y \end{array} \right\} \quad (27.1) \quad \{\text{eq:Bun-At1M1}\}$$

$$\begin{array}{ccccccc} & & u^1 & \xrightarrow{f_E} & u^2 & & \\ & & \downarrow i & & \downarrow i & & \\ U^1 & \xleftarrow{i} & U'^1 & \xrightarrow{f_E} & U'^2 & \xrightarrow{i} & U^2 \\ \downarrow \cong \phi^1 & & \downarrow \cong \phi^1 & & \downarrow \cong \phi'^2 & & \downarrow \cong \phi^2 \\ V^1 \times Y & \xleftarrow{i} & V'^1 \times Y & \xrightarrow{f_X \times f_Y} & \hat{V}'^2 \times Y & \xrightarrow{i} & V^2 \times Y \end{array}$$

Figure 17: Completed Bundle-atlas morphism

{fig:Bun-At1M1}

If $\mathbf{A}^1$ and $\mathbf{A}^2$ are maximal m-atlases then $\mathbf{f}$ is also a maximal $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\text{isAt1}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f})$.

The identity morphism of $(\mathbf{A}^i, \mathbf{B}^i)$ is

$$\text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) \quad (27.2) \quad \{\text{eq:Bun-At1M1}\}$$

This nomenclature will be justified later.

{def:Bun-ATLmorphequiv}

Definition 27.2 (Equivalence of bundle-atlas morphisms).

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is bundle equivalent to $\mathbf{g} \stackrel{\text{def}}{=} (g_E: E^1 \rightarrow E^2, g_X: X^1 \rightarrow X^2, g_Y: Y^1 \rightarrow Y^2, g_G: G^1 \rightarrow G^2)$ as a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$, abbreviated $\mathbf{f}$ and $\mathbf{g}$ are bundle equivalent $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphisms from $\mathbf{A}^1$ to $\mathbf{A}^2$, iff $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i), i = 1, 2$, is a protobundle, $\mathbf{A}^i$ is a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$, $\mathbf{f}$ and $\mathbf{g}$ are $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphisms from $\mathbf{A}^1$ to $\mathbf{A}^2$, $f_E = g_E, f_X = g_X$ and $f_G = g_G$,

By abuse of language we shall write $\mathbf{f}$ is bundle equivalent to $\mathbf{g}$ when the protobundles and bundle atlases are understood by context.

27.1.2 Proclamations on bundle-atlas morphisms

{lem:Bun-ATLmorph}

Lemma 27.3 (Bundle-atlas morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle and let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$.

Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ iff $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 and $(f_E, f_X \times f_Y)$ is a $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 .

Proof.

1. By lemma 26.3 (Bundle atlases) on page 162, $\mathbf{A}^i$ is an m-atlas of E^i in the coordinate space X^i, Y^i , every coordinate function preserves fibers and every chart of $\mathbf{A}^i$ is of the form $(U, V \times Y, \phi)$. {lem:Bun-Atl}
2. If $\mathbf{f}$ is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism then $f_X \times f_Y$ is a model function from X^1, Y^1 to X^2, Y^2 by lemma 24.11 (Morphisms of trivial G - ρ -model spaces) on page 155 and f_G is the function asserted to exist in eq. (24.3) (preservation of group action) of definition 24.3 (Morphisms of G - ρ -model spaces) on page 150, so $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 . {lem:Grhomodtrivmorph}
{def:Grhomorph}
3. If $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 and $(f_E, f_X \times f_Y)$ is an $\mathbf{E}^1$ - $\mathbf{E}^2$ m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 , then let $u^1 \in E^1$ be arbitrary, $u^2 \stackrel{\text{def}}{=} f_E(u^1)$ and $(U^i, V^i \times Y, \phi^i) \in \mathbf{A}^i$ be a chart at u^i . Let the chart

$(U'^1, V'^1 \times Y, \phi^1|_{V'^1 \times Y}) \in \mathbf{A}^1$ at u^1 , the open set $U'^2 \subseteq U^2$ and the chart $(U'^2, \hat{V}'^2 \times Y, \phi'^2|_{\hat{V}'^2 \times Y}) \in \mathbf{A}^2$ at $f_E(u^1)$ be as in definition 13.1, i.e., $f_E[U'^1] \subseteq U'^2$ and diagram (13.1) is commutative, as shown in fig. 7 (Completed m-atlas morphism) on page 62, with relabelling as in fig. 18 below.




Figure 18: Completed Bundle-atlas morphism with projection

All four functions are continuous: $f_E, f_X \times f_Y$ and f_G are continuous by definition 13.1 (M-atlas morphisms for model spaces) on page 61 and definition 24.3 (Morphisms of G - ρ -model spaces) on page 150. f_X and f_Y are continuous because $f_X \times f_Y$ is continuous.

f_G is a homomorphism: f_G is a homomorphism by definition 24.3 (Morphisms of G - ρ -model spaces) on page 150.

f commutes with π^i and ρ^i :

$\pi^2 \circ f_E = \pi_1 \circ \phi^2 \circ f_E = f_X \circ \pi_1 \circ \phi^1 = f_X \circ \pi^1$ on U'^1 . Since u is arbitray, $\pi^2 \circ f_E = f_X \circ \pi^1$ on all of E^1 .

$$(b) \left(\bigvee_{\substack{y \in Y^1 \\ g \in G^1}} f_Y(y) \star^2 f_G(g) = f_Y(y \star^1 g) \right) \text{ by eq. (24.3) in definition 24.3}$$

(Morphisms of G - ρ -model spaces) on page 150.

Commutation relation of charts: The commutativity of definition 27.1 follows from that of fig. 18.

□

Corollary 27.4 (Bundle-atlas morphisms). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2, 3$, be a protobundle, $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$ and $\mathbf{f}^i \stackrel{\text{def}}{=} (f_E^i: E^i \rightarrow E^{i+1}, f_X^i: X^i \rightarrow X^{i+1}, f_Y^i: Y^i \rightarrow Y^{i+1}, f_G^i: G^i \rightarrow G^{i+1})$ a $\mathbf{B}^i$ - $\mathbf{B}^{i+1}$ bundle-atlas morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$.

$f^2 \circ f^1$ is a bundle-atlas morphism from A^1 to A^3 .

Proof. $f_X^1 \times f_Y^1 \circ f_X^2 \times f_Y^2$ is a G^1 - G^3 - π^1 - π^3 morphism from X^1, Y^1 to X^3, Y^3 by item 4 of lemma 24.4 (Morphisms of G - ρ -model spaces) on page 151 and $(f_E^1, f_X^1 \times f_Y^1) \circ (f_E^2, f_X^2 \times f_Y^2)$ is a E^1 - E^3 m-atlas morphism from A^1 to A^3 in the coordinate space X^3, Y^3 by item 1 of lemma 13.13 (Composition of m-atlas morphisms) on page 79. $\square$

Lemma 27.5 (Composition of equivalent bundle-atlas morphisms). Let

1. $B^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle
2. A^i be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$
3. $f^i \stackrel{\text{def}}{=} (f_E^i: E^i \rightarrow E^{i+1}, f_X^i: X^i \rightarrow X^{i+1}, f_Y^i: Y^i \rightarrow Y^{i+1}, f_G^i: G^i \rightarrow G^{i+1})$ and $g^i \stackrel{\text{def}}{=} (g_E^i: E^i \rightarrow E^{i+1}, g_X^i: X^i \rightarrow X^{i+1}, g_Y^i: Y^i \rightarrow Y^{i+1}, g_G^i: G^i \rightarrow G^{i+1})$, $i = 1, 2$, be bundle equivalent B^1 - B^{i+1} bundle-atlas morphisms from A^i to A^{i+1}

Then $f^2 \circ f^1$ is bundle equivalent to $g^2 \circ g^1$ as a B^1 - B^3 bundle-atlas morphism from A^1 to A^3 .

Proof. $f_E^1 = g_E^1$, and $f_E^2 = g_E^2$, hence $f_E^2 \circ f_E^1 = g_E^2 \circ g_E^1$. $f_X^1 = g_X^1$, and $f_X^2 = g_X^2$, hence $f_X^2 \circ f_X^1 = g_X^2 \circ g_X^1$. $f_Y^1 = g_Y^1$, and $f_Y^2 = g_Y^2$, hence $f_Y^2 \circ f_Y^1 = g_Y^2 \circ g_Y^1$. $f_G^1 = g_G^1$, and $f_G^2 = g_G^2$, hence $f_G^2 \circ f_G^1 = g_G^2 \circ g_G^1$. $\square$

27.2 Categories of bundle atlases and functors

Definition 27.6 (Sets of bundle atlas morphisms). Let

$B^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $B \stackrel{\text{def}}{=} \{B^\alpha | \alpha < A\}$ be a set of protobundles.

$$\mathcal{Atl}_{\text{Ar}}^{\text{Bun}}(B^\alpha, B^\beta) \stackrel{\text{def}}{=} \left\{ (f, (A^\alpha, B^\alpha), (A^\beta, B^\beta)) \mid \text{isAtl}_{\text{Ar}}^{\text{Bun}}(A^\alpha, B^\alpha, A^\beta, B^\beta, f) \right\} \quad (27.3)$$

$$\mathcal{Atl}_{\text{Ar}}^{\text{Bun}}(B^\alpha) \stackrel{\text{def}}{=} \left\{ (f, (A^\alpha, B^\alpha), (A^\alpha, B^\alpha)) \mid \text{isAtl}_{\text{Ar}}^{\text{Bun}}(A^\alpha, B^\alpha, A^\alpha, B^\alpha, f) \right\} \quad (27.4)$$

$$\mathcal{Att}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \bigcup_{\substack{\mathbf{B}^\mu \in \mathbf{B} \\ \mathbf{B}^\nu \in \mathbf{B}}} \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\mu, \mathbf{B}^\nu) \quad (27.5)$$

{def:Bun-Atlcat}

Definition 27.7 (Categories of bundle atlases). Let

$\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then

$$\mathcal{Att}_{(\text{max})}^{\text{Bun}}(\mathbf{B}^\alpha) \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}^\alpha), \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}^\alpha, \mathbf{B}^\alpha), \overset{A}{\circ} \right) \quad (27.6)$$

$$\mathcal{Att}_{(\text{max})}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \left(\mathcal{Att}_{\text{Ob}}^{\text{Bun}}(\mathbf{B}), \mathcal{Att}_{\text{Ar}}^{\text{Bun}}(\mathbf{B}), \overset{A}{\circ} \right) \quad (27.7)$$

{lem:Bun-ATLiscat}

Lemma 27.8 ($\mathcal{Att}^{\text{Bun}} \mathbf{B}$ is a category). Let $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle and $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles. Then $\mathcal{Att}_{(\text{max})}^{\text{Bun}} \mathbf{B}$ is a category.

Let $(\mathbf{A}^\alpha, \mathbf{B}^\alpha) \in \mathcal{Att}_{\text{Ob}}^{\text{Bun}} \mathbf{B}$. Then $\text{Id}_{(\mathbf{A}^\alpha, \mathbf{B}^\alpha)}$ is the identity morphism for $(\mathbf{A}^\alpha, \mathbf{B}^\alpha)$.

Proof. Let $(\mathbf{A}^i, \mathbf{B}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ and let

$m^i \stackrel{\text{def}}{=} (f^i, (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^{i+1}, \mathbf{B}^{i+1}))$ be a morphism of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$. Then

Composition: $(f^2 \circ f^1, (\mathbf{A}^1, \mathbf{B}^1, \mathbf{C}^1), (\mathbf{A}^3, \mathbf{B}^3, \mathbf{C}^3))$ is a morphism of $\mathcal{Att}^{\text{Bun}} \mathbf{B}$ by corollary 27.4 (Bundle-atlas morphisms) on page 167. {cor:Bun-ATLmorph}

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Identity: $\text{Id}_{(\mathbf{A}^i, \mathbf{B}^i, \mathbf{C}^i)}$ is an identity morphism by lemma 1.19.

{lem:labcomp}

□

{def:FuncBuntoM}

Definition 27.9 (Functor from Bundle atlases to m-atlases). Let

$\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $i = 1, 2$, be a protobundle, let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$ and let

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \longrightarrow E^2, f_X: X^1 \longrightarrow X^2, f_Y: Y^1 \longrightarrow Y^2, f_G: G^1 \longrightarrow G^2)$ be a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$. Then

$$\mathcal{F}_{\text{Bun}, \text{M}}^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i) \stackrel{\text{def}}{=} \left(\mathbf{A}^i, \begin{matrix} E^i \\ X^i - \pi^i \text{triv} \end{matrix}, \begin{matrix} X^i, Y^i \\ G^i - \rho^i \text{triv} \end{matrix} \right) \quad (27.8)$$

$$\mathcal{F}_{\text{Bun}, \text{M}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} \left((f_E, f_X \times f_Y), \left(\mathbf{A}^1, \begin{matrix} E^1 \\ X^1 - \pi^1 \text{triv} \end{matrix}, \begin{matrix} C^1 \\ G^1 - \rho^1 \text{triv} \end{matrix} \right), \left(\mathbf{A}^2, \begin{matrix} E^2 \\ X^2 - \pi^2 \text{triv} \end{matrix}, \begin{matrix} C^2 \\ G^2 - \rho^2 \text{triv} \end{matrix} \right) \right) \quad (27.9)$$

{the:FuncBuntoM}

Theorem 27.10 (Functor from Bundle atlases to m-atlases). Let $\mathbf{E}$ be a set of topological spaces, $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha \in \mathbf{E}, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$, $\alpha < A$, be a protobundle, $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$ be a set of protobundles and $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha, Y^\alpha)_{G^\alpha - \rho^\alpha - \text{triv}}$. Then $\mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}}$ is a

functor from $\mathcal{Att}_{(\text{max})}^{\text{Bun}} \mathbf{B}$ to $\mathcal{Att}_{(\text{max})} \left(\frac{\mathbf{E}}{\mathcal{T}\text{riv}}, \frac{\mathbf{B}}{\text{Bun} - \mathcal{T}\text{riv}} \right)$ and a functor from

$\mathcal{Att}_{(\text{max})}^{\text{Bun}} \mathbf{B}$ to $\mathcal{Att}_{(\text{max})} \left(\frac{\mathbf{E}}{\mathcal{T}\text{riv}}, \frac{\mathbf{B}}{\text{Bun} - \text{prod} - \mathcal{T}\text{riv}} \right)$.

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$, $i \in [1, 3]$, be an object of $\mathcal{Att}^{\text{Bun}}(\mathbf{B})$, $o'^i \stackrel{\text{def}}{=} \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} o^i = \left(\mathbf{A}^i, \frac{E^i}{X^i - \pi^i \text{triv}}, \frac{C^i}{G^i - \rho^i \text{triv}} \right)$ be the corresponding object of $\mathcal{Att}_{(\text{max})} \left(\frac{\mathbf{E}}{\mathcal{T}\text{riv}}, \frac{\mathbf{B}}{\text{Bun} - \mathcal{T}\text{riv}} \right)$, $m^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o^i, o^{i+1})$, $i = 1, 2$, be a morphism of $\mathcal{Att}^{\text{Bun}}(\mathbf{B})$ from o^i to o^{i+1} and

$$\begin{aligned} m'^i &\stackrel{\text{def}}{=} \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} m^i \\ &= \left((f_E^i, f_X^i \times f_Y^i), \left(\mathbf{A}^i, \frac{E^i}{X^i - \pi^i \text{triv}}, \frac{C^i}{G^i - \rho^i \text{triv}} \right), \left(\mathbf{A}^{i+1}, \frac{E^{i+1}}{X^{i+1} - \pi^{i+1} \text{triv}}, \frac{C^2}{G^{i+1} - \rho^{i+1} \text{triv}} \right) \right) \\ &= ((f_E^i, f_X^i \times f_Y^i), o'^i, o'^{i+1}) \end{aligned}$$

be the corresponding morphism of $\mathcal{Att}_{(\text{max})} \left(\frac{\mathbf{E}}{\mathcal{T}\text{riv}}, \frac{\mathbf{B}}{\text{Bun} - \mathcal{T}\text{riv}} \right)$.

Preservation of endpoints: $\frac{E^i}{X^i - \pi^i \text{triv}} \stackrel{\text{Ob}}{\in} \frac{\mathbf{E}}{\mathcal{T}\text{riv}} \cdot \frac{C^i}{G^i - \rho^i \text{triv}} \stackrel{\text{Ar}}{\in} \frac{\mathbf{B}}{\text{Bun} - \text{prod} - \mathcal{T}\text{riv}}.$

By theorem 27.10 (Functor from Bundle atlases to m-atlases) on page 170, $(f_E, f_X \times f_Y)$ is a $\frac{E^i}{X^i - \pi^i \text{triv}} - \frac{E^{i+1}}{X^{i+1} - \pi^{i+1} \text{triv}}$ m-atlas morphism from $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\frac{X^i, Y^i}{G^i - \rho^i \text{triv}}, \frac{X^{i+1}, Y^{i+1}}{G^{i+1} - \rho^{i+1} \text{triv}}$. {the:FuncBuntoM}

$$\begin{aligned} \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} m^i &= \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} (\mathbf{f}^i, o^i, o^{i+1}) \\ &= ((f_E^i, f_X^i \times f_Y^i), o'^i, o'^{i+1}) \end{aligned}$$

Identity:

$$\begin{aligned} \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} &= \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), o^i, o^i) \\ &= ((\text{Id}_{\frac{E^i}{X^i - \pi^i \text{triv}}}, \text{Id}_{C^i}), o'^i, o'^i) \\ &= ((\text{Id}_{\frac{E^i}{X^i - \pi^i \text{triv}}}, \text{Id}_{C^i}), \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} o^i, \mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} o^i) \\ &= \text{Id}_{\mathcal{F}_{\text{Bun}, \mathbf{M}}^{\text{Bun}} o^i} \end{aligned}$$

Composition:

$$\begin{aligned}
m^2 \overset{A}{\circ} m^1 &= ((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), o^1, o^3) \\
\mathcal{F}_{\text{Bun}, M}^{\text{Bun}}(m^2 \circ m^1) &= ((f_E^2 \circ f_E^1, (f_X^2 \circ f_X^1) \times (f_Y^2 \circ f_Y^1)), o'^1, o'^3) \\
&= ((f_E^2 \circ f_E^1, (f_X^2 \times f_Y^2) \circ (f_X^1 \times f_Y^1)), o'^1, o'^3) \\
&= \mathcal{F}_{\text{Bun}, M}^{\text{Bun}} m^2 \overset{A}{\circ} \mathcal{F}_{\text{Bun}, M}^{\text{Bun}} m^1
\end{aligned}$$

□

{lem:Bun-Atlbasmorph}

Lemma 27.11 (Base space functions derived from bundle-atlas morphisms). Let $E^i, i = 1, 2$, be a model space, X^i, Y^i topological spaces, G^i a topological group, ρ^i an effective action of G^i on Y^i , $\mathbf{C}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i)$ a G^i - ρ^i model space of $X^i \times Y^i$, $f_C: \mathbf{C}^1 \rightarrow \mathbf{C}^2$ a G^1 - G^2 - ρ^1 - ρ^2 morphism of $X^1 \times Y^1$ to $X^2 \times Y^2$, i.e., a model function that preserves group action, A^i an m-atlas of E^i in the coordinate model space $\mathbf{C}^i$ and $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_C: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$.

Then there exists a unique function $f_X: X^1 \rightarrow X^2$ such that $f_X \circ \pi_1 = \pi_1 \circ f_C$.

Proof. Let x in X^1, y, y' in Y^1 , Then f_C preserves fibers, i.e.,

$\pi_1(f_C(x, y)) = \pi_1(f_C(x, y'))$. by lemma 24.4 (Morphisms of G - ρ -model spaces)

{lem:Grhomorph}

on page 150. Define $f_X(x) \stackrel{\text{def}}{=} \pi_1(f_C(x, y))$

□

Remark 27.12. There need not exist $f_Y: Y^1 \rightarrow Y^2$ such that $f_C = f_X \times f_Y$.

{def:FuncMtoBun}

Definition 27.13 (Functor from m-atlases to Bundle atlases). Let $\mathfrak{E}$ be a trivial model category, $\mathcal{C}$ be a trivial product coordinate model category with objects

$\{\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}\mathcal{Y}^\alpha) \mid \alpha < A\}$, G a group valued function on $\text{Ob}(\mathcal{C})$ and ρ a function

valued function on $\text{Ob}(\mathcal{C})$ such that for every $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}\mathcal{Y}^\alpha) \in \mathcal{C}$, $\rho(\mathbf{C}^\alpha)$ is an effective action of $G(\mathbf{C}^\alpha)$ on Y^α and $X^\alpha, Y^\alpha \stackrel{\text{def}}{=} \mathbf{C}^\alpha$.

$G(\mathbf{C}^\alpha) = \rho(\mathbf{C}^\alpha) = \text{triv}$

Let $E^i \stackrel{\text{def}}{=} (E^i, \mathcal{E}^i) \in \mathfrak{E}, i = 1, 2$, be a trivial model space, $\mathbf{C}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}\mathcal{Y}^i) \in \mathcal{C}$

$G^i \stackrel{\text{def}}{=} G(\mathbf{C}^i), \rho^i \stackrel{\text{def}}{=} \rho(\mathbf{C}^i), A^i$ a full m-atlas of E^i in the coordinate space $\mathbf{C}^i, \pi^i: E^i \rightarrow$

X^i the unique function asserted in lemma 26.11 and $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$. Then define

{lem:mGrho}

$$\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}(A^i, E^i, \mathbf{C}^i) \stackrel{\text{def}}{=} (A^i, \mathbf{B}^i) \quad (27.10)$$

Let $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_C \stackrel{\text{def}}{=} f_X \times f_Y: \mathbf{C}^1 \rightarrow \mathbf{C}^2)$ be an E^1 - E^2 m-atlas morphism of A^1 to A^2 in the coordinate spaces $\mathbf{C}^1, \mathbf{C}^2$ that preserves the group action,

$f_G: G^1 \longrightarrow G^2$ the unique function asserted to exist in eq. (24.3) (preservation of group action) of definition 24.3 (Morphisms of G - ρ -model spaces) on page 150. Then

$$\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{E}^1, \mathbf{C}^1), (\mathbf{A}^2, \mathbf{E}^2, \mathbf{C}^2)) \stackrel{\text{def}}{=} ((f_E, f_X, f_Y, f_G), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \quad (27.11)$$

Theorem 27.14 (Functor from m-atlases to bundle atlases). Let

1. $\mathcal{C}$ be a trivial product coordinate model category,
2. $\mathcal{E}$ be a model category,
3. G be a group valued function on $\text{Ob}(\mathcal{C})$
4. ρ be a function valued function on $\text{Ob}(\mathcal{C})$ such that for every $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (X^\alpha \times Y^\alpha, \mathcal{X}^\alpha \mathcal{Y}^\alpha) \in \mathcal{C}$, $\rho(\mathbf{C}^\alpha)$ is an effective action of $G(\mathbf{C}^\alpha)$ on Y^α and $\mathbf{C}^\alpha \stackrel{\text{def}}{=} \frac{X^\alpha, Y^\alpha}{G(\mathbf{C}^\alpha) - \rho(\mathbf{C}^\alpha) - \text{triv}}$.

5. π be the unique function valued function on $\mathcal{A}t\ell_{\text{full}}^{\text{Ob}}(\mathcal{E}, \mathcal{C})$ such that for every

$$(\mathbf{A}^\alpha, (E^\alpha, \mathcal{E}^\alpha), (\mathbf{C}^\alpha, \mathcal{C}^\alpha)) \in \mathcal{A}t\ell_{\text{full}}^{\text{Ob}}(\mathcal{E}, \mathcal{C}) \text{ and every } (U, V, \phi) \in \mathbf{A}^\alpha, \pi_1 \circ \phi = \pi(\mathbf{A}^\alpha) \downarrow_U$$

$$6. \mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, G(\mathbf{C}^\alpha), \pi(\mathbf{A}^\alpha), \rho(\mathbf{C}^\alpha)) \mid (\mathbf{A}^\alpha, (E^\alpha, \mathcal{E}^\alpha), (\mathbf{C}^\alpha \stackrel{\text{def}}{=} X^\alpha \times Y^\alpha, \mathcal{C}^\alpha)) \in \mathcal{A}t\ell_{\text{full}}^{\text{Ob}}(\mathcal{E}, \mathcal{C}) \right\}$$

Then $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}}$ is a functor from $\mathcal{A}t\ell_{\mathcal{T}\text{riv}, \text{Bun-prod-}\mathcal{T}\text{riv}}^{\text{Bun}}(\mathbf{E}, \mathbf{B})$ to $\mathcal{A}t\ell^{\text{Bun}}(\mathbf{B})$.

Proof. Let $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{E}^i, \mathbf{C}^i \stackrel{\text{def}}{=} (X^i \times Y^i, \mathcal{X}^i \mathcal{Y}^i)) \in \mathcal{M}$, $i \in [1, 3]$, $G^i \stackrel{\text{def}}{=} G(o^i)$, $\pi^i \stackrel{\text{def}}{=} \pi(o^i)$, $\rho^i \stackrel{\text{def}}{=} \rho(o^i)$, $\mathbf{A}^i$ an m-atlas of E^i in the coordinate model space $\mathbf{C}^i$, $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i)$, $o'^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$, $m^i \stackrel{\text{def}}{=} (\mathbf{f}^i, o^i, o^{i+1}) \stackrel{\text{Ar}}{\in} \mathcal{M}$, $i = 1, 2$, an E^i - E^{i+1} m-atlas morphism of $\mathbf{A}^i$ to $\mathbf{A}^{i+1}$ in the coordinate spaces $\mathbf{C}^i, \mathbf{C}^{i+1}$ that preserves the group action, $f_G: G^i \longrightarrow G^{i+1}$ the unique function asserted to exist in eq. (24.3) (preservation of group action) of definition 24.3 (Morphisms of G - ρ -model spaces) on page 150 and $\pi^i: E^i \rightarrow X^i$ the unique function asserted in lemma 26.11.

Let $f_G^i: G^i \longrightarrow G^{i+1}$ be the function asserted by eq. (24.3) (Morphisms of G - ρ -model spaces) on page 150; let $f_1^i = f_X^i \times f_Y^i$ be the decomposition given by definition 24.7 (Trivial G - ρ -model spaces) on page 152, and let $m'^i \stackrel{\text{def}}{=} \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^i = (\mathbf{f}^i, (f_0^i f_X^i f_Y^i f_G^i), o'^i, o'^{i+1})$.

Preservation of endpoints: E^i , X^i and Y^i are topological spaces.

G^i is a topological group,

$\pi^i: E^i \twoheadrightarrow X^i$ is a continuous surjection.

$\rho^i \stackrel{\text{def}}{=} \rho(\mathbf{C}^i)$ is an effective action of $G^i \stackrel{\text{def}}{=} G(\mathbf{C}^i)$ on Y^α .

Hence B^i is a protobundle.

Identity:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} &= \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)}((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), o^i, o^i) \\ &= ((\text{Id}_{\substack{E^i \\ X^i - \pi^i \text{triv}}}, \text{Id}_{C^i}), o^i, o^i) \\ &= ((\text{Id}_{\substack{E^i \\ X^i - \pi^i \text{triv}}}, \text{Id}_{C^i}), \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i) \\ &= \text{Id}_{\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i} \end{aligned}$$

Composition:

$\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} m^i$ is a morphism from $\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i$ to $\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^{i+1}$:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(m^i) &= \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(\mathbf{f}^i, o^i, o^{i+1}) \\ &= ((f_0^i, f_X^i, f_Y^i, f_G^i), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^{i+1}, \mathbf{B}^{i+1})) \\ &= ((f_0^i, f_X^i, f_Y^i, f_G^i), \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^{i+1}) \end{aligned}$$

$\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}$ maps identity functions to identity functions:

$$\begin{aligned} \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} \text{Id}_{o^i} &= \\ \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}((\text{Id}_{E^i}, \text{Id}_{C^i}), o^i, o^i) &= \\ ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) &= \\ ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i, \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i) &= \\ \text{Id}_{\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} o^i} \end{aligned} \tag{27.12}$$

$\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} m^2 \stackrel{A}{\circ} \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}} m^1 = \mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(m^2 \stackrel{A}{\circ} m^1)$:

1. $m^2 \stackrel{A}{\circ} m^1 = ((f_0^2 \circ f_0^1, f_1^2 \circ f_1^1), o^1, o^3)$
2. $\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(o^i) = (\mathbf{A}^i, \mathbf{B}^i)$
3. $\mathcal{F}_{\mathbf{M}-G-\rho, \text{Bun}}^{\text{Bun}}(m^i) = ((f_0^i, f_X^i, f_Y^i, f_G^i), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^{i+1}, \mathbf{B}^{i+1}))$

4. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^2 \overset{A}{\circ} \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} m^1 =$
 $((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), (A^1, B^1), (A^3, B^3))$
5. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (m^2 \circ m^1) =$
 $((f_0^2 \circ f_0^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), (A^1, B^1), (A^3, B^3))$

□

$$\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} \circ \mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} = \text{Id}.$$

Proof. Expanding the definitions, we have

1. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} (A^i, E^i, C^i) = (A^i, B^i)$
2. $\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} (A^i, B^i) = (A^i, \overset{E^i}{X^i - \pi^i \text{triv}}, \overset{X^i, Y^i}{G^i - \rho^i \text{triv}}) (A^i, E^i, C^i)$, since by definition 27.13, {def:FuncMtoBun}
 E^i and C^i are trivial.
3. $\mathcal{F}_{M-G-\rho, \text{Bun}}^{\text{Bun}} ((f_E^1, f_X^1 \times f_Y^1), (A^1, E^1, C^1), (A^2, E^2, C^2)) =$
 $((f_E^1, f_X^1, f_Y^1, f_G^1), (A^1, B^1), (A^2, B^2))$
4. $\mathcal{F}_{\text{Bun}, M}^{\text{Bun}} ((f_E^1, f_X^1, f_Y^1, f_G^1), (A^1, B^1), (A^2, B^2)) =$
 $((f_E^1, f_X^1 \times f_Y^1), (A^1, \overset{E^1}{X^1 - \pi^1 \text{triv}}, \overset{C^1}{G^1 - \rho^1 \text{triv}}), (A^2, \overset{E^2}{X^2 - \pi^2 \text{triv}}, \overset{C^2}{G^2 - \rho^2 \text{triv}})) =$
 $((f_E^1, f_X^1 \times f_Y^1), (A^1, E^1, C^1), (A^2, E^2, C^2))$, since by definition 27.13, E^i and C^i are trivial. {def:FuncMtoBun}

□

28 Fiber bundles

Conventionally a fiber bundle is different from its atlases, but definition 27.7 (Categories of bundle atlases) on page 169 encourages treating them on an equal footing. All of the results for maximal bundle atlases carry directly over to results for fiber bundles.

{sec:fib}
{def:Bun-Atlcat}
{def:Bun}

Definition 28.1 (fiber bundles). $(E, X, Y, \pi, G, \rho, A)$ is a weak fiber bundle iff

1. E, X and Y are topological spaces
2. $\pi: E \rightarrow X$ is surjective,
3. G is a topological group,
4. $\rho: Y \times G \rightarrow Y$ is an effective right action of G on Y
5. A is a π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$.

$(E, X, Y, \pi, G, \rho, \mathbf{A})$ is also a proper fiber bundle, or just fiber bundle, iff $\mathbf{A}$ is a maximal π - G - ρ -bundle atlas of E in the coordinate space $X \times Y$.

Let

1. $E^\alpha, X^\alpha, Y^\alpha$ be topological spaces
2. G^α be a topological group,
3. $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ be surjective
4. $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ be an effective right action of G^α on Y^α
5. $\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha) \mid \alpha < A \right\}$

Then

$$\text{Bun}_{\text{Ob}}^{\text{def}} \mathbf{B} \stackrel{\text{def}}{=} \mathcal{A} \mathcal{L}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \quad (28.1)$$

{lem:Bun}

Lemma 28.2 (fiber bundles). $(E, X, Y, \pi, G, \rho, \mathbf{A})$ is a weak fiber bundle iff

1. E, X, Y are topological spaces
2. G is a topological group,
3. $\pi: E \rightarrow X$ is surjective
4. $\rho: Y \times G \rightarrow Y$ an effective right action of G on Y
5. $\mathbf{A}$ is an m-atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\underset{G-\rho-\text{triv}}{X, Y}$
6. Every coordinate function in $\mathbf{A}$ preserves fibers.

Proof. The result follows from definition 28.1 and lemma 26.3 (Bundle atlases) on page 162. {def:Bun}Atl □

$(E, X, Y, \pi, G, \rho, \mathbf{A})$ is also a proper fiber bundle iff $\mathbf{A}$ is a maximal m-atlas of $\underset{\text{triv}}{E}$ in the coordinate space $\underset{G-\rho-\text{triv}}{X, Y}$.

Proof. The result follows from lemma 26.9 (Maximal bundle atlases) on page 163 {lem:Bun-ATLmax} □

29 Bundle maps, categories of fiber bundles and functors

{sec:Bunmap}

29.1 Bundle maps

{sub:Bunmap}

29.1.1 Definition of Bundle maps

{def:Bun-map}

Definition 29.1 (Bundle maps). Let

1. $E^\alpha, X^\alpha, Y^\alpha$ be topological spaces
2. G^α be a topological group,
3. $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ be surjective
4. $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ be an effective right action of G^α on Y^α
5. $\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha) \mid \alpha < A \right\}$

Then

$$\text{Bun}_{\text{Ar}} \mathbf{B} \stackrel{\text{def}}{=} \text{Att}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \quad (29.1)$$

$$\text{Bun} \mathbf{B} \stackrel{\text{def}}{=} (\text{Bun}_{\text{Ob}} \mathbf{B}, \text{Bun}_{\text{Ar}} \mathbf{B}, \circ) \quad (29.2)$$

Let $(\mathbf{A}^i, \mathbf{B}^i) \in \text{Bun}_{\text{Ob}} \mathbf{B}, i = 1, 2$. Then

$\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a bundle map from $(\mathbf{A}^1, \mathbf{B}^1)$ to $(\mathbf{A}^2, \mathbf{B}^2)$ iff it is a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from A^1 to A^2 . The identity morphism for $(\mathbf{A}^i, \mathbf{B}^i)$ is

$$\text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} \stackrel{\text{def}}{=} ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) \quad (29.3)$$

29.1.2 Proclamations on Bundle maps

{lem:Bun-map}

Lemma 29.2 (Bundle maps). Let $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i), i = 1, 2$, be a protobundle and let $\mathbf{A}^i$ be a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$. Then $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \rightarrow E^2, f_X: X^1 \rightarrow X^2, f_Y: Y^1 \rightarrow Y^2, f_G: G^1 \rightarrow G^2)$ is a bundle map from $(\mathbf{A}^1, \mathbf{B}^1)$ to $(\mathbf{A}^2, \mathbf{B}^2)$ iff $f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to G^1 - ρ^1 -triv

X^2, Y^2 and $(f_E, cf_X \times f_Y)$ is a X^1 - π^1 -triv X^2 - π^2 -triv m -atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 . G^1 - ρ^1 -triv G^2 - ρ^2 -triv

Proof. The result follows from definition 29.1 above and lemma 27.3 (Bundle-atlas morphisms) on page 166. □

29.2 Categories of fiber bundles and functors

sub:Fib-cat {the:Bun}

Theorem 29.3 (Categories of fiber bundles). Let

$\mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \mid \alpha < A \right\}$, where $E^\alpha, X^\alpha, Y^\alpha$ are topological spaces, G^α a topological group, $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ surjective and $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ an effective right action of G^α on Y^α . Then $\text{Bun}\mathbf{B}$ is a category and $\text{Id}_{\mathbf{B}^\alpha}$ is the identity morphism for $\mathbf{B}^\alpha$.

Proof. The result follows directly from $\stackrel{\text{def}}{\text{definition 28.1 (fiber bundles)}}$, $\stackrel{\text{def}}{\text{definition 29.1 (Bundle maps)}}$ and lemma 27.8 ($\text{Att}^{\text{Bun}}\mathbf{B}$ is a category) on page 169. $\square$ {def:Bun-map} {lem:Bun-ATLiscat}

29.2.1 Functors for fiber bundles

def:BunLCS

Definition 29.4 (Functor from fiber bundles to Local Coordinate Spaces). Let

1. $\mathbf{E}, \mathbf{X}$ and $\mathbf{Y}$ be sets of topological spaces
2. $\mathbf{G}$ be a set of topological groups
3. $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha \in \mathbf{E}, X^\alpha \in \mathbf{X}, Y^\alpha \in \mathbf{Y}, G^\alpha \in \mathbf{G}, \pi^\alpha, \rho^\alpha), \alpha < A$, be a protobundle
4. $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha \mid \alpha < A\}$ be a set of protobundles
5. $\mathbf{C}^\alpha \stackrel{\text{def}}{=} X^\alpha \times Y^\alpha$,
6. $\mathcal{E}, \mathcal{X}, \mathcal{Y}$ and $\mathcal{G}$ be model categories
7. $\mathcal{X}\mathcal{Y}\mathcal{G}\rho$ be a G - ρ -model category
8. $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{E}, \mathcal{X}\mathcal{Y}\mathcal{G}\rho, \mathcal{X}, \mathcal{Y}, \mathcal{G})$,
9. $\mathcal{M}_{\text{triv}} \stackrel{\text{def}}{=} \left(\mathbf{E}_{\mathcal{T}\text{riv}}, \mathbf{B}_{\text{Bun-}\mathcal{T}\text{riv}}, \mathbf{X}_{\mathcal{T}\text{riv}}, \mathbf{Y}_{\mathcal{T}\text{riv}}, \mathbf{G}_{\mathcal{T}\text{riv}} \right)$
10. $\mathcal{M}_{\text{triv}} \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}$
11. $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i) \in \mathbf{B}, i = 1, 2$. with
 - (a) $\star^i: G^i \times G^i \rightarrow G^i$ the group operation for G^i
 - (b) $\mathbf{A}^i$ a maximal π^i - G^i - ρ^i -bundle atlas of E^i in the coordinate space $\mathbf{C}^i \stackrel{\text{def}}{=} X^i \times Y^i$
 - (c) $\pi_X^i \stackrel{\text{def}}{=} \pi_1: X^i \times Y^i \rightarrow X^i$
 - (d) $\pi_Y^i \stackrel{\text{def}}{=} \pi_2: X^i \times Y^i \rightarrow Y^i$
 - (e) $\mathbf{o}^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$

- (f) $\mathbf{F}^i \stackrel{\text{def}}{=} (\pi^i, \pi_X^i, \pi_Y^i, \star^i, \rho^i)$
- (g) $\mathcal{M}_{\text{triv}}^i \stackrel{\text{def}}{=} \left(\begin{array}{c} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{array}, \begin{array}{c} X^i, Y^i \\ G^i - \rho^i - \text{triv} \\ \text{Sing} \end{array}, \begin{array}{c} X^i, Y^i, G^i \\ \text{Sing} \end{array} \right)$
- (h) $\mathbf{M}^i \stackrel{\text{def}}{=} \left(\begin{array}{c} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{array}, \begin{array}{c} X^i, Y^i \\ G^i - \rho^i - \text{triv} \\ \text{Sing} \end{array}, X^i, Y^i, G^i \right)$
- (i) $\Sigma \stackrel{\text{def}}{=} ((0, 2), (1, 2), (1, 3), (4, 4, 4), (3, 4, 3))$
- (j) $\mathbf{L}^i \stackrel{\text{def}}{=} \left(\begin{array}{c} \mathcal{M}^i \\ \text{triv} \\ \text{Sing} \end{array}, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma \right)$
- (k) $\mathbf{L}^i, \mathcal{M} \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma)$
- (l) $\mathbf{f} \stackrel{\text{def}}{=} (f_E: E^1 \longrightarrow E^2, f_X: X^1 \longrightarrow X^2, f_Y: Y^1 \longrightarrow Y^2, f_G: G^1 \longrightarrow G^2)$ a $\mathbf{B}^1$ - $\mathbf{B}^2$ bundle-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$

Then

$$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i) \stackrel{\text{def}}{=} \mathbf{L}^i \quad (29.4)$$

$$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} ((f_E, f_X \times f_Y, f_X, f_Y, f_G), \mathbf{L}^1, \mathbf{L}^2) \quad (29.5)$$

$$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{A}^i, \mathbf{B}^i) \stackrel{\text{def}}{=} \mathbf{L}^i, \mathcal{M} \quad (29.6)$$

$$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} ((f_E, f_X \times f_Y, f_X, f_Y, f_G), \mathbf{L}^1, \mathcal{M}, \mathbf{L}^2, \mathcal{M}) \quad (29.7)$$

$$\text{LCS}_{\text{Ob}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\mathbf{A}, (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha)) \mid (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \in \mathbf{B} \wedge \text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \right\} \quad (29.8)$$

$$\text{LCS}_{\text{Ar}}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \mid \mathbf{B}^i \in \mathbf{B} \wedge \text{isAtl}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f}) \right\} \quad (29.9)$$

$$\text{LCS}^{\text{Bun}} \mathbf{B} \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}^{\text{Bun}} \mathbf{B}, \text{LCS}_{\text{Ar}}^{\text{Bun}} \mathbf{B}, \circ^A) \quad (29.10)$$

$$\text{LCS}_{\text{Ob}}^{\text{Bun}, \mathcal{M}} \mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{A}, (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha)) \mid \right. \\ \left. (E^\alpha, X^\alpha, Y^\alpha, G^\alpha, \pi^\alpha, \rho^\alpha) \in \mathbf{B} \wedge \text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}, E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha) \right\} \quad (29.11)$$

$$\text{LCS}_{\text{Ar}}^{\text{Bun}, \mathcal{M}} \mathbf{B} \stackrel{\text{def}}{=} \left\{ \mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \mid \right. \\ \left. \mathbf{B}^i \in \mathbf{B} \wedge \text{isAtl}_{\text{Ar}}^{\text{Bun}}(\mathbf{A}^1, \mathbf{B}^1, \mathbf{A}^2, \mathbf{B}^2, \mathbf{f}) \right\} \quad (29.12)$$

$$\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B} \stackrel{\text{def}}{=} (\text{LCS}_{\text{Ob}}^{\text{Bun}, \mathcal{M}} \mathbf{B}, \text{LCS}_{\text{Ar}}^{\text{Bun}, \mathcal{M}} \mathbf{B}, \circ^A) \quad (29.13)$$

{the : BunLCS}

Theorem 29.5 (Functor from fiber bundles to Local Coordinate Spaces). Let

1. E, X and Y be sets of topological spaces, G be a set of topological groups,
2. $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha \in E, X^\alpha \in X, Y^\alpha \in Y, G^\alpha \in Y, \pi^\alpha, \rho^\alpha) \alpha < A$, be a protobundle with group operation $\star^\alpha: G^\alpha \times G^\alpha \rightarrow G^\alpha$
3. $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha \mid \alpha < A\}$ be a set of protobundles
4. $C^\alpha \stackrel{\text{def}}{=} X^\alpha \times Y^\alpha$, $\mathcal{E}, \mathcal{X}, \mathcal{Y}$ and $\mathcal{G}$ be model categories
5. $\mathcal{X}\mathcal{Y}\mathcal{G}\rho$ be a G - ρ -model category,
6. $\mathcal{M} \stackrel{\text{def}}{=} (\mathcal{E}, \mathcal{X}\mathcal{Y}\mathcal{G}\rho, \mathcal{X}, \mathcal{Y}, \mathcal{G})$, $\mathcal{M}_{\text{triv}} \stackrel{\text{def}}{=} (\mathbf{E}_{\text{triv}}, \mathbf{B}_{\text{Bun-Triv}}, \mathbf{X}_{\text{triv}}, \mathbf{Y}_{\text{triv}}, \mathbf{G}_{\text{triv}})$
7. $\mathcal{M}_{\text{triv}}^{\text{full-cat}} \subseteq \mathcal{M}$, $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i) i = 1, 2, 3$, be a protobundle in $\mathbf{B}$ with group operation $\star^i: G^i \times G^i \rightarrow G^i$
8. $\mathbf{A}^i$ be a maximal π^i - G^i - ρ^i -bundle atlas of E^i in the coordinate space $C^i \stackrel{\text{def}}{=} X^i \times Y^i$, i.e., $\text{isAtl}_{\text{Ob}}^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i)$
9. $o^i \stackrel{\text{def}}{=} (\mathbf{A}^i, \mathbf{B}^i)$ be the object corresponding to the fiber bundle $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i, \mathbf{A}^i)$
10. $\pi_X^i \stackrel{\text{def}}{=} \pi_1: X^i \times Y^i \rightarrow X^i$, $\pi_Y^i \stackrel{\text{def}}{=} \pi_2: X^i \times Y^i \rightarrow Y^i$

11. $F^i \stackrel{\text{def}}{=} (\pi^i, \pi_X^i, \pi_Y^i, \star^i, \rho^i)$
12. $\mathcal{M}_{\text{triv}}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i \\ G^i - \rho^i - \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i, G^i \\ \text{Sing} \end{smallmatrix} \right), \mathbf{M}^i \stackrel{\text{def}}{=} \left(\begin{smallmatrix} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i \\ G^i - \rho^i - \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i, G^i \\ \text{Sing} \end{smallmatrix} \right)$
13. $\Sigma \stackrel{\text{def}}{=} ((0, 2), (1, 2), (1, 3), (4, 4, 4), (3, 4, 3))$ be a signature
14. $L^i \stackrel{\text{def}}{=} \left(\mathcal{M}_{\text{triv}}^i, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma \right), L^{i, \mathcal{M}} \stackrel{\text{def}}{=} (\mathcal{M}, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma)$ be local coordinate spaces
15. $\mathbf{f}^i \stackrel{\text{def}}{=} (f_E^i: E^i \longrightarrow E^{i+1}, f_X^i: X^i \longrightarrow X^{i+1}, f_Y^i: Y^i \longrightarrow Y^{i+1}, f_G^i: G^i \longrightarrow G^{i+1})$
and $\mathbf{g}^i \stackrel{\text{def}}{=} (g_E^i: E^i \longrightarrow E^{i+1}, g_X^i: X^i \longrightarrow X^{i+1}, g_Y^i: Y^i \longrightarrow Y^{i+1}, g_G^i: G^i \longrightarrow G^{i+1})$,
 $i = 1, 2$, be bundle maps from $(\mathbf{A}^i, \mathbf{B}^i)$ to $(\mathbf{A}^{i+1}, \mathbf{B}^{i+1})$
16. $\mathbf{f}^{i, \text{LCS}} \stackrel{\text{def}}{=} (f_E^i, f_X^i \times f_Y^i, f_X^i, f_Y^i, f_G^i)$ and $\mathbf{g}^{i, \text{LCS}} \stackrel{\text{def}}{=} (g_E^i, g_X^i \times g_Y^i, g_X^i, g_Y^i, g_G^i), i = 1, 2$

Then:

L^i is a local $\mathcal{M}_{\text{triv}}^i$ - Σ coordinate space and $L^{i, \mathcal{M}}$ is a local $\mathcal{M}$ - Σ coordinate space.

Proof. They satisfy the criteria in definition 15.1 (Local $\mathbf{M}$ - Σ and $\mathcal{M}$ - Σ coordinate spaces) on page 114: {def:LCS}

1. $\mathcal{M}_{\text{triv}}^i$ and $\mathcal{M}$ are sequences of categories by construction.
2. $\mathbf{M}^i \in \mathcal{M}_{\text{triv}}^i \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}_{\text{triv}}$ by construction and $\mathcal{M}_{\text{triv}} \stackrel{\text{full-cat}}{\subseteq} \mathcal{M}$ by hypothesis.
3. $\begin{smallmatrix} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{smallmatrix}$ is a model category by definition 3.1 (Trivial model spaces) on page 21 {def:trivmod}
, $\begin{smallmatrix} X^i, Y^i \\ G^i - \rho^i - \text{triv} \\ \text{Sing} \end{smallmatrix}$ is a model category by definition 24.7 (Trivial G - ρ -model spaces) {def:Grhomodtriv}
on page 152, $\mathcal{E}$ is a model category by hypothesis and $\mathcal{X}\mathcal{Y}\mathcal{E}\rho$ is a model category by definition 24.6 (Categories of G - ρ -model spaces) on page 151. {def:Grhomodcat}
4. $\left(\mathcal{M}_{\text{triv}}^i, \mathbf{M}^i, \Sigma, \mathbf{F}^i \right)$ and $(\mathcal{M}, \mathbf{M}^i, \Sigma, \mathbf{F}^i)$ are prestructures: $\mathbf{F}^i$ has $\mathcal{M}_{\text{triv}}^i$ -signatures {def:trivmod}
 Σ and $\mathbf{F}^i$ has $\mathcal{M}$ -signatures Σ .

5. A^i is a maximal m-atlas of $\begin{smallmatrix} E^i \\ X^i - \pi^i - \text{triv} \end{smallmatrix}$ in $\begin{smallmatrix} X^i, Y^i \\ G^i - \rho^i - \text{triv} \end{smallmatrix}$ by lemma 27.3 (Bundle-atlas morphisms) on page 166. {lem:Bun-ATLmorph}

6. There are no constraint functions. □

$\text{LCS}^{\text{Bun}} \mathbf{B}$ and $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$ are categories and the identity morphism for L^i is $\text{Id}_{L^i} \stackrel{\text{def}}{=} ((\text{Id}_{E^i} \text{Id}_{X^i} \text{Id}_{X^i \times Y^i} \text{Id}_{Y^i} \text{Id}_{G^i}), L^i, L^i)$.

Proof. satisfy the definition of a category:

Composition: $f^2 \circ f^1$ is a bundle map from (A^1, B^1) to (A^1, B^1) by corollary 27.4 (Bundle-atlas morphisms) on page 167. Then {cor:Bun-ATLmorph}

$$\begin{aligned} \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(f^2, o^2, o^3) \circ \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(f^1, o^1, o^2) &= \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(f^2 \circ f^1, o^1, o^3) \\ \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(f^2, o^2, o^3) \circ \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(f^1, o^1, o^2) &= \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(f^2 \circ f^1, o^1, o^3) \end{aligned}$$

Associativity: Composition is associative by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

Unit: Id_{L^i} is an identity morphism by lemma 1.19 (Tuple composition for labeled morphisms) on page 13. {lem:labcomp}

□

$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(f^i, (A^i, B^i), (A^{i+1}, B^{i+1}))$ is a morphism from $\text{LCS}^{\text{Bun}} \mathbf{B}^i$ to $\text{LCS}^{\text{Bun}} \mathbf{B}^{i+1}$ and $\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(f^i, (A^i, B^i), (A^{i+1}, B^{i+1}))$ is a strict morphism from $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}^i$ to $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}^{i+1}$.

Proof.

1. Prestructure morphism:
 f^1 Σ -commutes with F^1, F^2 .

2. m-atlas morphism:
 (f_0, f_1) is an m-atlas morphism from $\begin{smallmatrix} E^1 \\ X^1 - \pi^1 - \text{triv} \end{smallmatrix}, \begin{smallmatrix} C^1 \\ G^1 - \rho^1 - \text{triv} \end{smallmatrix}$ to $\begin{smallmatrix} E^2 \\ X^2 - \pi^2 - \text{triv} \end{smallmatrix}, \begin{smallmatrix} C^2 \\ G^2 - \rho^2 - \text{triv} \end{smallmatrix}$ by lemma 27.3 (Bundle-atlas morphisms) on page 166 {lem:Bun-ATLmorph}

□

$\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}$ is a functor from $\text{Bun} \mathbf{B}$ to $\text{LCS}^{\text{Bun}} \mathbf{B}$ and $\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}$ is a functor from $\text{Bun} \mathbf{B}$ to $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$.

Proof. $\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}$ and $\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}$ satisfy the definition of a functor:

Preservation of endpoints:

$$\begin{aligned}
\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\mathbf{f}^i(A^i, \mathbf{B}^i), (A^{i+1}, \mathbf{B}^{i+1})) &= (\mathbf{f}^{i,\text{LCS}}, L^i, L^{i+1}) \\
\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{f}^i(A^i, \mathbf{B}^i), (A^{i+1}, \mathbf{B}^{i+1})) &= (\mathbf{f}^{i,\text{LCS}}, L^i, \mathcal{M}, L^{i+1}, \mathcal{M}) \\
\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(A^i, \mathbf{B}^i) &= L^i \\
\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(A^i, \mathbf{B}^i) &= L^i, \mathcal{M}
\end{aligned}$$

Composition: This follows from the proof above that $\text{LCS}^{\text{Bun}} \mathbf{B}$ and $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$ are categories.

Identity:

$$\begin{aligned}
\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(\text{Id}_{(A^i, \mathbf{B}^i)}) &= \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (A^i, \mathbf{B}^i), (A^i, \mathbf{B}^i)) \\
&= ((\text{Id}_{E^i}, \text{Id}_{X^i} \times \text{Id}_{Y^i}, \underset{\text{triv}}{X^i}, \underset{\text{triv}}{Y^i}, \underset{\text{triv}}{G^i}), L^i, L^i) \\
&= \text{Id}_{L^i} \\
&= \text{Id}_{\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}}(A^i, \mathbf{B}^i)} \\
&= \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(\text{Id}_{(A^i, \mathbf{B}^i)}) \\
&= \mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (A^i, \mathbf{B}^i), (A^i, \mathbf{B}^i)) = \\
&= ((\text{Id}_{E^i}, \text{Id}_{X^i} \times \text{Id}_{Y^i}, \underset{\text{triv}}{X^i}, \underset{\text{triv}}{Y^i}, \underset{\text{triv}}{G^i}), L^i, \mathcal{M}, L^i, \mathcal{M}) \\
&= \text{Id}_{L^i, \mathcal{M}} \\
&= \text{Id}_{\mathcal{F}_{\text{Fib,LCS}}^{\text{Bun}, \mathcal{M}}(A^i, \mathbf{B}^i)}
\end{aligned}$$

□

$\mathbf{f}^{i,\text{LCS}}$ is $(0, 2, 4)$ -equivalent to $\mathbf{f}^{i,\text{LCS}}$ iff $\mathbf{f}^i$ is bundle equivalent to $\mathbf{g}^i$.

Proof. Both statments are true iff $f_E^i = g_E^i$, $f_X^i = g_X^i$ and $f_G^i = g_G^i$ □

{def:LCS to Bun}

Definition 29.6 (Functor from local coordinate spaces to fiber bundles). Let $\mathcal{E}$ be a model category, $\mathcal{X}\mathcal{Y}\mathcal{G}\rho$ a G - ρ -model category, $\mathcal{X}$, $\mathcal{Y}$ and $\mathcal{G}$ topological categories,

$\mathcal{M}$ a sequence of categories, $\mathbf{M}^i$, $i = 1, 2$, a sequence, $E^i \in \mathcal{E}$, $\mathbf{C}^i = (C^i, \mathcal{C}^i) \in \mathcal{X}\mathcal{Y}\mathcal{G}\rho$,

$X^i \in \mathcal{X}$, $Y^i \in \mathcal{Y}$, $G^i \in \mathcal{G}$, $\pi^i: E^i \rightarrow X^i$, $\pi_X^i: C^i \rightarrow X^i$, $\pi_Y^i: C^i \rightarrow Y^i$, $\star^i: G^i \times G^i \rightarrow G^i$

$\rho^i: Y^i \times G^i \rightarrow Y^i$, $\mathcal{M}_{\text{triv}}^i$ a sequence of categories, $L^i \stackrel{\text{def}}{=} \left(\mathcal{M}_{\text{triv}}^i, \mathbf{M}^i, A^i, F^i, \Sigma \right)$ and

$L^{i, \mathcal{M}} \stackrel{\text{def}}{=} (\mathcal{M}^i, \mathbf{M}^i, A^i, F^i, \Sigma)$ local coordinate spaces, $\mathbf{B}^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, G^i, \pi^i, \rho^i)$ and

$\mathbf{f} \stackrel{\text{def}}{=} (f_\beta, \beta < \text{length}(\mathbf{M}^1))$ a morphism from L^1 to L^2 , satisfying

1. $\text{head}(\mathcal{M}, 5) = (\mathcal{E}, \mathcal{X}\mathcal{Y}\mathcal{G}\rho, \mathcal{X}, \mathcal{Y}, \mathcal{G})$
2. $\text{head}\left(\begin{smallmatrix} \mathcal{M}^i \\ \text{triv} \\ \text{Sing} \end{smallmatrix}, 5\right) = \left(\begin{smallmatrix} E^i \\ X^i - \pi^i \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i \\ G^i - \rho^i \text{triv} \\ \text{Sing} \end{smallmatrix}, \begin{smallmatrix} X^i, Y^i, G^i \\ \text{Sing} \end{smallmatrix}\right)$
3. $\text{head}(\mathbf{M}^i, 5) = (\mathbf{E}^i, \mathbf{C}^i, X^i, Y^i, G^i)$
4. $\mathbf{M}^i \overset{\circ}{\in} \mathcal{M}$
5. $\text{head}(\mathbf{F}^i, 5) = (\pi^i, \pi_X^i, \pi_Y^i, \star^i, \rho^i)$
6. $C^i = X^i \times Y^i$
7. G^i is a topological group with group operation $\star^i$. Subsequent references to G^i should be read as referring to the group rather than just the underlying topological space.
8. π^i is surjective.
9. $\pi_X^i = \pi_1: X^i \times Y^i \twoheadrightarrow X^i$.
10. $\pi_Y^i = \pi_2: X^i \times Y^i \twoheadrightarrow Y^i$.
11. ρ^i is an effective right action of G^i on Y^i .
12. $\text{head}(\Sigma, 5) = ((0, 2), (1, 2), (1, 3), (4, 4, 4), (3, 4, 3))$

Remark 29.7. Due to the commutation requirement, specifying $\mathcal{M}_4 = \mathbf{G}_{\text{group-}\mathcal{T}\text{riv}}$ is not necessary in order to ensure that f_G is a homomorphism.

Then

$$\mathcal{F}_{\text{LCS, Fib}}^{\text{Bun}} L^i \stackrel{\text{def}}{=} (E^i, X^i, Y^i, \pi^i, G^i, \rho^i, \mathbf{A}^i) \quad (29.14)$$

$$\mathcal{F}_{\text{LCS, Fib}}^{\text{Bun}}(\mathbf{f}, (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) \stackrel{\text{def}}{=} ((f_0, f_2, f_3, f_4), \mathcal{F}_{\text{LCS, Fib}}^{\text{Bun}} L^1, \mathcal{F}_{\text{LCS, Fib}}^{\text{Bun}} L^2) \quad (29.15) \quad \{\text{the :LCS to Bun}\}$$

Theorem 29.8 (Functor from local coordinate spaces to fiber bundles). Let

1. $\mathcal{E}$ be a model category
2. $\mathcal{X}\mathcal{Y}\mathcal{G}\rho$ a G - ρ -model category
3. $\mathcal{X}, \mathcal{Y}$ and $\mathcal{G}$ topological categories
4. $\mathcal{M}$ a sequence of categories
5. $\mathbf{E}^\alpha \stackrel{\text{def}}{=} (E^\alpha, \mathcal{E}^\alpha) \overset{\text{Ob}}{\in} \mathcal{E}, \alpha < A$
6. $\mathbf{C}^\alpha \stackrel{\text{def}}{=} (C^\alpha, \mathcal{C}^\alpha) \overset{\text{Ob}}{\in} \mathcal{X}\mathcal{Y}\mathcal{G}\rho$

7. $X^\alpha \in \mathcal{X}^{\text{Ob}}$
8. $Y^\alpha \in \mathcal{Y}^{\text{Ob}}$
9. $G^\alpha \in \mathcal{G}^{\text{Ob}}$ a topological group with group operation $\star^\alpha$
10. $\pi^\alpha: E^\alpha \rightarrow X^\alpha$ surjective
11. $\pi_X^\alpha: C^\alpha \rightarrow X^\alpha$,
12. $\pi_Y^\alpha: C^\alpha \rightarrow Y^\alpha$ surjective
13. $\star^\alpha: G^\alpha \times G^\alpha \rightarrow G^\alpha$ surjective
14. $\rho^\alpha: Y^\alpha \times G^\alpha \rightarrow Y^\alpha$ is an effective right action of G^α on Y^α
15. $\mathcal{M}^{\alpha}_{\text{triv Sing}}$ a sequence of categories
16. $\mathbf{M}^\alpha$ a sequence
17. $L^\alpha \stackrel{\text{def}}{=} \left(\mathcal{M}^{\alpha}_{\text{triv Sing}}, \mathbf{M}^\alpha, A^\alpha, F^\alpha, \Sigma \right)$ and $L^{\alpha, \mathcal{M}} \stackrel{\text{def}}{=} (\mathcal{M}^\alpha, \mathbf{M}^\alpha, A^\alpha, F^\alpha, \Sigma)$ local coordinate spaces
18. $\mathbf{B}^\alpha \stackrel{\text{def}}{=} (E^\alpha, X^\alpha, Y^\alpha, \pi^\alpha, G^\alpha, \rho^\alpha)$
19. $\mathbf{B} \stackrel{\text{def}}{=} \{\mathbf{B}^\alpha | \alpha < A\}$

such that

20. $\text{head}(\mathcal{M}, 5) = (\mathcal{E}, \mathcal{X} \mathcal{Y} \mathcal{G} \rho, \mathcal{X}, \mathcal{Y}, \mathcal{G})$
21. $\text{head} \left(\mathcal{M}^{\alpha}_{\text{triv Sing}}, 5 \right) = \left(E^\alpha, X^\alpha, Y^\alpha, X^\alpha, Y^\alpha, G^\alpha \right)_{\text{triv Sing } G^\alpha - \rho^\alpha - \text{triv Sing } \text{Sing } \text{Sing}}$
22. $\text{head}(\mathbf{M}^\alpha, 5) = (E^\alpha, \mathbf{C}^\alpha, X^\alpha, Y^\alpha, G^\alpha)$
23. $\mathbf{M}^\alpha \in \mathcal{M}^{()}$
24. $\text{head}(F^\alpha, 5) = (\pi^\alpha, \pi_X^\alpha, \pi_Y^\alpha, \star^\alpha, \rho^\alpha)$
25. $C^\alpha = X^\alpha \times Y^\alpha$
26. G^α is a topological group with group operation $\star^\alpha$ Subsequent references to G^α should be read as referring to the group rather than just the underlying topological space.

27. π^α is surjective.
28. $\pi_X^\alpha = \pi_1: X^\alpha \times Y^\alpha \rightarrow X^\alpha$.
29. $\pi_Y^\alpha = \pi_2: X^\alpha \times Y^\alpha \rightarrow Y^\alpha$.
30. ρ^α an effective right action of G^α on Y^α .
31. The coordinate functions in A^α preserve fibers.
32. $\text{head}(\Sigma, 5) = ((0, 2), (1, 2), (1, 3), (4, 4, 4), (3, 4, 3))$

Let

1. $\alpha^i < A, i = 1, 2$
2. $\mathbf{M}^i \stackrel{\text{def}}{=} \mathbf{M}^{\alpha^i}$
3. $\mathbf{E}^i = (E^i, \mathfrak{E}^i) \stackrel{\text{def}}{=} \mathbf{E}^{\alpha^i}$
4. $\mathbf{C}^i = (C^i, \mathcal{C}^i) \stackrel{\text{def}}{=} \mathbf{C}^{\alpha^i}$
5. $\mathbf{A}^i \stackrel{\text{def}}{=} \mathbf{A}^{\alpha^i}$
6. $X^i \stackrel{\text{def}}{=} X^{\alpha^i}$
7. $Y^i \stackrel{\text{def}}{=} Y^{\alpha^i}$
8. $G^i \stackrel{\text{def}}{=} G^{\alpha^i}$
9. $\pi^i \stackrel{\text{def}}{=} \pi^{\alpha^i}$
10. $\pi_X^i \stackrel{\text{def}}{=} \pi_X^{\alpha^i}$
11. $\pi_Y^i \stackrel{\text{def}}{=} \pi_Y^{\alpha^i}$
12. $\star^i \stackrel{\text{def}}{=} \star^{\alpha^i}$
13. $\rho^i \stackrel{\text{def}}{=} \rho^{\alpha^i}$
14. $\mathcal{M}_{\text{triv}}^i \stackrel{\text{def}}{=} \mathcal{M}_{\text{triv}}^{\alpha^i}$
 Sing Sing
15. $\mathbf{L}^i \stackrel{\text{def}}{=} \mathbf{L}^{\alpha^i}$
16. $\mathbf{L}^i, \mathcal{M} \stackrel{\text{def}}{=} \mathbf{L}^{\alpha^i}, \mathcal{M}$
17. $\mathbf{B}^i \stackrel{\text{def}}{=} \mathbf{B}^{\alpha^i}$

18. $f \stackrel{\text{def}}{=} (f_\beta, \beta < \text{length}(\mathcal{M}))$ a morphism from L^1 to L^2
19. $f_E \stackrel{\text{def}}{=} f_0$
20. $f_C \stackrel{\text{def}}{=} f_1$
21. $f_X \stackrel{\text{def}}{=} f_2$
22. $f_Y \stackrel{\text{def}}{=} f_3$
23. $f_G \stackrel{\text{def}}{=} f_4$

Then:

1. $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} L^i$ and $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} L^{\alpha^i}, \mathcal{M}$ are fiber bundles.

Proof. $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} L^i = (E^i, X^i, Y^i, \pi^i, G^i, \rho^i, A^i) = \mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} L^i, \mathcal{M}$ satisfy the conditions of definition 28.1 (fiber bundles) on page 174: {def:Bun}

C^i is a G^α - ρ^α model space of $X^\alpha \times Y^\alpha$: $\mathcal{X}\mathcal{Y}\mathcal{G}\rho$ is a G - ρ -model category and $C^i \in \mathcal{X}\mathcal{Y}\mathcal{G}\rho^{\text{Ob}}$ by hypothesis, and thus C^i is a G^α - ρ^α model space of $X^\alpha \times Y^\alpha$: by definition 24.6 (Categories of G - ρ -model spaces) on page 151. {def:Grhomodcat}

A^i is an m -atlas of E^i in the coordinate space C^i : By definition 15.1 (Local $M - \Sigma$ and $\mathcal{M} - \Sigma$ coordinate spaces) on page 114. {def:LCS}

A^i is an m -atlas of E^i in the coordinate space X, Y : By definition 24.7 (Trivial G - ρ -model spaces) on page 152. {def:Grhomodtriv}

A^i is a π - G - ρ -bundle atlas of E^i in the coordinate space $X^i \times Y^i$: By lemma 26.3 (Bundle atlases) on page 162. {lem:Bun-Atl}

□

2. $f_C = f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 . $G^1-\rho^1-\text{triv}$ $G^2-\rho^2-\text{triv}$

Proof. f_C satisfies the conditions of definition 24.3 (Morphisms of G - ρ -model spaces) on page 150: {def:Grhomorph}

$f_C = f_1$ is a model function: By definition 15.1 (Local $M - \Sigma$ and $\mathcal{M} - \Sigma$ coordinate spaces) on page 114. {def:LCS}

f_C is a model function that preserves the group action: f Σ -commutes with F^1, F^2 , as shown in fig. 5 (f Σ -commutes with the function sequences F^1, F^2) on page 43, by definition 15.1 (Local $M - \Sigma$ and $\mathcal{M} - \Sigma$ coordinate spaces) on page 114 and definition 10.7 (Morphisms of pre-structures) on page 46. {fig:fF1F2} {def:LCS} {def:premorph}

□

3. $\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}, \mathbf{L}^1, \mathbf{L}^2)$ and $\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}, \mathbf{L}^1, \mathcal{M}, \mathbf{L}^2, \mathcal{M})$ are bundle maps from $(\mathbf{A}^1, \mathbf{B}^1)$ to $(\mathbf{A}^2, \mathbf{B}^2)$.

Proof. $\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}, \mathbf{L}^1, \mathbf{L}^2)$ and $\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}, \mathbf{L}^1, \mathcal{M}, \mathbf{L}^2, \mathcal{M})$ satisfy the conditions of lemma 29.2 (Bundle maps) on page 176

{lem:Bun-map}

$\mathbf{B}^i$ is a protobundle: By hypothesis

$\mathbf{A}^i$ is a π^i - G^i - ρ^i -atlas of E^i in the coordinate space $C^i = X^i \times Y^i$: Shown above.

$f_X \times f_Y$ is a G^1 - G^2 - ρ^1 - ρ^2 morphism from X^1, Y^1 to X^2, Y^2 : Shown above.

$(f_E, f_X \times f_Y)$ is an E^1 - E^2 m-atlas morphism from $\mathbf{A}^1$ to $\mathbf{A}^2$ in the coordinate spaces X^1, Y^1 , X^2, Y^2 : By lemma 24.9 (Trivial G - ρ -model spaces) on page 153.

{lem:Grhomodtriv}

□

4. $\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}$ is a functor from $\text{LCS}^{\text{Bun}} \mathbf{B}$ to $\text{Bun} \mathbf{B}$ and is a functor from $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$ to $\text{Bun} \mathbf{B}$.

Proof.

Preservation of endpoints:

$$\begin{aligned} \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}^1, \mathbf{L}^1, \mathbf{L}^2) &= ((f_E^1, f_X^1, f_Y^1, f_G^1), \mathbf{B}^1, \mathbf{B}^2) \\ \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathcal{M}^i, \mathbf{M}^i, \mathbf{A}^i, \mathbf{F}^i, \Sigma) &= \mathbf{B}^i \end{aligned}$$

Composition:

$$\begin{aligned} \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}((\mathbf{f}^2, \mathbf{L}^2, \mathbf{L}^3) \overset{A}{\circ} (\mathbf{f}^1, \mathbf{L}^1, \mathbf{L}^2)) &= \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}((\mathbf{f}^2 \overset{0}{\circ} \mathbf{f}^1, \mathbf{L}^1, \mathbf{L}^3)) \\ &= ((f_E^2 \circ f_E^1, f_X^2 \circ f_X^1, f_Y^2 \circ f_Y^1, f_G^2 \circ f_G^1), \mathbf{B}^1, \mathbf{B}^3) \\ &= \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}^2, \mathbf{L}^2, \mathbf{L}^3) \overset{A}{\circ} \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\mathbf{f}^1, \mathbf{L}^1, \mathbf{L}^2) \end{aligned}$$

Identity:

$$\begin{aligned} \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\text{Id}_{L^i}) &= \mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}}(\text{Id}_{\mathbf{M}^i}, \mathbf{L}^i, \mathbf{L}^i) \\ &= ((\text{Id}_{E^i}, \text{Id}_{X^i}, \text{Id}_{Y^i}, \text{Id}_{G^i}), (\mathbf{A}^i, \mathbf{B}^i), (\mathbf{A}^i, \mathbf{B}^i)) \\ \text{Id}_{(\mathbf{A}^i, \mathbf{B}^i)} &= \text{Id}_{\mathcal{F}_{\text{LCS,Fib}}^{\text{Bun}} \mathbf{L}^i} \end{aligned}$$

The proof does not depend on the category, so it applies to $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$ as well. $\square$

5. $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} \circ \mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}}$ and $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} \circ \mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}$ are the identity functor on $\text{Bun} \mathbf{B}$.

Proof. Expanding the definitions, we have

- (a) $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}(\mathbf{A}^i, \mathbf{B}^i) = \mathbf{L}^i$
- (b) $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} \mathbf{L}^i = (\mathbf{A}^i, \mathbf{B}^i)$
- (c) $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}((f_E, f_X, f_Y, f_G), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) =$
 $((f_E, f_X \times f_Y, f_X, f_Y, f_G), \mathbf{L}^1, \mathbf{L}^2)$
- (d) $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}}((f_E, f_X \times f_Y, f_X, f_Y, f_G), \mathbf{L}^1, \mathbf{L}^2) =$
 $((f_E, f_X, f_Y, f_G), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2))$

The proof does not depend on the category, so it applies to $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$ as well. $\square$

6. $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}} \circ \mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}}$ is the identity functor on $\text{LCS}^{\text{Bun}} \mathbf{B}$ and $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}} \circ \mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}}$ is the identity functor on $\text{LCS}^{\text{Bun}, \mathcal{M}} \mathbf{B}$.

Proof. Expanding the definitions, we have

- (a) $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}} \mathbf{L}^i = (\mathbf{A}^i, \mathbf{B}^i)$
- (b) $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}}(\mathbf{A}^i, \mathbf{B}^i) = \mathbf{L}^i$
- (c) $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}}((f_E^1, f_X^1, f_Y^1, f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2)) =$
 $((f_E^1, f_X^1 \times f_Y^1, f_X^1, f_Y^1, f_G^1), \mathbf{L}^1, \mathbf{L}^2)$
- (d) $\mathcal{F}_{\text{LCS}, \text{Fib}}^{\text{Bun}}((f_E^1, f_X^1 \times f_Y^1, f_X^1, f_Y^1, f_G^1), \mathbf{L}^1, \mathbf{L}^2) =$
 $((f_E^1, f_X^1, f_Y^1, f_G^1), (\mathbf{A}^1, \mathbf{B}^1), (\mathbf{A}^2, \mathbf{B}^2))$

The same proof applies for $\mathcal{F}_{\text{Fib}, \text{LCS}}^{\text{Bun}, \mathcal{M}}$ with $\mathbf{L}^{i, \mathcal{M}}$ in place of $\mathbf{L}^i$. $\square$

Part XI

Future directions

{part: fut}

If this paradigm proves useful, it can be extended to include a set of admissible functions on the model neighborhoods of the charts, possibly using the language of sheaves. That might be desirable for coordinate spaces more general than Fréchet spaces.

Further work is needed to determine whether it is productive to define (semi-)strict m-atlas morphisms in terms of concrete categories over a category of model spaces rather than directly in terms of a model category.

Further work is needed to determine whether it is productive to allow a local coordinate space to have more than one atlas, e.g., for more than one bundle structure on the same base space.

The extension of paracompactness to model spaces is intended to be useful for partitions of unity on fiber bundles. Further work is needed to determine whether that is actually the case.

The definitions given here include some fairly strong conditions, e.g., AOC. Further work is needed to determine whether they should be relaxed for applications beyond manifolds and fiber bundles.

Further work is needed to determine whether the concepts of category-based atlases²¹ of model spaces and prestructures have general utility.

Further work is needed to determine where it is most fruitful to require full subcategories and where just subcategories.

Further work is needed to devise a definition of constraints that expresses global properties, e.g., compactness, and is both clear and rigorous.

Further work is needed to determine conditions for mappings associated with atlas morphisms to be model functions.

This paper uses the language of category theory as an organizing principle, but defines various notions concretely with sets. It may be desirable to abstract away some of the details, in the spirit of, e.g., topoi.

It may be desirable to define presequences in the language of natural transformations.

References

- [1] Jiří Adámek, Horst Herrlich, and George E. Strecker. *Abstract and Concrete Categories: The Joy of Cats*. John Wiley and Sons, Inc., 1990.
- [2] John L. Kelley. *General Topology*. D. Van Nostrand Company, 1955. First edition.
- [3] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of Differential Geometry, Volume I*. John Wiley and Sons, Inc., 1996.
- [4] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer-Verlag, 2nd edition, 1998.
- [5] Shmuel (Seymour J.) Metz. Towards a taxonomy of atlases and of morphisms between them. 2019. arXiv:1906.11690.
- [6] Norman Steenrod. *The Topology of Fibre Bundles*. Princeton University Press, 1999. Seventh printing.

²¹As opposed to pseudogroup based