


SUBLINEAR EXPANDERS

Shoham Letzter
UCL


BCC
July 2024

Joint with António Girão, Tao Jiang,
Abhishek Methuku, and Liana Yepremyan

Subdivisions :




K_4




K_4 -subdivision

Subdivisions:



K_4




K_4 -subdivision


Question (Mader '67 , Erdős-Hajnal '69): What is
 $\text{sub}(k) := \min \{d : \text{if } \underline{d}(G) \geq d \text{ then } G \text{ contains a } K_k\text{-subdivision}\}?$

↓
average degree of G

Subdivisions:



K_4




K_4 -subdivision

Question (Mader '67 , Erdős-Hajnal '69): What is
 $\text{sub}(k) := \min \{d : \text{if } \underline{d}(G) \geq d \text{ then } G \text{ contains a } K_k\text{-subdivision}\}?$


↓
average degree of G

- * $\text{sub}(k) = \mathcal{O}(k^2)$. (Consider $K_{t,t}$ for $t = \binom{k/2}{2} - 1$.)

Subdivisions:



K_4



K_4 -subdivision

Question (Mader '67 , Erdős-Hajnal '69): What is
 $\text{sub}(k) := \min \{d : \text{if } \underline{d}(G) \geq d \text{ then } G \text{ contains a } K_k\text{-subdivision}\}?$

↓
average degree of G

* $\text{sub}(k) = \mathcal{O}(k^2)$. (Consider $K_{t,t}$ for $t = \binom{k/2}{2} - 1$.)

Komlós-Szemerédi '96 / Bollobás-Thomason '96: $\text{sub}(k) = O(k^2)$.

BT: If G is $22k$ -connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

BT: If G is $22k$ -connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

BT: If G is $22k$ -Connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

An expander is a graph where all sets of vertices that are not too small or too large have large neighbourhoods.

BT: If G is $22k$ -Connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

An expander is a graph where all sets of vertices that are not too small or too large have large neighbourhoods.

* E.g. G is an α -expander if $|N(U)| \geq \alpha |U|$ for $U \subseteq V(G)$ with $|U| \leq \frac{|G|}{2}$.

\downarrow
constant

BT: If G is $22k$ -Connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

An expander is a graph where all sets of vertices that are not too small or too large have large neighbourhoods.

- * E.g. G is an α -expander if $|N(U)| \geq \alpha |U|$ for $U \subseteq V(G)$ with $|U| \leq \frac{|G|}{2}$.
 \downarrow
constant
- * Introduced in the 70's.

BT: If G is $22k$ -Connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

An expander is a graph where all sets of vertices that are not too small or too large have large neighbourhoods.

- * E.g. G is an α -expander if $|N(U)| \geq \alpha |U|$ for $U \subseteq V(G)$ with $|U| \leq \frac{|G|}{2}$.
 \downarrow
constant
- * Introduced in the 70's.
- * Have desirable properties, making them useful in combinatorics and related fields.

BT: If G is $22k$ -Connected, it is k -linked.

can join, vertex-disjointly, any k pairs of vertices

KS introduced a notion of sublinear expanders.

An expander is a graph where all sets of vertices that are not too small or too large have large neighbourhoods.

- * E.g. G is an α -expander if $|N(U)| \geq \alpha |U|$ for $U \subseteq V(G)$ with $|U| \leq \frac{|G|}{2}$.
constant
- * Introduced in the 70's.
- * Have desirable properties, making them useful in combinatorics and related fields.
- * Can be found in random graphs, with high probability.

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{\left(\log\left(\frac{tx}{t}\right)\right)^2}$.

$\xrightarrow[\text{as } x \rightarrow \infty]{}$ 0

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{(\log(\frac{tx}{t}))^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \ \exists \varepsilon > 0$ s.t. every G has an (ε, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{(\log(\frac{tx}{\varepsilon}))^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \exists \varepsilon > 0$ s.t. every G has an (ε, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

* Typically, ε is a constant. Then

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{(\log(\frac{tx}{\varepsilon}))^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \exists \varepsilon > 0$ s.t. every G has an (ε, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

* Typically, ε is a constant. Then

- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq O(t)$, expand linearly.

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{\left(\log\left(\frac{tx}{\varepsilon}\right)\right)^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \ \exists \varepsilon > 0$ s.t. every G has an (ε, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

* Typically, ε is a constant. Then

- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq O(t)$, expand linearly.
- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq \frac{n}{2}$, expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.

G is an (ε, t) -expander if $|N(U)| \geq p_{\varepsilon, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\varepsilon, t}(x) = \frac{\varepsilon}{\left(\log\left(\frac{tx}{\varepsilon}\right)\right)^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \ \exists \varepsilon > 0$ s.t. every G has an (ε, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

* Typically, ε is a constant. Then

- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq O(t)$, expand linearly.
- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq \frac{n}{2}$, expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.
- There is no information about sets of size $< \frac{t}{2}$.

G is an (Σ, t) -expander if $|N(U)| \geq p_{\Sigma, t}(|U|) \cdot |U|$ for $U \subseteq V(G)$

with $\frac{t}{2} \leq |U| \leq \frac{|G|}{2}$, where $p_{\Sigma, t}(x) = \frac{\Sigma}{\left(\log\left(\frac{18x}{\Sigma}\right)\right)^2}$.

$\xrightarrow{\text{as } x \rightarrow \infty} 0$

Theorem (Komlós-Szemerédi '96). $\forall \delta > 0 \ \exists \Sigma > 0$ s.t. every G has an (Σ, t) -expander $H \subseteq G$ with $d(H) \geq (1 - \delta)d(G)$.

* Typically, Σ is a constant. Then

- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq O(t)$, expand linearly.
- Sets $U \subseteq V(G)$, with $\frac{t}{2} \leq |U| \leq \frac{n}{2}$, expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.
- There is no information about sets of size $< \frac{t}{2}$.
(So every graph on $< \frac{t}{2}$ vertices is an (Σ, t) -expander).

- * Sometimes it makes sense to work with a weaker notion of expanders, where sets are only required to expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.

- * Sometimes it makes sense to work with a weaker notion of expanders, where sets are only required to expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.
- Moshkovitz-Shapira '18: Cannot guarantee an expander with expansion rate $\Omega\left(\frac{1}{(\log n)^{1-c}}\right)$.

- * Sometimes it makes sense to work with a weaker notion of expanders, where sets are only required to expand at rate $\Omega\left(\frac{1}{(\log n)^2}\right)$.
- Moshkovitz-Shapira '18: Cannot guarantee an expander with expansion rate $\Omega\left(\frac{1}{(\log n)^{1-c}}\right)$.
- Shapira-Sudakov '15: Can always find an expander with expansion rate $\Omega\left(\frac{1}{\log n (\log \log n)^2}\right)$.

Sketch of proof that $\text{sub}(t_k) = O(k^2)$:

Sketch of proof that $\text{sub}(k) = O(k^2)$:

* Need to show: $\exists c > 0$ s.t. if $d(G) \geq ck^2$ then G has R_k -subdivision.

Sketch of proof that $\text{sub}(k) = O(k^2)$:

- * Need to show: $\exists c > 0$ s.t. if $d(G) \geq ck^2$ then G has K_k -subdivision.
- * By Theorem, may assume G is an (ε, t) -expander.

Sketch of proof that $\text{sub}(k) = O(k^2)$:

- * Need to show: $\exists c > 0$ s.t. if $d(G) \geq ck^2$ then G has R_k -subdivision.
- * By Theorem, may assume G is an (ε, t) -expander.
- * Here it makes sense to take $t = c'k^2$.

Sketch of proof that $\text{sub}(k) = O(k^2)$:

- * Need to show: $\exists c > 0$ s.t. if $d(G) \geq ck^2$ then G has K_k -subdivision.
- * By Theorem, may assume G is an (ε, t) -expander.
- * Here it makes sense to take $t = c'k^2$.
- * If there are k vertices with large degree, join them one by one, vertex-disjointly, by shortest paths.

Sketch of proof that $\text{sub}(k) = O(k^2)$:

- * Need to show: $\exists c > 0$ s.t. if $d(G) \geq ck^2$ then G has K_k -subdivision.
- * By Theorem, may assume G is an (ε, t) -expander.
- * Here it makes sense to take $t = c'k^2$.
- * If there are k vertices with large degree, join them one by one, vertex-disjointly, by shortest paths.

This works because between every two vertices there is a path of length $O(\log n)^3$ avoiding any smallish set of vertices.

* Otherwise, may assume G is an expander with smallish max degree.

- * Otherwise, may assume G is an expander with smallish max degree.
- * Find $2k$ vertices with 'nice' structures around them, that have few overlaps.

- * Otherwise, may assume G is an expander with smallish max degree.
- * Find $2k$ vertices with 'nice' structures around them, that have few overlaps.
- * Join pairs one by one, omitting vertices whose structures become overused.

- * Otherwise, may assume G is an expander with smallish max degree.
- * Find $2k$ vertices with ‘nice’ structures around them, that have few overlaps.
- * Join pairs one by one, omitting vertices whose structures become overused.

This is a somewhat representative proof sketch using (Σ, ϵ) -expanders, with the definition of ‘nice’ varying.

- * Otherwise, may assume G is an expander with smallish max degree.
- * Find $2k$ vertices with ‘nice’ structures around them, that have few overlaps.
- * Join pairs one by one, omitting vertices whose structures become overused.

This is a somewhat representative proof sketch using (Σ, ϵ) -expanders, with the definition of ‘nice’ varying.

Best bounds: $(1+o(1)) \frac{9k^2}{64} \leq \text{sub}(k) \leq (1+o(1)) \frac{10k^2}{23}$.

bipartite random graphs \downarrow \downarrow Kühn–Osthus ‘06

More on subdivisions.

More on subdivisions.

- * Conjecture (Mader '99): $\exists c$ s.t. if $d(G) \geq ck$ and G is C_4 -free then G has a K_k -subdivision.

More on subdivisions.


* Conjecture (Mader '99): $\exists c$ s.t. if $d(G) \geq ck$ and G is C_4 -free then G has a K_k -subdivision.

Proved by Liu-Montgomery '17.

More on subdivisions.

* Conjecture (Mader '99): $\exists c$ s.t. if $d(G) \geq ck$ and G is C_4 -free then G has a K_k -subdivision.


Proved by Liu-Montgomery '17.



More on subdivisions.

* Conjecture (Mader '99): $\exists c$ s.t. if $d(G) \geq ck$ and G is C_4 -free then G has a K_k -subdivision.

Proved by Liu-Montgomery '17.



[Here it makes sense to take $t = c'k^2$, because second neighbourhoods have size $\mathcal{O}(k^2)$].

More on subdivisions.

- * Montgomery 'is: If $d(G) \geq \text{sub}(k) + \varepsilon$ then G has a K_k -subdivision on $O_\varepsilon(\log |G|)$ vertices.

More on subdivisions.

- * Montgomery '15: If $d(G) \geq \text{sub}(k) + \varepsilon$ then G has a K_k -subdivision on $O_\varepsilon(\log |G|)$ vertices.
 - Tight up to a constant factor.

More on subdivisions.

* Montgomery 'is: If $d(G) \geq \text{sub}(k) + \varepsilon$ then G has a K_k -subdivision on $O_\varepsilon(\log |G|)$ vertices.

- Tight up to a constant factor.
- Uses a different notion of expanders, because (ε, t) -expanders have diameter $\gg \log n$.


More on subdivisions.

* Montgomery '15: If $d(G) \geq \text{sub}(k) + \varepsilon$ then G has a K_k -subdivision on $O_\varepsilon(\log |G|)$ vertices.

- Tight up to a constant factor.
- Uses a different notion of expanders, because (ε, t) -expanders have diameter $\gg \log n$.
- Montgomery also proves an analogous result for minors, improving on Shapira-Sudakov '15.


More on subdivisions.

* Balanced subdivisions:



More on subdivisions.

* Balanced subdivisions:




Luan-Tang-Wang-Yang '23 /

Gil Fernandez-Hyde - Liu - Pikhurko - Wu '23:

$d(G) \geq ck^2 \Rightarrow$ balanced K_k -subdivision .

More on subdivisions.

* Balanced subdivisions:




Luan-Tang-Wang-Yang '23 /


Gil Fernandez-Hyde - Liu - Pikhurko - Wu '23:

$d(G) \geq ck^2 \Rightarrow$ balanced K_k -subdivision.

Uses ideas from Liu-Montgomery '23 resolving Erdős and Hajnal's odd cycle problem.





Simple adjuster





adjuster

Immersions: An immersion of H is obtained by replacing each edge uv in H by a path P_{uv} from u to v , s.t. these paths are edge-disjoint.





Immersions: An immersion of H is obtained by replacing each edge uv in H by a path P_{uv} from u to v , s.t. these paths are edge-disjoint.



$\text{imm}(k) := \min\{d : \text{if } d(G) \geq d \text{ then } G \text{ has a } k_k\text{-immersion?}\}$.


Immersions: An immersion of H is obtained by replacing each edge uv in H by a path P_{uv} from u to v , s.t. these paths are edge-disjoint.



$\text{imm}(k) := \min\{d : \text{if } d(G) \geq d \text{ then } G \text{ has a } k_k\text{-immersion}\}.$

DeVos - Dvořák - Fox - McDonald - Mohar - Scheide '14: $\text{imm}(k) = O(k)$.

Immersions: An immersion of H is obtained by replacing each edge uv in H by a path P_{uv} from u to v , s.t. these paths are edge-disjoint.



$\text{imm}(k) := \min\{d : \text{if } d(G) \geq d \text{ then } G \text{ has a } k_k\text{-immersion}\}.$

DeVos - Dvořák - Fox - McDonald - Mohar - Scheide '14: $\text{imm}(k) = O(k)$.

Dvořák - Yepremyan '18: $\delta(G) \geq 11k + 7 \Rightarrow K_k\text{-immersion}.$

Question: Does $\delta(G) \geq (1+\varepsilon)\ell$ imply G has a K_h -immersion?

Question: Does $\delta(G) \geq (1+\varepsilon)k$ imply G has a K_k -immersion?

- * Tight, because K_k -immersions have vertices of degree $\geq k-1$.

Question: Does $\delta(G) \geq (1+\varepsilon)k$ imply G has a K_h -immersion?

- * Tight, because K_k -immersions have vertices of degree $\geq k-1$.
- * Liu-Wang-Yang '22: Yes, if G is $K_{s,t}$ -free.

Question: Does $\delta(G) \geq (1+\varepsilon)k$ imply G has a K_k -immersion?

- * Tight, because K_k -immersions have vertices of degree $\geq k-1$.
- * Liu-Wang-Yang '22: Yes, if G is $K_{s,t}$ -free.

Question. If $d(G) \geq \text{imm}(k) + \varepsilon$, is there a K_k -immersion on $O_\varepsilon(\log |G|)$ vertices?


Question: Does $\delta(G) \geq (1+\varepsilon)k$ imply G has a K_k -immersion?

- * Tight, because K_k -immersions have vertices of degree $\geq k-1$.
- * Liu-Wang-Yang '22: Yes, if G is $K_{s,t}$ -free.



Question. If $d(G) \geq \text{imm}(k) + \varepsilon$, is there a K_k -immersion on $O_\varepsilon(\log |G|)$ vertices?

Question. Is there $c > 0$ s.t. if $d(G) \geq ck$ then G has a 'balanced' K_k -immersion?

A directed graph G is Eulerian if $d^+(v) = d^-(v)$ for every vertex v .





A directed graph G is Eulerian if $d^+(v) = d^-(v)$ for every vertex v .



Girão-L. '23: If G is an Eulerian directed graph with $\min \deg \geq ck$ then it has a \vec{K}_k -immersion.

↓
Complete digraph on k vertices

A directed graph G is Eulerian if $d^+(v) = d^-(v)$ for every vertex v .





Girão-L. '23: If G is an Eulerian directed graph with $\min \deg \geq ck$ then it has a \vec{K}_k -immersion.

↓
Complete digraph on k vertices

- * False without requiring G to be Eulerian, even for $k=3$.

A directed graph G is Eulerian if $d^+(v) = d^-(v)$ for every vertex v .





Girão-L. '23: If G is an Eulerian directed graph with $\min \deg \geq ck$ then it has a \vec{K}_k -immersion.

↓
Complete digraph on k vertices

- * False without requiring G to be Eulerian, even for $k=3$.
- * Rare use of expanders for directed graphs

A directed graph G is Eulerian if $d^+(v) = d^-(v)$ for every vertex v .



Girão-L. '23: If G is an Eulerian directed graph with $\min \deg \geq ck$ then it has a \vec{K}_k -immersion.

↓
Complete digraph on k vertices

- * False without requiring G to be Eulerian, even for $k=3$.
- * Rare use of expanders for directed graphs
[Eulerian + immersion helps].

Hypergraphs with no tight cycles


Hypergraphs with no tight cycles

An r-uniform hypergraph consists of vertices and edges which are sets of r vertices.

Hypergraphs with no tight cycles

An r-uniform hypergraph consists of vertices and edges which are sets of r vertices.

An r-uniform tight cycle of length l has vertices $\{v_1, \dots, v_l\}$ and edges $\{v_i \dots v_{i+r-1} \text{ (mod } l\} : i \in [l]\}$.




a tight 3-uniform cycle of length 10

Hypergraphs with no tight cycles

An r-uniform hypergraph consists of vertices and edges which are sets of r vertices.

An r-uniform tight cycle of length l has vertices $\{v_1, \dots, v_l\}$ and edges $\{v_i \dots v_{i+r-1} \text{ (mod } l\} : i \in [l]\}$.



a tight 3-uniform cycle of length 10

Question (Sós/Verstraëte): What is the maximum number of edges in an n-vertex r-uniform hypergraph with no tight cycles?

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

Sudakov-Tomon '22: $\leq n^{r-1+o(1)}.$

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

Sudakov-Tomon '22: $\leq n^{r-1+o(1)}.$

- * Used sublinear expanders (on 'r-line-graphs').

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

Sudakov-Tomon '22: $\leq n^{r-1+o(1)}.$

- * Used sublinear expanders (on 'r-line-graphs').
- * Showed that in an expander H every hyperedge can reach almost every other edge via a short tight path.

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

Sudakov-Tomon '22: $\leq n^{r-1+o(1)}.$

- * Used sublinear expanders (on ' r -line-graphs').
- * Showed that in an expander H every hyperedge can reach almost every other edge via a short tight path.
- * Ideally: split H randomly into H_1 and H_2 , show both H_i 's maintain above property, and then can find a cycle.

B. Janzer '20: $\geq \Omega\left(n^{r-1} \frac{\log n}{\log \log n}\right).$

Sudakov-Tomon '22: $\leq n^{r-1+o(1)}.$

- * Used sublinear expanders (on 'r-line-graphs').
- * Showed that in an expander H every hyperedge can reach almost every other edge via a short tight path.
- * Ideally: split H randomly into H_1 and H_2 , show both H_i 's maintain above property, and then can find a cycle.

L. '23: $\leq O(n^{r-1}(\log n)^s)$.

L. '23: $\leq O(n^{r-1}(\log n)^5)$.

* Strengthened above property to:

For every edge e in an expander H , can reach almost every edge via a short tight path so that no vertex (outside e) is overused.

Rainbow cycles and clique subdivisions

Rainbow cycles and clique subdivisions

A graph is properly-edge-coloured if intersecting edges have different colours.

Rainbow cycles and clique subdivisions

A graph is properly-edge-coloured if intersecting edges have different colours.

An edge-coloured graph is rainbow if its edges have distinct colours.

Rainbow cycles and clique subdivisions

A graph is properly-edge-coloured if intersecting edges have different colours.

An edge-coloured graph is rainbow if its edges have distinct colours.

Question: (Keevash-Mubayi-Sudakov-Verstraëte '06)

What is the maximum number of edges in a properly-edge-coloured n -vertex graph with no rainbow cycles?

Rainbow cycles and clique subdivisions

A graph is properly-edge-coloured if intersecting edges have different colours.

An edge-coloured graph is rainbow if its edges have distinct colours.

Question: (Keevash-Mubayi-Sudakov-Verstraëte '06)

What is the maximum number of edges in a properly-edge-coloured n -vertex graph with no rainbow cycles?

Question: Let $k \geq 3$. What is the maximum number of edges in a properly-edge-coloured n -vertex graph with no rainbow K_k -subdivision?

Rainbow cycles

KMSV '06: $\Omega(n \log n)$. [Hypercube.]

Rainbow cycles

KMSV '06: $\Omega(n \log n)$. [Hypercube.]

O. Janzer '22: $O(n(\log n)^4)$. [Counting homomorphisms.]

Rainbow clique subdivisions

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

- * Adjusted ST's approach to the rainbow setting.

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

- * Adjusted ST's approach to the rainbow setting.

In particular, showed:

- In a properly-edge-coloured expander for every vertex can reach almost every other vertex via short rainbow path.

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

- * Adjusted ST's approach to the rainbow setting.

In particular, showed:

- In a properly-edge-coloured expander for every vertex can reach almost every other vertex via short rainbow path.

Jiang - L. - Methuku - Yepremyan '24: $\leq O(n(\log n)^{53})$.

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

- * Adjusted ST's approach to the rainbow setting.

In particular, showed:

- In a properly-edge-coloured expander for every vertex can reach almost every other vertex via short rainbow path.

Jiang - L. - Methuku - Yepremyan '24: $\leq O(n(\log n)^{53})$.

- * Found almost regular expanders

Rainbow clique subdivisions

Jiang - Methuku - Yepremyan '23: $\leq n^{1+o(1)}$.

- * Adjusted ST's approach to the rainbow setting.

In particular, showed:

- In a properly-edge-coloured expander for every vertex can reach almost every other vertex via short rainbow path.

Jiang - L. - Methuku - Yepremyan '24: $\leq O(n(\log n)^{53})$.

- * Found almost regular expanders
- * Used the fast mixing time of random walks in expanders.

Tomon '22: $O(n(\log n)^{6+o(1)})$.

Tomon '22: $O(n(\log n)^{6+o(1)})$.

- * Used sampling trick to make ST's approach work, i.e. show random subhypergraphs maintain expansion properties.

Tomon '22: $O(n(\log n)^{6+o(1)})$.

- * Used sampling trick to make ST's approach work, i.e. show random subhypergraphs maintain expansion properties.

Wang '21: $O(n(\log n)^{2+o(1)})$.

Tomon '22: $O(n(\log n)^{6+o(1)})$.

- * Used sampling trick to make ST's approach work, i.e. show random subhypergraphs maintain expansion properties.

Wang '21: $O(n(\log n)^{2+o(1)})$.

- * Optimised Tomon's approach for this particular problem.

Back to rainbow cycles

Back to rainbow cycles

Tomon '22: $O(n(\log n)^{2+o(1)})$.

Back to rainbow cycles

Tomon '22: $O(n(\log n)^{2+o(1)})$.

Janzer-Sudakov '22 / Kim-Lee-Liu-Tran '22: $O(n(\log n)^2)$.

Back to rainbow cycles

Tomon '22: $O(n(\log n)^{2+o(1)})$.

Janzer-Sudakov '22 / Kim-Lee-Liu-Tran '22: $O(n(\log n)^2)$.

Alon-Bucić-Sauermann-Zakharov-Zamir '23: $O(n \log n \log \log n)$.

Back to rainbow cycles

Tomon '22: $O(n(\log n)^{2+o(1)})$.

Janzer-Sudakov '22 / Kim-Lee-Liu-Tran '22: $O(n(\log n)^2)$.

Alon-Bucić-Sauermann-Zakharov-Zamir '23: $O(n \log n \log \log n)$.

* Optimised notion of expansion.

Back to rainbow cycles

Tomon '22: $O(n(\log n)^{2+o(1)})$.

Janzer-Sudakov '22 / Kim-Lee-Liu-Tran '22: $O(n(\log n)^2)$.

Alon-Bucić-Sauermann-Zakharov-Zamir '23: $O(n \log n \log \log n)$.

- * Optimised notion of expansion.
- * New sampling trick.

More results

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.
- * Kim-Liu-Sharifzaadeh-Staden '17 / Cambie-Gao-Liu '23: counting Hamiltonian sets.

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.
- * Kim-Liu-Sharifzaadeh-Staden '17 / Cambie-Gao-Liu '23: counting Hamiltonian sets.
- * Bucić-Montgomery '22: cycle decompositions.
- * Z. '22: separating path systems. (Improved by Bonamy-Botler-Dross-Naia-Skokan '23.)

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.
- * Kim-Liu-Sharifzaadeh-Staden '17 / Cambie-Gao-Liu '23: counting Hamiltonian sets.
- * Bucić-Montgomery '22: cycle decompositions.
- * Z. '22: separating path systems. (Improved by Bonamy-Botler-Dross-Naia-Skokan '23.)
- * Haslegrave-Hyde-Kim-Liu '23: Ramsey goodness.

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.
- * Kim-Liu-Sharifzaadeh-Staden '17 / Cambie-Gao-Liu '23: counting Hamiltonian sets.
- * Bucić-Montgomery '22: cycle decompositions.
- * Z. '22: separating path systems. (Improved by Bonamy-Botler-Dross-Naia-Skokan '23.)
- * Haslegrave-Hyde-Kim-Liu '23: Ramsey goodness.
- * Montgomery '23: Latin squares.

More results

- * Im-Kim-Kim-Liu '22: K_k -subdivisions in graphs with large 'crux'.
- * Haslegrave-Kim-Liu '22: subdivisions of sparse graphs.
- * Gil Fernandez-Liu '23: pillars.
- * Gil Fernandez-Kim-Kim-Liu '22: nested cycles without crossings.
- * Draganić-Methuku-Munhá-Correia-Sudakov '23: cycles with many chords.
- * Chakraborti-Janzer-Methuku-Montgomery '24: edge-disjoint cycles with same vertex set.
- * Kim-Liu-Sharifzaadeh-Staden '17 / Cambie-Gao-Liu '23: counting Hamiltonian sets.
- * Bucić-Montgomery '22: cycle decompositions.
- * Z. '22: separating path systems. (Improved by Bonamy-Botler-Dross-Naia-Skokan '23.)
- * Haslegrave-Hyde-Kim-Liu '23: Ramsey goodness.
- * Montgomery '23: Latin squares.

Thank you for listening!