

Result: 100% 20 / 20	 100410 001182	Neptun ID: Draft papers: ☐ Name: _____
--	--	---

The test consists of **5 questions**. The available time to solve the quiz is **20 minutes**. Please use a blue coloured pen; the use of any other tools is forbidden.

- | | a | b | c | d | e | f |
|-----|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 1.) | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ |
| 2.) | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ |
| 3.) | ☐ | ☐ | ☐ | ☐ | | |
| 4.) | ☐ | ☐ | ☐ | ☐ | | |
| 5.) | (essay) | | | | | |

Please copy the multiple choice questions' solution in the attached table. For each correct answer marked, 2 points are awarded; for each incorrect answer marked, -1 point is awarded. There is no limit to the number of correct answers per question. A negative aggregate score will be interpreted as 0 points.

1.) Let $L^{(0)} = \langle LC, Con, Form \rangle$ be a zero-order language such that $Con = \{p, q\}$. Which string of symbols are membered in the set $Form$?

2 $\neg(\neg(p \vee q) \wedge q)$

2 $\neg(\neg p \supset \neg q)$

(c) $((p))$

(d) $((p \supset q) \neg p)$

2 $(p \equiv \neg q)$

(f) $(p + q)$

2.) Which statements are true for the formula below?

$$((\neg p \equiv q) \supset \neg(q \vee (\neg p \wedge q)))$$

2 $(\neg p \equiv q)$ is a subformula

(b) $(q \equiv \neg p)$ is a subformula

2 $((\neg p \equiv q) \supset \neg(q \vee (\neg p \wedge q)))$ is a subformula

(d) $\neg q$ is a subformula

2 the formula has 8 subformulae

(f) the formula has only one direct subformula

3.) Which interpretations are models of the following set of formulae?

$$\{ (q \supset (q \supset (q \vee p))), \neg(p \equiv q) \}$$

(a) $\varrho(p) = 0, \quad \varrho(q) = 0$

2 $\varrho(p) = 0, \quad \varrho(q) = 1$

2 $\varrho(p) = 1, \quad \varrho(q) = 0$

(d) $\varrho(p) = 1, \quad \varrho(q) = 1$

4.) Which interpretations are models of the following set of formulae?

$$\{ ((p \wedge q) \vee q), ((q \wedge p) \supset \neg q) \}$$

(a) $\varrho(p) = 0, \quad \varrho(q) = 0$

2 $\varrho(p) = 0, \quad \varrho(q) = 1$

(c) $\varrho(p) = 1, \quad \varrho(q) = 0$

(d) $\varrho(p) = 1, \quad \varrho(q) = 1$

5.) Define the semantic value of $(A \vee B)$!

Score: 2 / 2.