Next-Day Air Quality (AQI) Forecasting for Indian Cities

Problem Statement · DataTalks.Club ML Zoomcamp Midterm Project

1. Background: What is AQI?

The Air Quality Index (AQI) is a single number that tells the public how polluted the air is and what health risks it carries. India's National AQI, launched by the Central Pollution Control Board (CPCB) in 2014, combines up to eight pollutants: $PM_{2.5}$, PM_{10}, NO_2, SO_2, CO, O_3, NH_3, and lead. It reports the worst of their sub-indices as the overall AQI.

The index is grouped into six categories:

AQI Range	Category	Health Impact
0–50	**Good**	Minimal impact
51–100	**Satisfactory**	Minor breathing discomfort for sensitive people
101–200	**Moderate**	Discomfort for people with lung or heart disease, children, and older adults
201–300	**Poor**	Breathing discomfort for most people on prolonged exposure
301–400	**Very Poor**	Respiratory illness on prolonged exposure
401–500	**Severe**	Affects healthy people and seriously harms those with existing diseases

2. The Problem: Air Pollution in India

India has some of the most polluted cities in the world. Many cities in the Indo-Gangetic Plain, including Delhi, Ghaziabad, Noida, Lucknow, Patna, and Kanpur, regularly cross into the “Poor,” “Very Poor,” and “Severe” categories, especially during winter.

Main causes of pollution

- Vehicle emissions in dense, fast-growing cities
- Industrial activity and coal-based power plants
- Construction and road dust, a major source of PM_{10}
- Crop residue (stubble) burning in Punjab and Haryana during October–November
- Household burning of biomass, wood, and waste
- Firecrackers during festivals
- Winter weather, when low wind speeds, temperature inversion, and low mixing height trap pollutants close to the ground

3. Impact

Health

- Fine particulate matter ($PM_{2.5}$) is small enough to enter the lungs and bloodstream. It is linked to asthma, chronic obstructive pulmonary disease (COPD), lung cancer, heart attacks, strokes, and poor lung development in children.

- A 2020 study in *The Lancet Planetary Health* attributed about 1.67 million deaths in India in 2019 to air pollution, roughly 18% of all deaths that year.
- Children, older adults, pregnant women, outdoor workers, and people with existing heart or lung conditions are the most vulnerable.

Economic

- The same Lancet study estimated that lost output from pollution-related illness and early deaths cost India about 1.4% of its GDP in 2019.
- Pollution also increases healthcare spending, reduces worker productivity, and lowers tourism.

Social and daily life

- Schools in Delhi NCR have had to close or move online during severe pollution episodes.
- Construction bans and vehicle restrictions under Delhi's Graded Response Action Plan (GRAP) disrupt work and income, especially for daily-wage workers.
- Flights and road travel are affected by smog and low visibility.

4. The Gap: Why Forecasting Matters

Current AQI dashboards, such as CPCB's SAMEER app, mostly report what the air quality is right now. By the time AQI reaches “Severe,” people have already been exposed and authorities are reacting rather than preventing.

A reliable next-day forecast would allow:

- **Citizens** to plan outdoor activities, exercise, and commutes, and to take precautions such as masks and air purifiers
- **Schools** to decide on closures or indoor activities in advance
- **Hospitals** to prepare for more respiratory and cardiac cases
- **Local authorities** to trigger GRAP measures, such as construction bans, traffic restrictions, and water sprinkling, before pollution peaks rather than after

5. Project Objective

This project builds a machine learning model that predicts the **next day's AQI category** for an Indian city using:

- Pollutant readings from the current and previous days ($PM_{2.5}$, PM_{10}, NO_2, CO, SO_2, O_3, and others)
- Recent trends, using rolling averages of pollutant levels and AQI
- Seasonal and calendar information (month, day of week, season)
- The city

The task is framed as a **multi-class classification** problem whose target is one of the six AQI categories.

> **Why not predict today's AQI?** AQI is computed directly from pollutant readings, so predicting the current AQI from current pollutants would only re-learn a known formula. Forecasting the next day's AQI is a genuine prediction problem with real practical value.

6. Dataset

Air Quality Data in India (2015–2020), available on Kaggle and sourced from CPCB monitoring stations. It contains daily pollutant readings, AQI values, and AQI categories for 26 Indian cities.

7. Success Criteria

- The model should outperform a simple **persistence baseline**, which assumes tomorrow's AQI category equals today's.
- Because severe days are rare but the most important to catch, models are evaluated with **macro F1-score** and a **confusion matrix**, with particular attention to **recall** on the “Very Poor” and “Severe” classes.
- The final model is deployed as a web service (FastAPI/Flask in Docker) that returns the predicted AQI category and class probabilities for given inputs.