

CHAPTER - 21


CARBOXYLIC ACIDS

SYNOPSIS

Saturated monocarboxylic acids form a homologous series which has a general formula $C_nH_{2n+1}COOH$ or $C_nH_{2n}O_2$

In the IUPAC system, aliphatic carboxylic acids are named as 'alkanoic acid'.


Formula	Common Name	IUPAC name
1. CH_3-CH_2-COOH	Propionic acid	Propanoic acid
2. $CH_3-CH_2-CH_2-COOH$	Butyric acid	Butanoic acid
3. $CH_3 - \underset{\substack{ \\ CH_3}}{CH} - COOH$	Isobutyric acid	2-methyl propanoic acid
4. $CH_3-(CH_2)_3-COOH$	Valeric acid	Pentanoic acid
5. $CH_3-(CH_2)_4-COOH$	Caproic acid	Hexanoic acid
6. $HOOC-CH_2-COOH$	Malonic acid	Propane-1,3-dioic acid
7. $HOOC-(CH_2)_2-COOH$	Succinic acid	Butane-1,4-dioic acid
8. $HOOC-(CH_2)_3-COOH$	Glutaric acid	Pentane-1,5-dioic acid
9. $HOOC-(CH_2)_4-COOH$	Adipic acid	Hexane-1,6-dioic acid
10. C_6H_5-COOH	Benzoic acid	Benzene carboxylic acid
	OR (Benzoic acid)	

11. 	Phthalic acid	Benzene-1,2-dicarboxylic acid
---	---------------	-------------------------------

Structure & Isomerism

In $-COOH$ group, carboxyl carbon undergoes sp^2 -hybridisation. The geometry around the carboxyl carbon is trigonal planar and the bond angle is nearly 120° .

The carboxylic C is less electrophilic than carbonyl C due to resonance.





- The carboxylic acids exhibit chain, position, functional and stereo isomerism.
- Esters are functional isomers of carboxylic acids.
- Optical isomerism is observed in Lactic acid.
- Fumaric acid and maleic acids are geometrical isomers.

Methods of preparation

I. Oxidation of 1° Alcohols, aldehydes & ketones


1° ROH are readily oxidised to acids with KMnO_4 in neutral, acidic or basic medium or with $\text{K}_2\text{Cr}_2\text{O}_7$ and with CrO_3 in acidic medium




- Aldehydes are more easily oxidised to acids on treatment with oxidising agents such as HNO_3 , KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$ etc. Even mild oxidising agents like Tollens and Fehling's oxidise aldehyde.
- Ketones are oxidised by strong oxidising agents at elevated temperature.

II From alkyl benzenes :



Side chain oxidation of alkyl benzenes with chromic acid or acidic or alkaline KMnO_4 gives acids. The entire side chain is oxidised to $-\text{COOH}$ group. 1° and 2° alkyl groups are oxidised in the same manner, while 3° group is not affected.



III Hydrolysis of Nitriles




- Hydrolysis can be done with alkali also
 - HCN on hydrolysis forms formic acid
- #### IV. Hydrolysis of ester, acyl halides and anhydrides




V. Hydrolysis of Trihalides


All those compounds containing $-\text{CCl}_3$ group which is not directly attached to carbonyl group are hydrolysed with alkali to corresponding acid



VI. From R – Mg – X



VII. From Olefins (Koch reaction)




Physical properties :

1. Lower fatty acids upto C₁₀ are colourless liquids, higher members are colourless waxy solids.
2. The first three members have a sharp pungent odour, the middle C₄ to C₉ have an odour of rancid butter, whereas higher members are odourless
3. Lower members are completely miscible with water and the solubility decreases with increasing molecular weight
4. B.P. of carboxylic acids increase regularly with increase in mol. wt.. BP of RCOOH > R-OH

Acidic Nature

- Among organic compounds, carboxylic acids are most acidic and ionize in aqueous solution







$$K_a = \frac{[\text{RCOO}^-][\text{H}_3\text{O}^+]}{[\text{RCOOH}]}; \quad \text{p}K_a = -\log K_a$$

- The resulting carboxylate anion is stabilized by resonance, as negative charge is dispersed on both the oxygen atom
- Carboxylic acids are weaker than mineral acids, but they are stronger acids than alcohols, phenols and peroxy acids
- Electron withdrawing groups increase the acidity and electron donating groups decrease the acidity.
- All ortho - substituted acids are stronger than para and meta counterparts


Chemical properties

I Reactions involving cleavage of O-H bond




II Reactions involving cleavage of C-OH bond



a) Formation of anhydride






b) Esterification





c) Reaction with PCl₅, PCl₃ and SOCl₂



d) Reaction with NH_3





III. Reactions involving $-\text{COOH}$ group




Acid first forms Na, K, or Ca carboxylate salts which loses CO_2 to form carbanion. More stable the carbanion formed, more readily the alkane is formed

c) Kolbe's electrolysis:



IV Substitution reaction

a) H.V.Z Reaction




Although monocarboxylic acid themselves are not readily halogenated, their acid chlorides and anhydrides are halogenated readily

b) Ring substitution

An aromatic acid undergoes S.E. reactions in which the $-\text{COOH}$ group acts as a deactivating and meta directing group. It does not undergo F.C. reaction because the $-\text{COOH}$ group is deactivating and the






catalyst AlCl_3 gets bonded to the COOH group



PART-I (JEE MAIN)




SECTION-I (Straight objective type questions)

1. The final product in the following reaction sequence is







2. $\text{CH}_3 - \text{C} \equiv \text{C} - \text{H} \xrightarrow{\text{CH}_3\text{MgBr}} \text{CH}_4 + \text{A} \xrightarrow[\text{H}_3\text{O}^+]{\text{CO}_2} \text{B}$





Product B in the above conversion is




3. $\text{R}-\text{CH}_2-\text{CH}_2\text{OH}$ can be converted to $\text{R}-\text{CH}_2\text{CH}_2\text{COOH}$. The CORRECT sequence of reagents for the conversion is



4. Carboxylic acids do not give nucleophilic addition reaction even though they have $\text{—}\overset{\text{O}}{\parallel}{\text{C}}\text{—}$ group. It is due to
- 1) +I effect of R-group
 - 2) -I effect of —COOH group
 - 3) Hyperconjugation
 - 4) Resonance
5. Carboxylic acid group can be easily detected by using
- 1) NaHCO_3
 - 2) 2,4-DNP
 - 3) Iodoform test
 - 4) Fehling's test
6. The decreasing order of rate of formation of ester for the following compounds with acetic acid in presence of H^+ is



- 1) $\text{i} > \text{iii} > \text{ii} > \text{iv}$
 - 2) $\text{i} > \text{ii} > \text{iii} > \text{iv}$
 - 3) $\text{iv} > \text{ii} > \text{iii} > \text{i}$
 - 4) $\text{iv} > \text{iii} > \text{ii} > \text{i}$
7. What are A and B in the following sequence of reactions







- 1) $\text{CH}_3 - \underset{\text{Br}}{\text{CH}} - \text{COOH}$, $\text{CH}_2 = \text{CH} - \text{COOH}$
- 2) $\text{CH}_3 - \text{CH}_2 - \text{COBr}$, $\text{CH}_2 = \text{CH} - \text{COOH}$
- 3) $\text{CH}_2 - \underset{\text{Br}}{\text{CH}} - \text{COOH}$, $\text{CH}_2 = \text{CH} - \text{COOH}$
- 4) $\text{CH}_3 - \text{CH}_2 - \text{COBr}$, $\text{CH}_3 - \text{CH}_2 - \text{COOH}$

8. Assertion (A) : Compounds containing $-CHO$ groups are easily oxidised with mild oxidising agents to corresponding carboxylic acids






Reason (R) : Carboxylic acids can be reduced to alcohols by treatment with $LiAlH_4$

In the light of the above statements choose the correct option

- 1) Both A and R are correct but R is not the correct explanation of A
 - 2) A is false but R is true
 - 3) Both A and R are true and R is the correct explanation of A
 - 4) Both A and R are false
9. The major product obtained in the monobromination of phenyl benzoate is



10. Benzoic acid when treated with bromine in presence of $FeBr_3$ forms
- 1) o-Bromobenzoic acid
 - 2) Benzyl bromide
 - 3) Benzoyl bromide
 - 4) m-Bromobenzoic acid
11. End product of the following conversion is




SECTION-II - Numerical Type Questions

12. Phthalic acid $\xrightarrow{NH_3}$ [X] $\xrightarrow{\text{strong heating}}$ Y

Total number of heteroatoms present in a molecule of Y is —

13. Percentage of carbon in a carboxylic acid with molecular mass 146 is 49.3%. The number of carbon atoms present in the molecule is —

14. The number of moles of NaOH that would react with one mole of the following compound is —







15. The total number of compound (s) among the following that do(es) not undergo HVZ reaction is —
- 1) Formic acid
 - 2) Benzoic acid
 - 3) 2,2-dimethyl propanoic acid
 - 4) Ethanoic acid
 - 5) 2-methyl propanoic acid
 - 6) Salicylic acid





PART-II (JEE ADVANCED)

Section-III - Only one option correct type

16. Which of the following reactions is correct?



17. List the following esters in the order of decreasing reactivity towards nucleophilic substitution reaction




A) IV > I > III > II

B) IV > III > I > II

C) III > IV > I > II

D) II > I > III > IV

18. Consider the following reaction scheme:



Choose the correct set of reagents X and Y from the following

	X	Y
A)	Br_2	NaNH_2
B)	Br_2, P	NaOH
C)	Br_2, P	NH_3
D)	HBr	NH_3

19. Which of the following dicarboxylic acids gives a cyclic anhydride on heating?


A) Oxalic acid

B) Succinic acid

C) Adipic acid

D) Malonic acid

20. Consider the following reaction sequence.




Compound (F) on treatment with Br_2/Fe mainly gives



Section IV - One or more option correct type

21. The reagent (s) which can bring about the following transformation is/are



- (A) P_2O_5 (B) SOCl_2 (C) POCl_3 (D) PCl_5

22. Which of the following compounds on hydrolysis yield(s) acetic acid?

- A) $\text{CH}_3\text{CO}_2\text{MgX}$ B) $\text{CH}_3\text{C} \equiv \text{N}$ C) $\text{CH}_3\text{COOCH}_3$ D) $\text{CH}_3\text{CH}_2\text{CN}$


23. When methanoic acid is treated with $\text{NaH}^{14}\text{CO}_3$, the gas(es) evolved is/are

- A) H_2 B) ^{14}CO C) CO_2 D) $^{14}\text{CO}_2$


24. Which of the following is/are not correct for carboxylic acids?
- A) Lower acids are largely soluble in water
- B) The hydrogen bonding in acids is weaker than that in alcohols
- C) Carboxylic acids do not exist as dimer in the vapour phase
- D) Benzoic acid is insoluble in cold water where as soluble in benzene and ether

Section V - Numerical type questions

25. How many water molecules are eliminated in the following reaction ?



26. A mixture of $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OC}_2\text{H}_5$ and $\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OC}_2\text{H}_5$ is treated with $\text{C}_2\text{H}_5\text{ONa}$ in ethanol. How many different condensation products (excluding stereoisomers) are formed in the reaction?



The number of alcohol molecules produced in the above reaction is —

28. 59 g of an amide obtained from a carboxylic acid $\text{R}-\text{COOH}$, when heated with NaOH gave 17 g of NH_3 . Find the total number of carbon atoms present in the parent acid ($\text{R}-\text{COOH}$)

Section-VI - Matrix match type

29. Match the following

Column I (Compound)

(I) Acid chloride

(II) Acid anhydride

(III) Ester

(IV) Acid amide

A) I - QRS, II - PQS, III - PQS, IV - QS

C) I - PQRS, II - QS, III - QS, IV - S

Column II (Reactivity towards nucleophiles)

(P) More reactive than aldehyde

(Q) More reactive than carboxylic acid

(R) More reactive than ketone

(S) More reactive than carboxylate

B) I-PQRS, II - QS, III- PQS, IV - S

D) I - QRS, II - PQS, III - QS, IV - QS

30. Match the following

Column-I

(Compound)

(I) HCOOH

(II) CH_3COOH

(III) Benzoic acid

(IV) Cinnamic acid

A) I - PQ, II - PQR, III - PQ, IV - PQ

C) I - PQS, II - PQ, III - PQ, IV - PQR

Column-II

(Characteristic reaction)

(P) Effervescence of CO_2 with NaHCO_3

(Q) Evolve hydrogen with Na

(R) Decolourise Br_2 water

(S) Give silver mirror with Tollens reagent

B) I - PQS, II - PQR, III - PQR, IV - PQ

D) I - PQS, II - PQ, III - PQR, IV - PQR