

Prediction of anomalous blood viscosity in confined shear flow.

Supplementary Material.

Marine Thiébaud,^{1,*} Zaiyi Shen,¹ Jens Harting,^{2,3} and Chaouqi Misbah^{1,†}

*¹Université Grenoble I/CNRS, Laboratoire Interdisciplinaire
de Physique/UMR5588, Grenoble F-38041, France*

*²Department of Applied Physics, Eindhoven University of Technology,
P.O. Box 513, 5600 MB Eindhoven, The Netherlands*

*³Faculty of Science and Technology,
Mesa+ Institute, University of Twente,
7500 AE Enschede, The Netherlands*

Abstract

These supplementary material contains the observation of the three-file organization.

PACS numbers: 87.16.D- 83.50.Ha 87.17.Jj 83.80.Lz 87.19.rh

The three-file organization For smaller confinement $W = 7R_0$ and larger concentration $\phi = 22\%$, a three-file organization has also been observed, as showed by the Fig. 1.




Figure 1: (Color online) A typical vesicle configuration; $W = 7R_0$ and $\phi = 22\%$. (a): The vesicles in top (bottom) file are collectively moving to the right (left) and each vesicle center of mass undergoes oscillations upward and downward in the course of time, while the membrane tank-treads. (b): The configuration at a later time showing the sliding of each vesicle and the vertical motion of their center of mass. (c): The configuration at a later time showing the sliding of each vesicle and the symmetry between the top and bottom files.

* Electronic address: `marine.thiebaud@ujf-grenoble.fr`

† Electronic address: `chaouqi.misbah@ujf-grenoble.fr`