



MINISTRY OF SCIENCE AND
HIGHER EDUCATION OF
THE REPUBLIC OF KAZAKHSTAN





Түркі әлемі математиктерінің
VII Дүниежүзілік Конгресі
(TWMS Congress-2023)
**БАЯНДАМАЛАРЫНЫҢ
ТЕЗИСТЕРІ**

ABSTRACTS
of the VII World Congress of Turkic
World Mathematicians
(TWMS Congress-2023)

ТЕЗИСЫ ДОКЛАДОВ
VII Всемирного Конгресса математиков
тюркского мира
(TWMS Congress-2023)

20–23 қыркүйек, 2023, Түркістан, Қазақстан
September 20–23, 2023, Turkestan, Kazakhstan
20–23 сентября, 2023, Туркестан, Казахстан



Түркі әлемі математиктерінің
VII Дүниежүзілік Конгресі
(TWMS Congress-2023)
**БАЯНДАМАЛАРЫНЫҢ
ТЕЗИСТЕРІ**

ABSTRACTS
of the VII World Congress of Turkic
World Mathematicians
(TWMS Congress-2023)

ТЕЗИСЫ ДОКЛАДОВ
VII Всемирного Конгресса математиков
тюркского мира
(TWMS Congress-2023)

20–23 қыркүйек, 2023, Түркістан, Қазақстан
September 20–23, 2023, Turkestan, Kazakhstan
20–23 сентября, 2023, Туркестан, Казахстан

TWO-DIMENSIONAL PROBLEM OF ANOMALOUS TRANSPORT IN A TWO-ZONE FRACTAL POROUS MEDIUM

BAKHTIYOR KHUZHAYOROV¹, AZIZBEK USMONOV², FAKHRIDDIN KHOLLIEV³

^{1,2,3} Samarkand State University, Samarkand, Uzbekistan.

e-mail: b.khuzhayorov@mail.ru; a.usmonov.91@mail.ru; surxon88@bk.ru

Two-zone media are very common in a macroscopically inhomogeneous media. In such a media, the process of solute transport proceed with the manifestation of internal mass transfer between different zones. This significantly changes the overall pattern of filtration and mass transfer [1]. The transfer equations here, unlike the previous ones, have fractional derivatives. Therefore, the object can be considered as a macroscopically inhomogeneous fractal medium.

In this case, the equation in the two-dimensional case is written as

$$\theta_m \frac{\partial c_m}{\partial t} + \gamma \theta_{im} \frac{\partial^\alpha c_{im}}{\partial t^\alpha} = \theta_m \left(D_{mx} \frac{\partial^\beta c_m}{\partial x^\beta} + D_{my} \frac{\partial^\beta c_m}{\partial y^\beta} \right) - v_{mx} \theta_m \frac{\partial c_m}{\partial x} - v_{my} \theta_m \frac{\partial c_m}{\partial y}, \quad (1)$$

where θ_m , θ_{im} - porosity; c_m , c_{im} - volumetric concentrations of the substance $\left(\frac{m^3}{m^3}\right)$; D_{mx} , D_{my} are the hydrodynamic dispersion coefficients in the moving zone $\left(\frac{m^{\beta+1}}{c}\right)$; v_{mx} , v_{my} - average solution velocity (m/c), the index m corresponds to the mobile zone, and im - to the immobile zone.

The presence of a stagnant (immobile) zone is taken into account on the basis of the kinetic equation

$$\gamma \theta_{im} \frac{\partial^\alpha c_{im}}{\partial t^\alpha} = \omega (c_m - c_{im}), \quad (2)$$

where is the γ mass transfer coefficient $[\gamma] = T^{\alpha-1}$, $[\omega] = T^{-1}$, $0 < \alpha \leq 1$, $0 < \beta \leq 2$.

The fields of pressures and filtration rates are also determined. For this, an anomalous filtration equation is used and it is derived on the basis of the anomalous Darcy law

$$\frac{\partial p}{\partial t} = X_x \frac{\partial^{1+\delta_1} p}{\partial x^{1+\delta_1}} + X_y \frac{\partial^{1+\delta_2} p}{\partial y^{1+\delta_2}}, \quad (3)$$

where χ_x and χ_y are the piezoconductivities in the directions x and y .

For equations (1) - (3), the problem of filtration and solute transport is fixed and solved numerically. The anomalous transfer is characterized by the order of the derivative in the diffusion term of the transfer equation and the mass transfer kinetics equation. Decreasing the order of the derivative β in the diffusion terms of the transport equation in both zones leads to "fast diffusion".

Keywords: Anomalous solute transport, diffusion, fractional derivative, porous media.

AMS Subject Classification: 76-10

REFERENCES

- [1] Khuzhayorov B.Kh., Makhmudov JM Flow of suspension in two- dimensional porous media with mobile and immobile liquid zones, *Journal of Porous Media*, 2010. Vol. 13, No. 5. P. 423-437.

Empty corners phenomenon for model of repulsive discrete electrical charges within convex polygons <i>Sabina Tagaeva</i>	333
The problem of optimal design of a circular arch <i>Serdar Nazarov, Muhametberdi Rakhimov</i>	334
The nonlinear differential game in the case of the l-catch <i>Ulmasjon Soyibboev, Mohisanam Turgunboyeva</i>	335
Математикалық және компьютерлік модельдеу	
Mathematical and computer modeling	
Математическое и компьютерное моделирование	
Investigation of the theory of own and forced oscillatory processes of different types of flat elements <i>A. Seitmuratov, Z. Yergalauova, S. Menlikhozhayev, A. Ibrayeva, G. Yensebayeva1</i>	337
Mathematical model of optimal management of the customs clearance process <i>Abdusobir Saidov</i>	338
Algebraic methods for solving recognition problems with non-crossing classes <i>Anvar Kabulov, Alimdzhani Babadzhanov, Islambek Saymanov</i>	340
Models, methods and algorithms for monitoring environmental impact on agricultural production <i>Anvar Kabulov, Ibrokhimali Normatov, Sattar Baizhumanov, Inomjon Yarashov</i>	341
Construction of high order mathematical models of the magic square and their computer simulation <i>Asan Baizakov, Zamirgul Kazakbaeva</i>	342
On the stability of nonlinear difference equations and some of their applications <i>Bakhadir Utebaev, Khoshruy Tokaeva, Aziza Utebaeva</i>	343
Two-dimensional problem of anomalous transport in a two-zone fractal porous medium <i>Bakhtiyor Khuzhayorov, Azizbek Usmonov, Fakhridin Kholliiev</i>	344
Generalized relaxational fractional differential model of fluid filtration in a porous medium <i>Bakhtiyor Khuzhayorov, Tursunpulot Djiyanov, Muhammad Zokirov</i>	345
Anomalous solute transport in an element of a fractured-porous medium <i>Bakhtiyor Khuzhayorov, Tursunpulot Dzhiyanov, Zarifzhon Eshdavlatov</i>	346
Solid particle dynamics simulation based on tlbm-ibb <i>Bakytzhan Zhumagulov, Ainur Zhumali, Dauren Zhakebayev</i>	347
Numerical study of two-component suspension filtration in a porous medium with dynamical factors <i>Bekzodjon Fayziev</i>	348

