

# Matrix theory - Assignment 8

Shreeprasad Bhat  
AI20MTECH14011

**Abstract—**This document illustrates finding subspaces of a vector spaces

Download latex-tikz from

<https://github.com/shreeprasadbhat/matrix-theory/blob/master/assignment8/>

## 1 PROBLEM

- Prove that only subspace of  $\mathbb{R}^1$  are  $\mathbb{R}^1$  and the zero subspace
- Prove that a subspace of  $\mathbb{R}^2$  is  $\mathbb{R}^2$ , or the zero subspace, or consists of all scalar multiples of some fixed vector in  $\mathbb{R}^2$ . (The last type of subspace is, intuitively, a straight line through the origin.)
- Can you describe the subspaces of  $\mathbb{R}^3$  ?

## 2 SOLUTION

- Let  $W \neq 0$  be subspace of  $\mathbb{R}^1$ . Then  $W$  is a nonempty subset of  $\mathbb{R}^1$  and there exist  $w \in W$  such that  $w \neq 0$  which gives us that there exist  $w^{-1}$ .

Let  $x \in \mathbb{R}^1$ . Since  $W$  is in  $\mathbb{R}^1$  we have that it is closed under scalar multiplication which gives us that  $(xw^{-1})w = x(w^{-1}w) = x.1 = x \in W$

Hence  $\mathbb{R}^1 \subset W$  and therefore  $W = \mathbb{R}^1$

Thus the only subspace of  $\mathbb{R}^1$  distinct of 0 is  $\mathbb{R}^1$  and therefore only subspaces of  $\mathbb{R}^1$  are 0 and  $\mathbb{R}^1$ .

- Clearly, 0 and  $\mathbb{R}^2$  itself are subspaces of  $\mathbb{R}^2$ . If  $u \neq 0$  and  $u \in \mathbb{R}^2$  then  $\text{span}\{\mathbf{u}\} = \{c\mathbf{u} : c \in \mathbb{R}\} = \text{set of all scalar multiples of } \mathbf{u}$  is a subspace of  $\mathbb{R}^2$ .

To show that these are the only subspaces of  $\mathbb{R}^2$ , assume that  $W \subset \mathbb{R}^2$  is any subspace of  $\mathbb{R}^2$ . Since  $W \subset \mathbb{R}^2$  is a subspace of  $\mathbb{R}^2$ , we have

that  $\mathbf{0} \in W$ . If  $W \neq \mathbf{0}$  then there is a vector  $\mathbf{u} \neq \mathbf{0}$  and  $\mathbf{u} \in W$ , and hence  $W$  contains  $c\mathbf{u}$  for every  $c \in \mathbb{R}$ . If  $W \neq \text{span}\{\mathbf{u}\}$ , then there is a vector  $\mathbf{v} \in W$  so that  $\mathbf{v} \neq k\mathbf{u}$  for any  $k \in \mathbb{R}$ .

Then  $\mathbf{z} = c\mathbf{u} + d\mathbf{v} \in \text{span}\{\mathbf{u}, \mathbf{v}\}$  for any  $c, d \in \mathbb{R}$ . Since  $W$  is a subspace  $c\mathbf{u}$  and  $d\mathbf{v} \in W$  for any  $c, d \in \mathbb{R}$ , and hence so does  $\mathbf{z} = c\mathbf{u} + d\mathbf{v}$ . Thus  $\mathbf{z} \in \text{span}\{\mathbf{u}, \mathbf{v}\} \implies \mathbf{z} \in W$ , and so  $\text{span}\{\mathbf{u}, \mathbf{v}\} \subset W \subset \mathbb{R}^2$ .

Let  $\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  be any vector in  $\mathbb{R}^2$ , and let  $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  and let  $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ . We show that there are real numbers  $c$  and  $d$  so that  $c\mathbf{u} + d\mathbf{v} = \mathbf{x}$

$$\begin{pmatrix} cu_1 \\ cu_2 \end{pmatrix} + \begin{pmatrix} dv_1 \\ dv_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad (2.0.1)$$

$$\begin{pmatrix} u_1 & v_1 \\ u_2 & v_2 \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad (2.0.2)$$

Since  $\mathbf{v} \neq k\mathbf{u}$  for any  $k \in \mathbb{R}$  and since  $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  assume that  $u_1 \neq 0$ , and since  $k\mathbf{u} \neq \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  assume that  $v_2 \neq 0$ . Then

$$A = \begin{pmatrix} u_1 & v_1 \\ u_2 & v_2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (2.0.3)$$

Hence  $A$  is row equivalent to  $I_2$  and so  $A$  is invertible and so (2.0.2) has unique solution for  $c$  and  $d$ . Thus for any  $\mathbf{x} \in \mathbb{R}^2$  we can find real numbers  $c$  and  $d$  such that  $\mathbf{x} = c\mathbf{u} + d\mathbf{v}$ . Hence  $\mathbf{x} \in \mathbb{R}^2 \implies \mathbf{x} \in \text{span}\{\mathbf{u}, \mathbf{v}\}$ . Thus  $\mathbb{R}^2 \subset \text{span}\{\mathbf{u}, \mathbf{v}\} \subset W \subset \mathbb{R}^2$ .

Hence  $\text{span}\{\mathbf{u}, \mathbf{v}\} = W = \mathbb{R}^2$ , and so the only subspace of  $\mathbb{R}^2$  are  $\mathbf{0}$ ,  $\mathbb{R}^2$ , and  $L = \{c\mathbf{u} : \mathbf{u} \neq \mathbf{0}, c \in \mathbb{R}\}$ .

c. Origin and  $\mathbb{R}^3$  itself are trivial subspaces of  $\mathbb{R}^3$ .

Every line through origin is a subspace of  $\mathbb{R}^3$ .

The other subspaces of  $\mathbb{R}^3$  are the planes passing through the origin.