

Matrix theory - Assignment 14

Shreeprasad Bhat
AI20MTECH14011

Abstract—This document proves properties on double dual

Download latex-tikz from

<https://github.com/shreeprasadbhat/matrix-theory/blob/master/assignment14/>

1 PROBLEM

Let S be a set, F a field, and $V(S; F)$ the space of all functions from S into F :

$$(f + g)(x) = f(x) + g(x)$$

$$(cf)(x) = cf(x)$$

Let W be any n -dimensional subspace of $V(S, F)$. Show that there exist points x_1, x_2, \dots, x_n in S and functions f_1, \dots, f_n in W such that $f_i(x_j) = \delta_{ij}$

2 SOLUTION

Given	<p>S is a set F is a field $V(S, F)$ is a linear functional such that W be n-dim subspace of $V(S, F)$.</p> <p>Also, $\delta_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$</p>
To prove	<p>$f_i(x_j) = \delta_{ij}$</p> <p>where $x_1, x_2, \dots, x_n \in S$ and $f_1, f_2, \dots, f_n \in W$</p>

Proof	<p>Let $\phi_x : W \rightarrow F$</p> <p>Suppose $\phi_x(f) = 0 \quad \forall x \in S \quad \& \quad f \in W$ $\implies f(x) = 0$</p> <p>If $\forall x, \phi_x(f) \neq 0$ for some $f \in W$ If $n > 0 \exists \in S$ such that $\phi_x(f) \neq 0$ for some $f \in W$ $\implies f_1(x_1) \neq 0$</p> <p>By scaling we can have $f_1(x_1) = 1$</p> <p>Hence $f_i(x_j) = \delta_{ij}$</p>
-------	--

3 FIGURE

Consider $x_1 = 1, x_2 = 0$, where $\{x_1, x_2\} \in S$.
Let $f_1(x) = x, f_2(x) = 1 - x$ where $\{f_1, f_2\} \in W$.




Fig. 0: Pictorial representation of functions and points in vector space