


# Towards Geospatial Intelligence with AI

Shreya Sharma

Data Science Researcher, NEC Corporation

# What is Geospatial Intelligence?




# Why do we care?

- Currently only 25% of the Earth's surface is seen by ground sensors





- Remote sensing (Satellite Earth Observation) is 'Eye from the Sky'


# How we see earth?



# How EO sees earth?



# Applications



So many datasets...

# My Research Work

- I. Ship Classification
- II. Change Detection (brief)



Microwave Image =>

# Motivation

Ship classification is a key application in of maritime surveillance

Helps in quick identification of ships involved in illegal activities



Overfishing



Oil-spills



Garbage dump



Smuggling

No. of illegal fishing cases detected by Japan Coast Guard




©Yomiuri Shinbun,  
28/02/2018

Microwave image is very useful as it is ‘cloud-free’ and image even in night

# Existing SAR Ship Classification Methods

## 1. Hand-crafted Features (HCF)-based




### Pros

- Intuitive features

### Cons

- Requires expert knowledge of ships

## 2. Convolutional Neural Network (CNN)-based



### Pros

- Does not require expert knowledge

### Cons



- Requires huge training data

These methods classify a ship based on its appearance


# Problem

Appearance of a ship varies with satellite viewing angle

Labelled images are very few to learn all possible variations



$$\theta_2 = 40^\circ$$




Only image information is insufficient for robust recognition

Appearance information is not sufficient to achieve robust classification



# Proposed Method

Use viewing angle as an additional information in a NN

Helps the neural network follow the appearance changes by learning an implicit relationship



# Experimental Set-up


Dataset: OpenSARShip*		Specifications			
Container			Satellite Sentinel-1		
			Resolution 20m		
			Polarization HH		
Image size 128x128		No. images 200/class	Image size 128x128		
Ground truth AIS + Marine Traffic		No. images 200/class			
Conventional Methods					
HCF	10 Features + SVM				
CNN	w/o incident angle				

\*Huang, L et al., "OpenSARShip: A dataset dedicated to Sentinel-1 ship interpretation," IEEE Journal of Sel. Top. in App. Earth Obs. and Rem. Sen. 11(1), 195-208 (2018).


# Experimental Result

Proposed method outperforms the conventional method

4.2% improved f-measure



25% reduced training data requirement



# Demo

# Change Detection - Motivation

Change detection enables us to understand dynamics of our planet

SAR images provide reliable visual evidence to detect changes

Image 1






Image 2





Change Map



# Conventional Method


Based on pixel-to-pixel difference followed by classification



Pixel-to-pixel  
difference

Classification


Change Map



Problem: Many False Changes in high resolution images!

# Proposed Method

Transform the images into features and compute distance between features



# Experimental Evaluation

- **Application**

- Parking Lot Monitoring

- **Baselines:**

- PCA + Kmeans [1]
- Sparse AE + Kmeans [2]

- **Evaluation Metrics**


- f-measure
- ROC-AUC

[1] T. Celik: Unsupervised change detection in satellite images using principal component analysis and k-means clustering, IEEE Geoscience and Remote Sensing Letters, vol. 6, no. 4, pp. 772-776, 2009.

[2] M. Gong., H. Yang, and P. Zhang: Feature learning and change feature classification based on deep learning for ternary change detection in SAR images, ISPRS Journal of Photogr. and Remote Sensing, no.129, pp.212-225, 2017.

Specifications	
Satellite	TerraSAR-X
Resolution	1m
Polarization	HH
No. of parking sites	5
No. of images/site	10
No. of pairs/site	9
Patch sizes	10x10, 16x16
Ground Truth (GT)	Manual interpretation


# Dataset



# Result [1/2] : f-measure

Siamese networks outperforms the conventional methods


Comparison of f-measure for 2 test pairs




# Result [2/2] : Change Maps

Siamese networks produce visually better change maps


Test Pair 1




GT




PCA-K



SAE-K




Siamese-10




Proposed


Test Pair 2




GT




PCA-K



SAE-K



Siamese-10



Siamese-16

# Conclusion

# Contents

1. What is geospatial intelligence?
2. Motivation
3. Electromagnetic Spectrum [1/2]
4. Electromagnetic Spectrum [2/2]
5. Applications
6. 2 applications (just intro)
7. Ship recognition- motivation
8. Existing methods
9. Problem
10. Proposed method
11. Experimental Evaluation
12. Result
13. Demo
14. Change detection
15. Conventional method and problem
16. Proposed method
17. Experimental Evaluation
18. Dataset
19. Result – 1
20. Result – 2
21. Conclusion