

Why do birds use green nest material? A systematic review and meta-analysis of experiments

Shreya Dimri^{a,*}, Tuba Rizvi^a, Júlio M.G. Segovia^a, Meinolf Ottensmann^b, Alfredo Sánchez-Tójar^{a,c,d}

^a*Department of Evolutionary Biology, Bielefeld University, Germany,*

^b*Department of Animal Behaviour, Bielefeld University, Germany,*

^c*CNC, Center for Neuroscience and Cell Biology, University of Coimbra, Portugal,*

^d*CIBB, Center for Innovative Biomedicine and Biotechnology, University of Coimbra, Portugal,*

*Corresponding author

Email address: shreya.dimri@uni-bielefeld.de (Shreya Dimri)

Abstract

Many animals build nests, often considered extended phenotypes that shape survival and reproduction beyond the builder’s body. Birds are key examples of nest builders, and some species add fresh green plant material to their nests. Yet, the adaptive value of this behaviour remains debated. Non-mutually exclusive hypotheses propose roles in courtship signalling, parasite defence, and direct enhancement of offspring condition through pharmacological effects independent of parasite reduction. Here, we conducted a pre-registered systematic review and meta-analysis of 28 experimental studies (published and unpublished), spanning seven bird species and 274 effect sizes, to test whether green nest material enhances fitness and to evaluate competing functional explanations. Our synthesis shows that green nest material can increase fitness in birds that exhibit this behaviour. However, this effect varies depending on the fitness proxy investigated, and its functional role remains inconclusive. We found no clear evidence supporting the courtship, nest protection, or drug hypothesis. Instead, experimental design was the strongest moderator of effect size variation, challenging the role of aromatic compounds in the fitness benefits of green nest material and highlighting the need for further research into the underlying mechanisms.

1. Introduction

Nest-building is fundamental across a diverse range of animals, including arthropods (e.g., ants, spiders), mammals (e.g., platypuses, rodents), and perhaps more famously, birds. Nests provide structural support for eggs, parents, and offspring, and protect them against predators and other environmental pressures, and are thus key to reproductive success and survival [1, 2, 3]. Most birds use organic plant materials such as twigs, branches, grass, and moss to build their nests. Some even incorporate inorganic (e.g., mud, stones), animal-derived (e.g., feathers, snake skins), and anthropogenic materials (e.g., plastic, cigarette butts) [4, 5, 6, 7]. These additions are considered non-structural and suggest that nests are more than just physical shelters—they can be complex tools shaped by both natural and sexual selection for multiple functions. Indeed, nests can play a key role in sexual display, and features such as nest size have been shown to act as sexually selected signals of parental quality (e.g., [8, 9]). Nest characteristics may even convey information interspecifically, for example, when brood parasites use host nest traits as cues to host quality[10]. Nest-building behaviour is increasingly recognised as an expression of the extended phenotype (reviewed by [11]), and phylogeny has been shown to be an important factor explaining nest morphology [12]. Yet, despite hundreds of scientific articles on bird nests and nesting behaviour being published each year, much remains unknown [13].

An intriguing nesting behaviour observed in some bird species is the addition of fresh, often aromatic, plant

material to the nest. This green nest material (*hereafter GNM; also called green plant material or greenery*) often includes leaves, sprigs, herbs, and branches. It is typically added from the start of nest building until egg-laying, although some species like the blue tit (*Cyanistes caeruleus*) keep adding it afterwards. Unlike the structural nest materials such as twigs and mud, GNM is selected from a small, non-random subset of local flora [14, 15, 16]. The selected plants often contain high levels of aromatic volatile compounds (commonly used by songbirds, [17]) or secondary metabolites such as oils and resins (commonly used by raptors, [18]). This suggests that GNM is unlikely to be gathered opportunistically, and collecting and replenishing these uncommon plants, sometimes over weeks, requires extra time and, in some cases, flying outside the territory [19]. Thus, the general costs associated with nest-building [20], and the additional effort, time, and energy devoted to adding these non-structural materials suggest an adaptive function of GNM.

Multiple hypotheses have been proposed to explain the adaptive function of GNM (reviewed in [21, 22]). First, the courtship hypothesis suggests that males use GNM as a visual or olfactory signal to attract females, maintain established pairs [23], or advertise nest-site occupancy to other males (e.g., [24, 25, 26]), with olfactory signalling being particularly relevant in cavity-nesting species [27]. For instance, a seminal study by [28] showed that male European starlings (*Sturnus vulgaris*) carry fresh plant material to their nests while singing and displaying to females. Subsequent studies in European and spotless starlings (*Sturnus unicolor*) supported this view by showing that females often prefer nests with fresh GNM, that unpaired males conspicuously deliver GNM during courtship [26], and increase the addition of GNM in the presence of potential mates [24]. Altogether, these studies offered support for the courtship hypothesis, which was later extended to account for the observed GNM addition solely by females in blue tits [29].

Second, the most widely tested hypothesis is the nest protection hypothesis, which proposes that the volatile compounds in GNM reduce parasitic and pathogenic loads in nests, enhancing offspring survival. This hypothesis was first suggested by [30] in a comparative study in raptors (Falconiformes). Soon after, experimental work in European starlings showed a preference for GNM containing insect-repellent and antimicrobial volatiles, and [14] proposed three key criteria for evaluating the nest protection hypothesis: selective plant use, bioactive chemical properties, and demonstrable adaptive benefits. Observational and experimental studies in raptors [18, 31] and passerines [32, 33, 17, 34] have shown mixed results. The drug hypothesis was later proposed because GNM has been shown, in some cases, to improve offspring condition without reducing parasite abundance [35, 36]. This alternative hypothesis suggests that GNM volatiles directly enhance offspring immune function and/or health independently of parasite or pathogen control [37]. While both the nest protection and drug hypotheses suggest that GNM enhances offspring survival, the drug hypothesis attributes this effect to a pharmacological mechanism, whereas the nest protection hypothesis emphasises parasite and pathogen repellency. Importantly, these two hypotheses are not mutually exclusive, and their potential effects on offspring survival are difficult to disentangle because improved nestling condition

may result either directly from pharmacological action or indirectly through reduced parasite or pathogen load. Therefore, in our meta-analysis, we combined them under the “parental care hypothesis”.

Other less explored hypotheses include a closely related female protection hypothesis proposed in blue tits [38, 39], suggesting that plant volatiles may primarily benefit the incubating female by reducing her parasite exposure and infection risk [40]. Several hypotheses have also been proposed that do not invoke the aromatic properties of GNM, including enhancement of nest crypticity [41], thermal insulation or sun protection for eggs and offspring [42], and a structural function for the added material [43]. However, these alternative hypotheses have received comparatively little empirical attention and therefore fall outside the scope of the present meta-analysis.

Despite decades of research, abundant observational and experimental studies, and two narrative reviews [21, 22], the overall adaptive significance of GNM, its generality across bird species, and the relative importance of each of the suggested mechanisms are still unknown. To address this gap, we conducted a pre-registered systematic review and meta-analysis of experiments testing the effect of GNM on various fitness proxies across birds. Our meta-analysis explicitly evaluates the overall evidence for the adaptive significance of GNM, and assesses the relative importance of the main suggested hypotheses: the courtship hypothesis and parental care hypothesis. In addition, we tested potential moderators predicted to explain the observed context-dependency (i.e., high heterogeneity among effect sizes). Specifically, we tested whether the effect of GNM varied depending on the experimental design used, the timing of GNM addition, the focal species tested, the category of fitness proxy investigated (e.g., morphological, reproductive, etc.), and the type of parasite studied. Our meta-analysis of experimental studies is the first quantitative synthesis testing the causal relationship between GNM and fitness across birds.

2. Methods

2.1. Registration and reporting

We pre-registered our study protocol on the Open Science Framework [44] before exploring or analysing the data. The pre-registration details our objectives, hypotheses, predictions, methods, and planned analyses. Unless stated otherwise, we adhered to the registered protocol. We followed the PRISMA-EcoEvo guidelines [45] for reporting this study (Table S1; Figure S1).

2.2. Information source and search

We conducted a systematic search on Web of Science Core Collection and Scopus on the 7th of September, 2022, and updated it on the 12th of August, 2024. The keyword search query was designed as a three-block query including terms related to our treatment of interest ((“green*” OR “herb*” OR “aromatic*”)

AND “nest*”), population of interest (“bird*” OR “aves” OR “avian” OR “ornithol*” OR “passerine*” OR “passeriform*” OR “songbird*” OR list of all bird genera) and to limit the search results to experimental studies only (“experiment*” OR “manipulat*”) (see complete search string in Figure S15). Although our search terms were in English, we did not restrict eligibility by language. We planned to include studies in English, Spanish, Portuguese or Hindi if they were retrieved, but all studies identified were in English. We searched through titles, abstracts and keywords without restrictions on publication date or study type. We validated our search using an initial library of 15 relevant articles identified before conducting the searches (Figure S2), ensuring all were retrieved. Additionally, we used the R package ‘litsearchr’ v1.0.0 [46], which confirmed our search queries. After removing 165 duplicates using the R package ‘revtools’ v0.4.1 [47], we identified 436 unique records.

2.3. Eligibility criteria

All titles and abstracts were equally allocated and screened independently by at least two of the three screeners (SD, TR, JMGS) using ‘revtools’ following a pre-designed decision tree with all inclusion/exclusion criteria (Figure S3). We included all studies on birds that experimentally manipulated the GNM (both aromatics and non-aromatic). Forty articles passed the title-and-abstract screening and were further evaluated during the full-text screening, where each article was independently screened by a single screener, with another screener verifying the procedure. We used the same criteria as above (Figure S3) with an additional evaluation of whether the variable of interest studied in the article was in our list of fitness proxies of interest (Table S2). All conflicts in screening decisions (1% for title-and-abstract screening, and none for full-text screening) were discussed and resolved. Twenty-three articles passed the full-text screening, and we identified three additional eligible articles recommended by two colleagues. In all, 26 published articles were eligible for our meta-analysis. We also obtained two additional unpublished studies (more details below; see also Figure S1 and Figure S4).

2.4. Data extraction

Data from each article were extracted by one of three authors (SD, TR, JMGS), with another screener verifying the procedure, and four authors (SD, TR, JMGS, AST) collectively discussing and revising unclear cases. We prioritised extracting primary data (i.e., means, variances and sample size for each control-treatment comparison) from text and tables, and if that was not possible, from figures (e.g., boxplots) using the R package ‘metaDigitise’ v1.0.1 [48]. If primary data were not available, we extracted inferential statistics (e.g., t and F values, contingency tables) to transform them into effect sizes (see below). Last, if these steps failed but the article had openly archived data, we calculated the means, standard deviations and sample sizes from each group ourselves (n = 4 studies). We extracted the number of nests in each group as the sample size rather than the number of individuals wherever possible (discussed below).

To capture differences in the type of experimental setup used in the studies, we coded each control-treatment comparison into an ‘experimental design’ moderator, with the following three levels: (i) non-aromatic GNM vs. aromatic GNM, (ii) no material vs. aromatic GNM, and (iii) no material vs. non-aromatic GNM. Where “non-aromatic GNM” refers to nests containing non-aromatic material (e.g., grass, moss, etc.), “aromatic GNM” refers to nests containing aromatic material added by birds, researchers, or both; and “no material” refers to nests that had no material added by researchers or where any plant material added by birds was removed. Importantly, our conclusions remained robust to the exclusion of the no material vs. non-aromatic GNM category ($n = 3$ studies; Table S3), so we kept this category in our dataset following our pre-registered plan. A fourth experimental design compared enhanced supplementation of GNM by the researchers (i.e., researchers + birds) to GNM added by birds alone ($n = 2$ studies). This experimental design represents a fundamentally different comparison focusing on potential additive effects of GNM, comparing different amounts of aromatic GNM. Therefore, we did not include this fourth category in the meta-regression testing the ‘experimental design’ moderator. In studies where three groups were compared ($n = 5$), we extracted all possible pairwise comparisons, and accounted for (shared group) non-independence in our analyses (more below).

For each study, we also extracted information related to the study subjects (bird species, geographical location of the study population) and the source from which we extracted data (e.g., figures, raw data files). We categorised the reported fitness proxies into 6 broad categories: physiology, morphology, reproduction, behaviour, parasite and pathogenic load, and phenology. For studies testing whether GNM affects the number of parasites or pathogens found in offspring and/or nests, we categorised them into arthropods (e.g., fleas) or microorganisms (e.g., bacteria). For a complete list of variables extracted, see Figure S7. During data extraction, we encountered some traits (e.g., Leukocyte measures, parent mass; $k = 15$ effect sizes, $n = 5$ studies) that we did not include in any of our analyses because it was unclear from the study and our own judgement how these traits related to fitness (see Table S2 for the reasons of exclusion).

We contacted four corresponding authors to obtain missing data (e.g., non-reported results, missing sample sizes). All four authors responded to our email, but only two provided the data as requested; one did not have the data anymore, and one did not give a conclusive response. Additionally, to locate grey literature (e.g., unpublished data), we used a standardised email template (Figure S5) to write to the corresponding author(s) of all the 23 articles that passed full-text screening asking for any potential unpublished data that could be included in our meta-analysis. We obtained unpublished data for a study on blue tits and a study on common buzzards (*Buteo buteo*; see description in Figure S6).

2.5. Effect size calculation

Using the means, standard deviations, and sample sizes for each control-treatment comparison, we calculated two effect sizes: a log response ratio (lnRR) and a standardised mean difference with heteroskedasticity correction (SMDH), using the ‘escalc’ function (measure = ‘ROM’ and ‘SMDH’, respectively, vtype = ‘LS’) in the R package ‘metafor’ v4.8-0 [49]. We specified the treatment group as the numerator and the control group as the denominator for lnRR, and treatment minus control for SMDH, such that positive values indicate a positive effect of GNM on fitness. In cases such as measures of parasite load or prevalence, laying or hatching dates, courtship time, or scab scores, the effect size sign was inverted before analysis to ensure that a positive value consistently meant an increase in fitness across all fitness proxies (i.e., to ensure comparability across effect sizes). Since SMDH can only be calculated when the degrees of freedom (i.e., n_1+n_2-2) are greater than 1, we had to exclude nine effect sizes from one study that did not fulfil that requirement.

For the few cases for which we could not extract means and variances but inferential statistics such as t and F values ($k = 4$, $n = 2$), we estimated SMD values from those statistics [50]. For dichotomous variables (e.g., nest success, recorded as successful vs unsuccessful and analysed as counts per group, $k = 4$, $n = 2$), we calculated log odds ratios (lnOR) using the ‘escalc’ function (measure = OR) from ‘metafor’ and subsequently converted them to SMD [50]. Since lnRR cannot be calculated from inferential statistics, dichotomous variables, or traits including zero or negative values ($k = 7$), the number of effect sizes included in the lnRR subset ($k = 259$, $n = 28$) is smaller than that of SMDH ($k = 274$, $n = 28$).

When calculating each effect size’s sampling variance, we used the number of nests per group, rather than the number of individuals, whenever possible, to account for within-nest non-independence due to shared genetic background and early-life environment. In cases where the same control or treatment group was used in multiple comparisons, sample sizes were adjusted before calculating effect sizes by dividing them by the number of times each group was used in a comparison to account for shared group non-independence. For cases where the authors of an original article reported a statistically non-significant difference between two groups but without providing numerical values ($k = 16$, $n = 2$), we assigned zero as the effect size for that comparison and imputed its sampling variance using the missing-case approach described in [51] for lnRR, and $\frac{n_1+n_2}{n_1 \times n_2}$ for SMDH [52].

2.6. Data analysis

We performed all analyses in R v4.5.1 [53]. All multilevel meta-analytical models and meta-regressions were fitted using ‘metafor’ v4.8-0 [49]. Sampling variances were modelled as variance-covariance matrices assuming a 0.5 correlation (ρ) between sampling variances from the same study [54] using the ‘vcalc’ function from ‘metafor’ (see Supplementary Appendix Figure S9 for sensitivity analyses showing that conclusions remain the same when using different values of ρ).

We originally considered the following levels of non-independence for our analyses: multiple effect sizes collected from (a) same study (Paper ID), (b) bird species (Species ID), (c) geographical locations (Population ID), (d) sub-populations within the same location (Experiment ID; e.g., birds studied at the same location for multiple years), (e) groups of individuals within a population (Group ID; e.g., males and females in a population, first clutch and second clutch), and (f) repeated measurement of the same trait for the same individuals (Repeated trait ID; e.g., chick mass on day 7, 14 and 21). In the end, our random effect structure deviated from our pre-registration because Experiment ID and Group ID represented nearly the same grouping structure, so we decided to leave out the one with lesser resolution (i.e., Experiment ID). Additionally, few effect sizes were measured repeatedly, making Repeated trait ID almost identical to Observation ID (Figure S8), so we excluded Repeated trait ID. Last, we decided to add Population ID since we noticed after extracting the data, but before any analyses, that several studies were performed in the same geographical locations. Unless stated otherwise, we fitted the following random effects in all our models to account for non-independence: Paper ID, Group ID, Species ID, Population ID, and an observation-level random effect (Observation ID).

Meta-analytic mean and hypotheses: To estimate the effect of GNM on fitness estimates across bird species, we fitted two separate sets of models for each effect size: lnRR and SMD(H). First, we fitted multilevel intercept-only meta-analytical models to test the overall effect of GNM on fitness estimates, its uncertainty and heterogeneity. We used the ‘rma.mv’ function from the ‘metafor’ package (method = ‘REML’ and test = ‘t’ for calculating the test statistics and confidence intervals for the moderators). Although we excluded traits for which their link with fitness is debated in the literature (e.g., sex ratio [55]; yolk hormones levels, [56]; more in Table S2) from our main analysis, we conducted a sensitivity analysis including these traits that confirmed the robustness of our results (Table S3; $k = 21$, $n = 8$). For all intercept-only models, we estimated the total amount of absolute heterogeneity (σ^2 ; i.e., the total variation not explained by sampling error), its sources or relative heterogeneity (I^2), mean-standardised metric (CVH2) and variance-mean-standardised metric of heterogeneity (M2) following the pluralistic approach suggested by [57].

We ran several sensitivity analyses to test the robustness of the overall results to the: (i) use of pre-registered bivariate multilevel meta-analytic model including both effect sizes as response variables; (ii) inclusion of traits for which their relationship with fitness was debated in the literature (non-pre-registered); (iii) removal of effect sizes that were originally reported as simply statistically non-significant and we assumed them to be zero (non-pre-registered); (iv) both (ii) and (iii) (non-pre-registered); (v) removal of effect sizes from non-aromatic vs. aromatic experimental design (non-pre-registered); (vi) removal of effect sizes from non-aromatic vs. no added material experimental design (non-pre-registered); (vii) removal of effect sizes from aromatic vs. non-aromatic material experimental design (non-pre-registered); (viii) removal of effect sizes that were calculated by transforming inferential statistics (pre-registered); (ix) removal of effect sizes that failed the Geary’s Test for lnRR (non-pre-registered); (x) removal of unpublished data (non-pre-registered); and (xi) use

of different values of correlation (ρ) between sampling variance for variance-covariance matrix (pre-registered). The number of effect sizes included in those sensitivity analyses ranged from 131 to 274 (Table S3).

To test the relative importance of the two main functional hypotheses suggested for GNM, that is, the courtship and the parental care hypothesis (which combines the nest protection and drug hypotheses), we ran a multilevel meta-regression including the moderator ‘hypothesis’ (levels: courtship, parental care, both, where ‘both’ refers to fitness proxies not uniquely assigned to either mechanism because they may reflect both courtship and parental care, e.g., offspring morphology or survival). For all meta-regressions, we estimated R^2_{marginal} , which corresponds to the percentage of heterogeneity explained by the moderators [58].

Exploratory meta-regressions: We performed additional pre-registered multilevel meta-regressions for which we did not have clear directional predictions. To understand the effect of the type of experimental design (i.e., whether the authors compared non-aromatic vs. aromatic, no material vs. aromatic, no material vs. non-aromatic), we ran a meta-regression with the type of ‘experimental design’ as the moderator. To test the effect of time of addition of GNM on fitness (levels: before egg hatching, after egg hatching, continuously throughout the nesting phase), we ran a meta-regression with ‘time of addition of GNM’ as the moderator. We also ran a meta-regression using ‘bird species’ as the moderator (levels: *Buteo buteo*, *Cyanistes caeruleus*, *Passer cinnamomeus*, *Parus major*, *Sturnus unicolor*, *Sturnus vulgaris*, *Tachycineta bicolor*) to understand if and how the effect of GNM on fitness might differ among species. Furthermore, we explored whether the effect of GNM on fitness varied across the different ‘category of fitness proxy’ studied (levels: behaviour, morphology, parasite and pathogenic load, phenology, physiology, reproduction) by running a meta-regression with this moderator. Last, for the data subset on parasites and pathogens ($k = 90$, $n = 16$), we ran a meta-regression with ‘type of parasite’ as the moderator (levels: arthropod, micro-organism).

In addition, following recommendations by [59] and [60], we performed three additional pre-registered multilevel meta-regressions to explore differences in internal validity across the studies associated with our conclusions (i.e., Risk of Bias, or RoB). Specifically, we evaluated the effects of (i) blinding (whether outcome assessors were blinded to the experimental group), (ii) random assignment to treatment, and (iii) partial reporting (whether all measured outcomes were completely reported or only a subset). For each study, each of these three moderators were coded (levels: yes, no) and included in a separate meta-regression for both lnRR and SMD(H).

Publication Bias Test: Publication bias occurs when a subset of research findings, most often statistically non-significant ones, are less likely to be published or published later [61]. Two types of publication biases are commonplace in ecology and evolutionary biology: small-study (i.e., small sample size studies reporting larger overall effect sizes due to missing effect sizes) and decline effects (i.e., effect sizes decreasing over time) [62]. We tested them for both lnRR and SMD(H) separately following [63]. To test for small-study effects, we ran

a multilevel meta-regression using the ‘square-root of the inverse of the effective sample size’ as the moderator (i.e., $\sqrt{(n_1 + n_2)/(4n_1n_2)}$). To test for decline effects (also known as time-lag bias, [64]), we ran a multilevel meta-regression using ‘year of publication’ (mean-centered) as the moderator. Throughout, 95% CI and 95% PI refer to 95% confidence intervals and 95% prediction intervals, respectively.

3. Results

We obtained 274 effect sizes from 28 studies conducted on 7 species across 17 geographical locations (Figure 1). About 81% of the effect sizes came from only three bird species (*Cyanistes caeruleus*, $k = 77$; *Sturnus unicolor*, $k = 60$; *Sturnus vulgaris*, $k = 86$). Our dataset included studies published between 1988 and 2024. The mean ($\pm$ standard deviation) number of effect sizes calculated per study was 9.8 ± 6.4 (median: 8.5, range: 2–24) and the mean effective sample size (i.e., the number of nests/individuals per effect size after accounting for shared-group non-independence) was 32.3 ± 31.1 (median: 22.0, range: 3–210 nests/individuals; 31 data points had a sample size of less than 5 nests in each group).




Figure 1: **Geographical distribution of the 17 study populations included in the meta-analysis testing the adaptive value of green nest material in birds.** Point size reflects the number of effect sizes calculated from each location (range: 3–34), and colour denotes species (see legend), with an inset zooming in on a dense cluster of data points.

3.1. Meta-analytic mean

Both lnRR and SMD(H) intercept-only meta-analyses showed an average positive effect of GNM on the fitness proxies; however, this overall effect was only statistically significant for SMD(H). For lnRR, nests with an experimental increase in GNM showed an average statistically non-significant increase in fitness estimates of 1.9% (Figure 2 A; lnRR = 0.019, 95% CI: [-0.012, 0.050], 95% PI: [-0.281, 0.319], p-value = 0.223, k = 238, n = 26) but the heterogeneity among effect sizes was high. Total raw heterogeneity (σ^2) was 0.023 ($Q = 2277.7$, $p < 0.001$), the measures of relative ($I^2_{total} = 98.6\%$) and standardised heterogeneity (CVH2_{total} = 63.59; M2_{total} = 0.98) all suggested high heterogeneity across effect sizes, exceeding the 75th percentile of empirically derived estimates from meta-analyses in ecology and evolution [57].

For SMD(H), nests with an experimental increase in GNM showed an average statistically significant increase in fitness estimates (Figure 2 B; SMD(H) = 0.179, 95% CI: [0.048, 0.310], 95% PI: [-0.829, 1.188], p-value = 0.008, k = 253, n = 26), but the heterogeneity among effect sizes was moderate to high. Total raw heterogeneity (σ^2) was 0.258 ($Q = 1114.7$, $p < 0.001$), and both relative ($I^2_{total} = 66.1\%$) and standardised heterogeneity metrics (CVH2_{total} = 8.02 ; M2_{total} = 0.89) lay between the 50th and 75th percentile of empirically derived estimates [57].

3.2. Functional hypotheses

Our multilevel meta-regressions found no clear support for either of the two hypotheses separately (Figure 3). The estimates for the courtship hypothesis (lnRR_{courtship} = 0.039, 95% CI: [-0.172, 0.250], 95% PI: [-0.329, 0.407], p-value = 0.716, k = 8, n = 2; SMD(H)_{courtship} = 0.171, 95% CI: [-0.360, 0.702], 95% PI: [-0.964, 1.305], p-value = 0.527, k = 8, n = 2), and the “parental care hypothesis” (lnRR_{parental_care} = -0.010, 95% CI: [-0.065, 0.045], 95% PI: [-0.316, 0.296], p-value = 0.716, k = 106, n = 19; SMD(H)_{parental_care} = 0.109, 95% CI: [-0.060, 0.278], 95% PI: [-0.908, 1.126], p-value = 0.204, k = 115, n = 19) were statistically non-significant. However, the third level in this meta-regression (“Both”), which contained about half of all effect sizes that could not be unequivocally assigned to only one hypothesis, showed an average positive GNM effect on the fitness estimates, statistically significantly so for SMD(H) (lnRR_{both} = 0.028, 95% CI: [-0.006, 0.062], 95% PI: [-0.275, 0.331], p-value = 0.110, k = 124, n = 22; SMD(H)_{both} = 0.231, 95% CI: [0.079, 0.382], 95% PI: [-0.784, 1.245], p-value = 0.003, k = 130, n = 23). None of the three categories differed statistically significantly from each other (p-value_{lnRR} > 0.211 and p-value_{SMD(H)} > 0.178 in all cases), and the moderator explained little heterogeneity ($R^2_{\text{marginal}(\text{lnRR})} = 1.6\%$ and $R^2_{\text{marginal}(\text{SMD(H)})} = 1.4\%$).

3.3. Exploratory meta-regressions

First, the overall effect of GNM on fitness estimates differed largely depending on the experimental design used in each study (Figure 4 A,B). The overall effect size for the comparison between no material and aromatic GNM was positive and statistically significant (lnRR_{no-material-vs-aromatic} = 0.111, 95% CI: [0.021,

0.200], 95% PI: [-0.225, 0.446], p-value = 0.016, k = 90, n = 12 and $SMD(H)_{\text{no-material-vs-aromatic}} = 0.712$, 95% CI: [0.309, 1.114], 95% PI: [-0.826, 2.250], p-value = 0.001, k = 94, n = 12], whereas, the comparison between non-aromatic GNM and aromatic GNM was statistically non-significant (Table 1). The comparison between no material and non-aromatic GNM was negative and, although based on only 3 studies, statistically significant for $\ln RR$ ($\ln RR_{\text{no-material-vs-non-aromatic}} = -0.349$, 95% CI: [-0.504, -0.195], 95% PI: [-0.708, 0.009], p-value < 0.001, k = 10, n = 3; Table 1). All levels differed statistically significantly from each other except one (p-value $_{\ln RR} < 0.004$ and p-value $_{SMD(H)} < 0.001$ in all cases, except $SMD(H)_{\text{no-material-vs-aromatic}}$ vs. $SMD(H)_{\text{no-material-vs-non-aromatic}}$ p-value = 0.140), and the moderator experimental design explained a large proportion of heterogeneity ($R^2_{\text{marginal}(\ln RR)} = 27.8\%$ and $R^2_{\text{marginal}(SMD(H))} = 24.4\%$).

Second, the overall positive effect of GNM on fitness outcomes tended to be larger when GNM was added continuously throughout the nesting phase (Figure 4 C,D). The overall effect was positive and statistically significant for $SMD(H)$ when material was continuously and experimentally added to the nest ($\ln RR_{\text{continuously}} = 0.034$, 95% CI: [-0.014, 0.083], 95% PI: [-0.270, 0.338], p-value = 0.165, k = 95, n = 12; $SMD(H)_{\text{continuously}} = 0.302$, 95% CI: [0.069, 0.534], 95% PI: [-0.726, 1.329], p-value = 0.011, k = 105, n = 12), whereas the effect was small and statistically non-significant when added before egg hatching (Table 1). Last, the overall effect also tended to be positive for $SMD(H)$ when GNM was added after egg hatching only, although only marginally so ($SMD(H)_{\text{after_egg_hatching}} = 0.267$, 95% CI: [-0.024, 0.559], 95% PI: [-0.775, 1.310], p-value = 0.072, k = 57, n = 5; Table 1). The three categories did not differ statistically significantly from each other (p-value $_{\ln RR} > 0.307$ and p-value $_{SMD(H)} > 0.243$ in all cases). The timing of GNM addition explained little heterogeneity ($R^2_{\text{marginal}(\ln RR)} = 1.2\%$ and $R^2_{\text{marginal}(SMD(H))} = 4.5\%$).

Third, although none of the species-specific mean estimates were statistically significant, most were positive (Figure S13, Table 1), and bird species explained a small amount of heterogeneity ($R^2_{\text{marginal}(\ln RR)} = 5.3\%$ and $R^2_{\text{marginal}(SMD(H))} = 8.5\%$). Fourth, most fitness proxy categories had positive mean estimates, but only morphology was statistically significant, and only for $SMD(H)$ ($SMDH_{\text{morphology}} = 0.310$, 95% CI: [0.112, 0.509], 95% PI: [-0.723, 1.343], p-value = 0.002, k = 65, n = 15; Figure S13; Table 1) and fitness proxy category accounted for a small amount of heterogeneity ($R^2_{\text{marginal}(\ln RR)} = 3.7\%$ and $R^2_{\text{marginal}(SMD(H))} = 2.4\%$). Fifth, for the subset of outcomes relating to pathogen and parasitic load, estimates for arthropods and micro-organisms did not differ statistically significantly (Figure S13 ; Table 1) and the type of parasite or pathogen explained little heterogeneity ($R^2_{\text{marginal}(\ln RR)} = 0.4\%$ and $R^2_{\text{marginal}(SMD(H))} = 1.6\%$). Lastly, the evaluation of potential Risk of Bias categories (i.e., blinding, random assignment to each treatment, and partial reporting) was limited by the small number of studies in some categories (details in Table S4).

3.4. Publication Bias

We found no conclusive evidence for the existence of publication bias in the published literature (Figure S14). There was no statistical evidence for small-study effect for lnRR (lnRR_{intercept} = 0.003, 95% CI: [-0.066, 0.072], p-value = 0.926, lnRR_{slope} = 0.079, 95% CI: [-0.215; 0.373], p-value = 0.595, k = 221, n = 24, R²_{marginal(lnRR)} = 0.3%). For SMD(H), there was a tendency for effect sizes to be larger when sample size was smaller (SMDH_{slope} = 1.393, 95% CI: [-0.143, 2.929], p-value = 0.075, k = 235, n = 24, R²_{marginal(SMD(H))} = 8.3%) with the overall adjusted effect decreasing substantially (SMDH_{intercept} = -0.035, 95% CI: [-0.319, 0.249], p-value = 0.809). In addition, since published effect sizes did not change over time, our analyses suggest no clear evidence of decline effects in published effect sizes (see Figure S14).

4. Discussion

Our meta-analysis of experimental studies shows that green nest material (GNM) has a small positive effect on fitness in birds. Our results were generally consistent across two effect size metrics and robust to several sensitivity analyses. However, moderate-to-high heterogeneity among effect sizes indicates that the benefits of GNM are context-dependent. We found no clear support for either of the two functional hypotheses traditionally proposed, the courtship hypothesis and the parental care hypothesis (i.e., nest protection and drug hypotheses combined). While several studies investigate the courtship hypothesis, only a few fitness proxies could be exclusively linked to this mechanism (e.g., courtship time, male provisioning), and these showed no effect of GNM. Similarly, fitness proxies exclusively associated with the parental care hypothesis (e.g., parasite loads, nestling health), also showed no effect of GNM. This finding is particularly surprising because the parental care hypothesis has been most frequently tested. While studies testing the courtship hypothesis have suggested that the aromatic volatile compounds could help receivers of the signal detect the presence of GNM in dark cavity nests [26], it is the parental care hypothesis that has stressed the importance of these compounds [65, 66, 67], perhaps explaining their widespread consideration. Our findings challenge this assumption since experimental designs comparing aromatic with non-aromatic GNM, which account for more than half of all tests, did not show a clear fitness effect, whereas comparisons between aromatic GNM and no material did. By contrast, comparisons between non-aromatic GNM and no material suggested a negative effect of non-aromatic GNM on fitness, although this result should be interpreted cautiously because it was based on only three studies (k = 10 effect sizes). Notably, experimental design alone explained about one-quarter of the observed heterogeneity (R²_{marginal} 24-28%).

One possible explanation is that aromatic vs. non-aromatic isolates the effects of aromaticity or volatile compounds, which may be weak, context-dependent (e.g., dependent on parasite pressure, [14]), or short-lived due to rapid decay of volatiles [15] and may require repeated replenishment [68]. By contrast, aromatic vs. no material, capture the net effect of adding green material itself, including potential non-chemical and volatile

effects, which may therefore show a clearer overall benefit. Such benefits could also relate to plant morphology such as shape or colour. In addition, only proxies that could not be clearly assigned to either of the functional hypotheses (e.g., survival rate, nestling morphology), and were therefore categorized as being associated with “both” hypotheses, showed a clear positive effect. This ambiguity reflects the difficulty of linking many fitness proxies to an underlying function because they can be shaped by multiple overlapping processes. For example, nestling mass, a commonly measured fitness proxy, could be influenced by pathogen load or direct physiological effects of plant compounds, even though our synthesis found no consistent overall evidence for parasite reduction or health benefits. It could also respond to altered parental investment through differential allocation, making such broad proxies of fitness difficult to assign to a single function.

Heterogeneity in the overall effect of GNM was mainly associated with within-study variation, with little to no among-study variation. This suggests that, once methodological (e.g., experimental design) and biological factors (e.g., species identity, category of fitness proxy) are accounted for, the overall effect found would be potentially generalisable across studies. In addition to experimental design, we tested the timing of GNM addition ($R^2_{\text{marginal}} = 1\text{-}5\%$). Continuous supplementation throughout the nesting period had stronger positive effects than supplementation restricted to before or after hatching. Although not conclusive, this pattern is more consistent with the parental care hypothesis than with the courtship hypothesis. In blue tits the concentration of active compounds released by the aromatic plants can decrease within 24 hours [15], and females often replace experimentally removed herbs [39, 68]. This repeated addition may help maintain an environment rich in volatiles that is hostile to parasites. However, fresh GNM could also improve fitness through structural or microclimate benefits [42, 2]. In addition, regular replenishment of GNM by birds may help nest lining as material desiccates, thereby improving insulation and humidity buffering [69, 42] and diluting faecal matter or food debris in the nest (i.e., hygiene function, [70]). This microclimate stability may accelerate nestling development while reducing thermoregulatory costs for the parents [71, 72].

Among the biological moderators tested, species identity seemed the most important one ($R^2_{\text{marginal}} = 5\text{-}9\%$), providing some evidence for species-specific effects; however, note that three species were represented in only one study each. Such variation across species is not unexpected, since some (e.g., *Tachycineta bicolor*) do not naturally incorporate GNM in their nests, while the other species do. Differences across species in life-history traits, nest construction, or habitat condition (such as parasite load, availability of resources), and their interplay likely influence how GNM affects fitness. The second most important biological moderator was the fitness proxy category ($R^2_{\text{marginal}} = 2\text{-}4\%$), potentially indicating differences in validity across fitness proxies or reflecting differences among underlying mechanisms. Among all categories, morphological outcomes (e.g., nestling mass, tarsus length) showed the largest positive effects, suggesting that GNM has a more pronounced effect on offspring development or growth. Morphology was the second most measured fitness proxy across studies, likely because these traits can be obtained easily and non-invasively in the field as

compared to long-term demographic (e.g., survival, recruitment) or physiological assays (e.g., blood-based immune or endocrine measures). However, while widely used as an indicator of offspring condition, these morphological measures, like many other outcomes in the literature, remain indirect fitness proxies, and their relationship to recruitment or survival can be context dependent [73]. This could also reflect that morphology responds quickly to within-nest conditions, whereas effects on long-term survival may be diluted by subsequent environmental variations [74]. All other fitness proxy categories, including physiology, reproduction, phenology, and behaviour, showed small to moderate positive effects, whereas proxies related to parasite or pathogen load, despite being the most commonly investigated, showed weak and inconsistent effects. Parasite reduction is central to one of the two mechanisms within the parental care hypothesis (the nest protection hypothesis, [30, 14]). Our results further emphasise that despite its prominence, there is a lack of clear overall evidence for the parental care hypothesis. The other hypothesis categorised within the parental care hypothesis (i.e., the drug hypothesis, [35, 19]), proposes an increase in offspring survival, with volatile compounds directly enhancing offspring immune function and/or health independent of parasite or pathogen control. While the largest positive effects of morphology could support the drug hypothesis, it should be noted that (i) the mechanism linking volatiles to enhanced offspring immune function or health is unclear, (ii) our results suggest that non-aromatic GNM may play a role, and (iii) several other mechanisms could also be at play such as nest microclimate and incubation conditions [75] or parental investment.

When interpreting our results, it is important to acknowledge some constraints of the study and the available evidence. The overall effect of GNM differed between the two effect size metrics, which make distinct assumptions regarding the underlying biological processes and the statistical properties. These divergent results highlight the importance of choosing appropriate effect size metrics for a meta-analysis that align with the biological data at hand or in cases where several different types of measures are present, using both [76]. We found high reporting quality in the primary studies included in our meta-analysis, with only 0.06% partial reporting, but some evidence of small-study effects. After adjusting for this, the effect of GNM was no longer significant. To reduce the influence of small-study effects we contacted all authors for unpublished data, but could only find access to two studies. In addition, many studies had very low sample sizes (41% of the effect sizes were based on fewer than 10 nests per group), which reduces power and increases heterogeneity [77]. Furthermore, studies remain heavily biased towards just three bird species, limiting our ability to infer broad patterns of this behaviour. Expanding the scope beyond the currently well-studied taxa could be informative, but in practice can be limited by feasibility constraints such as access to nests. Targeting species with diverse nesting ecologies and habitat conditions can be useful in understanding the adaptive benefits of this behaviour.

5. Conclusion

After more than 40 years of research on this topic, ours is the first quantitative synthesis confirming an overall effect of green nest material on fitness in birds. Despite moderate-to-high heterogeneity, our results suggest that the effect may be generalisable across studies once methodological and biological factors within studies are accounted for. However, our small-study effects analysis, although non-conclusively, indicates that accounting for potentially missing studies could cause the observed effect to disappear, despite our efforts to include unpublished work. Moreover, our study clearly emphasises that the functions remain largely unclear and challenges the assumption that volatile compounds alone explain the relationship between green nest material and fitness. Future experimental designs should focus on decoupling volatiles from structural or microclimate effects, while quantifying mediators (e.g., nest microclimate, parasite load) and prioritising fitness proxies directly linked to fitness (e.g., fledging success, recruitment), as well as testing explicitly, rather than assuming, the role of volatiles. Replications across parasite and climatic gradients, together with cross-fostering experiments, could allow us to understand the context dependency and distinguish nest-level effects from parental quality. In addition, theoretical work could further solidify the field’s foundations and yield testable predictions. Cost-benefit models of GNM use to predict when net benefits could occur and why effects vary among species and environments could help generate predictions (e.g., benefits only above certain parasite pressure or under certain temperature or humidity conditions) that are directly testable with multi-site experiments. Altogether, our synthesis contributes to a more nuanced understanding of the adaptive significance of one of the most fascinating and most studied natural structures—bird nests, opening doors to several questions that remain unanswered.

5.1. Data and Code Availability Statement

All data and code associated with this project can be found on GitHub at [https://github.com/shreyadimri/Green_Nest_Material]. It is also available on Zenodo [**insert link later**].

5.2. Author contributions

SD: Conceptualisation, Data curation, Formal analysis, Investigation (lead), Methodology, Project administration, Software, Validation, Visualization, Writing - original draft (lead), review & editing; **TR:** Conceptualisation, Investigation (equal), Writing - review & editing; **JMGS:** Conceptualisation, Investigation (equal), Writing - review & editing; **MO:** Investigation (supporting), Validation; **AST:** Conceptualisation, Funding acquisition, Methodology, Supervision, Writing - original draft (supporting), review & editing

5.3. Conflict of interest

Authors declare no competing interests.

5.4. Acknowledgements

We thank Klaus Reinhold for his support. We also thank all the authors who replied to our emails and who addressed our questions. We are particularly grateful to those who kindly shared their data: Adèle Mennerat, Dave Shutler and Gustavo Tomás Gutiérrez.

5.5. Funding

SD and AST were funded by the DFG (Deutsche Forschungsgemeinschaft) as part of the SFB TRR 212 (NC³) Second Phase 2022-2025 (Project number 316099922), and MO by the First Phase 2018-2021 (Project number 396780709). AST was also partially supported by Portuguese national funds via Fundação para a Ciência e a Tecnologia (FCT) under projects LA/P/0058/2020, UID/PRR/4539/2025 and UID/04539/2025, and project EXCELSIOR, funded by the EU's Horizon Europe under Grant Agreement No. 101087416. JMGS was financially supported by a Walter Benjamin Fellowship (Project number 528240618) from the DFG. TR was financially supported by a Research Grant for Doctoral Programmes by the DAAD (Deutscher Akademischer Austauschdienst).

6. References

- [1] M. Hansell, Bird Nests and Construction Behaviour, Cambridge University Press, Cambridge, UK, 2000. doi:10.1017/CB09781139106788.
- [2] M. C. Mainwaring, I. R. Hartley, M. M. Lambrechts, D. C. Deeming, The design and function of birds' nests, Ecology and Evolution 4 (20) (2014) 3909–3928. doi:10.1002/ece3.1054.
- [3] D. C. Deeming, Nest construction in mammals: a review of the patterns of construction and functional roles, Philosophical Transactions of the Royal Society B: Biological Sciences 378 (1884) (2023) 20220138. doi:10.1098/rstb.2022.0138.
- [4] F. Sergio, J. Blas, G. Blanco, A. Tanferna, L. López, J. A. Lemus, F. Hiraldo, Raptor nest decorations are a reliable threat against conspecifics, Science 331 (6015) (2011) 327–330. doi:10.1126/science.1199422.
- [5] M. Suárez-Rodríguez, I. López-Rull, C. Macías García, Incorporation of cigarette butts into nests reduces nest ectoparasite load in urban birds: new ingredients for an old recipe?, Biology Letters 9 (1) (2013) 20120931. doi:10.1098/rsbl.2012.0931.
- [6] Z. Jagiello, S. J. Reynolds, J. Nagy, M. C. Mainwaring, J. D. Ibáñez-Álamo, Why do some bird species incorporate more anthropogenic materials into their nests than others?, Philosophical Transactions of the Royal Society B: Biological Sciences 378 (1884) (2023) 20220156. doi:10.1098/rstb.2022.0156.

- [7] C. Sheard, L. Stott, S. E. Street, S. D. Healy, S. Sugawara, K. N. Lala, Anthropogenic nest material use in a global sample of birds, *Journal of Animal Ecology* 93 (6) (2024) 691–704. doi:10.1111/1365-2656.14078.
- [8] J. Moreno, M. Soler, A. P. Møller, M. Lindén, The function of stone carrying in the black wheatear, *Oenanthe leucura*, *Animal Behaviour* 47 (6) (1994) 1297–1309. doi:10.1006/anbe.1994.1178.
- [9] H. Hoi, B. Schleicher, F. Valera, Nest size variation and its importance for mate choice in penduline tits, *Remiz pendulinus*, *Animal Behaviour* 51 (2) (1996) 464–466. doi:10.1006/anbe.1996.0046.
- [10] J. J. Soler, M. Soler, A. P. Møller, J. G. Martínez, Does the great spotted cuckoo choose magpie hosts according to their parenting ability?, *Behavioral Ecology and Sociobiology* 36 (3) (1995) 201–206. doi:10.1007/bf00177797.
- [11] J. Moreno, Avian nests and nest-building as signals, *Avian Biology Research* 5 (4) (2012) 238–251. doi:10.3184/175815512x13534385822786.
- [12] D. M. Perez, L. T. Manica, I. Medina, Variation in nest-building behaviour in birds: a multi-species approach, *Philosophical Transactions of the Royal Society B: Biological Sciences* 378 (1884) (2023) 20220145. doi:10.1098/rstb.2022.0145.
- [13] M. Jezierski, R. B. J. Benson, W. J. Smith, E. E. Saupe, S. M. Clegg, A wide range of abiotic and biotic variables leaves most variation in bird nest architecture unexplained, *Proceedings of the Royal Society B: Biological Sciences* 292 (2057) (2025) 20252013. doi:10.1098/rspb.2025.2013.
- [14] L. Clark, J. R. Mason, Use of nest material as insecticidal and anti-pathogenic agents by the European Starling, *Oecologia* 67 (2) (1985) 169–176. doi:10.1007/bf00384280.
- [15] C. Petit, M. Hossaert-McKey, P. Perret, J. Blondel, M. M. Lambrechts, Blue tits use selected plants and olfaction to maintain an aromatic environment for nestlings, *Ecology Letters* 5 (4) (2002) 585–589. doi:10.1046/j.1461-0248.2002.00361.x.
- [16] J. Garrido-Bautista, J. A. Ramos, S. Arce, P. Melero-Romero, R. Ferreira, C. Santos-Baena, H. R. Guímaro, C. Martín-Villegas, G. Moreno-Rueda, A. C. Norte, Is there a role for aromatic plants in blue tit *Cyanistes caeruleus* nests? Results from a correlational and an experimental study, *Behavioral Ecology and Sociobiology* 77 (10) (2023) 118. doi:10.1007/s00265-023-03393-9.
- [17] A. Mennerat, P. Mirleau, J. Blondel, P. Perret, M. M. Lambrechts, P. Heeb, Aromatic plants in nests of the blue tit *Cyanistes caeruleus* protect chicks from bacteria, *Oecologia* 161 (4) (2009) 849–855. doi:10.1007/s00442-009-1418-6.
- [18] D. Ontiveros, J. Caro, J. M. Pleguezuelos, Green plant material versus ectoparasites in nests of Bonelli's eagle, *Journal of Zoology* 274 (1) (2008) 99–104. doi:10.1111/j.1469-7998.2007.00364.x.

- [19] A. Mennerat, P. Perret, M. M. Lambrechts, Local individual preferences for nest materials in a passerine bird, *PLoS ONE* 4 (4) (2009) e5104. [doi:10.1371/journal.pone.0005104](https://doi.org/10.1371/journal.pone.0005104).
- [20] M. C. Mainwaring, I. R. Hartley, The energetic costs of nest building in birds, *Avian Biology Research* 6 (1) (2013) 12–17. [doi:10.3184/175815512x13528994072997](https://doi.org/10.3184/175815512x13528994072997).
- [21] A. Dubiec, I. Gózdź, T. D. Mazgajski, Green plant material in avian nests, *Avian Biology Research* 6 (2) (2013) 133–146. [doi:10.3184/175815513X13615363233558](https://doi.org/10.3184/175815513X13615363233558).
- [22] J. F. Scott-Baumann, E. R. Morgan, A review of the nest protection hypothesis: does inclusion of fresh green plant material in birds’ nests reduce parasite infestation?, *Parasitology* 142 (8) (2015) 1016–1023. [doi:10.1017/S0031182015000189](https://doi.org/10.1017/S0031182015000189).
- [23] J. P. Veiga, V. Polo, J. Viñuela, Nest green plants as a male status signal and courtship display in the spotless starling, *Ethology* 112 (2) (2006) 196–204. [doi:10.1111/j.1439-0310.2006.01148.x](https://doi.org/10.1111/j.1439-0310.2006.01148.x).
- [24] M. Eens, R. Pinxten, R. F. Verheyen, Function of the song and song repertoire in the European Starling *Sturnus vulgaris*: an Aviary Experiment, *Behaviour* 125 (1–2) (1993) 51–66. [doi:10.1163/156853993x00182](https://doi.org/10.1163/156853993x00182).
- [25] H. Gwinner, The function of green plants in nests of the European Starling *Sturnus vulgaris*, *Behaviour* 134 (5–6) (1997) 337–351. [doi:10.1163/156853997X00575](https://doi.org/10.1163/156853997X00575).
- [26] L. Brouwer, J. Komdeur, Green nesting material has a function in mate attraction in the European Starling, *Animal Behaviour* 67 (3) (2004) 539–548. [doi:10.1016/j.anbehav.2003.07.005](https://doi.org/10.1016/j.anbehav.2003.07.005).
- [27] S. Y. Beasley, R. Freire, M. N. Callan, M. Massaro, The influence of plant scents on nest box inspection by eastern rosella (*Platycercus eximius*), *Emu - Austral Ornithology* 124 (2) (2024) 165–176. [doi:10.1080/01584197.2024.2323920](https://doi.org/10.1080/01584197.2024.2323920).
- [28] B. Kessel, A study of the breeding biology of the European Starling *Sturnus vulgaris* in North America, *The American Midland Naturalist* 58 (2) (1957) 257–331. [doi:10.2307/2422615](https://doi.org/10.2307/2422615).
- [29] G. Tomás, S. Merino, J. Martínez-de la Puente, J. Moreno, J. Morales, J. Rivero-de Aguilar, Nest size and aromatic plants in the nest as sexually selected female traits in blue tits, *Behavioral Ecology* 24 (4) (2013) 926–934. [doi:10.1093/beheco/art015](https://doi.org/10.1093/beheco/art015).
- [30] P. H. Wimberger, [The use of green plant material in bird nests to avoid ectoparasites](#), *The Auk* 101 (3) (1984) 615–618.
URL <https://www.jstor.org/stable/4086620>

- [31] C. R. Dykstra, J. L. Hays, M. M. Simon, Selection of fresh vegetation for nest lining by red-shouldered hawks, *The Wilson Journal of Ornithology* 121 (1) (2009) 207–210. doi:10.1676/08-035.1.
- [32] L. Lafuma, M. M. Lambrechts, M. Raymond, Aromatic plants in bird nests as a protection against blood-sucking flying insects?, *Behavioural Processes* 56 (2) (2001) 113–120. doi:10.1016/s0376-6357(01)00191-7.
- [33] R. D. Dawson, Does fresh vegetation protect avian nests from ectoparasites? an experiment with tree swallows, *Canadian Journal of Zoology* 82 (7) (2004) 1005–1010. doi:10.1139/z04-076.
- [34] M. Quiroga, J. C. Rebores, A. H. Beltzer, Host use by *Philornis* sp. in a passerine community in central Argentina, *Revista Mexicana de Biodiversidad* 83 (1) (2012). doi:10.22201/ib.20078706e.2012.1.1137.
- [35] H. Gwinner, M. Oltrogge, L. Trost, U. Nienaber, Green plants in starling nests: effects on nestlings, *Animal Behaviour* 59 (2) (2000) 301–309. doi:10.1006/anbe.1999.1306.
- [36] H. Gwinner, S. Berger, European starlings: nestling condition, parasites and green nest material during the breeding season, *Journal of Ornithology* 146 (4) (2005) 365–371. doi:10.1007/s10336-005-0012-x.
- [37] A. Mennerat, P. Perret, P. Bourgault, J. Blondel, O. Gimenez, D. W. Thomas, P. Heeb, M. M. Lambrechts, Aromatic plants in nests of blue tits: positive effects on nestlings, *Animal Behaviour* 77 (3) (2009) 569–574. doi:10.1016/j.anbehav.2008.11.008.
- [38] R. J. Cowie, S. A. Hinsley, Timing of return with green vegetation by nesting blue tits *Parus caeruleus*, *Ibis* 130 (6) (1988) 553–555. doi:10.1111/j.1474-919x.1988.tb02722.x.
- [39] J. Banbura, J. Blondel, H. de Wilde-Lambrechts, P. Perret, Why do female blue tits *Parus caeruleus* bring fresh plants to their nests?, *Journal für Ornithologie* 136 (2) (1995) 217–221. doi:10.1007/bf01651244.
- [40] J. García-Campa, S. González-Braojos, J. Morales, Nest secondary plants and their associations with haemosporidian blood parasites in blue tit females, *Parasitology* 151 (10) (2024) 1126–1136. doi:10.1017/s0031182024000775.
- [41] A. F. Skutch, The incubation patterns of birds, *Ibis* 99 (1) (1957) 69–93. doi:10.1111/j.1474-919x.1957.tb01934.x.
- [42] J. A. L. Mertens, Thermal conditions for successful breeding in great tits *Parus major*, *Oecologia* 28 (1) (1977) 1–29. doi:10.1007/bf00346834.
- [43] D. M. Lyons, J. A. Mosher, Morphological growth, behavioral development, and parental care of Broad-Winged Hawks (crecimiento, desarrollo de patrones de conducta y cuidado parental en *Buteo platypterus*),

Journal of Field Ornithology 58 (3) (1987) 334–344.

URL <https://www.jstor.org/stable/4513249>

[44] S. Dimri, T. Rizvi, J. do Monte Gonzalez de Segovia, A. Sánchez-Tójar, The adaptive value of green nest material across birds: A systematic review and meta-analysis, preregistration (2024). [doi:10.17605/OSF.IO/S7J6Z](https://doi.org/10.17605/OSF.IO/S7J6Z).

[45] R. E. O’Dea, M. Lagisz, M. D. Jennions, J. Koricheva, D. W. A. Noble, T. H. Parker, J. Gurevitch, M. J. Page, G. Stewart, D. Moher, S. Nakagawa, Preferred reporting items for systematic reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension, Biological Reviews 96 (5) (2021) 1695–1722. [doi:10.1111/brv.12721](https://doi.org/10.1111/brv.12721).

[46] E. M. Grames, A. N. Stillman, M. W. Tingley, C. S. Elphick, An automated approach to identifying search terms for systematic reviews using keyword co-occurrence networks, Methods in Ecology and Evolution 10 (10) (2019) 1645–1654. [doi:10.1111/2041-210x.13268](https://doi.org/10.1111/2041-210x.13268).

[47] M. J. Westgate, revtools: An R package to support article screening for evidence synthesis, Research Synthesis Methods 10 (4) (2019) 606–614. [doi:10.1002/jrsm.1374](https://doi.org/10.1002/jrsm.1374).

[48] J. L. Pick, S. Nakagawa, D. W. A. Noble, Reproducible, flexible and high-throughput data extraction from primary literature: The metadigitise R package, Methods in Ecology and Evolution 10 (3) (2018) 426–431. [doi:10.1111/2041-210x.13118](https://doi.org/10.1111/2041-210x.13118).

[49] W. Viechtbauer, Conducting meta-analyses in R with the metafor package, Journal of Statistical Software 36 (3) (2010) 1–48. [doi:10.18637/jss.v036.i03](https://doi.org/10.18637/jss.v036.i03).

[50] H. Cooper, L. V. Hedges, J. C. Valentine, The Handbook of Research Synthesis and Meta-Analysis, Russell Sage Foundation, New York, NY, 2019. [doi:10.7758/9781610448864](https://doi.org/10.7758/9781610448864).

[51] S. Nakagawa, D. W. A. Noble, M. Lagisz, R. Spake, W. Viechtbauer, A. M. Senior, A robust and readily implementable method for the meta-analysis of response ratios with and without missing standard deviations, Ecology Letters 26 (2) (2022) 232–244. [doi:10.1111/ele.14144](https://doi.org/10.1111/ele.14144).

[52] M. J. Lajeunesse, Recovering missing or partial data from studies: a survey of conversions and imputations for meta-analysis, in: Handbook of Meta-analysis in Ecology and Evolution, Princeton University Press, 2013, pp. 195–206.

[53] R Core Team, R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria (2025).

URL <https://www.R-project.org/>

- [54] D. W. A. Noble, M. Lagisz, R. E. O’Dea, S. Nakagawa, Nonindependence and sensitivity analyses in ecological and evolutionary meta-analyses, *Molecular Ecology* 26 (9) (2017) 2410–2425. doi:10.1111/mec.14031.
- [55] S. Krackow, Why parental sex ratio manipulation is rare in higher vertebrates (invited article), *Ethology* 108 (12) (2002) 1041–1056. doi:10.1046/j.1439-0310.2002.00843.x.
- [56] L. Montesana, M. Hau, P. B. D’Amelio, N. M. Adreani, A. Sánchez-Tójar, Do egg hormones have fitness consequences in wild birds? A Systematic Review and Meta-Analysis, *Ecology Letters* 28 (3) (2025) e70100. doi:10.1111/ele.70100.
- [57] Y. Yang, D. W. A. Noble, R. Spake, A. M. Senior, M. Lagisz, S. Nakagawa, A pluralistic framework for measuring, interpreting and decomposing heterogeneity in meta-analysis, *Methods in Ecology and Evolution* 16 (11) (2025) 2710–2725. doi:10.1111/2041-210x.70155.
- [58] S. Nakagawa, H. Schielzeth, A general and simple method for obtaining R^2 from generalized linear mixed-effects models, *Methods in Ecology and Evolution* 4 (2) (2012) 133–142. doi:10.1111/j.2041-210x.2012.00261.x.
- [59] J. Stanhope, P. Weinstein, Critical appraisal in ecology: What tools are available, and what is being used in systematic reviews?, *Research Synthesis Methods* 14 (3) (2022) 342–356. doi:10.1002/jrsm.1609.
- [60] A. Culina, D. Foster, M. Grainger, R. E. O’Dea, A. Sánchez-Tójar, O. Pescott, R. Boyd, J. Koricheva, S. Nakagawa, G. Stewart, Elevating the importance of risk of bias assessment for ecology and evolution, *EcoEvoRxiv*, Preprint (2025). doi:10.32942/X2764C.
- [61] M. D. Jennions, C. J. Lortie, M. S. Rosenberg, H. R. Rothstein, Publication and related biases, in: J. Koricheva, J. Gurevitch, K. Mengersen (Eds.), *Handbook of Meta-analysis in Ecology and Evolution*, Princeton University Press, Princeton, NJ, 2013, pp. 207–236.
- [62] Y. Yang, A. Sánchez-Tójar, R. E. O’Dea, D. W. A. Noble, J. Koricheva, M. D. Jennions, T. H. Parker, M. Lagisz, S. Nakagawa, Publication bias impacts on effect size, statistical power, and magnitude (type m) and sign (type s) errors in ecology and evolutionary biology, *BMC Biology* 21 (1) (2023). doi:10.1186/s12915-022-01485-y.
- [63] S. Nakagawa, M. Lagisz, M. D. Jennions, J. Koricheva, D. W. A. Noble, T. H. Parker, A. Sánchez-Tójar, Y. Yang, R. E. O’Dea, Methods for testing publication bias in ecological and evolutionary meta-analyses, *Methods in Ecology and Evolution* 13 (1) (2022) 4–21. doi:10.1111/2041-210X.13724.
- [64] A. Sánchez-Tójar, S. Nakagawa, M. Sánchez-Fortún, D. A. Martin, S. Ramani, A. Girndt, V. Bókony, B. Kempnaers, A. Liker, D. F. Westneat, T. Burke, J. Schroeder, Meta-analysis challenges a textbook

example of status signalling and demonstrates publication bias, *eLife* 7 (2018) e37385. doi:10.7554/elife.37385.

[65] P. T. Fauth, D. G. Krementz, J. E. Hines, Ectoparasitism and the role of green nesting material in the European Starling, *Oecologia* 88 (1) (1991) 22–29. doi:10.1007/BF00328399.

[66] L. Clark, J. R. Mason, Effect of biologically active plants used as nests material and the derived benefit to starling nestlings, *Oecologia* 77 (2) (1988) 174–180. doi:10.1007/bf00379183.

[67] D. Shutler, A. A. Campbell, Experimental addition of greenery reduces flea loads in nests of a non-greenery using species, the tree swallow *Tachycineta bicolor*, *Journal of Avian Biology* 38 (1) (2007) 7–12. doi:10.1111/j.2007.0908-8857.04015.x.

[68] A. Mennerat, P. Perret, S. P. Caro, P. Heeb, M. M. Lambrechts, Aromatic plants in blue tit *Cyanistes caeruleus* nests: no negative effect on blood-sucking protocoelliphora blow fly larvae, *Journal of Avian Biology* 39 (2) (2008) 127–132. doi:10.1111/j.0908-8857.2008.04400.x.

[69] P. A. Taverner, Purple martins gathering leaves, *The Auk* 50 (1) (1933) 110–111. doi:10.2307/4076584.

[70] J. M. Peralta-Sanchez, A. P. Møller, A. M. Martin-Platero, J. J. Soler, Number and colour composition of nest lining feathers predict eggshell bacterial community in barn swallow nests: an experimental study, *Functional Ecology* 24 (2) (2010) 426–433. doi:10.1111/j.1365-2435.2009.01669.x.

[71] J.-Å. Nilsson, A. Nord, The use of the nest for parental roosting and thermal consequences of the nest for nestlings and parents, *Behavioral Ecology and Sociobiology* 71 (12) (2017). doi:10.1007/s00265-017-2400-7.

[72] D. M. Perez, J. L. Gardner, I. Medina, Climate as an evolutionary driver of nest morphology in birds: A Review, *Frontiers in Ecology and Evolution* 8 (2020). doi:10.3389/fevo.2020.566018.

[73] M. K. Labocha, J. P. Hayes, Morphometric indices of body condition in birds: a review, *Journal of Ornithology* 153 (1) (2011) 1–22. doi:10.1007/s10336-011-0706-1.

[74] B. Pires, A. D. Belo, F. Diamantino, J. Rabaça, S. Merino, Development of nestling blue tits (*Cyanistes caeruleus*) is affected by experimental addition of aromatic plants, *Avian Biology Research* 13 (3) (2020) 44–48. doi:10.1177/1758155920921075.

[75] A. Corregidor-Castro, O. R. Jones, The effect of nest temperature on growth and survival in juvenile great tits *Parus major*, *Ecology and Evolution* 11 (12) (2021) 7346–7353. doi:10.1002/ece3.7565.

[76] Y. Yang, C. Williams, A. M. Senior, K. Morrison, L. Ricolfi, J. Pan, M. Lagisz, S. Nakagawa, Bivariate multilevel meta-analysis of log response ratio and standardized mean difference for robust and reproducible environmental and biological sciences, *bioRxiv*, Preprint (2024). doi:10.1101/2024.05.13.594019.

663 [77] J. IntHout, J. P. A. Ioannidis, G. F. Borm, J. J. Goeman, Small studies are more heterogeneous
664 than large ones: a meta-meta-analysis, Journal of Clinical Epidemiology 68 (8) (2015) 860–869. doi:
665 10.1016/j.jclinepi.2015.03.017.

666 7. Supplementary Appendix

667 7.1. PRISMA Checklist

Table 1: {#tblsupp-checklist}

Checklist item	Item score	Sub-item number	Sub-item	Reported by authors?	Notes
Title and abstract	100%	1.1	Identify the review as a systematic review, meta-analysis, or both	Yes	Identified in title and in abstract as both a systematic review and a meta-analysis
		1.2	Summarise the aims and scope of the review	Yes	In the abstract
		1.3	Describe the data set	Yes	In the abstract
		1.4	State the results of the primary outcome	Yes	In the abstract
		1.5	State conclusions	Yes	In the abstract
		1.6	State limitations	Yes	In the abstract
Aims and questions	100%	2.1	Provide a rationale for the review	Yes	Rationale explained in the Introduction in detail
		2.2	Reference any previous reviews or meta-analyses on the topic	Yes	Clearly stated twice in the introduction

2.3	State the aims and scope of the review (including its generality)	Yes	Scope (only birds and experimental studies) explained along with the aims in the last paragraph of Introduction
2.4	State the primary questions the review addresses (e.g., which moderators were tested)	Yes	Moderators tested are clearly stated in the methods in the data analysis section.
2.5	Describe whether effect sizes were derived from experimental and/or observational comparisons	Yes	Only experimental studies were selected, described in the last paragraph of introduction as well as the methods

Review registration	100%	3.1	Register review aims, hypotheses (if applicable), and methods in a time-stamped and publicly accessible archive and provide a link to the registration in the methods section of the manuscript. Ideally registration occurs before the search, but it can be done at any stage before data analysis.	Yes	https://doi.org/10.17605/OSF.IO/S7J6Z
		3.2	Describe deviations from the registered aims and methods	Yes	Described in methods wherever applicable
		3.3	Justify deviations from the registered aims and methods	Yes	When applicable, described in the methods

Eligibility criteria	100%	4.1	Report the specific criteria used for including or excluding studies when screening titles and/or abstracts, and full texts, according to the aims of the systematic review (e.g., study design, taxa, data availability)	Yes	Described briefly in the methods, but mainly in the decision tree (see Supplementary Materials)
		4.2	Justify criteria, if necessary (i.e. not obvious from aims and scope.	Yes	Described along with the decision tree (see Supplementary Materials)
Finding studies	100%	5.1	Define the type of search (e.g.,comprehensive search, representative sample)	Yes	Comprehensive search conducted, described in methods along with search string (complete search string provided in the supplementary materials)
		5.2	State what sources of information were sought (e.g.,published and unpublished studies, personal communications)	Yes	Published, unpublished as well as personal communications. Described in detail in the Eligibility Criteria and Data Extraction section of Methods Section

Study selection	100%	5.3	Include, for each database searched, the exact search strings used, with keyword combinations and Boolean operators	Yes	See complete search string in the Supplementary Materials
		5.4	Provide enough information to repeat the equivalent search (if possible), including the timespan covered (start and end dates)	Yes	See complete search string details in the Supplementary Materials. Search dates are described in the methods.
		6.1	Describe how studies were selected for inclusion at each stage of the screening process (e.g., use of decision trees, screening software)	Yes	Abstract-Title screening was conducted using the R package revtools GUI and Fulltext screening was conducted in excel manually downloading the articles. Described in detail in the pre-registration and methods section. The decision tree is provided in the supplementary materials.
		6.2	Report the number of people involved and how they contributed (e.g., independent parallel screening)	Yes	Described in detail in the methods section. All screening decision files are shared as .xls/.csv files with notes on the decisions made.

Data collection process	100%	7.1	Describe where in the reports data were collected from (e.g.,text or figures)	Yes	In all files after the data extraction (shared along with the project), the column data_location contains this information
		7.2	Describe how data were collected (e.g.,software used to digitize figures, external data sources)	Yes	Described in detail in the Data extraction section.
		7.3	Describe moderator variables that were constructed from collected data (e.g., number of generations calculated from years and average generation time)	Yes	All moderators are described in detail in the metadata file of the overall dataset. Please see data_cleaning.Rmd and data_preparation.Rmd files for any cleaning done to the files. All codes are provided along with the dataset.

		7.4	Report how missing or ambiguous information was dealt with during data collection (e.g., authors of original studies were contacted for missing descriptive statistics, and/or effect sizes were calculated from test statistics)	Yes	Described in detail in the data extraction and effect size calculation section in the methods.
		7.5	Report who collected data	Yes	The extractor_ID column in the data file has information on the ID of who collected the data. The methods section also describes the ID of the data collector.
		7.6	State the number of extractions that were checked for accuracy by co-authors		Described in the methods section along with % of disagreement and process of resolution. The data_checker_ID column in the data files has information on the ID of who checked the data.
Data items	100%	8.1	Describe the key data sought from each study	Yes	The priority order of data extraction is described in detail in the Data extraction section of the methods.

		8.2	Describe items that do not appear in the main results, or which could not be extracted due to insufficient information	Yes	All information that is excluded from analysis is described in detail in the methods. Some proxies that could not be included in the analysis for various reasons are reported in the supplementary material along with the reason.
		8.3	Describe main assumptions or simplifications that were made (e.g., categorising both ‘length’ and ‘mass’ as ‘morphology’)	Yes	All major decisions are described in the methods section, for any remaining description please check the notes column in the data files.
		8.4	Describe the type of replication unit (e.g., individuals, broods, study sites)	Yes	Described in the methods, we prioritised nests as the unit of replication whenever possible.
		9.1	Describe whether the quality of studies included in the systematic review or meta-analysis was assessed (e.g., blinded data collection, reporting quality, experimental versus observational)	Yes	The risk of bias assessment performed for the individual studies is described in the methods and results reported in the results section (See supplementary material for all model output and figures). The risk of bias categories we evaluated were blinding, random assignment to each treatment and partial reporting
Assessment of individual study quality	100%				

		9.2	Describe how information about study quality was incorporated into analyses (e.g., meta-regression and/or sensitivity analysis)	Yes	We conducted three meta-regressions using blinding, random assignment to each treatment and partial reporting as moderators in each analysis. We reported the outcome briefly in the Results (See supplementary material for all model output and figures)
Effect size measures	100%	10.1	Describe effect size(s) used	Yes	Described in Effect Size Calculation section in Methods
		10.2	Provide a reference to the equation of each calculated effect size (e.g., standardised mean difference, log response ratio) and (if applicable) its sampling variance	Yes	Described in the Effect Size Calculation section in Methods. Check <code>data_preparation.Rmd</code> in Code to see the exact function, its parameters and any equations that were used to estimate the effect sizes.
		10.3	If no reference exists, derive the equations for each effect size and state the assumed sampling distribution(s)	Yes	Does not apply. Any custom equations used from books etc. are clearly mentioned as well as referenced in the Methods section.

Missing data	100%	11.1	Describe any steps taken to deal with missing data during analysis (e.g., imputation, complete case, subset analysis)	Yes	All imputations conducted for the missing data are stated clearly in the Effect Size Calculations in Methods Section
		11.2	Justify the decisions made to deal with missing data	Yes	We decided to impute the missing data but also conducted the sensitivity analysis excluding the imputed value. All justifications are clearly stated in the methods.
Meta-analytic model description	100%	12.1	Describe the models used for synthesis of effect sizes	Yes	Described all models in Data Analysis section of the Methods

		12.2	The most common approach in ecology and evolution will be a random-effects model, often with a hierarchical/multilevel structure. If other types of models are chosen (e.g., common/fixed effects model, unweighted model), provide justification for this choice	Yes	Not applicable, Multilevel meta analytical model and meta-regressions were conducted. No justification needed
Software	100%	13.1	Describe the statistical platform used for inference (e.g., R)	Yes	R v4.5.1 was used for the analysis and is clearly stated in the Methods.
		13.2	Describe the packages used to run models	Yes	All packages and their versions are described in the Methods.
		13.3	Describe the functions used to run models	Yes	All important functions used for running the models are stated in the Methods.
		13.4	Describe any arguments that differed from the default settings	Yes	We have clearly stated all the important arguments that differed from the default settings. However, see data_analysis.Rmd for full description of the arguments for each model.

Non-independence	100%	13.5	Describe the version numbers of all software used	Yes	All packages and their versions are described in the Methods.
		14.1	Describe the types of non-independence encountered (e.g., phylogenetic, spatial, multiple measurements over time)	Yes	See data analysis for more details on how non-independence was handled.
		14.2	Describe how non-independence has been handled	Yes	See data analysis for more details on how non-independence was handled.
		14.3	Justify decisions made	Yes	See data analysis for more details on how non-independence was handled.
Meta-regression and model selection	100%	15.1	Provide a rationale for the inclusion of moderators (covariates) that were evaluated in meta-regression models	Yes	All the moderators that were evaluated were pre-registered with the rationale behind it clearly stated. See Data analysis for more details (but also see pre-registration).

Publication bias and sensitivity analysis	100%	15.2	Justify the number of parameters estimated in models, in relation to the number of effect sizes and studies (e.g., interaction terms were not included due to insufficient sample sizes)	Yes	We described the random effects that were not included in the models in the Methods section
		15.3	Describe any process of model selection	Yes	No model selection was conducted
		16.1	Describe assessments of the risk of bias due to missing results (e.g., publication, time-lag, and taxonomic biases)	Yes	Time-lag bias and small study bias are reported in the methods, results and discussion.
		16.2	Describe any steps taken to investigate the effects of such biases (if present)	Yes	We discussed the effects of potential biases and conducted a sensitivity analysis accounting these effects for the two main models.

		16.3	Describe any other analyses of robustness of the results, e.g., due to effect size choice, weighting or analytical model assumptions, inclusion or exclusion of subsets of the data, or the inclusion of alternative moderator variables in meta-regressions	Yes	All sensitivity analysis are mentioned in the methods, results and discussed where appropriate. All the results from the sensitivity analysis/robustness checks are reported as a table in the supplementary material.
Clarification of post hoc analyses	100%	17.1	When hypotheses were formulated after data analysis, this should be acknowledged.	Yes	All analysis are pre-registered and exploratory analysis are clearly stated under exploratory analysis both in the pre-registration and final article
Metadata, data, and code	100%	18.1	Share metadata (i.e. data descriptions)	Yes	See data and code availability statement for details.
		18.2	Share data required to reproduce the results presented in the manuscript	Yes	See data and code availability statement for details.

Results of study selection process	100%	18.3	Share additional data, including information that was not presented in the manuscript (e.g., raw data used to calculate effect sizes, descriptions of where data were located in papers)	Yes	See data and code availability statement for details.
		18.4	Share analysis scripts (or, if a software package with graphical user interface (GUI) was used, then describe full model specification and fully specify choices)	Yes	See data and code availability statement for details.
		19.1	Report the number of studies screened	Yes	Details in method and the PRISMA Flowchart in supplementary material
		19.2	Report the number of studies excluded at each stage of screening	Yes	Details in method and the PRISMA Flowchart in supplementary material
		19.3	Report brief reasons for exclusion from the full text stage	Yes	Details in the PRISMA Flowchart in supplementary material

Sample sizes and study characteristics	100%	19.4	Present a Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)-like flowchart (www.prisma-statement.org).	Yes	Figure S1 in supplementary materials is a PRISMA Flowchart
		20.1	Report the number of studies and effect sizes for data included in meta-analyses	Yes	Details in method and supplementary tables
		20.2	Report the number of studies and effect sizes for subsets of data included in meta-regressions	Yes	Details in method and supplementary tables
		20.3	Provide a summary of key characteristics for reported outcomes (either in text or figures; e.g., one quarter of effect sizes reported for vertebrates and the rest invertebrates)	Yes	All figures contain effect sizes (k) and unique articles (n) for different moderator levels. Summary of reported outcomes is also in the first paragraph of the Results.

		20.4	Provide a summary of limitations of included moderators (e.g., collinearity and overlap between moderators)	NA	All moderators tested were categorical and only publication year was a continuous moderator. Therefore, no collinearity or overlap is possible.
		20.5	Provide a summary of characteristics related to individual study quality (risk of bias)	Yes	This data is presented in dataset in columns blinding, missing_data, random_assignment
Meta-analysis	100%	21.1	Provide a quantitative synthesis of results across studies, including estimates for the mean effect size, with confidence/credible intervals	Yes	See results and supplementary material for detail of all the synthesis.
Heterogeneity	100%	22.1	Report indicators of heterogeneity in the estimated effect (e.g., I ² , tau ² and other variance components)	Yes	All estimates of heterogeneity are reported in the results and model table. Also see supplementary material for the results not reported in the main text.

Meta-regression	100%	23.1	Provide estimates of meta-regression slopes (i.e. regression coefficients) and confidence/credible intervals	Yes	See results and supplementary material for detail of all the synthesis.
		23.2	Include estimates and confidence/credible intervals for all moderator variables that were assessed (i.e. complete reporting)	Yes	See results and supplementary material for detail of all the synthesis.
		23.3	Report interactions, if they were included	NA	There were no interactions included
		23.4	Describe outcomes from model selection, if done (e.g., R ² and AIC)	NA	No model selection was conducted.

Outcomes of publication bias and sensitivity analysis	100%	24.1	Provide results for the assessments of the risks of bias (e.g., Egger's regression, funnel plots)	Yes	Risk of bias was assessed using meta-regression models with moderator levels for blinding, missing data, random assignment in experiments and reported in results (and see supplementary material for complete details). We also reported meta-regressions for publication bias (time-lag bias and small-study effects) in results (and see supplementary material for complete details)
		24.2	Provide results for the robustness of the review's results (e.g., subgroup analyses, meta-regression of study quality, results from alternative methods of analysis, and temporal trends)	Yes	All robustness checks/sensitivity analysis conducted are reported in full detail in the supplementary material and mentioned in the results clearly.
Discussion	100%	25.1	Summarise the main findings in terms of the magnitude of effect	Yes	

25.2	Summarise the main findings in terms of the precision of effects (e.g., size of confidence intervals, statistical significance)	Yes	We use confidence intervals, prediction intervals and statistical significance to discuss the precision of the effects in the results and discussion section.
25.3	Summarise the main findings in terms of their heterogeneity	Yes	We report and discuss the heterogeneity and explore some of the possible causes of it in our results and discussion.
25.4	Summarise the main findings in terms of their biological/practical relevance	Yes	We have discussed all important findings in terms of their biological and practical implications
25.5	Compare results with previous reviews on the topic, if available	Yes	We have discussed the results comparing them to other published literature (including two narrative reviews that summarise the field) but no quantitative review on the field exists to compare the results with.

Contributions and funding	100%	25.6	Consider limitations and their influence on the generality of conclusions, such as gaps in the available evidence (e.g., taxonomic and geographical research biases)	Yes	We discuss the limitations of the evidence.
		26.1	Provide names, affiliations, and funding sources of all co-authors	Yes	See section Fundings
		26.2	List the contributions of each co-author	Yes	See section Author Contributions
		26.3	Provide contact details for the corresponding author	Yes	See manuscript Author details: Shreya Dimri - shreya.dimri@uni-bielefeld.de
References	100%	26.4	Disclose any conflicts of interest	Yes	See section Conflict of interest
		27.1	Provide a reference list of all studies included in the systematic review or meta-analysis	Yes	See supplementary material for a list.

27.2	List included studies as referenced sources (e.g., rather than listing them in a table or supplement)	Yes	We have included all the studies in our meta-analysis within the text of the article in appropriate places.
------	---	-----	---

- 668 7.2. *PRISMA Flowchart*
- 669 7.3. *Initial library used for search validation*
- 670 7.4. *Decision Tree*
- 671 7.5. *Fitness proxies included in or excluded from the meta-analysis*

Table 2: {#tblsupp-proxies}

Courtship			
Hypothesis			
Fitness	Effect	Definition (as	Expected Direction (as described
Proxy	Size(studies)	measures/provided	or expected by authors of primary
		by authors of primary	study)
		study)	

Sex Ratio*	9(2)	Sex Ratio is the hatching broods sex ratio measures as the proportion of male offspring in a breeding attempt.	In these experiments, adding fresh GNM is treated as a courtship signal. Under sex-allocation as described by the authors, females paired to more attractive males (that add more GNM), are expected to bias investment towards the sex with higher potential fitness returns, here assumed to be sons. So the predicted effect is a more male-biased sex ratio in nests with GNM (Polo et al., 2004; Veiga et al., 2008). However, we only include them in a sensitivity analysis since the evidence for whether birds can change sex ratio is debated (see Krackow, S. (2002), Why Parental Sex Ratio Manipulation is Rare in Higher Vertebrates (Invited Article). <i>Ethology</i>, 108: 1041-1056. https://doi.org/10.1046/j.1439-0310.2002.00843.x).
------------	------	--	---

Females Testosterone*	4(2)	Circulating (plasma) testosterone concentration of breeding females (during laying or early incubation), quantified from a blood sample collected during the breeding attempt and assayed using ELISA.	In these experiments author predicted higher female testosterone in GNM-added nests. Experimental GNM addition is treated as increasing the perceived attractiveness/quality of the male (i.e., a courtship signal). The authors predict (and test) that this should elevate female testosterone because (i) females may increase competitive/aggressive capacity to maintain access to preferred males or nest sites under heightened female–female competition, and/or (ii) female endocrine state may shift as part of differential allocation when paired to (perceived) attractive males. This is described as a mechanistic mediator that would show competitive state of the females, which could indirectly affect reproductive outcomes (e.g., maintaining access to higher quality males thus benefiting the offspring or altering maternal investment strategies). (López-Rull & Gil, 2009; Polo et al., 2010). However, we only include them in a sensitivity analysis because in these studies, only one measure of testosterone was taken per female and the values of testosterone can considerably change within the time of the day or phase of breeding, for consistency with other physiological measures like blood glucose, which we only test in a sensitivity analysis, we also excluded it. Furthermore, an increase in female testosterone does not have clear fitness benefits associated with it.
--------------------------	------	---	---

Courtship Time	3(1)	Interval between first appearance of greenery to first egg.	Under the courtship hypothesis (greenery as a sexually selected ornament/mate-attraction cue), the expected relationship is that GNM should facilitate mate attraction and progression to laying, so addition would be expected to shorten courtship time (i.e., faster progression to laying). Early laying of eggs can then have several benefits for the offspring. (Brouwer & Komdeur, 2004)
Male Provisioning Rate	2(1)	Male provisioning rate is the male's feeding visit rate to nestlings, measured as the number of visits per hour to the nest box while provisioning chicks.	A higher male provisioning rates in nests supplied with aromatic plants (GNM) is expected. In this experimental context, the amount of aromatic plant material in the nest is treated as a female sexual signal that could elicit differential allocation by males. If males interpret aromatic plant addition as indicating higher female quality or higher expected reproductive value of the brood, it would predict increased male parental investment, measured here as increased provisioning rate. (Tomás et al., 2013)

Male Visit	2(1)	A participation or attendance measure indicating whether the male was present and entered the nest-box to provision nestlings.	A higher male visit in nests supplied with aromatic plants (GNM) is expected. In this experimental context, the amount of aromatic plant material in the nest is treated as a female sexual signal that could elicit differential allocation by males. If males interpret aromatic plant addition as indicating higher female quality or higher expected reproductive value of the brood, it would predict increased male parental investment, measured here as increased male visit to the nest. (Tomás et al., 2013)
------------	------	---	---

Yolk Hormones*	2(1)	Yolk androgen concentrations measured from eggs (e.g., testosterone (T), androstenedione (A4))	A higher yolk androgen deposition is expected in experimental nests with added green plant material (GNM). Under the courtship hypothesis, females paired to more attractive males (who bring more GNM) are predicted to increase pre-laying maternal investment, including transferring higher androgen levels to eggs. (López-Rull & Gil, 2009). However, we only include them in a sensitivity analysis because, according to a meta-analysis on yolk hormone and fitness, there is no strong evidence for such a relationship. (Mentesana, L., Hau, M., D'Amelio, P.B., Adreani, N.M. and Sánchez-Tójar, A. (2025), Do Egg Hormones Have Fitness Consequences in Wild Birds? A Systematic Review and Meta-Analysis. <i>Ecology Letters</i>, 28: e70100. https://doi.org/10.1111/ele.70100)
Male Risk Taking	1(1)	Measured as a composite index of male parental risk-taking/investment, constructed as the mean of many binomial behavioural components spanning the nestling period. Higher scores indicate greater willingness to incur risk to attend the nest under disturbance or predation cues.	The authors interpret entering the nest-box under potential danger as risky and treat male risk-taking as an index of male parental investment. Under the courtship hypothesis, aromatic plant supplementation (a female trait/signal in this case) is expected to elicit greater male reproductive investment, expressed here as increased risk-taking. (Tomás et al., 2013)

Parental Care Fitness Proxy	Hypothesis Effect Size(studies)	Definition (as measures/provided by authors of primary study)	Expected Direction (as described or expected by authors of primary study)
Ectoparasite Abu- dance/Load	Mites: 22(7) Blowflies number: 8(6) Fleas number: 6(6) Carnusflies load: 5(2) Blackflies number: 2(1) Midges number: 2(1) Beetles: 1(1) Blowfly Parasitoids Number: 1(1) Booklice And Barklice: 1(1) Butterflies And Moths: 1(1) Clown Beetles: 1(1) Flies: 1(1) Lice Score: 1(1) Rove Beetles: 1(1) Skin Beetles: 1(1) Springtails: 1(1) Tick Load: 1(1) Ticks: 1(1) True Bugs: 1(1) Wasps, Bees, And Ants: 1(1)	These are the different measures of the amount of ectoparasites found in the nests or on the nestlings. Many different ectoparasites have been measured in different studies.	It is expected that the addition of GNM would reduce the amount/abundance/load of ectoparasites in the nests, thereby increasing the fitness of the nestlings. These are considered the most direct measures of the nest protection hypothesis.

Microorganism Abundance/Load/Community metrics	Bacterial Load: 20(4) Bacterial Increase: 2(1) Bacterial Richness: 2(1) Leucocytozoon Prevalence: 3(1)	Different measures of microbial load, including increase or richness, are present in the nests or on the nestlings.	It is expected that the addition of GNM would reduce the diversity/abundance/load of microbes in the nests, thereby increasing the fitness of the nestlings. These are considered the most direct measures of the nest protection hypothesis.
Hemoglobin Measure	9(6)	Hemoglobin concentration in blood is a measure of oxygen-carrying capacity and the general physiological condition of nestlings.	It is expected that the addition of GNM would increase the Hemoglobin Measure. Increased hemoglobin (improved physiological condition) under the parental care hypothesis, because of reduced parasitism/pathogen pressure and/or pharmacological benefits, would support a better physiological state.
Hematocrit %	1(1)	The percentage of blood volume composed of red blood cells is another indicator of oxygen transport and the physiological condition of the nestlings.	Increased hematocrit (improved physiological condition) under the parental care hypothesis, because of reduced parasitism/pathogen pressure and/or pharmacological benefits, would support a better physiological state.
Scab Score	5(3)	Score quantifying skin lesions/scabs on nestlings, typically interpreted as damage from ectoparasites.	With increased GNM we expect reduced scab score because of reduced ectoparasite load or reduced inflammation from secondary infection.

Telomere Measure	4(1)	Measure of telomere length (but in one case, telomere dynamics).	We expect a positive relation between telomere length and GNM. Telomere length shortens as the individual ages and the cell dies once there is not enough protection provided by the telomere cap. Therefore, the shorter the telomere, the worse they are in their fitness. We expect a negative relationship between telomere dynamics and GNM. Telomere dynamics are calculated as the rank difference in telomere length.
Nestling Immunity	2(2)	A direct measure of immune function (varies by study; e.g., PHA swelling response) intended to reflect immune performance/competence.	Increased immune response when GNM is added through the direct pharmacological effect of GNM, thus enabling a better immune response.

Glucose Measure*	2(2)	Blood glucose concentration, a metabolic indicator reflecting energetic state and/or physiological stress.	Authors of the primary studies expect the addition of GNM to decrease the value of Glucose concentration in the blood, i.e., it is closer to the baseline level, showing less stress. However, we only include them in a sensitivity analysis, as Glucose Measure is highly context-dependent. We do not consider glucose concentration to be a meaningful fitness proxy because (1) the blood samples of the birds were collected at random time of the day without any strict protocol for between samplin differences. Blood glucose levels could change very rapidly depending on the consumption of food and there could be great between individual variation in the population. Circadian rhythm could also influence the blood sugar levels. (2) Authors themselves call this a less robust proxy - “The level of blood glucose is considered a reverse (lower concentration of glucose is related to higher body condition) and a less robust (more variable) indicator of body condition than haemoglobin but is also useful in studies of ecophysiology”
For data from	Tachycineta bicolor (Because addition	of GNM is not shown by this species,	these proxies were treated as parental-care proxies only)

Nestling Mass	1(1)	Body mass of nestlings measured at a different ages in <i>Tachycineta bicolor</i> .	Increased nestling mass would be expected with addition of GNM in <i>Tachycineta bicolor</i> . Since this bird does not show this behaviour in wild, the effect of GNM cannot be attributed to courtship hypothesis.
Nestling Size	1(1)	A composite size index measured in nestlings using other structural size metrics like tarsus length, wing length.	Increased nestling size would be expected with addition of GNM in <i>Tachycineta bicolor</i> . Since this bird does not show this behaviour in wild, the effect of GNM cannot be attributed to courtship hypothesis.
Number of Fledglings	1(1)	Count of young that fledge per nesting attempt.	If GNM reduces parasite/pathogen pressure and improves nestling condition, we should see an increased survival to fledgeling with GNM addition.
Number of Nestlings	1(1)	Count of nestlings present/survive to a certain age.	If GNM reduces early mortality, we expect more chicks to remain alive during the nestling period.
Reproductive Success	3(1)	Percentage of nests with complete reproductive success in <i>Tachycineta bicolor</i> .	With the addition of GNM, we should see increased reproductive success through reduced mortality.
Clutch Size	1(1)	Number of eggs laid in a clutch in <i>Tachycineta bicolor</i> .	With the addition of GNM, we should see an increased clutch size.

Both

Hypothesis

**Fitness
Proxy**

**Effect
Size(studies)**

**Definition (as
measures/provided
by authors of primary
study)**

**Expected Direction (as described
or expected by authors of primary
study)**

Growth, size or condition of offspring	Nestling Mass:	These proxies capture	We expect GNM to have a positive effect on growth, size and body condition of the offspring. This could be because of reduced pathogens or drug effect, but also because of investment of parents through various mechanisms. Therefore, it is not possible to categorise this into a clear mechanism.
	20(9)	various measurements of	
	Chick Mass: 16(4)	offspring growth,	
	Nestling Tarsus	structural size and	
	Length: 13(5)	development condition	
	Fledgeling Mass:	by measuring body mass,	
	7(2)	skeletal measures (e.g.,	
	Nestling Wing	tarsus/wing length),	
	Length: 2(2)	integrated body	
	Fledgeling	condition indices, fat	
	Tarsus–Metatarsus	reserves, and feather	
	Lengths (Females):	development, all of which	
	1(1)	reflect the developmental	
	Fledgeling	quality of the nestling at	
	Tarsus–Metatarsus	various measurement	
	Lengths (Males):	ages.	
	1(1)		
	Body Condition		
	Estimate: 3(1)		
	Fat Score: 1 (1)		
	Mouth Colouration:		
	3 (1)		
	Chick Feather		
	Development: 2 (1)		

Reproductive Output/Suc- cess	Clutch Size: 16(8) Number Of Nestling: 7(5) Fledgling Success: 4 (4) Hatching Success: 4 (3) Local Recruitment Probability: 4 (1) Survival To Fledgeling: 3(1) Nest Mortality: 1 (1) Percent Survived: 1(1) Survival Per Brood: 1(1) Survival Rate Nestling: 1(1)	These proxies quantify the reproductive output and survival (hatching, fledgling, nest failure and recruitment) across different stages of breeding.	GNM is predicted to increase productivity and survival and decrease mortality. These proxies of fitness could improve via increased parental investment, better genetic makeup or through lower parasites in the nests. Therefore, it is difficult to categorise these proxies into a particular mechanism.
Laying Date	3(3)	The date of the clutch initiation (first egg laid), expressed as a relative count from the reference date.	Since earlier laying is often associated with higher reproductive output, we expected GNM would lead to advancement in laying date (i.e., smaller values of laying date proxy).
Hatching Date	1(1)	The date of the first egg hatched, expressed as a relative count from the reference date.	Earlier hatching can have several benefits due to the availability of better quality food; we expected GNM would lead to advancement in hatching date (i.e., smaller values of hatching date proxy).

Clutch Development Score	2(1)	It is a measure of how far embryonic development has progressed (egg, wet hatchling, dry hatchling), at a standardized time late in incubation.	Earlier development on a fixed day is interpreted as a benefit due to reduced exposure time to risks during incubation and improved postnatal condition. Therefore, it is expected that GNM would increase the clutch development score.
Egg Size	1(1)	As a measure of egg volume, derived from the length and width of the egg calculated as $0.45 \times L \times W^2$.	It is expected that GNM would lead to females laying larger eggs.
Feeding Rate*	1(1)	Measure of parental provisioning, quantified from nest-box video as the total number of feeding events standardised per hour during the recording period.	We only test this in a sensitivity analysis, because the relation of this to fitness can be argued. A higher feeding rate can indicate better care and resource collection, which might suggest higher fitness of the parents or brood. However, if feeding is high, it could also imply poor quality of food availability or compensation for the poor health of the nestling.

Time Spent Begging By Broods*	1(1)	Total begging duration, summed across nestlings, divided by brood size, and expressed as seconds of begging per hour of recording (s/h).	If GNM reduces parasite/pathogen burden and/or improves nestling condition, nestlings are expected to show reduced need, expressed as less begging time. However, we only test this in a sensitivity analysis because it is difficult and contentious to interpret. Begging is generally a signal that chicks are hungry and in need of care. More begging might indicate a greater demand for resources and therefore potentially less fitness in terms of immediate brood condition, but studies also suggest that begging intensity is an evolved behaviour to maximise parental care and could imply it is not related to fitness but rather a strategy to elicit care.
-------------------------------------	------	---	---

Fitness proxies	excluded from the	meta-analysis	
Fitness	Effect		Reason for exclusion
Proxy	Size(studies)		
Male Mass	3(1)		Variables such as adult male or female mass as a fitness proxy are tricky. On one hand, adults in better condition could provide better parental care for their offspring and be more likely to survive to reproduce the following season, thereby increasing their overall fitness. However, whether adult mass is directly related to fitness remains unclear.

Female Body Condition 2(1)

The authors used female body condition simply to control for its effect. We are excluding it, as the relationship between adult body condition and GNM is rather tricky to interpret and difficult to describe in terms of a true direction of effect.

Female Provisioning Rate 2(1)

We will exclude this variable from all our analyses. Since females add the GNM as noted by the authors, it is the male that should adjust his investment accordingly. One could argue that female investment will be an indirect reflection of the male’s investment; however, this relationship is very indirect, and the direction of this proxy in relation to the GNM addition can be argued both ways. That is, if females use GNM as an honest signal to increase male investment, their own investment would not be directly related to fitness. Alternatively, they might increase investment to compensate for a lack of male investment, or it could be negatively related to fitness because they invested both in bringing in the GNM and in higher provisioning. Therefore, we excluded it.

Measures of WBC (White Blood Cells)	Number of Basophils: 2(1) Number of Lymphocytes: 2(1) Eosinophils of Nestlings: 1(1) Heterophils of Nestlings: 1(1) Nestling Leukocytes: 1(1) WBC measures of Nestlings: 1(1)	It is very difficult to assign a direction of the relationship. Does an increase in WBC imply better or worse fitness? We consulted several researchers in the field and there was no clear consensus on this. Therefore, we decided to exclude it. For further details, see the discussion provided in the project materials.
---	--	---

672 7.6. Studies included in the meta-analysis

Table 3: {#tblsupp-proxies}

Paper ID	Title	Authors	Journal	Year	DOI
GNM_001	Consequences of experimental addition of fresh, aromatic plants into nests of blue tits (<i>Cyanistes caeruleus</i>) on the physiological condition of nestlings	Gładalski M; Kaliński A; Wawrzyniak J; Bańbura M; Markowski M; Skwarska J; Bańbura J	Behavioral Ecology and Sociobiology	2020	10.1007/s00265- 020-2812-7
GNM_002	Development of nestling blue tits (<i>Cyanistes caeruleus</i>) is affected by experimental addition of aromatic plants	Pires BA; Belo ADF; Diamantino F; Rabaça JE; Merino S	Avian Biology Research	2020	10.1177/1758155920921075

Paper ID	Title	Authors	Journal	Year	DOI
GNM_010	‘Green incubation’: avian offspring benefit from aromatic nest herbs through improved parental incubation behaviour	Gwinner H; Capilla-Lasheras P; Cooper C; Helm B	Proceedings of the Royal Society B: Biological Sciences	2018	10.1098/rspb.2018.0376
GNM_016	Green plants in nests reduce offspring recruitment rates in the spotless starling	Polo V; Rubalcaba JG; Veiga JP	Behavioral Ecology	2015	10.1093/be- heco/arv056
GNM_018	Nest size and aromatic plants in the nest as sexually selected female traits in blue tits	Tomás G; Merino S; Martínez-de la Puente J; Moreno J; Morales J; Rivero-de Aguilar J	Behavioral Ecology	2013	10.1093/be- heco/art015
GNM_023	Experimental Addition of Green Plants to the Nest Increases Testosterone Levels in Female Spotless Starlings	Polo V; López-Rull I; Gil D; Veiga JP	Ethology	2010	10.1111/j.1439- 0310.2009.01724.x
GNM_024	Do female spotless starlings (<i>Sturnus</i> unicolor) adjust maternal investment according to male attractiveness?	López-Rull I; Gil D	Journal of Avian Biology	2009	10.1111/j.1600- 048X.2009.04553.x
GNM_028	Age dependent sex allocation in the polygynous spotless starling	Veiga JP; Polo V; Cordero PJ	Evolutionary Ecology	2008	10.1007/s10682- 007-9166-8

Paper ID	Title	Authors	Journal	Year	DOI
GNM_032	Female starlings adjust primary sex ratio in response to aromatic plants in the nest	Polo V; Veiga JP; Cordero PJ; Viñuela J; Monaghan P	Proceedings of the Royal Society of London. Series B: Biological Sciences	2004	10.1098/rspb.2004.2801
GNM_069	Telomere length and dynamics of spotless starling nestlings depend on nest-building materials used by parents	Soler JJ; Ruiz-Castellano C; Figuerola J; Martín-Vivaldi M; Martínez-de la Puente J; Ruiz-Rodríguez M; Tomás G	Animal Behaviour	2017	10.1016/j.anbehav.2017.01.018
GNM_101	Interacting effects of aromatic plants and female age on nest-dwelling ectoparasites and blood-sucking flies in avian nests	Tomás G; Merino S; Martínez-de la Puente J; Moreno J; Morales J; Lobato E; Rivero-de Aguilar J; del Cerro S	Behavioural Processes	2012	10.1016/j.beproc.2012.02.003
GNM_121	Aromatic plants in nests of the blue tit <i>Cyanistes caeruleus</i> protect chicks from bacteria	Mennerat A; Mirleau P; Blondel J; Perret P; Lambrechts MM; Heeb P	Oecologia	2009	10.1007/s00442-009-1418-6

Paper ID	Title	Authors	Journal	Year	DOI
GNM_123	Aromatic plants in nests of blue tits: positive effects on nestlings	Mennerat A; Perret P; Bourgault P; Blondel J; Gimenez O; Thomas DW; Heeb P; Lambrechts MM	Animal Behaviour	2009	10.1016/j.anbehav.2008.11.008
GNM_128	Aromatic plants in blue tit <i>Cyanistes caeruleus</i> nests: no negative effect on blood-sucking <i>Protocalliphora</i> blow fly larvae	Mennerat A; Perret P; Caro SP; Heeb P; Lambrechts MM	Journal of Avian Biology	2008	10.1111/j.0908-8857.2008.04400.x
GNM_137	Experimental addition of greenery reduces flea loads in nests of a non-greenery using species, the tree swallow <i>Tachycineta bicolor</i>	Shutler D; Campbell AA	Journal of Avian Biology	2007	10.1111/j.2007.0908-8857.04015.x
GNM_143	European starlings: nestling condition, parasites and green nest material during the breeding season	Gwinner H; Berger S	Journal of Ornithology	2005	10.1007/s10336-005-0012-x
GNM_147	Does fresh vegetation protect avian nests from ectoparasites? An experiment with tree swallows	Dawson RD	Canadian Journal of Zoology	2004	10.1139/Z04-076

Paper ID	Title	Authors	Journal	Year	DOI
GNM_152	Green nesting material has a function in mate attraction in the European starling	Brouwer L; Komdeur J	Animal Behaviour	2004	10.1016/j.anbehav.2003.07.005
GNM_185	Effects of nest-box environment on fledgling success rate and pathogen load	Scott-Baumann JF; Morgan ER; Cogan TA	Parasitology	2022	10.1017/S0031182022000695
GNM_237	Nest Material Shapes Eggs Bacterial Environment	Ruiz-Castellano C; Tomás G; Ruiz-Rodríguez M; Martín-Gálvez D; Soler JJ	PLOS ONE	2016	10.1371/journal.pone.0148894
GNM_349	Ectoparasitism and the role of green nesting material in the European starling	Fauth PT; Krementz DG; Hines JE	Oecologia	1991	10.1007/BF00328399
GNM_355	Effect of biologically active plants used as nest material and the derived benefit to starling nestlings	Clark L; Mason JR	Oecologia	1988	10.1007/BF00379183
GNM_001_add	Is there a role for aromatic plants in blue tit (<i>Cyanistes caeruleus</i>) nests? Results from a correlational and an experimental study	Garrido-Bautista J; Ramos JA; Arce SI; Melero-Romero P; Ferreira R; Santos-Baena C; Guímaro HR; Martín-Villegas C; Moreno-Rueda G; Norte AC	Behavioral Ecology and Sociobiology	2023	10.1007/s00265-023-03393-9

Paper ID	Title	Authors	Journal	Year	DOI
GNM_001_rep	Effects of experimental nest treatment with herbs on ectoparasites and body condition of nestlings	Gładalski M; Norte AC; Bartos M; Demeško I; Kaliński A; Markowski M; Skwarska J; Wawrzyniak J; Zieliński P; Bańbura J	Behavioral Ecology	2025	10.1093/beheco/arae103
GNM_002_add	Green plants in starling nests: effects on nestlings	Gwinner H; Oltrogge M; Trost L; Nienaber U	Animal Behaviour	2000	10.1006/anbe.1999.1306
GNM_003_add	Sparrows use a medicinal herb to defend against parasites and increase offspring condition	Yang C; Ye P; Huo J; Møller AP; Liang W; Feeney WE	Current Biology	2020	10.1016/j.cub.2020.10.021
GNM_1_Unpublished	Greenery experiment – Common buzzard	Meinolf Ottensmann	Unpublished Data (Not peer-reviewed)	NA	NA
GNM_2_Unpublished	Aromatic plants in nests of blue tits (Cyanistes caeruleus)	Anna Pantazi	Master Thesis (Not peer-reviewed)	2024	https://hdl.handle.net/11250/3172874

7.7. Standardized Email Template

We included two unpublished studies in our meta-analysis:

7.8.0.1. GNM_1_Unpublished:.

Greenery manipulation experiment was conducted over two breeding seasons in 2018 and 2019. There were four treatment groups: Control, Spruce, Plastic (Branches of an artificial Christmas tree) and Greenery removal. Nests assigned to the spruce and plastic treatments, were lined with approximately ten twigs in such a way that the entire nest platform was covered. At nests assigned to the greenery removal treatment, all fresh greenery was carefully removed while the nest structure was left untouched. Greenery removal treatment

was conditioned on the existence of fresh greenery at the first sampling occasion. Control nests were handled similarly but the greenery composition was not altered. For the purpose of the meta-analysis we did not include the plastic treatment, only, addition or removal of green material and control group. Sampling interval was 5 to 19 days (mean: 7.72 ± 0.54 , $n = 42$ nests).

Sample sizes 42 nests in total: 18 nests allocated to the spruce treatment, 9 greenery removal, 10 plastic and 5 controls

Fitness proxies included in the meta-analysis:

1. Body condition of the nestling
2. Ectoparasite load (*Carnus hemapterus*): Semi-quantitative measure with five levels, ranging from 0 (no infestation on any of the four locations) to 4 (*C. hemapterus* larvae or fresh biting marks present on all four examined body locations)
3. Endoparasite load (*Leucocytozoon*): Five levels 0 – 4

Data used for the meta-analysis can be found data/02_data_extraction/data_extraction_MO_checkedSD.xlsx

7.8.0.2. GNM_2_Unpublished.:

An experiment on Blue Tits conducted in Bergen (Norway), adding lavender in nests and comparing nestling growth & survival with controls. 5 fresh leaves of lavender were added / replaced every 2nd day in broods in the “aromatic” group, from day 5 to day 13 after hatching. In the control group nothing was added, but the broods were otherwise treated similarly.

The following variables were provided in the dataset:

1. Weight D5, D7, D9, D11, D13 - Weight of nestlings at 5, 7, 9, 11, 13 days after hatching of first egg
2. Weight D14-17 - Final weight measure on nestlings at days 14-17 post-hatching, seen as fledging weight
3. TarsusLength D14-17 - Tarsus length measured on nestlings at days 14-17 post-hatching, 0.1mm accuracy
4. Percent Survived - Percentage of all chicks initially hatched in each group that survived until fledging (therefore no SD)
5. Survival Per Brood - Percentage of chicks in each brood that survived until fledging (the unit is the brood here)

This data was part of a Master thesis “**Aromatic plants in nests of blue tits (*Cyanistes caeruleus*)**” and all the details can be found at <https://hdl.handle.net/11250/3172874>.

Data used for the meta-analysis can be found data/02_data_extraction/data_extraction_UnpublishedThesis.xls

7.9. Complete List of Variables extracted from each study

- **Authors** Complete information of all authors. Each author name to be separated by ;. Example: Dave Shutler and Adam A. Campbell → “Shutler D; Campbell A A”.
- **Year of Publication** The year the article was published.
- **Population Location** Geographical location of the study population extracted as provided by the authors along with any coordinates provided. If multiple sites are studied in a location, extract them as PopulationName_SiteA.
- **Experiment ID** An identifier given to the estimates coming from the same sub-population within a location.
- **Group ID** An identifier given to the estimates coming from the same group of experimental units (e.g., males and females in a population, first clutch and second clutch).
- **Repeated Trait ID** An identifier given to the repeated measurement of the same trait within the same individuals (e.g., chick mass on day 7, 14, and 21).
- **Bird Species** Scientific name of the bird species studied.
- **Treatment Group Plant Species** Scientific or common name of the plant species used as the green nest material (treatment group).
- **Control Group Plant Species** Scientific name of the plant(s) species used as control treatment(s) in the study. If the scientific name is not given, extract it as reported by the authors. If there is no material used for control, note “blank”.
- **Comparison Type** Experimental design used to code the comparison between the two groups: 1 = non-aromatic (control) vs. aromatic (treatment) 2 = no material (control) vs. aromatic (treatment) 3 = no material (control) vs. non-aromatic (treatment)
- **CH Estimates** related to the courtship hypothesis — yes (1) or no (0).
- **PCH Estimates** related to the nest protection or drug hypothesis — yes (1) or no (0).

- Measure of Central Tendency (treatment group) Estimated value of central tendency for the samples of the treatment group.
- Type of Measure of Central Tendency (treatment group) Type of central tendency value reported for the treatment group (e.g., Mean, Median, Mode).
- Measure of Dispersal (treatment group) Value of the measure of variation for the treatment group.
- Type of Measure of Dispersal (treatment group) Type of variation of the data (e.g., SE, SD, CI95) reported for the treatment group.
- Sample Size (treatment group) Sample size for the treatment group.
- Measure of Central Tendency (control group) Estimated value of central tendency for the samples of the control group.
- Type of Measure of Central Tendency (control group) Type of central tendency value reported for the control group (e.g., Mean, Median, Mode).
- Measure of Dispersal (control group) Value of the measure of variation for the control group.
- Type of Measure of Dispersal (control group) Type of variation of the data (e.g., SE, SD, CI95) reported for the control group.
- Sample Size (control group) Sample size for the control group.
- Fitness Proxy Measured Variable name used in the study as a proxy of fitness. Extracted as presented in the article and later standardized across studies.
- Category/Type of Fitness Proxy Measured Type of trait (i.e., fitness proxy) studied: behaviour, morphology, parasite and pathogenic load, phenology, physiology, or reproduction.
- Fitness Proxy Sign Sign assigned in relation to fitness: +1 if fitness increases with proxy (e.g., survival rate, body mass). -1 if fitness decreases with proxy (e.g., mortality, parasite load).
- Statistics Type Type of statistics used to address the effect of GNM on fitness when primary data were not reported (e.g., t-test, ANOVA). Extracted only when central tendency was not reported.
- Test Statistics Type Type of test statistic used (e.g., t, F).
- Statistics Value Value of the test statistic used.
- P Value Reported p-value (e.g., $p = 0.003$).

- Sign Relationship Positive when treatment increases the trait value and negative when it decreases the trait value. Used when only the statistical value was provided.
- Total Sample Size Total sample size used to calculate the statistical value.
- Degree of Freedom Degrees of freedom for the statistics used.
- Data Location The part of the text or figure from which the data were extracted.
- Parasite Type Type of parasite when fitness proxy measured parasite/pathogen load (Arthropod or Micro-organism).
- Time of Green Nest Material Addition Time of green nest material addition by authors or birds [continuous (c), before (b), or after (a) egg hatching].
- Blinding Whether blinding was conducted — yes (y) or no (n).
- Random Assignment Whether nests were assigned to treatment or control randomly — yes (y) or no (n).
- Missing Data Whether data were only partially or not reported — yes (y) or no (n).
- Shared Treatment Group Used to identify non-independence between shared treatment groups. Value = 1 if individuals were compared only once as a treatment group.
- Shared Control Group Used to identify non-independence between shared control groups. Value = 1 if individuals were compared only once as a control group.

7.10. Random Effects

7.11. Sensitivity analysis of sampling correlation values (ρ) in variance-covariance structures

7.12. All sensitivity-analysis/robustness check

The overall results and conclusion were robust to several sensitivity analyses including the bivariate multilevel meta-analytic model including both effect sizes as response variables ($\ln\text{RR} = -0.015$, $p\text{-value} = 0.624$ and $\text{SMD(H)} = 0.250$, $p\text{-value} = 0.069$).

7.13. Plots without cropping the axis (showing complete X-axis)

7.14. Risk of Bias Test

The risk of bias categories we evaluated (i.e., blinding, random assignment to each treatment and partial reporting) were not statistically significant for $\ln\text{RR}$ (Table S4). SMD(H) was positive and statistically significant for studies without blinding ($\text{SMD(H)}_{\text{No Blinding}} = 0.184$, 95% CI: [0.051, 0.316], 95% PI: [-0.828, 1.195], $p\text{-value} = 0.007$, $k = 242$, $n = 25$; Table S4), with random assignment to treatment ($\text{SMD(H)}_{\text{Random Assignment}}$

= 0.205, 95% CI: [0.049, 0.360], 95% PI: [-0.813, 1.223], p-value = 0.010, k = 225, n = 23; Table S4) and where no data was missing ($SMD(H)_{No\ Partial\ Reporting} = 0.196$, 95% CI: [0.055, 0.337], 95% PI: [-0.817, 1.209], p-value = 0.007, k = 216, n = 22; Table S4). Notably, other levels of each category included fewer than four studies, indicating that the observed significance likely reflects the higher statistical power in the larger sample level rather than any systematic bias. All the three categories for risk of bias explained very little heterogeneity ($R^2_{marginal(lnRR)} > 0.8\%$; $R^2_{marginal(SMD(H))} > 0.7\%$).

7.15. Publication Bias

Since published effect sizes did not seem to change considerably over time, our analyses suggest no clear evidence of decline effects (i.e., time-lag bias) in published effect sizes ($lnRR_{intercept} = 0.021$, 95% CI: [-0.011; 0.053], p-value = 0.196, $lnRR_{slope} = -0.002$, 95% CI: [-0.005 to 0.001], p-value = 0.225, k = 221, n = 24, $R^2_{marginal(lnRR)} = 1.8\%$; $SMDH_{intercept} = 0.193$, 95% CI: [0.058 to 0.329], p-value = 0.005, $SMDH_{slope} = -0.008$, 95% CI: [-0.023 to 0.006], p-value = 0.264, k = 235, n = 24, $R^2_{marginal(SMD(H))} = 2.9\%$).

The search was conducted twice, the first time on 07.09.2022 and then repeated on 12.08.2024 to ensure it was the most up to date and did not miss any additional articles on the topic that were published meanwhile. For both the searches, we did not use any time limitation i.e. it included all the published literature till that given date. We used **Advance Search** option for both Web of Science Core Collection Database and Scopus Database using the search query given below.

Search string for the Web of Science Core Collection

Databases used in the search included

-Science Citation Index Expanded (SCI-EXPANDED) [1900-present]

-Social Sciences Citation Index (SSCI) [1900-present]

-Arts & Humanities Citation Index (AHCI) [1975 - present]

-Conference Proceedings Citation Index - Science (CPCI-S) [1991 - present]

-Conference Proceedings Citation Index - Social Science & Humanities (CPCI-SSH) [1991 - present]

-Emerging Sources Citation Index (ESCI) [2017- present]

We used advance search option and Title (TI), Abstract (AB) and Author Keywords (AK) to conduct our systematic search using the following string

<TI=(((“green*” OR “herb*” OR “aromatic*”) AND “nest*” AND (“bird*” OR “aves” OR “avian” OR “ornithol*” OR “passerine*” OR “passeriform*” OR “songbird*” OR “Accipiter” OR “Aegypius” OR “Aquila” OR “Aviceda” OR “Busarellus” OR “Butastur” OR “Buteo” OR “Buteogallus” OR “Chondrohierax” OR “Circaetus” OR “Circus” OR “Clanga” OR “Elanoides” OR “Elanus” OR “Gampsonyx” OR “Geranoaetus”

826 OR "Geranospiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR "Haliastur" OR
 827 "Harpagus" OR "Harpia" OR "Hieraetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco" OR "Leptodon" OR
 828 "Leucopternis" OR "Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus" OR "Necrosyrtes" OR
 829 "Neophron" OR "Nisaetus" OR "Parabuteo" OR "Pernis" OR "Pithecophaga" OR "Polemaetus" OR "Pseudas-
 830 tur" OR "Rostrhamus" OR "Rupornis" OR "Sarcogyps" OR "Spilornis" OR "Spizaetus" OR "Stephanoaetus"
 831 OR "Terathopius" OR "Torgos" OR "Trigonocephalus" OR "Cathartes" OR "Coragyps" OR "Gymnogyps" OR
 832 "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix" OR "Alopochen" OR "Amazonetta"
 833 OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR "Bucephala" OR "Cairina" OR
 834 "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloeophaga" OR "Clangula" OR "Coscoroba"
 835 OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR "Histrionicus" OR "Hymeno-
 836 laimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca" OR "Marmaronetta" OR
 837 "Melanitta" OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neochen" OR "Netta" OR "Nettapus"
 838 OR "Nomonyx" OR "Oxyura" OR "Plectropterus" OR "Polysticta" OR "Pteronetta" OR "Sarkidiornis"
 839 OR "Sibirionetta" OR "Somateria" OR "Spatula" OR "Tachyeres" OR "Tadorna" OR "Thalassornis" OR
 840 "Anhima" OR "Chauna" OR "Anseranas" OR "Aegotheles" OR "Aerodramus" OR "Aeronautus" OR "Apus"
 841 OR "Chaetura" OR "Collocalia" OR "Cypseloides" OR "Cypsiurus" OR "Hirundapus" OR "Panyptila" OR
 842 "Streptoprocne" OR "Tachornis" OR "Tachymarptis" OR "Hemiprocne" OR "Abeillia" OR "Adelomyia"
 843 OR "Aglaeactis" OR "Aglaiocercus" OR "Amazilia" OR "Androdon" OR "Anopetia" OR "Anthocephala"
 844 OR "Anthracothorax" OR "Archilochus" OR "Atthis" OR "Augastes" OR "Avocettula" OR "Basilinna" OR
 845 "Boissonneaua" OR "Calliphlox" OR "Calypte" OR "Campylopterus" OR "Chaetocercus" OR "Chalcostigma"
 846 OR "Chalybura" OR "Chlorestes" OR "Chlorostilbon" OR "Chrysolampis" OR "Chrysornis" OR "Clyto-
 847 laema" OR "Coeligena" OR "Colibri" OR "Cyanophaea" OR "Cyananthus" OR "Discosura" OR "Doricha"
 848 OR "Doryfera" OR "Elvira" OR "Ensifera" OR "Eriocnemis" OR "Eugenes" OR "Eulampis" OR "Eulidia"
 849 OR "Eupetomena" OR "Eupherusa" OR "Eutoxeres" OR "Florisuga" OR "Glaucis" OR "Goethalsia" OR
 850 "Goldmania" OR "Haplophaedia" OR "Heliactin" OR "Heliangelus" OR "Heliodoxa" OR "Heliomaster"
 851 OR "Heliophryx" OR "Hylocharis" OR "Juliamyia" OR "Klais" OR "Lafresnaya" OR "Lampornis" OR
 852 "Lamprolaima" OR "Lepidopyga" OR "Lesbia" OR "Leucippus" OR "Leucochloris" OR "Lophornis" OR
 853 "Mellisuga" OR "Metallura" OR "Microchera" OR "Myrmia" OR "Myrtis" OR "Ocreatus" OR "Opistho-
 854 prora" OR "Oreonympha" OR "Oreotrochilus" OR "Orthorhynchus" OR "Oxyechus" OR "Pantherpe" OR
 855 "Patagona" OR "Phaeochroa" OR "Phaethornis" OR "Phlogophilus" OR "Polyonymus" OR "Polytmus" OR
 856 "Pterophanes" OR "Ramphodon" OR "Ramphomicron" OR "Rhodopsis" OR "Sappho" OR "Schistes" OR
 857 "Selasphorus" OR "Sephanoides" OR "Stephanoxis" OR "Sternoclyta" OR "Taphrosphilus" OR "Thalurania"
 858 OR "Thaumatocorys" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR
 859 "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR
 860 "Rhabdoterrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus"

861 OR "Upupa" OR "Antrostomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus"
 862 OR "Hydropsalis" OR "Lurocalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus"
 863 OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius"
 864 OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariama" OR "Casuarius" OR "Dromaius" OR
 865 "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cererhinca" OR "Fratricula" OR
 866 "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus"
 867 OR "Charadrius" OR "Elseyornis" OR "Erythrogonyx" OR "Hoploxypterus" OR "Oreopholus" OR "Pel-
 868 tohyas" OR "Phegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR
 869 "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR
 870 "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR
 871 "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelochelidon" OR "Gygis" OR "Hydrocoloeus" OR
 872 "Hydroprogne" OR "Ichthyaetus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR
 873 "Pagophila" OR "Phaetusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula"
 874 OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR
 875 "Recurvirostra" OR "Nycticryphes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR
 876 "Calidris" OR "Coenocorypha" OR "Gallinago" OR "Limnodromus" OR "Limosa" OR "Lymnocyptes" OR
 877 "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagis"
 878 OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR
 879 "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chal-
 880 cophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula"
 881 OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR
 882 "Gymnophaps" OR "Hemiphaga" OR "Henicophaps" OR "Leptotila" OR "Leptotrygon" OR "Leucosarcia"
 883 OR "Lopholaimus" OR "Macropygia" OR "Metriopelia" OR "Ocyphaps" OR "Oena" OR "Otidiphaps"
 884 OR "Patagioenas" OR "Petrophassa" OR "Pezophaps" OR "Phapitreron" OR "Phaps" OR "Ptilinopus"
 885 OR "Raphus" OR "Reinwardtoena" OR "Spilopelia" OR "Streptopelia" OR "Treron" OR "Trugon" OR
 886 "Turacoena" OR "Turtur" OR "Uropelia" OR "Zenaida" OR "Zentrygon" OR "Actenoides" OR "Alcedo" OR
 887 "Caridonax" OR "Ceryle" OR "Ceyx" OR "Chloroceryle" OR "Cittura" OR "Corythornis" OR "Dacelo" OR
 888 "Halcyon" OR "Ispidina" OR "Lacedo" OR "Megaceryle" OR "Melidora" OR "Syma" OR "Tanyptera" OR
 889 "Todiramphus" OR "Atelornis" OR "Brachypteracias" OR "Geobiastes" OR "Coracias" OR "Eurystomus"
 890 OR "Merops" OR "Nyctyornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes"
 891 OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR
 892 "Chrysococcyx" OR "Clamator" OR "Coccyzua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus"
 893 OR "Dasylophus" OR "Dromococcyx" OR "Eudynamis" OR "Geococcyx" OR "Guira" OR "Hierococcyx"
 894 OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicophaeus" OR "Piaya" OR "Rhinortha"
 895 OR "Scythrops" OR "Surniculus" OR "Tapera" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR

896 “Anomalopteryx” OR “Emeus” OR “Euryapteryx” OR “Eurypyga” OR “Rhynochetos” OR “Caracara” OR
 897 “Daptrius” OR “Falco” OR “Herpetotheres” OR “Ibycter” OR “Micrastur” OR “Microhierax” OR “Milvago”
 898 OR “Phalcoboenus” OR “Polihierax” OR “Spiziapteryx” OR “Aburria” OR “Chamaepetes” OR “Crax” OR
 899 “Mitu” OR “Nothocrax” OR “Oreophasis” OR “Ortalis” OR “Pauxi” OR “Penelope” OR “Penelopina” OR
 900 “Pipile” OR “Alectura” OR “Megapodius” OR “Acryllium” OR “Guttera” OR “Numida” OR “Callipepla”
 901 OR “Colinus” OR “Cyrtonyx” OR “Dendrortyx” OR “Odontophorus” OR “Oreortyx” OR “Philortyx” OR
 902 “Ptilopachus” OR “Rhynchortyx” OR “Alectoris” OR “Ammoperdix” OR “Arborophila” OR “Argusianus”
 903 OR “Bambusicola” OR “Bonasa” OR “Caloperdix” OR “Centrocercus” OR “Chrysolophus” OR “Coturnix”
 904 OR “Crossoptilon” OR “Dendragapus” OR “Excalfactoria” OR “Falcipennis” OR “Francolinus” OR “Gallus”
 905 OR “Haematoryx” OR “Ithaginis” OR “Lagopus” OR “Lerwa” OR “Lophophorus” OR “Lophura” OR
 906 “Lyrurus” OR “Meleagris” OR “Pavo” OR “Peliperdix” OR “Perdix” OR “Phasianus” OR “Polyplectron” OR
 907 “Pternistis” OR “Pucrasia” OR “Rhizothera” OR “Rollulus” OR “Scleroptila” OR “Syrmaticus” OR “Tetrao”
 908 OR “Tetraogallus” OR “Tetraophasis” OR “Tragopan” OR “Tympanuchus” OR “Gavia” OR “Aramus”
 909 OR “Anthropoides” OR “Antigone” OR “Balearica” OR “Bugeranus” OR “Grus” OR “Leucogeranus” OR
 910 “Heliornis” OR “Psophia” OR “Aenigmatolimnas” OR “Amaurolimnas” OR “Amaurornis” OR “Anurolimnas”
 911 OR “Aramides” OR “Atlantisia” OR “Coturnicops” OR “Crex” OR “Dryolimnas” OR “Eulabeornis” OR
 912 “Fulica” OR “Gallicrex” OR “Gallinula” OR “Gallirallus” OR “Laterallus” OR “Lewinia” OR “Micropygia”
 913 OR “Neocrex” OR “Nesoclopeus” OR “Paragallinula” OR “Pardirallus” OR “Porphyrio” OR “Porphyriops”
 914 OR “Porzana” OR “Rallacula” OR “Rallina” OR “Rallus” OR “Tribonyx” OR “Canirallus” OR “Sarothrura”
 915 OR “Leptosomus” OR “Corythaeola” OR “Corythaixoides” OR “Crinifer” OR “Musophaga” OR “Tauraco”
 916 OR “Opisthocomus” OR “Afrotis” OR “Ardeotis” OR “Chlamydotis” OR “Neotis” OR “Otis” OR “Tetrax”
 917 OR “Acanthisitta” OR “Pachyplichas” OR “Traversia” OR “Xenicus” OR “Acanthiza” OR “Acanthornis” OR
 918 “Aethomyias” OR “Aphelocephala” OR “Calamanthus” OR “Gerygone” OR “Hylacola” OR “Neosericornis”
 919 OR “Oreoscopus” OR “Origma” OR “Pachycare” OR “Pycnoptilus” OR “Pyrrholaemus” OR “Sericornis”
 920 OR “Smicrornis” OR “Acrocephalus” OR “Arundinax” OR “Calamonastides” OR “Hippolais” OR “Iduna”
 921 OR “Nesillas” OR “Aegithalos” OR “Leptopoecile” OR “Psaltiriparus” OR “Aegithina” OR “Alaemon” OR
 922 “Alauda” OR “Alaudala” OR “Ammomanes” OR “Calandrella” OR “Calendulauda” OR “Chersomanes”
 923 OR “Eremophila” OR “Eremopterix” OR “Galerida” OR “Lullula” OR “Melanocorypha” OR “Mirafraga” OR
 924 “Pinarocorys” OR “Spizocorys” OR “Artamus” OR “Gymnorhina” OR “Melloria” OR “Bombycilla” OR
 925 “Buphagus” OR “Calcarius” OR “Plectrophenax” OR “Rhynchophanes” OR “Callaeas” OR “Heteralocha” OR
 926 “Philesturnus” OR “Calyptophilus” OR “Campephaga” OR “Ceblepyris” OR “Coracina” OR “Edolisoma” OR
 927 “Lalage” OR “Malindangia” OR “Pericrocotus” OR “Amaurospiza” OR “Cardinalis” OR “Caryothraustes” OR
 928 “Chlorothraupis” OR “Cyanocompsa” OR “Cyanoloxia” OR “Granatellus” OR “Habia” OR “Passerina” OR
 929 “Periporphyrus” OR “Pheucticus” OR “Piranga” OR “Spiza” OR “Certhia” OR “Salpornis” OR “Abroscopus”
 930 OR “Cettia” OR “Horornis” OR “Phyllergates” OR “Tesia” OR “Urosphena” OR “Chloropsis” OR “Cinclus”

931 OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR
 932 "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia"
 933 OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR
 934 "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR
 935 "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR
 936 "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus"
 937 OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus"
 938 OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR
 939 "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias"
 940 OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR
 941 "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza"
 942 OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR
 943 "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis"
 944 OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga"
 945 OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calypomena" OR "Cymbirhynchus" OR
 946 "Eurylaimus" OR "Psarisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR
 947 "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chlo-
 948 ris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra"
 949 OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorhous" OR "Hemignathus" OR
 950 "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR
 951 "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR
 952 "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza"
 953 OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR
 954 "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus"
 955 OR "Certhiasomus" OR "Certhiaxis" OR "Cinclodes" OR "Clibanornis" OR "Coryphistera" OR "Cranioleuca"
 956 OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR
 957 "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geocerthia" OR "Geositta" OR "Glyphorhynchus"
 958 OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR
 959 "Limnortites" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR
 960 "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor"
 961 OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR
 962 "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus"
 963 OR "Spartonoica" OR "Sylviorthorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonimus" OR
 964 "Thripadectes" OR "Thriphophaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes"
 965 OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hylopezus" OR "Myrmothera" OR "Alopocheli-

966 don" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon"
 967 OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psolidoprocne"
 968 OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR
 969 "Tachycineta" OR "Hylia" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Am-
 970 blycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives"
 971 OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus"
 972 OR "Lamprospiza" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR
 973 "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar"
 974 OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura"
 975 OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus"
 976 OR "Leiothrix" OR "Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochalopecton" OR "Turdoides" OR
 977 "Bradypterus" OR "Cincloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR
 978 "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR
 979 "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malaconotus" OR "Nilaus" OR "Rhodophoneus" OR
 980 "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR
 981 "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia"
 982 OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomy-
 983 zon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes"
 984 OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis"
 985 OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius"
 986 OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cinclocerthia" OR "Dumetella" OR
 987 "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR
 988 "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis"
 989 OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus"
 990 OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR
 991 "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Campicoloides" OR "Cercotrichas" OR "Chamaetylas"
 992 OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR
 993 "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia"
 994 OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiomela" OR "Myioparus" OR "Myophonus"
 995 OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola"
 996 OR "Sheppardia" OR "Sholicola" OR "Stiphornis" OR "Tarsiger" OR "Thamnolaea" OR "Vauriella" OR
 997 "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR
 998 "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkinogramma" OR "Leptocoma" OR "Nectarinia"
 999 OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleadryas" OR "Oriolus" OR
 1000 "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR

1001 "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR
 1002 "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR
 1003 "Cyanistes" OR "Lophophanes" OR "Machlolophus" OR "Melaniparus" OR "Melanochlora" OR "Pardali-
 1004 parus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus"
 1005 OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros"
 1006 OR "Leiothlypis" OR "Limnothlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis"
 1007 OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR
 1008 "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisiospiza"
 1009 OR "Atlapetes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza"
 1010 OR "Melozone" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR
 1011 "Pezopetes" OR "Pipilo" OR "Pooecetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizel-
 1012 loides" OR "Xenospiza" OR "Zonotrichia" OR "Carpospiza" OR "Gymnoris" OR "Hypocryptadius" OR
 1013 "Montifringilla" OR "Onychostruthus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR
 1014 "Gampsorhynchus" OR "Illadopsis" OR "Jabouilleia" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR
 1015 "Malacopteron" OR "Napothena" OR "Pellorneum" OR "Rimtor" OR "Trichastoma" OR "Amalocichla"
 1016 OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca"
 1017 OR "Monachella" OR "Pachycephalopsis" OR "Peneoenanthe" OR "Peneothello" OR "Petroica" OR "Poe-
 1018 cilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR
 1019 "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR
 1020 "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR
 1021 "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyrannetes" OR "Xenopipo"
 1022 OR "Erythropitta" OR "Hydrornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalor-
 1023 nis" OR "Euplectes" OR "Malimbus" OR "Plocepasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR
 1024 "Pnoepyga" OR "Microbates" OR "Polioptila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR
 1025 "Cinclosoma" OR "Ptilorrhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR
 1026 "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla"
 1027 OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Crimiger" OR "Eurillas" OR
 1028 "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus"
 1029 OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus"
 1030 OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pterop-
 1031 tochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR
 1032 "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynch" OR "Culicicapa" OR "Elminia" OR "Acridotheres"
 1033 OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora"
 1034 OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotonis" OR "Leucopsar"
 1035 OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR

1036 “Sarcops” OR “Scissirostrum” OR “Speculipastor” OR “Spodiopsar” OR “Streptocitta” OR “Sturnia” OR
 1037 “Sturnus” OR “Chamaea” OR “Chleuasicus” OR “Chloropeta” OR “Chrysomma” OR “Conostoma” OR
 1038 “Fulvetta” OR “Lioparus” OR “Paradoxornis” OR “Pseudoalcippe” OR “Psittiparus” OR “Rhopophilus” OR
 1039 “Sinosuthora” OR “Suthora” OR “Sylvia” OR “Teretistris” OR “Akletos” OR “Ampelornis” OR “Aprosi-
 1040 tornis” OR “Batara” OR “Cercomacra” OR “Cercomacroides” OR “Cymbilaimus” OR “Dichrozona” OR
 1041 “Drymophila” OR “Dysithamnus” OR “Epinecrophylla” OR “Euchrepomis” OR “Formicivora” OR “Freder-
 1042 ickena” OR “Gymnocichla” OR “Gymnopithys” OR “Hafferia” OR “Herpsilochmus” OR “Hylophylax” OR
 1043 “Hypocnemis” OR “Hypocnemoides” OR “Hypoedaleus” OR “Isleria” OR “Mackenziaena” OR “Megastictus”
 1044 OR “Microrhopias” OR “Myrmeciza” OR “Myrmelastes” OR “Myrmoborus” OR “Myrmochanes” OR “Myr-
 1045 moderus” OR “Myrmophylax” OR “Myrmorchilus” OR “Myrmornis” OR “Myrmotherula” OR “Neotantes”
 1046 OR “Oneillornis” OR “Percnostola” OR “Phaenostictus” OR “Phlegopsis” OR “Pithys” OR “Poliocrania”
 1047 OR “Pygiptila” OR “Pyriglena” OR “Rhegmatorhina” OR “Rhopias” OR “Rhopornis” OR “Sakesphorus”
 1048 OR “Sciaphylax” OR “Sclateria” OR “Sipia” OR “Taraba” OR “Thamnistes” OR “Thamnomanes” OR
 1049 “Thamnophilus” OR “Willisornis” OR “Xenornis” OR “Anisognathus” OR “Bangsia” OR “Buthraupis”
 1050 OR “Calochaetes” OR “Catamblyrhynchus” OR “Catamenia” OR “Charitospiza” OR “Chlorochrysa” OR
 1051 “Chlorophanes” OR “Chlorornis” OR “Chrysothlypis” OR “Cissopis” OR “Cnemoscopus” OR “Coereba”
 1052 OR “Compsospiza” OR “Compsothraupis” OR “Conirostrum” OR “Conothraupis” OR “Coryphospingus”
 1053 OR “Creurgops” OR “Cyanerpes” OR “Cyanicterus” OR “Cypsnagra” OR “Dacnis” OR “Delothraupis”
 1054 OR “Diglossa” OR “Diuca” OR “Dolospingus” OR “Donacospiza” OR “Dubusia” OR “Emberizoides” OR
 1055 “Embernagra” OR “Eucometis” OR “Euneornis” OR “Geospiza” OR “Gubernatrix” OR “Haplospiza” OR
 1056 “Hemispingus” OR “Hemithraupis” OR “Heterospingus” OR “Incasppiza” OR “Iridophanes” OR “Iridosornis”
 1057 OR “Lanio” OR “Lophospingus” OR “Loxigilla” OR “Loxipasser” OR “Melanodera” OR “Melanospiza” OR
 1058 “Melopyrrha” OR “Nemosia” OR “Nephelornis” OR “Oreomanes” OR “Oryzoborus” OR “Parkerthraustes”
 1059 OR “Paroaria” OR “Phrygilus” OR “Piezorina” OR “Pipraeidea” OR “Poospiza” OR “Pyrrhocomma” OR
 1060 “Ramphocelus” OR “Rhodospingus” OR “Saltator” OR “Saltatricula” OR “Schistochlamys” OR “Sericossypha”
 1061 OR “Sicalis” OR “Sporophila” OR “Stephanophorus” OR “Tachyphonus” OR “Tangara” OR “Tersina” OR
 1062 “Thlypopsis” OR “Thraupis” OR “Tiaris” OR “Trichothraupis” OR “Volatinia” OR “Wetmorethraupis” OR
 1063 “Xenodacnis” OR “Xenospingus” OR “Tichodroma” OR “Macronus” OR “Pomatorhinus” OR “Spelaeornis”
 1064 OR “Stachyridopsis” OR “Stachyris” OR “Iodopleura” OR “Laniisoma” OR “Laniocera” OR “Myiobius”
 1065 OR “Onychorhynchus” OR “Oxyruncus” OR “Pachyramphus” OR “Schiffornis” OR “Terenotriccus” OR
 1066 “Tityra” OR “Xenopsaris” OR “Campylorhynchus” OR “Cantorchilus” OR “Catherpes” OR “Cinnycerthia”
 1067 OR “Cistothorus” OR “Cyphorhinus” OR “Henicorhina” OR “Microcerculus” OR “Odontorchilus” OR
 1068 “Pheugopedius” OR “Salpinctes” OR “Thryomanes” OR “Thryophilus” OR “Thryothorus” OR “Troglodytes”
 1069 OR “Uropsila” OR “Catharus” OR “Cichlopsis” OR “Entomodestes” OR “Geokichla” OR “Hylocichla” OR
 1070 “Ixoreus” OR “Myadestes” OR “Neocossyphus” OR “Ridgwayia” OR “Sialia” OR “Stizorhina” OR “Turdus”

1071 OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola"
 1072 OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Capsiempis" OR "Casiornis" OR "Cnemarchus" OR
 1073 "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR
 1074 "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR
 1075 "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis"
 1076 OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR
 1077 "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus"
 1078 OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myio-
 1079 dynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiotriccus" OR
 1080 "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR
 1081 "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes"
 1082 OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis"
 1083 OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR
 1084 "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis"
 1085 OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR
 1086 "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis"
 1087 OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR
 1088 "Urocynchramus" OR "Artamella" OR "Calicalicus" OR "Cyanolanius" OR "Euryceros" OR "Falculea"
 1089 OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR
 1090 "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris"
 1091 OR "Anomalospiza" OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR
 1092 "Vireo" OR "Vireolanius" OR "Zeledonia" OR "Apalopteron" OR "Dasycrotapha" OR "Heleia" OR "Lopho-
 1093 zosterops" OR "Sterrhoptilus" OR "Yuhina" OR "Zosterops" OR "Zosterornis" OR "Agamia" OR "Ardea" OR
 1094 "Ardeola" OR "Botaurus" OR "Bubulcus" OR "Butorides" OR "Cochlearius" OR "Dupetor" OR "Egretta"
 1095 OR "Gorsachius" OR "Ixobrychus" OR "Nyctanassa" OR "Nycticorax" OR "Ptilerodius" OR "Syrigma" OR
 1096 "Tigriornis" OR "Tigrisoma" OR "Zebrius" OR "Balaeniceps" OR "Pelecanus" OR "Scopus" OR "Bostrychia"
 1097 OR "Eudocimus" OR "Geronticus" OR "Lophotibis" OR "Mesembrinibis" OR "Nipponia" OR "Phimosus"
 1098 OR "Platalea" OR "Plegadis" OR "Pseudibis" OR "Theristicus" OR "Threskiornis" OR "Phaethon" OR
 1099 "Phoeniconaias" OR "Phoenicoparrus" OR "Phoenicopterus" OR "Bucco" OR "Chelidoptera" OR "Hapalop-
 1100 tila" OR "Hypnelus" OR "Malacoptila" OR "Micromonacha" OR "Monasa" OR "Nonnula" OR "Notharchus"
 1101 OR "Nystalus" OR "Capito" OR "Eubucco" OR "Brachygalba" OR "Galbalcyrrhynchus" OR "Galbula" OR
 1102 "Jacamaralcyon" OR "Jacamerops" OR "Indicator" OR "Melichneutes" OR "Malignomon" OR "Prodotiscus"
 1103 OR "Buccanodon" OR "Gymnobucco" OR "Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus"
 1104 OR "Psilopogon" OR "Blythipicus" OR "Campephilus" OR "Campethera" OR "Celeus" OR "Chloropicus"
 1105 OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Dendrocopos" OR "Dendrocytes" OR

1106 “Dendropicos” OR “Dinopium” OR “Dryobates” OR “Dryocopus” OR “Gecinulus” OR “Geocolaptes” OR
 1107 “Jynx” OR “Leiopicus” OR “Leuconotopicus” OR “Meiglyptes” OR “Melanerpes” OR “Micropternus” OR
 1108 “Mulleripicus” OR “Nesotites” OR “Picoides” OR “Piculus” OR “Picumnus” OR “Picus” OR “Reinwardtipi-
 1109 cus” OR “Sasia” OR “Sphyrapicus” OR “Veniliornis” OR “Yungipicus” OR “Andigena” OR “Aulacorhynchus”
 1110 OR “Pteroglossus” OR “Ramphastos” OR “Selenidera” OR “Semnornis” OR “Aechmophorus” OR “Podiceps”
 1111 OR “Podilymbus” OR “Poliiocephalus” OR “Rollandia” OR “Tachybaptus” OR “Diomedea” OR “Phoebastria”
 1112 OR “Phoebetria” OR “Thalassarche” OR “Hydrobates” OR “Oceanodroma” OR “Fregetta” OR “Garrodia”
 1113 OR “Nesofregetta” OR “Oceanites” OR “Pelagodroma” OR “Aphrodroma” OR “Ardena” OR “Bulweria”
 1114 OR “Calonectris” OR “Daption” OR “Fulmarus” OR “Halobaena” OR “Macronectes” OR “Pachyptila” OR
 1115 “Pagodroma” OR “Pelecanoides” OR “Procellaria” OR “Pseudobulweria” OR “Pterodroma” OR “Puffinus” OR
 1116 “Thalassioica” OR “Cacatua” OR “Callocephalon” OR “Calyptorhynchus” OR “Eolophus” OR “Lophochroa”
 1117 OR “Nymphicus” OR “Probosciger” OR “Alipiopsitta” OR “Amazona” OR “Anodorhynchus” OR “Ara” OR
 1118 “Aratinga” OR “Bolborhynchus” OR “Brotogeris” OR “Conuropsis” OR “Cyanoliseus” OR “Cyanopsitta” OR
 1119 “Derophtus” OR “Diopsittaca” OR “Enicognathus” OR “Eupsittula” OR “Forpus” OR “Graydidascalus” OR
 1120 “Guarouba” OR “Guaruba” OR “Hapalopsittaca” OR “Leptosittaca” OR “Myiopsitta” OR “Nannopsittaca”
 1121 OR “Orthopsittaca” OR “Pionites” OR “Pionopsitta” OR “Pionus” OR “Poicephalus” OR “Primolius”
 1122 OR “Psilopsiagon” OR “Psittacara” OR “Psittacus” OR “Pyralia” OR “Pyrrhura” OR “Rhynchopsitta”
 1123 OR “Thectocercus” OR “Touit” OR “Triclaria” OR “Agapornis” OR “Alisterus” OR “Aprosmictus” OR
 1124 “Barnardius” OR “Bolbopsittacus” OR “Chalcopsitta” OR “Charmosyna” OR “Coracopsis” OR “Cyanoram-
 1125 phus” OR “Cyclopsitta” OR “Eclectus” OR “Eos” OR “Eunymphicus” OR “Geoffroyus” OR “Glossopsitta”
 1126 OR “Lathamus” OR “Loriculus” OR “Lorius” OR “Melopsittacus” OR “Micropsitta” OR “Neophema” OR
 1127 “Neopsephotus” OR “Neopsittacus” OR “Northiella” OR “Oreopsittacus” OR “Parvipsitta” OR “Pezoporus”
 1128 OR “Phigys” OR “Platycercus” OR “Polytelis” OR “Prioniturus” OR “Prosopiea” OR “Psephotellus” OR
 1129 “Psephotus” OR “Pseudeos” OR “Psittacella” OR “Psittacula” OR “Psittaculirostris” OR “Psitteuteles” OR
 1130 “Psittinus” OR “Psittichas” OR “Purpureicephalus” OR “Tanygnathus” OR “Trichoglossus” OR “Vini” OR
 1131 “Nestor” OR “Strigops” OR “Pterocles” OR “Syrrhaptes” OR “Rhea” OR “Aptenodytes” OR “Eudytes” OR
 1132 “Eudiptula” OR “Megadyptes” OR “Pygoscelis” OR “Spheniscus” OR “Aegolius” OR “Asio” OR “Athene”
 1133 OR “Bubo” OR “Glaucidium” OR “Ketupa” OR “Lophostrix” OR “Megascops” OR “Micrathene” OR “Ninox”
 1134 OR “Otus” OR “Pseudoscops” OR “Psiloscops” OR “Pulsatrix” OR “Sceloglaux” OR “Scotopelia” OR “Strix”
 1135 OR “Surnia” OR “Xenoglaux” OR “Phodilus” OR “Tyto” OR “Aepyornis” OR “Mullerornis” OR “Struthio”
 1136 OR “Anhinga” OR “Fregata” OR “Leucocarbo” OR “Microcarbo” OR “Nannopterum” OR “Phalacrocorax”
 1137 OR “Morus” OR “Sula” OR “Crypturellus” OR “Eudromia” OR “Nothocercus” OR “Nothoprocta” OR
 1138 “Nothura” OR “Rhynchotus” OR “Tinamotis” OR “Tinamus” OR “Apalharpactes” OR “Apaloderma” OR
 1139 “Harpactes” OR “Pharomachrus” OR “Trogon” OR “Tetrastes” OR “Hesperiphona”) AND (“experiment*”
 1140 OR “manipulat*”))) OR AB= (((“green*” OR “herb*” OR “aromatic*”) AND “nest*” AND (“bird*” OR

1141 “aves” OR “avian” OR “ornithol*” OR “passerine*” OR “passeriform*” OR “songbird*” OR “Accipiter” OR
 1142 “Aegyptius” OR “Aquila” OR “Aviceda” OR “Busarellus” OR “Butastur” OR “Buteo” OR “Buteogallus” OR
 1143 “Chondrohierax” OR “Circaetus” OR “Circus” OR “Clanga” OR “Elanoides” OR “Elanus” OR “Gampsonyx”
 1144 OR “Geranoaetus” OR “Geranospiza” OR “Gypaetus” OR “Gypohierax” OR “Gyps” OR “Haliaeetus” OR
 1145 “Haliastur” OR “Harpagus” OR “Harpia” OR “Hieraetus” OR “Ictinaetus” OR “Ictinia” OR “Kaupifalco”
 1146 OR “Leptodon” OR “Leucopternis” OR “Lophotriorchis” OR “Milvus” OR “Morphnarchus” OR “Morphnus”
 1147 OR “Necrosyrtes” OR “Neophron” OR “Nisaetus” OR “Parabuteo” OR “Pernis” OR “Pithecophaga” OR
 1148 “Polemaetus” OR “Pseudastur” OR “Rostrhamus” OR “Rupornis” OR “Sarcogyps” OR “Spilornis” OR
 1149 “Spizaetus” OR “Stephanoaetus” OR “Terathopius” OR “Torgos” OR “Trigonoceps” OR “Cathartes” OR
 1150 “Coragyps” OR “Gymnogyps” OR “Sarcoramphus” OR “Vultur” OR “Pandion” OR “Sagittarius” OR “Aix”
 1151 OR “Alopochen” OR “Amazonetta” OR “Anas” OR “Anser” OR “Asarcornis” OR “Aythya” OR “Branta” OR
 1152 “Bucephala” OR “Cairina” OR “Callonetta” OR “Cereopsis” OR “Chen” OR “Chenonetta” OR “Chloephaga”
 1153 OR “Clangula” OR “Coscoroba” OR “Cyanochen” OR “Cygnus” OR “Dendrocygna” OR “Heteronetta” OR
 1154 “Histrionicus” OR “Hymenolaimus” OR “Lophodytes” OR “Lophonetta” OR “Malacorhynchus” OR “Mareca”
 1155 OR “Marmaronetta” OR “Melanitta” OR “Merganetta” OR “Mergellus” OR “Mergus” OR “Neochen” OR
 1156 “Netta” OR “Nettapus” OR “Nomonyx” OR “Oxyura” OR “Plectropterus” OR “Polysticta” OR “Pteronetta”
 1157 OR “Sarkidiornis” OR “Sibirionetta” OR “Somateria” OR “Spatula” OR “Tachyeres” OR “Tadorna” OR
 1158 “Thalassornis” OR “Anhima” OR “Chauna” OR “Anseranas” OR “Aegothales” OR “Aerodramus” OR “Aero-
 1159 nautes” OR “Apus” OR “Chaetura” OR “Collocalia” OR “Cypseloides” OR “Cypsiurus” OR “Hirundapus”
 1160 OR “Panyptila” OR “Streptoprocne” OR “Tachornis” OR “Tachymarptis” OR “Hemiprocne” OR “Abeillia”
 1161 OR “Adelomyia” OR “Aglaeactis” OR “Aglaiocercus” OR “Amazilia” OR “Androdon” OR “Anopetia” OR
 1162 “Anthocephala” OR “Anthracothorax” OR “Archilochus” OR “Atthis” OR “Augastes” OR “Avocettula” OR
 1163 “Basilinna” OR “Boissonneaua” OR “Calliphlox” OR “Calypte” OR “Campylopterus” OR “Chaetocercus” OR
 1164 “Chalcostigma” OR “Chalybura” OR “Chlorestes” OR “Chlorostilbon” OR “Chrysolampis” OR “Chrysuronia”
 1165 OR “Clytolaema” OR “Coeligena” OR “Colibri” OR “Cyanophaea” OR “Cynanthus” OR “Discosura” OR
 1166 “Doricha” OR “Doryfera” OR “Elvira” OR “Ensifera” OR “Eriocnemis” OR “Eugenes” OR “Eulampis” OR
 1167 “Eulidia” OR “Eupetomena” OR “Eupherusa” OR “Eutoxeres” OR “Florisuga” OR “Glaucis” OR “Goethalsia”
 1168 OR “Goldmania” OR “Haplophaedia” OR “Heliactin” OR “Heliangelus” OR “Heliodoxa” OR “Heliomaster”
 1169 OR “Heliothryx” OR “Hylocharis” OR “Juliamyia” OR “Klais” OR “Lafresnaya” OR “Lampornis” OR
 1170 “Lamprolaima” OR “Lepidopyga” OR “Lesbia” OR “Leucippus” OR “Leucochloris” OR “Lophornis” OR
 1171 “Mellisuga” OR “Metallura” OR “Microchera” OR “Myrmia” OR “Myrtis” OR “Ocreatus” OR “Opistho-
 1172 prora” OR “Oreonympha” OR “Oreotrochilus” OR “Orthorhynchus” OR “Oxypogon” OR “Panterpe” OR
 1173 “Patagona” OR “Phaeochroa” OR “Phaethornis” OR “Phlogophilus” OR “Polyonymus” OR “Polytmus” OR
 1174 “Pterophanes” OR “Ramphodon” OR “Ramphomicron” OR “Rhodopis” OR “Sappho” OR “Schistes” OR
 1175 “Selasphorus” OR “Sephanoides” OR “Stephanoxis” OR “Sternoclyta” OR “Taphrosphilus” OR “Thalurania”

1176 OR "Thaumastura" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR
1177 "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR
1178 "Rhabdotorrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus"
1179 OR "Upupa" OR "Antrostomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus"
1180 OR "Hydropsalis" OR "Lurocalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus"
1181 OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius"
1182 OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariama" OR "Casuarius" OR "Dromaius" OR
1183 "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cerorhinca" OR "Fratricula" OR
1184 "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus"
1185 OR "Charadrius" OR "Elseyornis" OR "Erythrogonys" OR "Hoploxypterus" OR "Oreopholus" OR "Pel-
1186 tohyas" OR "Phegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR
1187 "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR
1188 "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR
1189 "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelocheleon" OR "Gygis" OR "Hydrocoloeus" OR
1190 "Hydroprogne" OR "Ichthyaeus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR
1191 "Pagophila" OR "Phaetusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula"
1192 OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR
1193 "Recurvirostra" OR "Nycticryphes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR
1194 "Calidris" OR "Coenocorypha" OR "Gallinago" OR "Limnodromus" OR "Limosa" OR "Lymnocyptes" OR
1195 "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagis"
1196 OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR
1197 "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chal-
1198 cophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula"
1199 OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR
1200 "Gymnophaps" OR "Hemiphaga" OR "Henicophaps" OR "Leptotila" OR "Leptotrygon" OR "Leucosarcia"
1201 OR "Lopholaimus" OR "Macropygia" OR "Metriopelia" OR "Ocyphaps" OR "Oena" OR "Otidiphaps"
1202 OR "Patagioenas" OR "Petrophassa" OR "Pezophaps" OR "Phapitreron" OR "Phaps" OR "Ptilinopus"
1203 OR "Raphus" OR "Reinwardtoena" OR "Spilopelia" OR "Streptopelia" OR "Treron" OR "Trugon" OR
1204 "Turacoena" OR "Turtur" OR "Uropelia" OR "Zenaida" OR "Zentrygon" OR "Actenoides" OR "Alcedo" OR
1205 "Caridonax" OR "Ceryle" OR "Ceyx" OR "Chloroceryle" OR "Cittura" OR "Corythornis" OR "Dacelo" OR
1206 "Halcyon" OR "Ispidina" OR "Lacedo" OR "Megaceryle" OR "Melidora" OR "Syma" OR "Tanyptera" OR
1207 "Todiramphus" OR "Atelornis" OR "Brachypteracias" OR "Geobiastes" OR "Coracias" OR "Eurystomus"
1208 OR "Merops" OR "Nyctyornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes"
1209 OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR
1210 "Chrysococcyx" OR "Clamator" OR "Coccyua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus"

1211 OR "Dasylophus" OR "Dromococcyx" OR "Eudynamys" OR "Geococcyx" OR "Guira" OR "Hierococcyx"
1212 OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicopterus" OR "Piaya" OR "Rhinortha"
1213 OR "Scythrops" OR "Surniculus" OR "Tapera" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR
1214 "Anomalopteryx" OR "Emeus" OR "Euryapteryx" OR "Eurypyga" OR "Rhynochetos" OR "Caracara" OR
1215 "Daptrius" OR "Falco" OR "Herpetotheres" OR "Ibycter" OR "Micrastur" OR "Microhierax" OR "Milvago"
1216 OR "Phalcoboenus" OR "Polihierax" OR "Spiziapteryx" OR "Aburria" OR "Chamaepetes" OR "Crax" OR
1217 "Mitu" OR "Nothocrax" OR "Oreophasis" OR "Ortalis" OR "Pauxi" OR "Penelope" OR "Penelopina" OR
1218 "Pipile" OR "Alectura" OR "Megapodius" OR "Acryllium" OR "Guttera" OR "Numida" OR "Callipepla"
1219 OR "Colinus" OR "Cyrtonyx" OR "Dendrortyx" OR "Odontophorus" OR "Oreortyx" OR "Philortyx" OR
1220 "Ptilopachus" OR "Rhynchortyx" OR "Alectoris" OR "Ammoperdix" OR "Arborophila" OR "Argusianus"
1221 OR "Bambusicola" OR "Bonasa" OR "Caloperdix" OR "Centrocercus" OR "Chrysolophus" OR "Coturnix"
1222 OR "Crossoptilon" OR "Dendragapus" OR "Excalfactoria" OR "Falcipennis" OR "Francolinus" OR "Gallus"
1223 OR "Haematortyx" OR "Ithaginis" OR "Lagopus" OR "Lerwa" OR "Lophophorus" OR "Lophura" OR
1224 "Lyrurus" OR "Meleagris" OR "Pavo" OR "Peliperdix" OR "Perdix" OR "Phasianus" OR "Polyplectron" OR
1225 "Pternistis" OR "Pucrasia" OR "Rhizothera" OR "Rollulus" OR "Scleroptila" OR "Syrmaticus" OR "Tetrao"
1226 OR "Tetraogallus" OR "Tetraophasis" OR "Tragopan" OR "Tympanuchus" OR "Gavia" OR "Aramus"
1227 OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR
1228 "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amauornis" OR "Anurolimnas"
1229 OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR
1230 "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia"
1231 OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrio" OR "Porphyriops"
1232 OR "Porzana" OR "Rallacula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura"
1233 OR "Leptosomus" OR "Corythaeola" OR "Corythaixoides" OR "Crimifer" OR "Musophaga" OR "Tauraco"
1234 OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax"
1235 OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR
1236 "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis"
1237 OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis"
1238 OR "Smicrornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna"
1239 OR "Nesillas" OR "Aegithalos" OR "Leptopoecile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR
1240 "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes"
1241 OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafraga" OR
1242 "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR
1243 "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhynchophanes" OR "Callaeas" OR "Heteralocha" OR
1244 "Philesturnus" OR "Calyptophilus" OR "Campephaga" OR "Ceblepyris" OR "Coracina" OR "Edolisoma" OR
1245 "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amaurospiza" OR "Cardinalis" OR "Ceryle" OR

1246 "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Passerina" OR
 1247 "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus"
 1248 OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus"
 1249 OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR
 1250 "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia"
 1251 OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR
 1252 "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR
 1253 "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR
 1254 "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus"
 1255 OR "Pyrrhonorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus"
 1256 OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR
 1257 "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias"
 1258 OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR
 1259 "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza"
 1260 OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR
 1261 "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis"
 1262 OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga"
 1263 OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calypomena" OR "Cymbirhynchus" OR
 1264 "Eurylaimus" OR "Psarisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR
 1265 "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chlo-
 1266 ris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra"
 1267 OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorhous" OR "Hemignathus" OR
 1268 "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR
 1269 "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR
 1270 "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza"
 1271 OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR
 1272 "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus"
 1273 OR "Certhiasomus" OR "Certhiaxis" OR "Cinclodes" OR "Clibanornis" OR "Coryphistera" OR "Cranioleuca"
 1274 OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR
 1275 "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geocerthia" OR "Geositta" OR "Glyphorhynchus"
 1276 OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR
 1277 "Limnortites" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR
 1278 "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor"
 1279 OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR
 1280 "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus"

1281 OR "Spartonoica" OR "Sylviorthorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonomus" OR
 1282 "Thripadectes" OR "Thripophaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes"
 1283 OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hylopezus" OR "Myrmothera" OR "Alopocheli-
 1284 don" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon"
 1285 OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psolidoprocne"
 1286 OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR
 1287 "Tachycineta" OR "Hylia" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Am-
 1288 blycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives"
 1289 OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus"
 1290 OR "Lamprospiza" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR
 1291 "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar"
 1292 OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura"
 1293 OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus"
 1294 OR "Leiothrix" OR "Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochalopteron" OR "Turdoides" OR
 1295 "Bradypterus" OR "Cincloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR
 1296 "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR
 1297 "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malaconotus" OR "Nilaus" OR "Rhodophoneus" OR
 1298 "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR
 1299 "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia"
 1300 OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomy-
 1301 zon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes"
 1302 OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis"
 1303 OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius"
 1304 OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cinclocerthia" OR "Dumetella" OR
 1305 "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR
 1306 "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis"
 1307 OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus"
 1308 OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR
 1309 "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Campicoloides" OR "Cercotrichas" OR "Chamaetylas"
 1310 OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR
 1311 "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia"
 1312 OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiomela" OR "Myioparus" OR "Myophonus"
 1313 OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola"
 1314 OR "Sheppardia" OR "Sholicola" OR "Stiphornis" OR "Tarsiger" OR "Thamnolaea" OR "Vauriella" OR
 1315 "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR

1316 “Cyanomitra” OR “Deleornis” OR “Hedydipna” OR “Kurochkinogramma” OR “Leptocoma” OR “Nectarinia”
 1317 OR “Daphoenositta” OR “Nesospingus” OR “Nicator” OR “Notiomystis” OR “Aleadryas” OR “Oriolus” OR
 1318 “Pitohui” OR “Sphecotheres” OR “Turnagra” OR “Colluricincla” OR “Falcunculus” OR “Melanorectes” OR
 1319 “Pachycephala” OR “Pseudorectes” OR “Panurus” OR “Cicinnurus” OR “Diphyllodes” OR “Manucodia” OR
 1320 “Paradisaea” OR “Ptiloris” OR “Paramythia” OR “Pardalotus” OR “Baeolophus” OR “Cephalopyrus” OR
 1321 “Cyanistes” OR “Lophophanes” OR “Machlolophus” OR “Melaniparus” OR “Melanochlora” OR “Pardali-
 1322 parus” OR “Parus” OR “Periparus” OR “Poecile” OR “Pseudopodoces” OR “Sittiparus” OR “Sylviparus”
 1323 OR “Basileuterus” OR “Cardellina” OR “Catharopeza” OR “Dendroica” OR “Geothlypis” OR “Helmitheros”
 1324 OR “Leiothlypis” OR “Limnethlypis” OR “Mniotilta” OR “Myioborus” OR “Myiothlypis” OR “Oporornis”
 1325 OR “Oreothlypis” OR “Parkesia” OR “Protonotaria” OR “Seiurus” OR “Setophaga” OR “Vermivora” OR
 1326 “Aimophila” OR “Ammodramus” OR “Amphispiza” OR “Arremon” OR “Arremonops” OR “Artemisiospiza”
 1327 OR “Atlapetes” OR “Calamospiza” OR “Chlorospingus” OR “Chondestes” OR “Junco” OR “Melospiza”
 1328 OR “Melozone” OR “Oreothraupis” OR “Oriturus” OR “Passerculus” OR “Passerella” OR “Peucaea” OR
 1329 “Pezopetes” OR “Pipilo” OR “Pooecetes” OR “Pselliophorus” OR “Rhynchospiza” OR “Spizella” OR “Spizel-
 1330 loides” OR “Xenospiza” OR “Zonotrichia” OR “Carpospiza” OR “Gymnoris” OR “Hypocryptadius” OR
 1331 “Montifringilla” OR “Onychostruthus” OR “Passer” OR “Petronia” OR “Pyrgilauda” OR “Alcippe” OR
 1332 “Gampsorhynchus” OR “Illadopsis” OR “Jabouilleia” OR “Kenopia” OR “Laticilla” OR “Malacocincla” OR
 1333 “Malacopteron” OR “Napothera” OR “Pellorneum” OR “Rimator” OR “Trichastoma” OR “Amalocichla”
 1334 OR “Drymodes” OR “Eopsaltria” OR “Eugerygone” OR “Heteromyias” OR “Melanodryas” OR “Microeca”
 1335 OR “Monachella” OR “Pachycephalopsis” OR “Peneoenanthe” OR “Peneothello” OR “Petroica” OR “Poe-
 1336 cilodryas” OR “Tregellasia” OR “Peucedramus” OR “Microligea” OR “Phaenicophilus” OR “Xenoligea” OR
 1337 “Phylloscopus” OR “Picathartes” OR “Antilophia” OR “Ceratopipra” OR “Chiroxiphia” OR “Chloropipo” OR
 1338 “Corapipo” OR “Cryptopipo” OR “Heterocercus” OR “Ilicura” OR “Lepidothrix” OR “Machaeropterus” OR
 1339 “Manacus” OR “Masius” OR “Neopelma” OR “Pipra” OR “Pseudopipra” OR “Tyranneutes” OR “Xenopipo”
 1340 OR “Erythropitta” OR “Hydrornis” OR “Pitta” OR “Batis” OR “Platysteira” OR “Anaplectes” OR “Bubalor-
 1341 nis” OR “Euplectes” OR “Malimbus” OR “Plocepasser” OR “Ploceus” OR “Quelea” OR “Sporopipes” OR
 1342 “Pnoepyga” OR “Microbates” OR “Polioptila” OR “Ramphocaenus” OR “Garritornis” OR “Prunella” OR
 1343 “Cinclosoma” OR “Ptilorrhoa” OR “Phainopepla” OR “Phainoptila” OR “Ptiliogonys” OR “Ailuroedus” OR
 1344 “Amblyornis” OR “Ptilonorhynchus” OR “Sericulus” OR “Alophoixus” OR “Andropadus” OR “Arizelocichla”
 1345 OR “Atimastillas” OR “Baeopogon” OR “Bleda” OR “Chlorocichla” OR “Criniger” OR “Eurillas” OR
 1346 “Hemixos” OR “Hypsipetes” OR “Iole” OR “Ixos” OR “Neolestes” OR “Phyllastrephus” OR “Pycnonotus”
 1347 OR “Spizixos” OR “Stelgidillas” OR “Tricholestes” OR “Regulus” OR “Anthoscopus” OR “Auriparus”
 1348 OR “Remiz” OR “Rhagologus” OR “Eleoscytalopus” OR “Liosceles” OR “Psilorhamphus” OR “Pterop-
 1349 tochos” OR “Scelorchilus” OR “Scytalopus” OR “Teledromas” OR “Rhipidura” OR “Rhodinocichla” OR
 1350 “Sapayoa” OR “Sitta” OR “Spindalis” OR “Chelidorhynx” OR “Culicicapa” OR “Elminia” OR “Acridotheres”

1351 OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora"
 1352 OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotornis" OR "Leucopsar"
 1353 OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR
 1354 "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR
 1355 "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR
 1356 "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR
 1357 "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprosi-
 1358 tornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR
 1359 "Drymophila" OR "Dysithamnus" OR "Epinecrophylla" OR "Euchrepomis" OR "Formicivora" OR "Freder-
 1360 ickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR
 1361 "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus"
 1362 OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myr-
 1363 moderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes"
 1364 OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania"
 1365 OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus"
 1366 OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR
 1367 "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis"
 1368 OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Charitospiza" OR "Chlorochrysa" OR
 1369 "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba"
 1370 OR "Compsospiza" OR "Compsothraupis" OR "Conirostrum" OR "Conothraupis" OR "Coryphospingus"
 1371 OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis"
 1372 OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR
 1373 "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR
 1374 "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Incaspiza" OR "Iridophanes" OR "Iridosornis"
 1375 OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR
 1376 "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes"
 1377 OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocomma" OR
 1378 "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha"
 1379 OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR
 1380 "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR
 1381 "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatorhinus" OR "Spelaeornis"
 1382 OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius"
 1383 OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR
 1384 "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia"
 1385 OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR

1386 “Pheugopedius” OR “Salpinctes” OR “Thryomanes” OR “Thryophilus” OR “Thryothorus” OR “Trogodytes”
 1387 OR “Uropsila” OR “Catharus” OR “Cichlopsis” OR “Entomodestes” OR “Geokichla” OR “Hylocichla” OR
 1388 “Ixoreus” OR “Myadestes” OR “Neocossyphus” OR “Ridgwayia” OR “Sialia” OR “Stizorhina” OR “Turdus”
 1389 OR “Zoothera” OR “Agriornis” OR “Alectrurus” OR “Anairetes” OR “Aphanotriccus” OR “Arundinicola”
 1390 OR “Atalotriccus” OR “Attila” OR “Camptostoma” OR “Casiornis” OR “Cnemarchus” OR
 1391 “Cnemotriccus” OR “Cnipodectes” OR “Colonia” OR “Colorhamphus” OR “Conopias” OR “Contopus” OR
 1392 “Corythopsis” OR “Culicivora” OR “Deltarhynchus” OR “Elaenia” OR “Empidonax” OR “Empidonomus” OR
 1393 “Euscarthmus” OR “Fluvicola” OR “Griseotyrannus” OR “Gubernetes” OR “Hemitriccus” OR “Heteroxolmis”
 1394 OR “Hirundinea” OR “Hymenops” OR “Inezia” OR “Knipolegus” OR “Lathrotriccus” OR “Legatus” OR
 1395 “Leptopogon” OR “Lessonia” OR “Lophotriccus” OR “Machetornis” OR “Mecocerculus” OR “Megarynchus”
 1396 OR “Mionectes” OR “Mitrephanes” OR “Muscigralla” OR “Muscisaxicola” OR “Myiarchus” OR “Myio-
 1397 dynastes” OR “Myiopagis” OR “Myiophobus” OR “Myiornis” OR “Myiotheretes” OR “Myiotriccus” OR
 1398 “Myiozetetes” OR “Neopipo” OR “Neoxolmis” OR “Nephelomyias” OR “Ochthoeca” OR “Ochthornis” OR
 1399 “Oncostoma” OR “Ornithion” OR “Phaeomyias” OR “Philohydor” OR “Phyllomyias” OR “Phylloscartes”
 1400 OR “Piprites” OR “Pitangus” OR “Platyrinchus” OR “Poecilotriccus” OR “Pogonotriccus” OR “Polioxolmis”
 1401 OR “Polystictus” OR “Pseudelaenia” OR “Pseudocolopteryx” OR “Pseudotriccus” OR “Pyrocephalus” OR
 1402 “Pyrrhomyias” OR “Ramphotrigon” OR “Rhynchocyclus” OR “Rhytipterna” OR “Satrapa” OR “Sayornis”
 1403 OR “Serpophaga” OR “Silvicultrix” OR “Sirystes” OR “Stigmatura” OR “Sublegatus” OR “Suiriri” OR
 1404 “Tachuris” OR “Taeniotriccus” OR “Todiostrostrum” OR “Tolmomyias” OR “Tumbezia” OR “Tyrannopsis”
 1405 OR “Tyrannulus” OR “Tyrannus” OR “Uromyias” OR “Xenotriccus” OR “Xolmis” OR “Zimmerius” OR
 1406 “Urocynchramus” OR “Artamella” OR “Calicalicus” OR “Cyanolanius” OR “Euryceros” OR “Falculea”
 1407 OR “Hypositta” OR “Leptopterus” OR “Mystacornis” OR “Newtonia” OR “Oriolia” OR “Philentoma” OR
 1408 “Prionops” OR “Pseudobias” OR “Schetba” OR “Tephrodornis” OR “Tylas” OR “Vanga” OR “Xenopirostris”
 1409 OR “Anomalospiza” OR “Vidua” OR “Cyclarhis” OR “Erpornis” OR “Hylophilus” OR “Pteruthius” OR
 1410 “Vireo” OR “Vireolanius” OR “Zeledonia” OR “Apalopteron” OR “Dasycrotapha” OR “Heleia” OR “Lopho-
 1411 zosterops” OR “Sternhoptilus” OR “Yuhina” OR “Zosterops” OR “Zosterornis” OR “Agamia” OR “Ardea” OR
 1412 “Ardeola” OR “Botaurus” OR “Bubulcus” OR “Butorides” OR “Cochlearius” OR “Dupetor” OR “Egretta”
 1413 OR “Gorsachius” OR “Ixobrychus” OR “Nyctanassa” OR “Nycticorax” OR “Ptilherodius” OR “Syrigma” OR
 1414 “Tigriornis” OR “Tigrisoma” OR “Zebrius” OR “Balaeniceps” OR “Pelecanus” OR “Scopus” OR “Bostrychia”
 1415 OR “Eudocimus” OR “Geronticus” OR “Lophotibis” OR “Mesembrinibis” OR “Nipponia” OR “Phimosus”
 1416 OR “Platalea” OR “Plegadis” OR “Pseudibis” OR “Theristicus” OR “Threskiornis” OR “Phaethon” OR
 1417 “Phoeniconaias” OR “Phoenicoparrus” OR “Phoenicopterus” OR “Bucco” OR “Chelidoptera” OR “Hapalop-
 1418 tila” OR “Hypnelus” OR “Malacoptila” OR “Micromonacha” OR “Monasa” OR “Nonnula” OR “Notharchus”
 1419 OR “Nystalus” OR “Capito” OR “Eubucco” OR “Brachygalba” OR “Galbalcyrrhynchus” OR “Galbula” OR
 1420 “Jacamaralcyon” OR “Jacamerops” OR “Indicator” OR “Melichneutes” OR “Malignomon” OR “Prodotiscus”

1421 OR "Buccanodon" OR "Gymnobucco" OR "Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus"
1422 OR "Psilopogon" OR "Blythipicus" OR "Campephilus" OR "Campethera" OR "Celeus" OR "Chloropicus"
1423 OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Dendrocopos" OR "Dendrocoptes" OR
1424 "Dendropicos" OR "Dinopium" OR "Dryobates" OR "Dryocopus" OR "Gecinulus" OR "Geocolaptes" OR
1425 "Jynx" OR "Leiopicus" OR "Leuconotopicus" OR "Meiglyptes" OR "Melanerpes" OR "Micropternus" OR
1426 "Mulleripicus" OR "Nesocittes" OR "Picoides" OR "Piculus" OR "Picumnus" OR "Picus" OR "Reinwardtipi-
1427 cus" OR "Sasia" OR "Sphyrapicus" OR "Veniliornis" OR "Yungipicus" OR "Andigena" OR "Aulacorhynchus"
1428 OR "Pteroglossus" OR "Ramphastos" OR "Selenidera" OR "Semnornis" OR "Aechmophorus" OR "Podiceps"
1429 OR "Podilymbus" OR "Poliocephalus" OR "Rollandia" OR "Tachybaptus" OR "Diomedea" OR "Phoebastria"
1430 OR "Phoebetria" OR "Thalassarche" OR "Hydrobates" OR "Oceanodroma" OR "Fregetta" OR "Garrodia"
1431 OR "Nesofregetta" OR "Oceanites" OR "Pelagodroma" OR "Aphrodroma" OR "Ardenna" OR "Bulweria"
1432 OR "Calonectris" OR "Daption" OR "Fulmarus" OR "Halobaena" OR "Macronectes" OR "Pachyptila" OR
1433 "Pagodroma" OR "Pelecanoides" OR "Procellaria" OR "Pseudobulweria" OR "Pterodroma" OR "Puffinus" OR
1434 "Thalassoica" OR "Cacatua" OR "Callocephalon" OR "Calyptorhynchus" OR "Eolophus" OR "Lophochroa"
1435 OR "Nymphicus" OR "Probosciger" OR "Alipiopsitta" OR "Amazona" OR "Anodorhynchus" OR "Ara" OR
1436 "Aratinga" OR "Bolborhynchus" OR "Brotogeris" OR "Conuropsis" OR "Cyanoliseus" OR "Cyanopsitta" OR
1437 "Derophtus" OR "Diopsittaca" OR "Enicognathus" OR "Eupsittula" OR "Forpus" OR "Graydidascalus" OR
1438 "Guarouba" OR "Guaruba" OR "Hapalopsittaca" OR "Leptosittaca" OR "Myiopsitta" OR "Nannopsittaca"
1439 OR "Orthopsittaca" OR "Pionites" OR "Pionopsitta" OR "Pionus" OR "Poicephalus" OR "Primolius"
1440 OR "Psilopsiagon" OR "Psittacara" OR "Psittacus" OR "Pyrilia" OR "Pyrrhura" OR "Rhynchopsitta"
1441 OR "Thectocercus" OR "Touit" OR "Triclaria" OR "Agapornis" OR "Alisterus" OR "Aprosmictus" OR
1442 "Barnardius" OR "Bolbopsittacus" OR "Chalcopsitta" OR "Charmosyna" OR "Coracopsis" OR "Cyanoram-
1443 phus" OR "Cyclopsitta" OR "Eclectus" OR "Eos" OR "Eunymphicus" OR "Geoffroyus" OR "Glossopsitta"
1444 OR "Lathamus" OR "Loriculus" OR "Lorius" OR "Melopsittacus" OR "Micropsitta" OR "Neophema" OR
1445 "Neopsephotus" OR "Neopsittacus" OR "Northiella" OR "Oreopsittacus" OR "Parvipsitta" OR "Pezoporus"
1446 OR "Phigys" OR "Platycercus" OR "Polytelis" OR "Prioniturus" OR "Prosopiea" OR "Psephotellus" OR
1447 "Psephotus" OR "Pseudeos" OR "Psittacella" OR "Psittacula" OR "Psittaculirostris" OR "Psitteuteles" OR
1448 "Psittinus" OR "Psittichas" OR "Purpureicephalus" OR "Tanygnathus" OR "Trichoglossus" OR "Vini" OR
1449 "Nestor" OR "Strigops" OR "Pterocles" OR "Syrrhaptes" OR "Rhea" OR "Aptenodytes" OR "Eudiptes" OR
1450 "Eudiptula" OR "Megadyptes" OR "Pygoscelis" OR "Spheniscus" OR "Aegolius" OR "Asio" OR "Athene"
1451 OR "Bubo" OR "Glaucidium" OR "Ketupa" OR "Lophotrix" OR "Megascops" OR "Micrathene" OR "Ninox"
1452 OR "Otus" OR "Pseudoscops" OR "Psiloscops" OR "Pulsatrix" OR "Sceloglaux" OR "Scotopelia" OR "Strix"
1453 OR "Surnia" OR "Xenoglaux" OR "Phodilus" OR "Tyto" OR "Aepyornis" OR "Mullerornis" OR "Struthio"
1454 OR "Anhinga" OR "Fregata" OR "Leucocarbo" OR "Microcarbo" OR "Nannopterum" OR "Phalacrocorax"
1455 OR "Morus" OR "Sula" OR "Crypturellus" OR "Eudromia" OR "Nothocercus" OR "Nothoprocta" OR

1456 “Nothura” OR “Rhynchotus” OR “Tinamotis” OR “Tinamus” OR “Apalharpactes” OR “Apaloderma” OR
 1457 “Harpactes” OR “Pharomachrus” OR “Trogon” OR “Tetrastes” OR “Hesperiphona”) AND (“experiment*”
 1458 OR “manipulat*”))) OR AK= (((“green*” OR “herb*” OR “aromatic*”) AND “nest*” AND (“bird*” OR
 1459 “aves” OR “avian” OR “ornithol*” OR “passerine*” OR “passeriform*” OR “songbird*” OR “Accipiter” OR
 1460 “Aegypius” OR “Aquila” OR “Aviceda” OR “Busarellus” OR “Butastur” OR “Buteo” OR “Buteogallus” OR
 1461 “Chondrohierax” OR “Circaetus” OR “Circus” OR “Clanga” OR “Elanoides” OR “Elanus” OR “Gampsonyx”
 1462 OR “Geranoaetus” OR “Geranospiza” OR “Gypaetus” OR “Gypohierax” OR “Gyps” OR “Haliaeetus” OR
 1463 “Haliastur” OR “Harpagus” OR “Harpia” OR “Hieraaetus” OR “Ictinaetus” OR “Ictinia” OR “Kaupifalco”
 1464 OR “Leptodon” OR “Leucopternis” OR “Lophotriorchis” OR “Milvus” OR “Morphnarchus” OR “Morphnus”
 1465 OR “Necrosyrtes” OR “Neophron” OR “Nisaetus” OR “Parabuteo” OR “Pernis” OR “Pithecophaga” OR
 1466 “Polemaetus” OR “Pseudastur” OR “Rostrhamus” OR “Rupornis” OR “Sarcogyps” OR “Spilornis” OR
 1467 “Spizaetus” OR “Stephanoaetus” OR “Terathopius” OR “Torgos” OR “Trigonoceps” OR “Cathartes” OR
 1468 “Coragyps” OR “Gymnogyps” OR “Sarcoramphus” OR “Vultur” OR “Pandion” OR “Sagittarius” OR “Aix”
 1469 OR “Alopochen” OR “Amazonetta” OR “Anas” OR “Anser” OR “Asarcornis” OR “Aythya” OR “Branta” OR
 1470 “Bucephala” OR “Cairina” OR “Callonetta” OR “Cereopsis” OR “Chen” OR “Chenonetta” OR “Chloephaga”
 1471 OR “Clangula” OR “Coscoroba” OR “Cyanochen” OR “Cygnus” OR “Dendrocygna” OR “Heteronetta” OR
 1472 “Histrionicus” OR “Hymenolaimus” OR “Lophodytes” OR “Lophonetta” OR “Malacorhynchus” OR “Mareca”
 1473 OR “Marmaronetta” OR “Melanitta” OR “Merganetta” OR “Mergellus” OR “Mergus” OR “Neochen” OR
 1474 “Netta” OR “Nettapus” OR “Nomonyx” OR “Oxyura” OR “Plectropterus” OR “Polysticta” OR “Pteronetta”
 1475 OR “Sarkidiornis” OR “Sibirionetta” OR “Somateria” OR “Spatula” OR “Tachyeres” OR “Tadorna” OR
 1476 “Thalassornis” OR “Anhima” OR “Chauna” OR “Anseranas” OR “Aegotheles” OR “Aerodramus” OR “Aero-
 1477 nautes” OR “Apus” OR “Chaetura” OR “Collocalia” OR “Cypseloides” OR “Cypsiurus” OR “Hirundapus”
 1478 OR “Panyptila” OR “Streptoprocne” OR “Tachornis” OR “Tachymarptis” OR “Hemiprocne” OR “Abeillia”
 1479 OR “Adelomyia” OR “Aglaeactis” OR “Aglaiocercus” OR “Amazilia” OR “Androdon” OR “Anopetia” OR
 1480 “Anthocephala” OR “Anthracothorax” OR “Archilochus” OR “Atthis” OR “Augustes” OR “Avocettula” OR
 1481 “Basilinna” OR “Boissonneaua” OR “Calliphlox” OR “Calypte” OR “Campylopterus” OR “Chaetocercus” OR
 1482 “Chalcostigma” OR “Chalybura” OR “Chlorestes” OR “Chlorostilbon” OR “Chrysolampis” OR “Chrysuronia”
 1483 OR “Clytolaema” OR “Coeligena” OR “Colibri” OR “Cyanophaia” OR “Cynanthus” OR “Discosura” OR
 1484 “Doricha” OR “Doryfera” OR “Elvira” OR “Ensifera” OR “Eriocnemis” OR “Eugenes” OR “Eulampis” OR
 1485 “Eulidia” OR “Eupetomena” OR “Eupherusa” OR “Eutoxeres” OR “Florisuga” OR “Glaucis” OR “Goethalsia”
 1486 OR “Goldmania” OR “Haplophaedia” OR “Heliactin” OR “Heliangelus” OR “Heliodoxa” OR “Heliomaster”
 1487 OR “Heliothryx” OR “Hylocharis” OR “Juliamyia” OR “Klais” OR “Lafresnaya” OR “Lampornis” OR
 1488 “Lamprolaima” OR “Lepidopyga” OR “Lesbia” OR “Leucippus” OR “Leucochloris” OR “Lophornis” OR
 1489 “Mellisuga” OR “Metallura” OR “Microchera” OR “Myrmia” OR “Myrtis” OR “Ocreatus” OR “Opistho-
 1490 prora” OR “Oreonympha” OR “Oreotrochilus” OR “Orthorhyncus” OR “Oxypogon” OR “Panterpe” OR

1491 “Patagona” OR “Phaeochroa” OR “Phaethornis” OR “Phlogophilus” OR “Polyonymus” OR “Polytmus” OR
 1492 “Pterophanes” OR “Ramphodon” OR “Ramphomicron” OR “Rhodopis” OR “Sappho” OR “Schistes” OR
 1493 “Selasphorus” OR “Sephanoides” OR “Stephanoxis” OR “Sternoclyta” OR “Taphrospilus” OR “Thalurania”
 1494 OR “Thaumastura” OR “Threnetes” OR “Topaza” OR “Trochilus” OR “Urochroa” OR “Urosticte” OR
 1495 “Apteryx” OR “Anthracoceros” OR “Buceros” OR “Bycanistes” OR “Lophoceros” OR “Penelopides” OR
 1496 “Rhabdotorrhinus” OR “Rhyticeros” OR “Tockus” OR “Bucorvus” OR “Phoeniculus” OR “Rhinopomastus”
 1497 OR “Upupa” OR “Antrostomus” OR “Caprimulgus” OR “Chordeiles” OR “Eleothreptus” OR “Eurostopodus”
 1498 OR “Hydropsalis” OR “Lurocalis” OR “Lyncornis” OR “Nyctidromus” OR “Nyctiphrynus” OR “Nyctipolus”
 1499 OR “Nyctiprogne” OR “Phalaenoptilus” OR “Setopagis” OR “Systellura” OR “Uropsalis” OR “Nyctibius”
 1500 OR “Batrachostomus” OR “Podargus” OR “Steatornis” OR “Cariama” OR “Casuarius” OR “Dromaius” OR
 1501 “Aethia” OR “Alca” OR “Alle” OR “Brachyramphus” OR “Cepphus” OR “Cerorhinca” OR “Fratricula” OR
 1502 “Pinguinus” OR “Ptychoramphus” OR “Synthliboramphus” OR “Uria” OR “Burhinus” OR “Anarhynchus”
 1503 OR “Charadrius” OR “Elseyornis” OR “Erythrogonys” OR “Hoploxypterus” OR “Oreopholus” OR “Pel-
 1504 tohyas” OR “Phegornis” OR “Pluvialis” OR “Thinornis” OR “Vanellus” OR “Chionis” OR “Dromas” OR
 1505 “Cursorius” OR “Glareola” OR “Rhinoptilus” OR “Stiltia” OR “Haematopus” OR “Actophilornis” OR
 1506 “Hydrophasianus” OR “Irediparra” OR “Jacana” OR “Metopidius” OR “Microparra” OR “Anous” OR
 1507 “Chlidonias” OR “Chroicocephalus” OR “Creagrus” OR “Gelochelidon” OR “Gygis” OR “Hydrocoloeus” OR
 1508 “Hydroprogne” OR “Ichthyaetus” OR “Larosterna” OR “Larus” OR “Leucophaeus” OR “Onychoprion” OR
 1509 “Pagophila” OR “Phaetusa” OR “Rhodostethia” OR “Rissa” OR “Rynchops” OR “Sterna” OR “Sternula”
 1510 OR “Thalasseus” OR “Xema” OR “Pedionomus” OR “Pluvianellus” OR “Pluvianus” OR “Himantopus” OR
 1511 “Recurvirostra” OR “Nycticryphes” OR “Rostratula” OR “Actitis” OR “Arenaria” OR “Bartramia” OR
 1512 “Calidris” OR “Coenocorypha” OR “Gallinago” OR “Limnodromus” OR “Limosa” OR “Lymnocyptes” OR
 1513 “Numenius” OR “Phalaropus” OR “Scolopax” OR “Tringa” OR “Xenus” OR “Stercorarius” OR “Attagis”
 1514 OR “Thinocorus” OR “Turnix” OR “Ciconia” OR “Ephippiorhynchus” OR “Jabiru” OR “Leptoptilos” OR
 1515 “Mycteria” OR “Colius” OR “Urocolius” OR “Alectroenas” OR “Alopecoenas” OR “Caloenas” OR “Chal-
 1516 cophaps” OR “Claravis” OR “Columba” OR “Columbina” OR “Didunculus” OR “Drepanoptila” OR “Ducula”
 1517 OR “Ectopistes” OR “Gallicolumba” OR “Geopelia” OR “Geophaps” OR “Geotrygon” OR “Goura” OR
 1518 “Gymnophaps” OR “Hemiphaga” OR “Henicophaps” OR “Leptotila” OR “Leptotrygon” OR “Leucosarcia”
 1519 OR “Lopholaimus” OR “Macropygia” OR “Metriopelia” OR “Ocyphaps” OR “Oena” OR “Otidiphaps”
 1520 OR “Patagioenas” OR “Petrophassa” OR “Pezophaps” OR “Phapitreron” OR “Phaps” OR “Ptilinopus”
 1521 OR “Raphus” OR “Reinwardtoena” OR “Spilopelia” OR “Streptopelia” OR “Treron” OR “Trugon” OR
 1522 “Turacoena” OR “Turtur” OR “Uropelia” OR “Zenaida” OR “Zentrygon” OR “Actenoides” OR “Alcedo” OR
 1523 “Caridonax” OR “Ceryle” OR “Ceyx” OR “Chloroceryle” OR “Cittura” OR “Corythornis” OR “Dacelo” OR
 1524 “Halcyon” OR “Ispidina” OR “Lacedo” OR “Megaceryle” OR “Melidora” OR “Syma” OR “Tanyptera” OR
 1525 “Todiramphus” OR “Atelornis” OR “Brachypteracias” OR “Geobiastes” OR “Coracias” OR “Eurystomus”

1526 OR "Merops" OR "Nyctyornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes"
1527 OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR
1528 "Chrysococcyx" OR "Clamator" OR "Coccyua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus"
1529 OR "Dasylophus" OR "Dromococcyx" OR "Eudynamys" OR "Geococcyx" OR "Guira" OR "Hierococcyx"
1530 OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicophaeus" OR "Piaya" OR "Rhinortha"
1531 OR "Scythrops" OR "Surniculus" OR "Tapera" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR
1532 "Anomalopteryx" OR "Emeus" OR "Euryapteryx" OR "Eurypyga" OR "Rhynochetos" OR "Caracara" OR
1533 "Daptrius" OR "Falco" OR "Herpetotheres" OR "Ibycter" OR "Micrastur" OR "Microhierax" OR "Milvago"
1534 OR "Phalcoboenus" OR "Polihierax" OR "Spizapteryx" OR "Aburria" OR "Chamaepetes" OR "Crax" OR
1535 "Mitu" OR "Nothocrax" OR "Oreophasis" OR "Ortalis" OR "Pauxi" OR "Penelope" OR "Penelopina" OR
1536 "Pipile" OR "Alectura" OR "Megapodius" OR "Acryllium" OR "Guttera" OR "Numida" OR "Callipepla"
1537 OR "Colinus" OR "Cyrtonyx" OR "Dendrortyx" OR "Odontophorus" OR "Oreortyx" OR "Philortyx" OR
1538 "Ptilopachus" OR "Rhynchortyx" OR "Alectoris" OR "Ammoperdix" OR "Arborophila" OR "Argusianus"
1539 OR "Bambusicola" OR "Bonasa" OR "Caloperdix" OR "Centrocercus" OR "Chrysolophus" OR "Coturnix"
1540 OR "Crossoptilon" OR "Dendragapus" OR "Excalfactoria" OR "Falcipennis" OR "Francolinus" OR "Gallus"
1541 OR "Haematortyx" OR "Ithaginis" OR "Lagopus" OR "Lerwa" OR "Lophophorus" OR "Lophura" OR
1542 "Lyrurus" OR "Meleagris" OR "Pavo" OR "Peliperdix" OR "Perdix" OR "Phasianus" OR "Polyplectron" OR
1543 "Pternistis" OR "Pucrasia" OR "Rhizothera" OR "Rollulus" OR "Scleroptila" OR "Syrmaticus" OR "Tetrao"
1544 OR "Tetraogallus" OR "Tetraophasis" OR "Tragopan" OR "Tympanuchus" OR "Gavia" OR "Aramus"
1545 OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR
1546 "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amauornis" OR "Anurolimnas"
1547 OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR
1548 "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia"
1549 OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrio" OR "Porphyriops"
1550 OR "Porzana" OR "Rallacula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura"
1551 OR "Leptosomus" OR "Corythaeola" OR "Corythaixoides" OR "Crinifer" OR "Musophaga" OR "Tauraco"
1552 OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax"
1553 OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR
1554 "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis"
1555 OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis"
1556 OR "Smicrornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna"
1557 OR "Nesillas" OR "Aegithalos" OR "Leptopoeile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR
1558 "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes"
1559 OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafraga" OR
1560 "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR

1561 “Buphagus” OR “Calcarius” OR “Plectrophenax” OR “Rhynchophanes” OR “Callaeas” OR “Heteralocha” OR
 1562 “Philesturnus” OR “Calyptophilus” OR “Campephaga” OR “Ceblepyris” OR “Coracina” OR “Edolisoma” OR
 1563 “Lalage” OR “Malindangia” OR “Pericrocotus” OR “Amaurospiza” OR “Cardinalis” OR “Caryothraustes” OR
 1564 “Chlorothraupis” OR “Cyanocompsa” OR “Cyanoloxia” OR “Granatellus” OR “Habia” OR “Passerina” OR
 1565 “Periporphyrus” OR “Pheucticus” OR “Piranga” OR “Spiza” OR “Certhia” OR “Salpornis” OR “Abroscopus”
 1566 OR “Cettia” OR “Horornis” OR “Phyllergates” OR “Tesia” OR “Urosphena” OR “Chloropsis” OR “Cinclus”
 1567 OR “Apalis” OR “Artisornis” OR “Camaroptera” OR “Cisticola” OR “Eremomela” OR “Heliolais” OR
 1568 “Hypergerus” OR “Neomixis” OR “Oreolais” OR “Orthotomus” OR “Phragmacia” OR “Poliolais” OR “Prinia”
 1569 OR “Schistolais” OR “Urolais” OR “Urorhipis” OR “Climacteris” OR “Cormobates” OR “Conopophaga” OR
 1570 “Struthidea” OR “Aphelocoma” OR “Calocitta” OR “Cissa” OR “Coloeus” OR “Corvus” OR “Cyanocitta” OR
 1571 “Cyanocorax” OR “Cyanolyca” OR “Cyanopica” OR “Dendrocitta” OR “Garrulus” OR “Gymnorhinus” OR
 1572 “Nucifraga” OR “Perisoreus” OR “Pica” OR “Platylophus” OR “Podoces” OR “Psilorhinus” OR “Ptilostomus”
 1573 OR “Pyrrhocorax” OR “Urocissa” OR “Ampelioides” OR “Ampelion” OR “Carpornis” OR “Cephalopterus”
 1574 OR “Conioptilon” OR “Cotinga” OR “Doliornis” OR “Gymnoderus” OR “Haematoderus” OR “Lipaugus” OR
 1575 “Perissocephalus” OR “Phoenicircus” OR “Phytotoma” OR “Pipreola” OR “Porphyrolaema” OR “Procnias”
 1576 OR “Pyroderus” OR “Querula” OR “Rupicola” OR “Snowornis” OR “Xipholena” OR “Zaratornis” OR
 1577 “Dasyornis” OR “Dicaeum” OR “Prionochilus” OR “Dicrurus” OR “Donacobius” OR “Dulus” OR “Emberiza”
 1578 OR “Amadina” OR “Amandava” OR “Cryptospiza” OR “Erythrura” OR “Estrilda” OR “Euodice” OR
 1579 “Euschistospiza” OR “Lagonosticta” OR “Lonchura” OR “Mandingoa” OR “Neochmia” OR “Nesocharis”
 1580 OR “Nigrita” OR “Ortygospiza” OR “Parmoptila” OR “Pyrenestes” OR “Pytilia” OR “Spermophaga”
 1581 OR “Stagonopleura” OR “Taeniopygia” OR “Uraeginthus” OR “Calyptomena” OR “Cymbirhynchus” OR
 1582 “Eurylaimus” OR “Psarisomus” OR “Serilophus” OR “Smithornis” OR “Chamaeza” OR “Formicarius” OR
 1583 “Acanthis” OR “Agraphospiza” OR “Akialoa” OR “Bucanetes” OR “Carduelis” OR “Carpodacus” OR “Chlo-
 1584 ris” OR “Chlorodrepanis” OR “Chlorophonia” OR “Chrysocorythus” OR “Coccothraustes” OR “Crithagra”
 1585 OR “Drepanis” OR “Eophona” OR “Euphonia” OR “Fringilla” OR “Haemorhous” OR “Hemignathus” OR
 1586 “Himatione” OR “Leucosticte” OR “Linaria” OR “Linurgus” OR “Loxia” OR “Loxioides” OR “Loxops” OR
 1587 “Magnumma” OR “Melamprosops” OR “Oreomystis” OR “Paroreomyza” OR “Pinicola” OR “Procarduelis” OR
 1588 “Pseudonestor” OR “Psittirostra” OR “Pyrrhoplectes” OR “Pyrrhula” OR “Rhodopechys” OR “Rhodospiza”
 1589 OR “Serinus” OR “Spinus” OR “Telespiza” OR “Anabacerthia” OR “Anabazenops” OR “Ancistrops” OR
 1590 “Anumbius” OR “Aphrastura” OR “Asthenes” OR “Automolus” OR “Berlepschia” OR “Campylorhamphus”
 1591 OR “Certhiasomus” OR “Certhiaxis” OR “Cinclodes” OR “Clibanornis” OR “Coryphistera” OR “Cranioleuca”
 1592 OR “Deconychura” OR “Dendrexetastes” OR “Dendrocincla” OR “Dendrocolaptes” OR “Dendroplex” OR
 1593 “Drymornis” OR “Drymotoxeres” OR “Furnarius” OR “Geocerthia” OR “Geositta” OR “Glyphorhynchus”
 1594 OR “Heliobletus” OR “Hellmayrea” OR “Hylexetastes” OR “Lepidocolaptes” OR “Leptasthenura” OR
 1595 “Limnortites” OR “Limnornis” OR “Lochmias” OR “Margarornis” OR “Mazaria” OR “Megaxenops” OR

1596 “Metopothrix” OR “Microxenops” OR “Nasica” OR “Ochetorhynchus” OR “Phacellodomus” OR “Philydor”
 1597 OR “Phleocryptes” OR “Premnoplex” OR “Premnornis” OR “Pseudasthenes” OR “Pseudocolaptes” OR
 1598 “Pseudoseisura” OR “Pygarrichas” OR “Roraimia” OR “Schoeniophylax” OR “Sclerurus” OR “Sittasomus”
 1599 OR “Spartonoica” OR “Sylviorthorhynchus” OR “Synallaxis” OR “Syndactyla” OR “Tarphonimus” OR
 1600 “Thripadectes” OR “Thripophaga” OR “Upucerthia” OR “Xenerpestes” OR “Xenops” OR “Xiphocolaptes”
 1601 OR “Xiphorhynchus” OR “Grallaria” OR “Grallaricula” OR “Hylopezus” OR “Myrmothera” OR “Alopocheli-
 1602 don” OR “Atticora” OR “Cecropis” OR “Delichon” OR “Haplochelidon” OR “Hirundo” OR “Neochelidon”
 1603 OR “Notiochelidon” OR “Orochelidon” OR “Petrochelidon” OR “Phedina” OR “Progne” OR “Psolidoprocne”
 1604 OR “Pseudhirundo” OR “Pseudochelidon” OR “Ptyonoprogne” OR “Riparia” OR “Stelgidopteryx” OR
 1605 “Tachycineta” OR “Hylia” OR “Hypocolius” OR “Agelaioides” OR “Agelaius” OR “Agelasticus” OR “Am-
 1606 blycercus” OR “Amblyramphus” OR “Anumara” OR “Cacicus” OR “Chrysomus” OR “Curaeus” OR “Dives”
 1607 OR “Dolichonyx” OR “Euphagus” OR “Gnorimopsar” OR “Gymnomystax” OR “Hypopyrrhus” OR “Icterus”
 1608 OR “Lamprosar” OR “Leistes” OR “Macroagelaius” OR “Molothrus” OR “Nesopsar” OR “Oreopsar” OR
 1609 “Psarocolius” OR “Pseudoleistes” OR “Quiscalus” OR “Sturnella” OR “Xanthocephalus” OR “Xanthopsar”
 1610 OR “Icteria” OR “Ifrita” OR “Irena” OR “Corvinella” OR “Eurocephalus” OR “Lanius” OR “Actinodura”
 1611 OR “Argya” OR “Garrulax” OR “Grammatoptila” OR “Heterophasia” OR “Ianthocincla” OR “Laniellus”
 1612 OR “Leiothrix” OR “Liocichla” OR “Minla” OR “Pterorhinus” OR “Trochalopecton” OR “Turdoides” OR
 1613 “Bradypterus” OR “Cincloramphus” OR “Helopsaltes” OR “Locustella” OR “Megalurus” OR “Poodytes” OR
 1614 “Machaerirhynchus” OR “Graueria” OR “Hylia” OR “Macrosphenus” OR “Melocichla” OR “Sylvietta” OR
 1615 “Chlorophoneus” OR “Dryoscopus” OR “Laniarius” OR “Malaconotus” OR “Nilaus” OR “Rhodophoneus” OR
 1616 “Tchagra” OR “Telophorus” OR “Amytornis” OR “Chenorhamphus” OR “Clytomyias” OR “Malurus” OR
 1617 “Stipiturus” OR “Melampitta” OR “Melanocharis” OR “Oedistoma” OR “Toxorhamphus” OR “Melanopareia”
 1618 OR “Acanthagenys” OR “Acanthorhynchus” OR “Anthochaera” OR “Anthornis” OR “Caligavis” OR “Entomy-
 1619 zon” OR “Epthianura” OR “Glycifolia” OR “Gymnomyza” OR “Lichmera” OR “Manorina” OR “Melidectes”
 1620 OR “Melilestes” OR “Meliphaga” OR “Melipotes” OR “Melithreptus” OR “Myzomela” OR “Nesoptilotis”
 1621 OR “Philemon” OR “Phylidonyris” OR “Prothemadera” OR “Ptiloprora” OR “Ptilotula” OR “Pycnopygius”
 1622 OR “Timeliopsis” OR “Xanthotis” OR “Menura” OR “Allenia” OR “Cinclocerthia” OR “Dumetella” OR
 1623 “Margarops” OR “Melanoptila” OR “Melanotis” OR “Mimus” OR “Oreoscoptes” OR “Ramphocinclus” OR
 1624 “Toxostoma” OR “Lamprospiza” OR “Mitrospingus” OR “Moho” OR “Mohoua” OR “Arses” OR “Carterornis”
 1625 OR “Clytorhynchus” OR “Grallina” OR “Hypothymis” OR “Monarcha” OR “Myiagra” OR “Symposiachrus”
 1626 OR “Terpsiphone” OR “Trochocercus” OR “Anthus” OR “Macronyx” OR “Motacilla” OR “Alethe” OR
 1627 “Anthipes” OR “Brachypteryx” OR “Calliope” OR “Campicoloides” OR “Cercotrichas” OR “Chamaetylas”
 1628 OR “Copsychus” OR “Cossypha” OR “Cyanoptila” OR “Cyornis” OR “Enicurus” OR “Erithacus” OR
 1629 “Eumyias” OR “Ficedula” OR “Fraseria” OR “Irania” OR “Larvivora” OR “Leonardina” OR “Luscinia”
 1630 OR “Melaenornis” OR “Monticola” OR “Muscicapa” OR “Myiomela” OR “Myioparus” OR “Myophonus”

1631 OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola"
 1632 OR "Sheppardia" OR "Sholicola" OR "Stiphrornis" OR "Tarsiger" OR "Thamnolaea" OR "Vauriella" OR
 1633 "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR
 1634 "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkinogramma" OR "Leptocoma" OR "Nectarinia"
 1635 OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleadryas" OR "Oriolus" OR
 1636 "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR
 1637 "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR
 1638 "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR
 1639 "Cyanistes" OR "Lophophanes" OR "Machlolophus" OR "Melaniparus" OR "Melanochlora" OR "Pardali-
 1640 parus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus"
 1641 OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros"
 1642 OR "Leiothlypis" OR "Limnothlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis"
 1643 OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR
 1644 "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisiospiza"
 1645 OR "Atlapetes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza"
 1646 OR "Melozone" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR
 1647 "Pezopetes" OR "Pipilo" OR "Pooecetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizel-
 1648 loides" OR "Xenospiza" OR "Zonotrichia" OR "Carpospiza" OR "Gymnoris" OR "Hypocryptadius" OR
 1649 "Montifringilla" OR "Onychostruthus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR
 1650 "Gampsorhynchus" OR "Illadopsis" OR "Jabouilleia" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR
 1651 "Malacopteron" OR "Napothea" OR "Pellorneum" OR "Rimtor" OR "Trichastoma" OR "Amalocichla"
 1652 OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca"
 1653 OR "Monachella" OR "Pachycephalopsis" OR "Peneoenanthe" OR "Peneothello" OR "Petroica" OR "Poe-
 1654 cilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR
 1655 "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR
 1656 "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR
 1657 "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyrannetes" OR "Xenopipo"
 1658 OR "Erythropitta" OR "Hydornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalor-
 1659 nis" OR "Euplectes" OR "Malimbus" OR "Plocepasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR
 1660 "Pnoepyga" OR "Microbates" OR "Polioptila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR
 1661 "Cinclosoma" OR "Ptilorrhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR
 1662 "Amblyornis" OR "Ptilonorhynchus" OR "Sericultus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla"
 1663 OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Crimiger" OR "Eurillas" OR
 1664 "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus"
 1665 OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus"

1666 OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pterop-
1667 tochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR
1668 "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynx" OR "Culicicapa" OR "Elminia" OR "Acridotheres"
1669 OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora"
1670 OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotornis" OR "Leucopsar"
1671 OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR
1672 "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR
1673 "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR
1674 "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR
1675 "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprosi-
1676 tornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR
1677 "Drymophila" OR "Dysithamnus" OR "Epinecrophylla" OR "Euchrepomis" OR "Formicivora" OR "Freder-
1678 ickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR
1679 "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus"
1680 OR "Microrhops" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myr-
1681 moderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes"
1682 OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania"
1683 OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus"
1684 OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR
1685 "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis"
1686 OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Charitospiza" OR "Chlorochrysa" OR
1687 "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba"
1688 OR "Compsospiza" OR "Compsothraupis" OR "Conirostrum" OR "Conothraupis" OR "Coryphospingus"
1689 OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis"
1690 OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR
1691 "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR
1692 "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Incasppiza" OR "Iridophanes" OR "Iridosornis"
1693 OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR
1694 "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes"
1695 OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocomma" OR
1696 "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha"
1697 OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR
1698 "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR
1699 "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatorhinus" OR "Spelaeornis"
1700 OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius"

1701 OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR
 1702 "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia"
 1703 OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR
 1704 "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Troglodytes"
 1705 OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR
 1706 "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus"
 1707 OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola"
 1708 OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Capsiempis" OR "Casiornis" OR "Cnemarchus" OR
 1709 "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR
 1710 "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR
 1711 "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis"
 1712 OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR
 1713 "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus"
 1714 OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myio-
 1715 dynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiotriccus" OR
 1716 "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR
 1717 "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes"
 1718 OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis"
 1719 OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR
 1720 "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis"
 1721 OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR
 1722 "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis"
 1723 OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR
 1724 "Urocynchramus" OR "Artamella" OR "Calicalicus" OR "Cyanolanius" OR "Euryceros" OR "Falculea"
 1725 OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR
 1726 "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris"
 1727 OR "Anomalospiza" OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR
 1728 "Vireo" OR "Vireolanius" OR "Zeledonia" OR "Apalopteron" OR "Dasycrotapha" OR "Heleia" OR "Lopho-
 1729 zosterops" OR "Sternhoptilus" OR "Yuhina" OR "Zosterops" OR "Zosterornis" OR "Agamia" OR "Ardea" OR
 1730 "Ardeola" OR "Botaurus" OR "Bubulcus" OR "Butorides" OR "Cochlearius" OR "Dupetor" OR "Egretta"
 1731 OR "Gorsachius" OR "Ixobrychus" OR "Nyctanassa" OR "Nycticorax" OR "Ptilerodius" OR "Syrigma" OR
 1732 "Tigriornis" OR "Tigrisoma" OR "Zebrilus" OR "Balaeniceps" OR "Pelecanus" OR "Scopus" OR "Bostrychia"
 1733 OR "Eudocimus" OR "Geronticus" OR "Lophotibis" OR "Mesembrinibis" OR "Nipponia" OR "Phimosus"
 1734 OR "Platalea" OR "Plegadis" OR "Pseudibis" OR "Theristicus" OR "Threskiornis" OR "Phaethon" OR
 1735 "Phoeniconaias" OR "Phoenicoparrus" OR "Phoenicopterus" OR "Bucco" OR "Chelidoptera" OR "Hapalop-

1736 tila" OR "Hypnelus" OR "Malacoptila" OR "Micromonacha" OR "Monasa" OR "Nonnula" OR "Notharchus"
1737 OR "Nystalus" OR "Capito" OR "Eubucco" OR "Brachygalba" OR "Galbalcyrhynchus" OR "Galbula" OR
1738 "Jacamaralcyon" OR "Jacamerops" OR "Indicator" OR "Melichneutes" OR "Malignomon" OR "Prodotiscus"
1739 OR "Buccanodon" OR "Gymnobucco" OR "Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus"
1740 OR "Psilopogon" OR "Blythipicus" OR "Campephilus" OR "Campethera" OR "Celeus" OR "Chloropicus"
1741 OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Dendrocopos" OR "Dendrocoptes" OR
1742 "Dendropicos" OR "Dinopium" OR "Dryobates" OR "Dryocopus" OR "Gecinulus" OR "Geocolaptes" OR
1743 "Jynx" OR "Leiopicus" OR "Leuconotopicus" OR "Meiglyptes" OR "Melanerpes" OR "Micropternus" OR
1744 "Mulleripicus" OR "Nesocites" OR "Picoides" OR "Piculus" OR "Picumnus" OR "Picus" OR "Reinwardtipi-
1745 cus" OR "Sasia" OR "Sphyrapicus" OR "Veniliornis" OR "Yungipicus" OR "Andigena" OR "Aulacorhynchus"
1746 OR "Pteroglossus" OR "Ramphastos" OR "Selenidera" OR "Semnornis" OR "Aechmophorus" OR "Podiceps"
1747 OR "Podilymbus" OR "Poliiocephalus" OR "Rollandia" OR "Tachybaptus" OR "Diomedea" OR "Phoebastria"
1748 OR "Phoebetria" OR "Thalassarche" OR "Hydrobates" OR "Oceanodroma" OR "Fregetta" OR "Garrodia"
1749 OR "Nesofregetta" OR "Oceanites" OR "Pelagodroma" OR "Aphrodroma" OR "Ardena" OR "Bulweria"
1750 OR "Calonectris" OR "Daption" OR "Fulmarus" OR "Halobaena" OR "Macronektes" OR "Pachyptila" OR
1751 "Pagodroma" OR "Pelecanoides" OR "Procellaria" OR "Pseudobulweria" OR "Pterodroma" OR "Puffinus" OR
1752 "Thalassoica" OR "Cacatua" OR "Callocephalon" OR "Calyptorhynchus" OR "Eolophus" OR "Lophochroa"
1753 OR "Nymphicus" OR "Probosciger" OR "Alipiopsitta" OR "Amazona" OR "Anodorhynchus" OR "Ara" OR
1754 "Aratinga" OR "Bolborhynchus" OR "Brotogeris" OR "Conuropsis" OR "Cyanoliseus" OR "Cyanopsitta" OR
1755 "Derophtus" OR "Diopsittaca" OR "Enicognathus" OR "Eupsittula" OR "Forpus" OR "Graydidascalus" OR
1756 "Guarouba" OR "Guaruba" OR "Hapalopsittaca" OR "Leptosittaca" OR "Myiopsitta" OR "Nannopsittaca"
1757 OR "Orthopsittaca" OR "Pionites" OR "Pionopsitta" OR "Pionus" OR "Poicephalus" OR "Primolius"
1758 OR "Psilopsiagon" OR "Psittacara" OR "Psittacus" OR "Pyrilia" OR "Pyrrhura" OR "Rhynchopsitta"
1759 OR "Thectocercus" OR "Touit" OR "Triclaria" OR "Agapornis" OR "Alisterus" OR "Aprosmictus" OR
1760 "Barnardius" OR "Bolbopsittacus" OR "Chalcopsitta" OR "Charmosyna" OR "Coracopsis" OR "Cyanoram-
1761 phus" OR "Cyclopsitta" OR "Eclectus" OR "Eos" OR "Eunymphicus" OR "Geoffroyus" OR "Glossopsitta"
1762 OR "Lathamus" OR "Loriculus" OR "Lorius" OR "Melopsittacus" OR "Micropsitta" OR "Neophema" OR
1763 "Neopsephotus" OR "Neopsittacus" OR "Northiella" OR "Oreopsittacus" OR "Parvipsitta" OR "Pezoporus"
1764 OR "Phigys" OR "Platycercus" OR "Polytelis" OR "Prioniturus" OR "Prosopeia" OR "Psephotellus" OR
1765 "Psephotus" OR "Pseudeos" OR "Psittacella" OR "Psittacula" OR "Psittaculirostris" OR "Psitteuteles" OR
1766 "Psittinus" OR "Psittichas" OR "Purpureicephalus" OR "Tanygnathus" OR "Trichoglossus" OR "Vini" OR
1767 "Nestor" OR "Strigops" OR "Pterocles" OR "Syrrhaptes" OR "Rhea" OR "Aptenodytes" OR "Eudypetes" OR
1768 "Eudypetula" OR "Megadyptes" OR "Pygoscelis" OR "Spheniscus" OR "Aegolius" OR "Asio" OR "Athene"
1769 OR "Bubo" OR "Glaucidium" OR "Ketupa" OR "Lophotrix" OR "Megascops" OR "Micrathene" OR "Ninox"
1770 OR "Otus" OR "Pseudoscops" OR "Psilosops" OR "Pulsatrix" OR "Sceloglaux" OR "Scotopelia" OR "Strix"

1771 OR "Surnia" OR "Xenoglaux" OR "Phodilus" OR "Tyto" OR "Aepyornis" OR "Mullerornis" OR "Struthio"
 1772 OR "Anhinga" OR "Fregata" OR "Leucocarbo" OR "Microcarbo" OR "Nannopterum" OR "Phalacrocorax"
 1773 OR "Morus" OR "Sula" OR "Crypturellus" OR "Eudromia" OR "Nothocercus" OR "Nothoprocta" OR
 1774 "Nothura" OR "Rhynchotus" OR "Tinamotis" OR "Tinamus" OR "Apalharpactes" OR "Apaloderma" OR
 1775 "Harpactes" OR "Pharomachrus" OR "Trogon" OR "Tetrastes" OR "Hesperiphona") AND ("experiment*" OR
 1776 OR "manipulat*"))))>

1777

1778 The search was also conducted on SCOPUS using Advance Search Option and Title Abstract and Keyword
 1779 (TITLE-ABS-KEY) on the same dates using the string

1780

1781 <TITLE-ABS-KEY (((("green*" OR "herb*" OR "aromatic*") AND "nest*" AND ("bird*" OR "aves" OR
 1782 "avian" OR "ornithol*" OR "passerine*" OR "passeriform*" OR "songbird*" OR "Accipiter" OR "Aegypius"
 1783 OR "Aquila" OR "Aviceda" OR "Busarellus" OR "Butastur" OR "Buteo" OR "Buteogallus" OR "Chon-
 1784 drohierax" OR "Circaetus" OR "Circus" OR "Clanga" OR "Elanoides" OR "Elanus" OR "Gampsonyx"
 1785 OR "Geranoaetus" OR "Geranospiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR
 1786 "Haliastur" OR "Harpagus" OR "Harpia" OR "Hieraaetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco"
 1787 OR "Leptodon" OR "Leucopternis" OR "Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus"
 1788 OR "Necrosyrtes" OR "Neophron" OR "Nisaetus" OR "Parabuteo" OR "Pernis" OR "Pithecophaga" OR
 1789 "Polemaetus" OR "Pseudastur" OR "Rostrhamus" OR "Rupornis" OR "Sarcogyps" OR "Spilornis" OR
 1790 "Spizaetus" OR "Stephanoaetus" OR "Terathopius" OR "Torgos" OR "Trigonoceps" OR "Cathartes" OR
 1791 "Coragyps" OR "Gymnogyps" OR "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix"
 1792 OR "Alopochen" OR "Amazonetta" OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR
 1793 "Bucephala" OR "Cairina" OR "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloephaga"
 1794 OR "Clangula" OR "Coscoroba" OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR
 1795 "Histrionicus" OR "Hymenolaimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca"
 1796 OR "Marmaronetta" OR "Melanitta" OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neochen" OR
 1797 "Netta" OR "Nettapus" OR "Nomonyx" OR "Oxyura" OR "Plectropterus" OR "Polysticta" OR "Pteronetta"
 1798 OR "Sarkidiornis" OR "Sibirionetta" OR "Somateria" OR "Spatula" OR "Tachyeres" OR "Tadorna" OR
 1799 "Thalassornis" OR "Anhima" OR "Chauna" OR "Anseranas" OR "Aegotheles" OR "Aerodramus" OR "Aero-
 1800 nautes" OR "Apus" OR "Chaetura" OR "Collocalia" OR "Cypseloides" OR "Cypsiurus" OR "Hirundapus"
 1801 OR "Panyptila" OR "Streptoprocne" OR "Tachornis" OR "Tachymarptis" OR "Hemiprocne" OR "Abeillia"
 1802 OR "Adelomyia" OR "Aglaeactis" OR "Aglaiocercus" OR "Amazilia" OR "Androdon" OR "Anopetia" OR
 1803 "Anthocephala" OR "Anthracothorax" OR "Archilochus" OR "Atthis" OR "Augastes" OR "Avocettula" OR
 1804 "Basilinna" OR "Boissonneaua" OR "Calliphlox" OR "Calypte" OR "Campylopterus" OR "Chaetocercus" OR
 1805 "Chalcostigma" OR "Chalybura" OR "Chlorestes" OR "Chlorostilbon" OR "Chrysolampis" OR "Chrysuronia"

1806 OR "Clytolaema" OR "Coeligena" OR "Colibri" OR "Cyanophaia" OR "Cynanthus" OR "Discosura" OR
 1807 "Doricha" OR "Doryfera" OR "Elvira" OR "Ensifera" OR "Eriocnemis" OR "Eugenes" OR "Eulampis" OR
 1808 "Eulidia" OR "Eupetomena" OR "Eupherusa" OR "Eutoxeres" OR "Florisuga" OR "Glaucis" OR "Goethalsia"
 1809 OR "Goldmania" OR "Haplophaedia" OR "Heliactin" OR "Heliangelus" OR "Heliodoxa" OR "Heliomaster"
 1810 OR "Heliothryx" OR "Hylocharis" OR "Juliamyia" OR "Klais" OR "Lafresnaya" OR "Lampornis" OR
 1811 "Lamprolaima" OR "Lepidopyga" OR "Lesbia" OR "Leucippus" OR "Leucochloris" OR "Lophornis" OR
 1812 "Mellisuga" OR "Metallura" OR "Microchera" OR "Myrmia" OR "Myrtis" OR "Ocreatus" OR "Opistho-
 1813 prora" OR "Oreonympha" OR "Oreotrochilus" OR "Orthorhyncus" OR "Oxypogon" OR "Panterpe" OR
 1814 "Patagona" OR "Phaeochroa" OR "Phaethornis" OR "Phlogophilus" OR "Polyonymus" OR "Polytmus" OR
 1815 "Pterophanes" OR "Ramphodon" OR "Ramphomicron" OR "Rhodopis" OR "Sappho" OR "Schistes" OR
 1816 "Selasphorus" OR "Sephanoides" OR "Stephanoxis" OR "Sternoclyta" OR "Taphrospilus" OR "Thalurania"
 1817 OR "Thaumastura" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR
 1818 "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR
 1819 "Rhabdotorrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus"
 1820 OR "Upupa" OR "Antrostomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus"
 1821 OR "Hydropsalis" OR "Lurocalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus"
 1822 OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius"
 1823 OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariama" OR "Casuarius" OR "Dromaius" OR
 1824 "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cerorhinca" OR "Fratricula" OR
 1825 "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus"
 1826 OR "Charadrius" OR "Elseyornis" OR "Erythrogonyx" OR "Hoploxypterus" OR "Oreopholus" OR "Pel-
 1827 tohyas" OR "Phegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR
 1828 "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR
 1829 "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR
 1830 "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelocheilidon" OR "Gygis" OR "Hydrocoloeus" OR
 1831 "Hydroprogne" OR "Ichthyaelus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR
 1832 "Pagophila" OR "Phaetusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula"
 1833 OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR
 1834 "Recurvirostra" OR "Nycticryphes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR
 1835 "Calidris" OR "Coenocorypha" OR "Gallinago" OR "Limnodromus" OR "Limosa" OR "Lymnocyptes" OR
 1836 "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagis"
 1837 OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR
 1838 "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chal-
 1839 cophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula"
 1840 OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR

1841 “Gymnophaps” OR “Hemiphaga” OR “Henicophaps” OR “Leptotila” OR “Leptotrygon” OR “Leucosarcia”
 1842 OR “Lopholaimus” OR “Macropygia” OR “Metriopelia” OR “Ocyphaps” OR “Oena” OR “Otidiphaps”
 1843 OR “Patagioenas” OR “Petrophassa” OR “Pezophaps” OR “Phapitreron” OR “Phaps” OR “Ptilinopus”
 1844 OR “Raphus” OR “Reinwardtoena” OR “Spilopelia” OR “Streptopelia” OR “Treron” OR “Trugon” OR
 1845 “Turacoena” OR “Turtur” OR “Uropelia” OR “Zenaida” OR “Zentrygon” OR “Actenoides” OR “Alcedo” OR
 1846 “Caridonax” OR “Ceryle” OR “Ceyx” OR “Chloroceryle” OR “Cittura” OR “Corythornis” OR “Dacelo” OR
 1847 “Halcyon” OR “Isipidina” OR “Lacedo” OR “Megaceryle” OR “Melidora” OR “Syma” OR “Tanyptera” OR
 1848 “Todiramphus” OR “Atelornis” OR “Brachypteracias” OR “Geobiastes” OR “Coracias” OR “Eurystomus”
 1849 OR “Merops” OR “Nyctornis” OR “Baryphthengus” OR “Electron” OR “Eumomota” OR “Hylomanes”
 1850 OR “Momotus” OR “Todus” OR “Cacomantis” OR “Carpococcyx” OR “Centropus” OR “Cercococcyx” OR
 1851 “Chrysococcyx” OR “Clamator” OR “Coccyua” OR “Coccyzus” OR “Coua” OR “Crotophaga” OR “Cuculus”
 1852 OR “Dasylophus” OR “Dromococcyx” OR “Eudynamys” OR “Geococcyx” OR “Guira” OR “Hierococcyx”
 1853 OR “Morococcyx” OR “Neomorphus” OR “Pachycoccyx” OR “Phaenicophaeus” OR “Piaya” OR “Rhinortha”
 1854 OR “Scythrops” OR “Surniculus” OR “Tapera” OR “Urodynamis” OR “Zanclostomus” OR “Dinornis” OR
 1855 “Anomalopteryx” OR “Emeus” OR “Euryapteryx” OR “Eurypyga” OR “Rhynochetos” OR “Caracara” OR
 1856 “Daptrius” OR “Falco” OR “Herpetotheres” OR “Ibycter” OR “Micrastur” OR “Microhierax” OR “Milvago”
 1857 OR “Phalcoboenus” OR “Polihierax” OR “Spiziapteryx” OR “Aburria” OR “Chamaepetes” OR “Crax” OR
 1858 “Mitu” OR “Nothocrax” OR “Oreophasis” OR “Ortalis” OR “Pauxi” OR “Penelope” OR “Penelopina” OR
 1859 “Pipile” OR “Alectura” OR “Megapodius” OR “Acryllium” OR “Guttera” OR “Numida” OR “Callipepla”
 1860 OR “Colinus” OR “Cyrtonyx” OR “Dendrortyx” OR “Odontophorus” OR “Oreortyx” OR “Philortyx” OR
 1861 “Ptilopachus” OR “Rhynchortyx” OR “Alectoris” OR “Ammoperdix” OR “Arborophila” OR “Argusianus”
 1862 OR “Bambusicola” OR “Bonasa” OR “Caloperdix” OR “Centrocercus” OR “Chrysolophus” OR “Coturnix”
 1863 OR “Crossoptilon” OR “Dendragapus” OR “Excalfactoria” OR “Falcipennis” OR “Francolinus” OR “Gallus”
 1864 OR “Haematortyx” OR “Ithaginis” OR “Lagopus” OR “Lerwa” OR “Lophophorus” OR “Lophura” OR
 1865 “Lyrurus” OR “Meleagris” OR “Pavo” OR “Peliperdix” OR “Perdix” OR “Phasianus” OR “Polyplectron” OR
 1866 “Pternistis” OR “Pucrasia” OR “Rhizothera” OR “Rollulus” OR “Scleroptila” OR “Syrmaticus” OR “Tetrao”
 1867 OR “Tetraogallus” OR “Tetraophasis” OR “Tragopan” OR “Tympanuchus” OR “Gavia” OR “Aramus”
 1868 OR “Anthropoides” OR “Antigone” OR “Balearica” OR “Bugeranus” OR “Grus” OR “Leucogeranus” OR
 1869 “Heliornis” OR “Psophia” OR “Aenigmatolimnas” OR “Amaurolimnas” OR “Amauornis” OR “Anurolimnas”
 1870 OR “Aramides” OR “Atlantisia” OR “Coturnicops” OR “Crex” OR “Dryolimnas” OR “Eulabeornis” OR
 1871 “Fulica” OR “Gallicrex” OR “Gallinula” OR “Gallirallus” OR “Laterallus” OR “Lewinia” OR “Micropygia”
 1872 OR “Neocrex” OR “Nesoclopeus” OR “Paragallinula” OR “Pardirallus” OR “Porphyrio” OR “Porphyriops”
 1873 OR “Porzana” OR “Rallacula” OR “Rallina” OR “Rallus” OR “Tribonyx” OR “Canirallus” OR “Sarothrura”
 1874 OR “Leptosomus” OR “Corythaeola” OR “Corythaixoides” OR “Crinifer” OR “Musophaga” OR “Tauraco”
 1875 OR “Opisthocomus” OR “Afrotis” OR “Ardeotis” OR “Chlamydotis” OR “Neotis” OR “Otis” OR “Tetrax”

1876 OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR
 1877 "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis"
 1878 OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis"
 1879 OR "Smicrornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna"
 1880 OR "Nesillas" OR "Aegithalos" OR "Leptopoecile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR
 1881 "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes"
 1882 OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafr" OR
 1883 "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR
 1884 "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhynchophanes" OR "Callaeas" OR "Heteralocha" OR
 1885 "Philesturnus" OR "Calyptophilus" OR "Campephaga" OR "Ceblepyris" OR "Coracina" OR "Edolisoma" OR
 1886 "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amaurospiza" OR "Cardinalis" OR "Caryothraustes" OR
 1887 "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Passerina" OR
 1888 "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus"
 1889 OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus"
 1890 OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR
 1891 "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia"
 1892 OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR
 1893 "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR
 1894 "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR
 1895 "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus"
 1896 OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus"
 1897 OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR
 1898 "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias"
 1899 OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR
 1900 "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza"
 1901 OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR
 1902 "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis"
 1903 OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga"
 1904 OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calyptomena" OR "Cymbirhynchus" OR
 1905 "Eurylaimus" OR "Psarisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR
 1906 "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chlo-
 1907 ris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra"
 1908 OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorhous" OR "Hemignathus" OR
 1909 "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR
 1910 "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR

1911 “Pseudonestor” OR “Psittirostra” OR “Pyrrhoptes” OR “Pyrrhula” OR “Rhodopechys” OR “Rhodospiza”
 1912 OR “Serinus” OR “Spinus” OR “Telespiza” OR “Anabacerthia” OR “Anabazenops” OR “Ancistrops” OR
 1913 “Anumbius” OR “Aphrastura” OR “Asthenes” OR “Automolus” OR “Berlepschia” OR “Campylorhamphus”
 1914 OR “Certhiasomus” OR “Certhiaxis” OR “Cinclodes” OR “Clibanornis” OR “Coryphistera” OR “Cranioleuca”
 1915 OR “Deconychura” OR “Dendrexetastes” OR “Dendrocincla” OR “Dendrocolaptes” OR “Dendroplex” OR
 1916 “Drymornis” OR “Drymotoxeres” OR “Furnarius” OR “Geocerthia” OR “Geositta” OR “Glyphorhynchus”
 1917 OR “Heliobletus” OR “Hellmayrea” OR “Hylexetastes” OR “Lepidocolaptes” OR “Leptasthenura” OR
 1918 “Limnortyx” OR “Limnornis” OR “Lochmias” OR “Margarornis” OR “Mazaria” OR “Megaxenops” OR
 1919 “Metopothrix” OR “Microxenops” OR “Nasica” OR “Ochetorhynchus” OR “Phacellodomus” OR “Philydor”
 1920 OR “Phleocryptes” OR “Premnoplex” OR “Premnornis” OR “Pseudasthenes” OR “Pseudocolaptes” OR
 1921 “Pseudoseisura” OR “Pygarrhichas” OR “Roraimia” OR “Schoeniophylax” OR “Sclerurus” OR “Sittasomus”
 1922 OR “Spartonoica” OR “Sylviorhynchus” OR “Synallaxis” OR “Syndactyla” OR “Tarphonornis” OR
 1923 “Thripadectes” OR “Thripophaga” OR “Upucerthia” OR “Xenerpestes” OR “Xenops” OR “Xiphocolaptes”
 1924 OR “Xiphorhynchus” OR “Grallaria” OR “Grallaricula” OR “Hylopezus” OR “Myrmothera” OR “Alopocheli-
 1925 don” OR “Atticora” OR “Cecropis” OR “Delichon” OR “Haplochelidon” OR “Hirundo” OR “Neochelidon”
 1926 OR “Notiochelidon” OR “Orochelidon” OR “Petrochelidon” OR “Phedina” OR “Progne” OR “Psolidoprocne”
 1927 OR “Pseudhirundo” OR “Pseudochelidon” OR “Ptyonoprogne” OR “Riparia” OR “Stelgidopteryx” OR
 1928 “Tachycineta” OR “Hylia” OR “Hypocolius” OR “Agelaioides” OR “Agelaius” OR “Agelasticus” OR “Am-
 1929 blycercus” OR “Amblyramphus” OR “Anumara” OR “Cacicus” OR “Chrysomus” OR “Curaeus” OR “Dives”
 1930 OR “Dolichonyx” OR “Euphagus” OR “Gnorimopsar” OR “Gymnomystax” OR “Hypopyrrhus” OR “Icterus”
 1931 OR “Lamprolaima” OR “Leistes” OR “Macroagelaius” OR “Molothrus” OR “Nesopsar” OR “Oreopsar” OR
 1932 “Psarocolius” OR “Pseudoleistes” OR “Quiscalus” OR “Sturnella” OR “Xanthocephalus” OR “Xanthopsar”
 1933 OR “Icteria” OR “Ifrita” OR “Irena” OR “Corvinella” OR “Eurocephalus” OR “Lanius” OR “Actinodura”
 1934 OR “Argya” OR “Garrulax” OR “Grammatoptila” OR “Heterophasia” OR “Ianthocincla” OR “Laniellus”
 1935 OR “Leiothrix” OR “Liocichla” OR “Minla” OR “Pterorhinus” OR “Trochaloxypterus” OR “Turdoides” OR
 1936 “Bradypterus” OR “Cincloramphus” OR “Helopsaltes” OR “Locustella” OR “Megalurus” OR “Poodytes” OR
 1937 “Machaerirhynchus” OR “Graueria” OR “Hylia” OR “Macrosphenus” OR “Melocichla” OR “Sylvietta” OR
 1938 “Chlorophoneus” OR “Dryoscopus” OR “Laniarius” OR “Malaconotus” OR “Nilaus” OR “Rhodophoneus” OR
 1939 “Tchagra” OR “Telophorus” OR “Amytornis” OR “Chenorhamphus” OR “Clytomyias” OR “Malurus” OR
 1940 “Stipiturus” OR “Melampitta” OR “Melanocharis” OR “Oedistoma” OR “Toxorhamphus” OR “Melanopareia”
 1941 OR “Acanthagenys” OR “Acanthorhynchus” OR “Anthochaera” OR “Anthornis” OR “Caligavis” OR “Entomy-
 1942 zon” OR “Epthianura” OR “Glycifolia” OR “Gymnomyza” OR “Lichmera” OR “Manorina” OR “Melidectes”
 1943 OR “Melilestes” OR “Meliphaga” OR “Melipotes” OR “Melithreptus” OR “Myzomela” OR “Nesoptilotis”
 1944 OR “Philemon” OR “Phylidonyris” OR “Prothemadon” OR “Ptiloprora” OR “Ptilotula” OR “Pycnopygius”
 1945 OR “Timeliopsis” OR “Xanthotis” OR “Menura” OR “Allenia” OR “Cinclocerthia” OR “Dumetella” OR

1946 “Margarops” OR “Melanoptila” OR “Melanotis” OR “Mimus” OR “Oreoscoptes” OR “Ramphocinclus” OR
 1947 “Toxostoma” OR “Lamprospiza” OR “Mitrospingus” OR “Moho” OR “Mohoua” OR “Arses” OR “Carterornis”
 1948 OR “Clytorhynchus” OR “Grallina” OR “Hypothymis” OR “Monarcha” OR “Myiagra” OR “Symposiachrus”
 1949 OR “Terpsiphone” OR “Trochocercus” OR “Anthus” OR “Macronyx” OR “Motacilla” OR “Alethe” OR
 1950 “Anthipes” OR “Brachypteryx” OR “Calliope” OR “Campicoloides” OR “Cercotrichas” OR “Chamaetylas”
 1951 OR “Copsychus” OR “Cossypha” OR “Cyanoptila” OR “Cyornis” OR “Enicurus” OR “Erithacus” OR
 1952 “Eumyias” OR “Ficedula” OR “Fraseria” OR “Irania” OR “Larvivora” OR “Leonardina” OR “Luscinia”
 1953 OR “Melaenornis” OR “Monticola” OR “Muscicapa” OR “Myiomela” OR “Myioparus” OR “Myophonus”
 1954 OR “Myrmecocichla” OR “Niltava” OR “Oenanthe” OR “Phoenicurus” OR “Pogonocichla” OR “Saxicola”
 1955 OR “Sheppardia” OR “Sholicola” OR “Stiphornis” OR “Tarsiger” OR “Thamnolaea” OR “Vauriella” OR
 1956 “Aethopyga” OR “Anabathmis” OR “Anthreptes” OR “Arachnothera” OR “Chalcomitra” OR “Cinnyris” OR
 1957 “Cyanomitra” OR “Deleornis” OR “Hedydipna” OR “Kurochkinogramma” OR “Leptocoma” OR “Nectarinia”
 1958 OR “Daphoenositta” OR “Nesospingus” OR “Nicator” OR “Notiomystis” OR “Aleadryas” OR “Oriolus” OR
 1959 “Pitohui” OR “Sphecotheres” OR “Turnagra” OR “Colluricincla” OR “Falcunculus” OR “Melanorectes” OR
 1960 “Pachycephala” OR “Pseudorectes” OR “Panurus” OR “Cicinnurus” OR “Diphyllodes” OR “Manucodia” OR
 1961 “Paradisaea” OR “Ptiloris” OR “Paramythia” OR “Pardalotus” OR “Baeolophus” OR “Cephalopyrus” OR
 1962 “Cyanistes” OR “Lophophanes” OR “Machlolophus” OR “Melaniparus” OR “Melanochlora” OR “Pardali-
 1963 parus” OR “Parus” OR “Periparus” OR “Poecile” OR “Pseudopodoces” OR “Sittiparus” OR “Sylviparus”
 1964 OR “Basileuterus” OR “Cardellina” OR “Catharopeza” OR “Dendroica” OR “Geothlypis” OR “Helmitheros”
 1965 OR “Leiothlypis” OR “Limnethlypis” OR “Mniotilta” OR “Myioborus” OR “Myiothlypis” OR “Oporornis”
 1966 OR “Oreothlypis” OR “Parkesia” OR “Protonotaria” OR “Seiurus” OR “Setophaga” OR “Vermivora” OR
 1967 “Aimophila” OR “Ammodramus” OR “Amphispiza” OR “Arremon” OR “Arremonops” OR “Artemisiospiza”
 1968 OR “Atlapetes” OR “Calamospiza” OR “Chlorospingus” OR “Chondestes” OR “Junco” OR “Melospiza”
 1969 OR “Melozone” OR “Oreothraupis” OR “Oriturus” OR “Passerculus” OR “Passerella” OR “Peucaea” OR
 1970 “Pezopetes” OR “Pipilo” OR “Pooecetes” OR “Pselliophorus” OR “Rhynchospiza” OR “Spizella” OR “Spizel-
 1971 loides” OR “Xenospiza” OR “Zonotrichia” OR “Carpospiza” OR “Gymnoris” OR “Hypocryptadius” OR
 1972 “Montifringilla” OR “Onychostruthus” OR “Passer” OR “Petronia” OR “Pyrgilauda” OR “Alcippe” OR
 1973 “Gampsorhynchus” OR “Illadopsis” OR “Jabouilleia” OR “Kenopia” OR “Laticilla” OR “Malacocincla” OR
 1974 “Malacopteron” OR “Napothera” OR “Pellorneum” OR “Rimator” OR “Trichastoma” OR “Amalocichla”
 1975 OR “Drymodes” OR “Eopsaltria” OR “Eugerygone” OR “Heteromyias” OR “Melanodryas” OR “Microeca”
 1976 OR “Monachella” OR “Pachycephalopsis” OR “Peneoenanthe” OR “Peneothello” OR “Petroica” OR “Poe-
 1977 cilodryas” OR “Tregellasia” OR “Peucedramus” OR “Microligea” OR “Phaenicophilus” OR “Xenoligea” OR
 1978 “Phylloscopus” OR “Picathartes” OR “Antilophia” OR “Ceratopipra” OR “Chiroxiphia” OR “Chloropipo” OR
 1979 “Corapipo” OR “Cryptopipo” OR “Heterocercus” OR “Ilicura” OR “Lepidothrix” OR “Machaeropterus” OR
 1980 “Manacus” OR “Masius” OR “Neopelma” OR “Pipra” OR “Pseudopipra” OR “Tyrannutes” OR “Xenopipo”

1981 OR "Erythropitta" OR "Hydrornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalor-
 1982 nis" OR "Euplectes" OR "Malimbus" OR "Plocepasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR
 1983 "Pnoepyga" OR "Microbates" OR "Polioptila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR
 1984 "Cinclosoma" OR "Ptilorrhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR
 1985 "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla"
 1986 OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Criniger" OR "Eurillas" OR
 1987 "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus"
 1988 OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus"
 1989 OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pterop-
 1990 tochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR
 1991 "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynx" OR "Culicicapa" OR "Elminia" OR "Acridotheres"
 1992 OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora"
 1993 OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotonis" OR "Leucopsar"
 1994 OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR
 1995 "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR
 1996 "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR
 1997 "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR
 1998 "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprosi-
 1999 tornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR
 2000 "Drymophila" OR "Dysithamnus" OR "Epinecrophylla" OR "Euchrepomis" OR "Formicivora" OR "Freder-
 2001 ickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR
 2002 "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus"
 2003 OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myr-
 2004 moderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes"
 2005 OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania"
 2006 OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus"
 2007 OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR
 2008 "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis"
 2009 OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Charitospiza" OR "Chlorochrysa" OR
 2010 "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba"
 2011 OR "Compsospiza" OR "Compsothraupis" OR "Conirostrum" OR "Conothraupis" OR "Coryphospingus"
 2012 OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis"
 2013 OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR
 2014 "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR
 2015 "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Incasiza" OR "Iridophanes" OR "Iridosornis"

2016 OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR
 2017 "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes"
 2018 OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocoma" OR
 2019 "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha"
 2020 OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR
 2021 "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR
 2022 "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatorhinus" OR "Spelaeornis"
 2023 OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius"
 2024 OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR
 2025 "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia"
 2026 OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR
 2027 "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Trogodytes"
 2028 OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR
 2029 "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus"
 2030 OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola"
 2031 OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Casiornis" OR "Cnemarchus" OR
 2032 "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR
 2033 "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR
 2034 "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis"
 2035 OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR
 2036 "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus"
 2037 OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myio-
 2038 dynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiotriccus" OR
 2039 "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR
 2040 "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes"
 2041 OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis"
 2042 OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR
 2043 "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis"
 2044 OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR
 2045 "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis"
 2046 OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR
 2047 "Urocynchramus" OR "Artamella" OR "Calicalicus" OR "Cyanolanius" OR "Euryceros" OR "Falculea"
 2048 OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR
 2049 "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris"
 2050 OR "Anomalospiza" OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR

2051 “Vireo” OR “Vireolanius” OR “Zeledonia” OR “Apalopteron” OR “Dasycrotapha” OR “Heleia” OR “Lopho-
 2052 zosterops” OR “Sterrhoptilus” OR “Yuhina” OR “Zosterops” OR “Zosterornis” OR “Agamia” OR “Ardea” OR
 2053 “Ardeola” OR “Botaurus” OR “Bubulcus” OR “Butorides” OR “Cochlearius” OR “Dupetor” OR “Egretta”
 2054 OR “Gorsachius” OR “Ixobrychus” OR “Nyctanassa” OR “Nycticorax” OR “Pilierodius” OR “Syrrigma” OR
 2055 “Tigriornis” OR “Tigrisoma” OR “Zebrius” OR “Balaeniceps” OR “Pelecanus” OR “Scopus” OR “Bostrychia”
 2056 OR “Eudocimus” OR “Geronticus” OR “Lophotibis” OR “Mesembrinibis” OR “Nipponia” OR “Phimosus”
 2057 OR “Platalea” OR “Plegadis” OR “Pseudibis” OR “Theristicus” OR “Threskiornis” OR “Phaethon” OR
 2058 “Phoeniconaias” OR “Phoenicoparrus” OR “Phoenicopterus” OR “Bucco” OR “Chelidoptera” OR “Hapalop-
 2059 tila” OR “Hypnelus” OR “Malacoptila” OR “Micromonacha” OR “Monasa” OR “Nonnula” OR “Notharchus”
 2060 OR “Nystalus” OR “Capito” OR “Eubucco” OR “Brachygalba” OR “Galbalcyrhynchus” OR “Galbula” OR
 2061 “Jacamaralcyon” OR “Jacamerops” OR “Indicator” OR “Melichneutes” OR “Malignomon” OR “Prodotiscus”
 2062 OR “Buccanodon” OR “Gymnobucco” OR “Lybius” OR “Pogoniulus” OR “Stactolaema” OR “Trachyphonus”
 2063 OR “Psilopogon” OR “Blythipicus” OR “Campephilus” OR “Campethera” OR “Celeus” OR “Chloropicus”
 2064 OR “Chrysocolaptes” OR “Chrysophlegma” OR “Colaptes” OR “Dendrocopos” OR “Dendrocytes” OR
 2065 “Dendropicus” OR “Dinopium” OR “Dryobates” OR “Dryocopus” OR “Gecinulus” OR “Geocolaptes” OR
 2066 “Jynx” OR “Leiopicus” OR “Leuconotopicus” OR “Meiglyptes” OR “Melanerpes” OR “Micropternus” OR
 2067 “Mulleripicus” OR “Nesocites” OR “Picoides” OR “Piculus” OR “Picumnus” OR “Picus” OR “Reinwardtipi-
 2068 cus” OR “Sasia” OR “Sphyrapicus” OR “Veniliornis” OR “Yungipicus” OR “Andigena” OR “Aulacorhynchus”
 2069 OR “Pteroglossus” OR “Ramphastos” OR “Selenidera” OR “Semnornis” OR “Aechmophorus” OR “Podiceps”
 2070 OR “Podilymbus” OR “Polioccephalus” OR “Rollandia” OR “Tachybaptus” OR “Diomedea” OR “Phoebastria”
 2071 OR “Phoebetria” OR “Thalassarche” OR “Hydrobates” OR “Oceanodroma” OR “Fregetta” OR “Garrodia”
 2072 OR “Nesofregetta” OR “Oceanites” OR “Pelagodroma” OR “Aphrodroma” OR “Ardena” OR “Bulweria”
 2073 OR “Calonectris” OR “Daption” OR “Fulmarus” OR “Halobaena” OR “Macronectes” OR “Pachyptila” OR
 2074 “Pagodroma” OR “Pelecanoides” OR “Procellaria” OR “Pseudobulweria” OR “Pterodroma” OR “Puffinus” OR
 2075 “Thalassoica” OR “Cacatua” OR “Callocephalon” OR “Calyptorhynchus” OR “Eolophus” OR “Lophochroa”
 2076 OR “Nymphicus” OR “Probosciger” OR “Alipiopsitta” OR “Amazona” OR “Anodorhynchus” OR “Ara” OR
 2077 “Aratinga” OR “Bolborhynchus” OR “Brotogeris” OR “Conuropsis” OR “Cyanoliseus” OR “Cyanopsitta” OR
 2078 “Derophtus” OR “Diopsittaca” OR “Enicognathus” OR “Eupsittula” OR “Forpus” OR “Graydidascalus” OR
 2079 “Guarouba” OR “Guaruba” OR “Hapalopsittaca” OR “Leptosittaca” OR “Myiopsitta” OR “Nannopsittaca”
 2080 OR “Orthopsittaca” OR “Pionites” OR “Pionopsitta” OR “Pionus” OR “Poicephalus” OR “Primolius”
 2081 OR “Psilopsiagon” OR “Psittacara” OR “Psittacus” OR “Pyrrhura” OR “Pyrrhura” OR “Rhynchopsitta”
 2082 OR “Thectocercus” OR “Touit” OR “Triclaria” OR “Agapornis” OR “Alisterus” OR “Aprosmictus” OR
 2083 “Barnardius” OR “Bolbopsittacus” OR “Chalcopsitta” OR “Charmosyna” OR “Coracopsis” OR “Cyanoram-
 2084 phus” OR “Cyclopsitta” OR “Eclectus” OR “Eos” OR “Eunymphicus” OR “Geoffroyus” OR “Glossopsitta”
 2085 OR “Lathamus” OR “Loriculus” OR “Lorius” OR “Melopsittacus” OR “Micropsitta” OR “Neophema” OR

2086 “Neopsephotus” OR “Neopsittacus” OR “Northiella” OR “Oreopsittacus” OR “Parvipsitta” OR “Pezoporus”
 2087 OR “Phigys” OR “Platycercus” OR “Polytelis” OR “Prioniturus” OR “Prosopeia” OR “Psephotellus” OR
 2088 “Psephotus” OR “Pseudeos” OR “Psittacella” OR “Psittacula” OR “Psittaculirostris” OR “Psitteuteles” OR
 2089 “Psittinus” OR “Psittirichas” OR “Purpureicephalus” OR “Tanygnathus” OR “Trichoglossus” OR “Vini” OR
 2090 “Nestor” OR “Strigops” OR “Pterocles” OR “Syrnhaptes” OR “Rhea” OR “Aptenodytes” OR “Eudytes” OR
 2091 “Eudiptula” OR “Megadyptes” OR “Pygoscelis” OR “Spheniscus” OR “Aegolius” OR “Asio” OR “Athene”
 2092 OR “Bubo” OR “Glaucidium” OR “Ketupa” OR “Lophotrix” OR “Megascops” OR “Micrathene” OR “Ninox”
 2093 OR “Otus” OR “Pseudoscops” OR “Psiloscops” OR “Pulsatrix” OR “Sceloglaux” OR “Scotopelia” OR “Strix”
 2094 OR “Surnia” OR “Xenoglaux” OR “Phodilus” OR “Tyto” OR “Aepyornis” OR “Mullerornis” OR “Struthio”
 2095 OR “Anhinga” OR “Fregata” OR “Leucocarbo” OR “Microcarbo” OR “Nannopterum” OR “Phalacrocorax”
 2096 OR “Morus” OR “Sula” OR “Crypturellus” OR “Eudromia” OR “Nothocercus” OR “Nothoprocta” OR
 2097 “Nothura” OR “Rhynchotus” OR “Tinamotis” OR “Tinamus” OR “Apalharpactes” OR “Apaloderma” OR
 2098 “Harpactes” OR “Pharomachrus” OR “Trogon” OR “Tetrastes” OR “Hesperiphona”) AND (“experiment*”
 2099 OR “manipulat*”)))>




Figure 2: Overall effect of green nest material on all fitness proxies combined. The meta-analytic mean (black circle), 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along with the data points (coloured circles) scaled by precision (inverse of standard error). Results are shown as an orchard plot for the intercept-only model (A) using log response ratio (lnRR), and (B) standardised mean difference (SMDH). k : number of effect sizes (number of studies is shown in brackets). X-axes are truncated for legibility, which partially explains the apparent asymmetry (see Figure S10).




Figure 3: There is no clear evidence for the courtship or the parental care hypothesis separately as drivers of green nest material fitness effects. Orchard plot of the multilevel meta-regression for (A) log response ratio (lnRR) and (B) standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars), and prediction intervals (thinner bars) are shown along with the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k : number of effect sizes (number of studies is shown in brackets). X-axes are truncated for legibility, which partially explains the apparent asymmetry (see Figure S11).




Figure 4: Experimental designs comparing nests with aromatic green nest material against nests with no green material, and those in which green nest material was experimentally added continuously throughout the nesting phase, lead to the largest fitness estimates. Orchard plots of the multilevel meta-regressions showing the overall effect of green nest material depending on experimental design (A: log response ratio (lnRR); B: standardised mean difference (SMD(H)), and time of green nest material addition (C: lnRR; D: SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along the individual data points (transparantly coloured circles) scaled by precision (inverse of standard error). k : number of effect sizes (number of studies is shown in brackets). X-axes are truncated for legibility, which partially explains the apparent asymmetry (see Figure S12).

Multilevel Meta Regressions for Exploratory Analyses												
Level	lnRR					SMDH						
	Estimate	P-val	95% CI	95% PI	k	n	Estimate	P-val	95% CI	95% PI	k	n
1. Experimental Design												
Non-aromatic vs. Aromatic	-0.019	0.677	[-0.111, 0.072]	[-0.356, 0.317]	122	16	-0.100	0.606	[-0.479, 0.280]	[-1.632, 1.432]	132	16
No added material vs. Aromatic	0.111	0.016	[0.021, 0.200]	[-0.225, 0.446]	90	12	0.712	0.001	[0.309, 1.114]	[-0.826, 2.250]	94	12
No added material vs. Non-aromatic	-0.349	<0.001	[-0.504, -0.195]	[-0.708, 0.009]	10	3	-0.581	0.085	[-1.244, 0.081]	[-2.207, 1.044]	11	3
2. Time of GNM Addition												
After egg hatching	0.028	0.425	[-0.040, 0.095]	[-0.280, 0.335]	52	5	0.267	0.072	[-0.024, 0.559]	[-0.775, 1.310]	57	5
Before egg hatching	-0.002	0.948	[-0.051, 0.048]	[-0.306, 0.303]	91	9	0.061	0.531	[-0.130, 0.251]	[-0.958, 1.080]	91	9
Continuously through nesting phase	0.034	0.165	[-0.014, 0.083]	[-0.270, 0.338]	95	12	0.302	0.011	[0.069, 0.534]	[-0.726, 1.329]	105	12
3. Bird Species												
Buteo buteo	-0.109	0.724	[-0.714, 0.497]	[-0.844, 0.626]	9	1	-0.251	0.732	[-1.691, 1.189]	[-2.263, 1.761]	9	1
Cyanistes caeruleus	0.006	0.969	[-0.288, 0.299]	[-0.504, 0.515]	68	10	0.087	0.866	[-0.926, 1.100]	[-1.646, 1.819]	74	10
Parus major	-0.037	0.815	[-0.347, 0.273]	[-0.556, 0.482]	19	1	0.651	0.275	[-0.521, 1.822]	[-1.178, 2.480]	22	1
Passer cinnamomeus	0.228	0.192	[-0.115, 0.570]	[-0.312, 0.767]	6	1	0.955	0.193	[-0.485, 2.396]	[-1.057, 2.968]	6	1
Sturnus unicolor	-0.011	0.941	[-0.308, 0.286]	[-0.523, 0.500]	44	5	0.069	0.896	[-0.967, 1.105]	[-1.676, 1.815]	45	5
Sturnus vulgaris	0.039	0.794	[-0.255, 0.333]	[-0.471, 0.549]	83	6	0.279	0.590	[-0.742, 1.300]	[-1.457, 2.016]	84	6
Tachycineta bicolor	0.092	0.582	[-0.237, 0.422]	[-0.439, 0.623]	9	2	0.204	0.716	[-0.899, 1.308]	[-1.582, 1.991]	13	2
4. Trait Category												
Behaviour	0.034	0.755	[-0.179, 0.247]	[-0.339, 0.406]	8	2	0.153	0.574	[-0.382, 0.688]	[-0.993, 1.299]	8	2
Morphology	0.036	0.109	[-0.008, 0.080]	[-0.273, 0.345]	62	15	0.310	0.002	[0.112, 0.509]	[-0.723, 1.343]	65	15
Parasitic and Pathogen Related	-0.036	0.328	[-0.107, 0.036]	[-0.350, 0.278]	86	16	0.112	0.242	[-0.076, 0.299]	[-0.919, 1.143]	90	16
Phenology	0.004	0.965	[-0.156, 0.163]	[-0.341, 0.348]	4	3	0.081	0.779	[-0.484, 0.645]	[-1.080, 1.241]	4	3
Physiology	0.026	0.521	[-0.053, 0.105]	[-0.290, 0.342]	22	9	0.157	0.288	[-0.133, 0.448]	[-0.897, 1.212]	23	9
Reproduction	0.010	0.727	[-0.045, 0.065]	[-0.301, 0.320]	56	16	0.153	0.109	[-0.034, 0.340]	[-0.878, 1.184]	63	19
5. Parasite Type												
Arthropod	0.013	0.898	[-0.186, 0.212]	[-0.780, 0.806]	56	13	0.144	0.292	[-0.126, 0.414]	[-1.107, 1.395]	59	13
Micro-organism	-0.041	0.720	[-0.268, 0.186]	[-0.842, 0.760]	26	6	-0.025	0.890	[-0.385, 0.335]	[-1.299, 1.248]	27	6

Table 1: Results from multilevel meta-regressions testing how several factors moderate the effect of green nest material on fitness estimates across birds. Columns show model estimates, p-values (P-val), 95% confidence intervals (CI), 95% prediction intervals (PI), number of effect sizes (k), and number of studies (n) for both effect size metrics: log response ratio (lnRR) and standardized mean difference (SMD(H)). Statistically significant results (p-values < 0.05) are shown in bold.

Completed PRISMA-EcoEvo Checklist [45]: This checklist shows how the systematic review and meta-analysis on green nest material and fitness outcomes complies with reporting standards in ecology and evolutionary biology

Table S1




Figure S1: PRISMA flowchart illustrating the steps and number of studies included in the systematic review and meta-analysis on green nest material and fitness outcomes.

Article Title	Authors	DOI	Journal	Year of Publication
Green incubation: avian offspring benefit from aromatic nest herbs through improved parental incubation behaviour	Gwinner, Helga and Capilla-Lasheras, Pablo and Cooper, Caren and Helm, Barbara	10.1098/rspb.2018.0376	Proceedings of the Royal Society B- Biological Sciences	2018
Telomere length and dynamics of spotless starling nestlings depend on nest-building materials used by parents	Soler, Juan J. and Ruiz-Castellano, Cristina and Figuerola, Jordi and Martin-Vivaldi, Manuel and Martinez-de la Puente, Josue and Ruiz-Rodriguez, Magdalena and Tomas, Gustavo	10.1016/j.anbehav.2017.01.018	Animal Behaviour	2017
Green plants in nests reduce offspring recruitment rates in the spotless starling	Polo, Vicente and Rubalcaba, Juan G. and Veiga, Jose P.	10.1093/beheco/arn056	Behavioural Ecology	2015
Nest size and aromatic plants in the nest as sexually selected female traits in blue tits	Tomas, Gustavo and Merino, Santiago and Martínez-de la Puente, Josue and Moreno, Juan and Morales, Judith and Rivero-de Aguilar, Juan	10.1093/beheco/art015	Behavioural Ecology	2013
Interacting effects of aromatic plants and female age on nest-dwelling ectoparasites and blood-sucking flies in avian nests	Tomas, G. and Merino, S. and Martínez-de la Puente, J. and Moreno, J. and Morales, J. and Lobato, E. and Rivero-de Aguilar, J. and del Cerro, S.	10.1016/j.beproc.2012.02.003	Behavioural Ecology	2012
Experimental Addition of Green Plants to the Nest Increases Testosterone Levels in Female Spotless Starlings	Polo, Vicente and Lopez-Rull, Isabel and Gil, Diego and Veiga, Jose P.	10.1111/j.1439-0310.2009.01724.x	Ethology	2010
Aromatic plants in nests of the blue tit <i>Cyanistes caeruleus</i> protect chicks from bacteria	Mennerat, Adele and Mirleau, Pascal and Blondel, Jacques and Perret, Philippe and Lambrechts, Marcel M. and Heeb, Philipp	10.1007/s00442-009-1418-6	Oecologia	2009
Do female spotless starlings <i>Sturnus unicolor</i> adjust maternal investment according to male attractiveness?	Lopez-Rull, Isabel and Gil, Diego	10.1111/j.1600-048X.2009.04553.x	Journal of Avian Biology	2009
Aromatic plants in nests of blue tits: positive effects on nestlings	Mennerat, Adele and Perret, Philippe and Bourgauff, Patrice and Blondel, Jacques and Gimenez, Olivier and Thomas, Don W. and Heeb, Philipp and Lambrechts, Marcel M.	10.1016/j.anbehav.2008.11.008	Animal Behaviour	2009
Blue tits (<i>Cyanistes caeruleus</i>) respond to an experimental change in the aromatic plant odour composition of their nest	Mennerat, A.	10.1016/j.beproc.2008.07.003	Behavioural Process	2008
Experimental addition of greenery reduces flea loads in nests of a non-greenery using species, the tree swallow <i>Tachycineta bicolor</i>	Shutler, Dave and Campbell, Adam A.	10.1111/j.2007.0908-8857.04015.x	Journal of Avian Biology	2007
European starlings: nestling condition, parasites and green nest material during the breeding season	Gwinner, H and Berger, S	10.1007/s10336-005-0012-x	Journal of Ornithology	2005
Does fresh vegetation protect avian nests from ectoparasites? An experiment with tree swallows	Dawson, RD	10.1139/Z04-076	Canadian Journal of Zoology- Revue Canadienne De Zoologie	2004
Ectoparasitism and the role of green nest material in the European Starling	Fauth, P T and Krementz, D G and Hines, J E	10.1007/BF00328399	Oecologia	1991
Aromatic plants in blue tit <i>Cyanistes caeruleus</i> nests: no negative effect on blood-sucking <i>Protophila</i> blow fly larvae	Mennerat, Adele and Perret, Philippe and Caro, Samuel P. and Heeb, Philipp and Lambrechts, Marcel M.	10.1111/j.0908-8857.2008.04400.x	Journal of Avian Biology	2008

Figure S2: Validation of the search using our own initial library of 15 relevant articles: We validated our search strategy by testing whether it retrieved a pre-compiled set of 15 known studies on green nest material. Searches conducted successfully recovered all 15 articles (100% retrieval), confirming the effectiveness of our search query




Figure S3: **Decision tree used for screening studies during title–abstract and full-text screening process:** The diagram illustrates the stepwise inclusion and exclusion criteria applied to each article during screenings. At each decision point, reviewers selected *include/uncertain* or *exclude*, following pre-registered criteria to ensure consistency.

Summary of all fitness proxies included in the meta-analysis: Each column represents a hypothesis tested in the dataset—Courtship Hypothesis, Parental Care Hypothesis, and Both Hypotheses (studies addressing both). Entries list each unique fitness proxy followed by the number of effect sizes (k, numbers without brackets) and unique articles (n, numbers within brackets) contributing data, formatted as Proxy: k(n). Proxies marked with an asterisk (*) were included only in sensitivity analyses and not in the main meta-analytic models. Summary of fitness proxies completely removed from the meta-analysis or used only in sensitivity analyses: Fitness proxies were excluded when they could not be assigned a consistent directional link to fitness or when the literature on the validity of a proxy for fitness is highly debated. “Completely excluded” proxies were omitted from all models, whereas “Sensitivity-analysis only” proxies were tested separately to evaluate the robustness of the meta-analytic results. Values denote the number of effect sizes (k) and number of studies (n) for each proxy.

Table S2

Summary of all studies included in the meta-analysis: For each study, the table reports the unique paper identifier used in this meta-analysis (**Paper ID**), the full article title (**Title**), the study authors (**Authors**), the journal in which the study was published (**Journal**), the publication year (**Year**), and the digital object identifier (**DOI**). The table includes both published studies and unpublished unpublished dataset and thesis.

Figure S4




Figure S5: **Standardized email template sent to all the authors whose article was included in the meta-analysis:** In cases where we needed the incomplete information, the fitness proxies of interest were mentioned. In all cases, we requested for any unpublished experimental data that could be used for the purpose of our meta-analysis

7.8. Description of Unpublished studies

Figure S6

Complete list of variables extracted from each study: comprehensive overview of all variables recorded during data extraction.

Figure S7




Figure S8: **Alluvial plot showing the overlap among random effects extracted for the meta-analytic dataset:** Each vertical bar represents a random-effect term i.e. Paper ID (28 levels), Experiment ID (48), Group ID (52), Repeated Trait ID (225), Observation ID (273), Bird Species (7), and Population ID (20). Connections between bars indicate shared values across levels. The figure shows that Experiment ID and Group ID were nearly identical, and that Repeated Trait ID had many levels almost as many as the Observation ID, and Repeated Trait ID was excluded from the final models. In contrast, Population ID captured additional clustering across studies and was therefore added to the random effects (not pre-registered).

Sensitivity to sampling correlation (ρ)										
Results for lnRR and SMD(H) Meta-analytic models										
ρ	Overall effect	SE	p	95% CI		LogLik	AIC	BIC	AICc	Effect.size
				CI low	CI high					
0.2	0.0168	0.0131	0.198	-0.0088	0.0424	-15.5399	43.0798	63.8881	43.4450	lnRR
0.3	0.0186	0.0138	0.180	-0.0086	0.0457	-17.4753	46.9505	67.7589	47.3157	lnRR
0.5	0.0190	0.0156	0.222	-0.0115	0.0495	-26.0231	64.0462	84.8546	64.4114	lnRR
0.7	0.0147	0.0181	0.416	-0.0207	0.0501	-43.3232	98.6464	119.4548	99.0117	lnRR
0.8	0.0086	0.0203	0.670	-0.0311	0.0483	-58.3988	128.7976	149.6060	129.1628	lnRR
0.2	0.1330	0.0501	0.008	0.0349	0.2311	-297.6150	607.2301	628.4066	607.5729	SMD(H)
0.3	0.1479	0.0556	0.008	0.0390	0.2568	-304.1710	620.3420	641.5186	620.6849	SMD(H)
0.5	0.1793	0.0665	0.007	0.0489	0.3098	-324.4948	660.9896	682.1661	661.3324	SMD(H)
0.7	0.2560	0.1180	0.030	0.0248	0.4872	-355.0793	722.1585	743.3351	722.5014	SMD(H)
0.8	0.2936	0.1508	0.051	-0.0019	0.5891	-372.2800	756.5600	777.7365	756.9028	SMD(H)

Figure S9: **Sensitivity of overall effect estimates to the assumed sampling correlation (ρ):** The table shows model estimates of the overall effect, standard errors (SE), p -values, 95% confidence intervals (CI), and fit statistics (log-likelihood, AIC, BIC, and AICc) across a range of assumed sampling correlations (ρ) used for the variance-covariance matrices.

Summary of Sensitivity Analyses for lnRR and SMD(H) Intercept-only models							
Effect Size	Data used in model	Estimate	95% CI	95% PI	P-value	k (No. of Effect Sizes)	n (No. of Studies)
lnRR	Including published data only	0.020	[-0.013, 0.052]	[-0.290, 0.329]	0.230	221	24
lnRR	Excluding Experimental Design Non-Aromatic Vs. No added material	0.027	[-0.009, 0.063]	[-0.203, 0.256]	0.147	228	26
lnRR	Excluding Experimental Design Aromatic Vs. Non-Aromatic	0.032	[-0.024, 0.088]	[-0.379, 0.443]	0.258	131	15
lnRR	Including flagged proxies	0.024	[-0.006, 0.053]	[-0.272, 0.319]	0.113	259	28
lnRR	Without those cases where author reported no effect of the treatment	0.019	[-0.013, 0.050]	[-0.291, 0.328]	0.238	222	26
lnRR	Including flagged proxies and without the cases with no effect of the treatment	0.024	[-0.006, 0.054]	[-0.280, 0.328]	0.121	243	28
lnRR	Excluding effect sizes where lnRR failed Geary's Test	0.029	[-0.003, 0.060]	[-0.271, 0.328]	0.074	182	26
SMD(H)	Including published data only	0.192	[0.056, 0.327]	[-0.849, 1.232]	0.006	235	24
SMD(H)	Excluding Experimental Design Non-Aromatic Vs. No added material	0.148	[0.019, 0.278]	[-0.816, 1.112]	0.025	242	26
SMD(H)	Excluding Experimental Design Aromatic Vs. Non-Aromatic	0.220	[0.020, 0.419]	[-0.972, 1.411]	0.031	136	15
SMD(H)	Including flagged proxies	0.191	[0.067, 0.315]	[-0.798, 1.181]	0.003	274	28
SMD(H)	Without those cases where author reported no effect of the treatment	0.186	[0.048, 0.324]	[-0.896, 1.268]	0.008	237	26
SMD(H)	Including flagged proxies and without the cases with no effect of the treatment	0.199	[0.069, 0.329]	[-0.854, 1.251]	0.003	258	28
SMDH	Without the cases where SMDH was calculated from inferential statistics	0.179	[0.044, 0.314]	[-0.856, 1.215]	0.009	245	26

Table S3: Results of sensitivity analyses assessing the robustness of the overall meta-analytic results for lnRR and SMD(H) intercept-only models. We tested whether the main conclusions were sensitive to alternative analytical decisions, including the use of published data only, inclusion of flagged proxies, exclusion of cases reported as having no treatment effect, exclusion of specific experimental-design categories, exclusion of effect sizes calculated from inferential statistics or failing Geary's test. Across analyses, the number of effect sizes ranged from 131 to 274, representing 15 to 28 studies. Overall, the results were qualitatively robust across these alternative model specifications.




Figure S10: Overall effect of green nest material on all fitness proxies combined. The meta-analytic mean (black circle) with 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along with the data points (coloured circles) scaled by precision (inverse of standard error). Results are shown as an orchard plot for the intercept-only model (A) using log response ratio (lnRR), and (B) standardised mean difference (SMD(H)). k : number of effect sizes (number of studies is shown in brackets).




Figure S11: There is no clear evidence to favour either the courtship or the parental care hypothesis as drivers of potential fitness benefits of green nest material. Orchard plot of the multilevel meta-regression for (A) log response ratio (lnRR) and (B) standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars), and prediction intervals (thinner bars) are shown along with the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k : number of effect sizes (number of studies is shown in brackets).




Figure S12: Experimental designs comparing nests with aromatic green nest material against nests without any green material, and those adding the material continuously, lead to the largest fitness estimates. Orchard plots of the multilevel meta-regressions showing the overall effect of green nest material depending on experimental design for (A: log response ratio (lnRR); B: standardised mean difference (SMD(H)), and time of green nest material addition (C: log response ratio (lnRR); D: standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k : number of effect sizes (number of studies is shown in brackets).




Figure S13: Orchard plot of the multilevel meta-regression for (A) log response ratio (lnRR) and (B) standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars), and prediction intervals (thinner bars) are shown along with the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k: number of effect sizes (number of studies is shown in brackets).

Multilevel Meta Regressions for Risk of Bias													
lnRR							SMDH						
Level	Estimate	P-val	95% CI	95% PI	k	n	Estimate	P-val	95% CI	95% PI	k	n	
1. Blinding													
No	0.022	0.168	[-0.009, 0.053]	[-0.278, 0.322]	227	25	0.184	0.007	[0.051, 0.316]	[-0.828, 1.195]	242	25	
Yes	-0.042	0.555	[-0.182, 0.098]	[-0.372, 0.288]	11	1	0.016	0.970	[-0.816, 0.847]	[-1.287, 1.318]	11	1	
2. Random assignment to treatment													
No	0.002	0.958	[-0.080, 0.084]	[-0.309, 0.313]	22	3	0.070	0.705	[-0.292, 0.431]	[-0.999, 1.138]	28	3	
Yes	0.022	0.199	[-0.011, 0.055]	[-0.280, 0.323]	216	23	0.205	0.010	[0.049, 0.360]	[-0.813, 1.223]	225	23	
3. Partial Reporting													
No	0.021	0.205	[-0.011, 0.053]	[-0.280, 0.322]	201	22	0.196	0.007	[0.055, 0.337]	[-0.817, 1.209]	216	22	
Yes	-0.001	0.989	[-0.110, 0.109]	[-0.320, 0.318]	37	4	0.071	0.695	[-0.286, 0.429]	[-0.994, 1.136]	37	4	

Table S4




Figure S14: Tests for publication bias in the effect of green nest material on fitness outcomes. Plots show relationships between effect sizes and (A–B) the inverse of effective sample size ($\sqrt{1/ff}$) testing for small-study effects, and (C–D) publication year (mean-centered) testing for time-lag bias, for log response ratio (lnRR, A and C) and standardized mean difference (SMD(H), B and D). Points represent individual effect sizes, scaled by precision (1/SE). Solid lines show meta-regression slopes, with 95% confidence intervals (dashed lines) and 95% prediction intervals (dotted lines). k : number of effect sizes included in each analysis.

7.16. Systematic Search Information

Figure S15