

Supplementary Appendix

Why do birds use green nest material? A systematic review and meta-analysis of experiments

Shreya Dimri^{1*}, Tuba Rizvi¹, Júlio M.G. Segovia¹, Meinolf Ottensmann², Alfredo Sánchez-Tójar^{1,3,4}

¹Department of Evolutionary Biology, Bielefeld University, Germany

²Department of Animal Behaviour, Bielefeld University, Germany

³CNC, Center for Neuroscience and Cell Biology, University of Coimbra, Portugal

⁴CIBB, Center for Innovative Biomedicine and Biotechnology, University of Coimbra, Portugal

Journal: Proceedings of the Royal Society B

DOI : 10.1098/rspb.2026-0777

Index

Item	Figure/Table	Brief description
Table S1	PRISMA checklist	Completed PRISMA-EcoEvo reporting checklist.
Figure S1	PRISMA flowchart	The steps and number of studies included in the study identification, screening, exclusion, and inclusion process.
Figure S2	Search-validation library	Validation of the search strategy using 15 known relevant articles (initial library).
Figure S3	Screening decision tree	Inclusion and exclusion criteria used during screening.
Table S2	Fitness proxies	Fitness proxies included, excluded, or used only in sensitivity analyses.
Table S3	Included studies	Published and unpublished studies included in the meta-analysis.
Figure S4	Author email template	Standardised email used to request missing or unpublished data.
Text S1	Unpublished studies	Description of the two unpublished datasets included in the meta-analysis.
Text S2	Extracted variables	Details of all the variables recorded during data extraction.
Figure S5	Random effects	Overlap among all possible random-effect terms.
Table S4	Sampling-correlation sensitivity	Sensitivity of model estimates to assumed sampling correlations.
Table S5	Robustness checks	Sensitivity analyses for the intercept-only models.
Figure S6	Overall-effect plots	Overall lnRR and SMD(H) estimates without axis cropping.
Figure S7	Hypothesis models	Moderator models comparing functional hypotheses.
Figure S8	Design and timing models	Moderator models for experimental design and timing of GNM addition.
Figure S9	Additional moderator models	Moderator models for parasite/pathogen type, bird species, and fitness proxy category.
Table S6	Risk-of-bias analyses	Effects of blinding, random assignment, and partial reporting.
Figure S10	Publication-bias plots	Small-study and decline-effect analyses.
Text S3	Search information	Search dates, databases, and complete search strings.

PRISMA Checklist

Completed PRISMA-EcoEvo Checklist (O'Dea et al., 2021): This checklist shows how the systematic review and meta-analysis on green nest material and fitness outcomes complies with reporting standards in ecology and evolutionary biology

Table S 1.

Checklist item	Item score	Sub-item number	Sub-item	Reported by authors?	Notes
Title and abstract	100%	1.1	Identify the review as a systematic review, meta-analysis, or both	Yes	Identified in title and in abstract as both a systematic review and a meta-analysis
		1.2	Summarise the aims and scope of the review	Yes	In the abstract
		1.3	Describe the data set	Yes	In the abstract
		1.4	State the results of the primary outcome	Yes	In the abstract
		1.5	State conclusions	Yes	In the abstract
		1.6	State limitations	Yes	In the abstract
Aims and questions	100%	2.1	Provide a rationale for the review	Yes	Rationale explained in the Introduction in detail
		2.2	Reference any previous reviews or meta-analyses on the topic	Yes	Clearly stated twice in the introduction
		2.3	State the aims and scope of the review (including its generality)	Yes	Scope (only birds and experimental studies) explained along with the aims in the last paragraph of Introduction
		2.4	State the primary questions the review addresses (e.g., which moderators were tested)	Yes	Moderators tested are clearly stated in the methods in the data analysis section.

		2.5	Describe whether effect sizes were derived from experimental and/or observational comparisons	Yes	Only experimental studies were selected, described in the last paragraph of introduction as well as the methods
Review registration	100%	3.1	Register review aims, hypotheses (if applicable), and methods in a time-stamped and publicly accessible archive and provide a link to the registration in the methods section of the manuscript. Ideally registration occurs before the search, but it can be done at any stage before data analysis.	Yes	https://doi.org/10.17605/OSF.IO/S7J6Z
		3.2	Describe deviations from the registered aims and methods	Yes	Described in methods wherever applicable
		3.3	Justify deviations from the registered aims and methods	Yes	When applicable, described in the methods
Eligibility criteria	100%	4.1	Report the specific criteria used for including or excluding studies when screening titles and/or abstracts, and full texts, according to the aims of the systematic review (e.g., study design, taxa, data availability)	Yes	Described briefly in the methods, but mainly in the decision tree (see Supplementary Materials)
		4.2	Justify criteria, if necessary (i.e. not obvious from aims and scope).	Yes	Described along with the decision tree (see Supplementary Materials)
Finding studies	100%	5.1	Define the type of search (e.g., comprehensive search, representative sample)	Yes	Comprehensive search conducted, described in methods along with search string (complete search string provided in the supplementary materials)
		5.2	State what sources of information were sought (e.g., published and unpublished studies, personal communications)	Yes	Published, unpublished as well as personal communications. Described in detail in the Eligibility Criteria and Data Extraction section of Methods Section
		5.3	Include, for each database searched, the exact search strings used, with keyword combinations and Boolean operators	Yes	See complete search string in the Supplementary Materials
		5.4	Provide enough information to repeat the equivalent search (if possible), including the timespan covered (start and end dates)	Yes	See complete search string details in the Supplementary Materials. Search dates are described in the methods.
Study selection	100%	6.1	Describe how studies were selected for inclusion at each stage of the screening process (e.g., use of decision trees, screening software)	Yes	Abstract-Title screening was conducted using the R package revtools GUI and Fulltext screening was conducted in excel manually downloading the articles. Described in detail in the pre-registration and methods section. The decision tree is provided in the supplementary materials.
		6.2	Report the number of people involved and how they contributed (e.g., independent parallel screening)	Yes	Described in detail in the methods section. All screening decision files are shared as .xls/.csv files with notes on the decisions made.
Data collection process	100%	7.1	Describe where in the reports data were collected from (e.g., text or figures)	Yes	In all files after the data extraction (shared along with the project), the column data_location contains this information
		7.2	Describe how data were collected (e.g., software used to digitize figures, external data sources)	Yes	Described in detail in the Data extraction section.
		7.3	Describe moderator variables that were constructed from collected data (e.g., number of generations calculated from years and average generation time)	Yes	All moderators are described in detail in the metadata file of the overall dataset. Please see data_cleaning.Rmd and data_preparation.Rmd files for any cleaning done to the files. All codes are provided along with the dataset.
		7.4	Report how missing or ambiguous information was dealt with during data collection (e.g., authors of original studies were contacted for missing descriptive statistics, and/or effect sizes were calculated from test statistics)	Yes	Described in detail in the data extraction and effect size calculation section in the methods.
		7.5	Report who collected data	Yes	The extractor_ID column in the data file has information on the ID of who collected the data. The methods section also describes the ID of the data collector.
		7.6	State the number of extractions that were checked for accuracy by co-authors		Described in the methods section along with % of disagreement and process of resolution. The data_checker_ID column in the data files has information on the ID of who checked the data.
Data items	100%	8.1	Describe the key data sought from each study	Yes	The priority order of data extraction is described in detail in the Data extraction section of the methods.
		8.2	Describe items that do not appear in the main results, or which could not be extracted due to insufficient information	Yes	All information that is excluded from analysis is described in detail in the methods. Some proxies that could not be included in the analysis for various reasons are reported in the supplementary material along with the reason.
		8.3	Describe main assumptions or simplifications that were made (e.g., categorising both 'length' and 'mass' as 'morphology')	Yes	All major decisions are described in the methods section, for any remaining description please check the notes column in the data files.
		8.4	Describe the type of replication unit (e.g., individuals, broods, study sites)	Yes	Described in the methods, we prioritised nests as the unit of replication whenever possible.
Assessment of individual study quality	100%	9.1	Describe whether the quality of studies included in the systematic review or meta-analysis was assessed (e.g., blinded data collection, reporting quality, experimental versus observational)	Yes	The risk of bias assessment performed for the individual studies is described in the methods and results reported in the results section (See supplementary material for all model output and figures). The risk of bias categories we evaluated were blinding, random assignment to each treatment and partial reporting

		9.2	Describe how information about study quality was incorporated into analyses (e.g., meta-regression and/or sensitivity analysis)	Yes	We conducted three meta-regressions using blinding, random assignment to each treatment and partial reporting as moderators in each analysis. We reported the outcome briefly in the Results (See supplementary material for all model output and figures)
Effect size measures	100%	10.1	Describe effect size(s) used	Yes	Described in Effect Size Calculation section in Methods
		10.2	Provide a reference to the equation of each calculated effect size (e.g., standardised mean difference, log response ratio) and (if applicable) its sampling variance	Yes	Described in the Effect Size Calculation section in Methods. Check data_preparation.Rmd in Code to see the exact function, its parameters and any equations that were used to estimate the effect sizes.
		10.3	If no reference exists, derive the equations for each effect size and state the assumed sampling distribution(s)	Yes	Does not apply. Any custom equations used from books etc. are clearly mentioned as well as referenced in the Methods section.
Missing data	100%	11.1	Describe any steps taken to deal with missing data during analysis (e.g., imputation, complete case, subset analysis)	Yes	All imputations conducted for the missing data are stated clearly in the Effect Size Calculations in Methods Section
		11.2	Justify the decisions made to deal with missing data	Yes	We decided to impute the missing data but also conducted the sensitivity analysis excluding the imputed value. All justifications are clearly stated in the methods.
Meta-analytic model description	100%	12.1	Describe the models used for synthesis of effect sizes	Yes	Described all models in Data Analysis section of the Methods
		12.2	The most common approach in ecology and evolution will be a random-effects model, often with a hierarchical/multilevel structure. If other types of models are chosen (e.g., common/fixed effects model, unweighted model), provide justification for this choice	Yes	Not applicable, Multilevel meta analytical model and meta-regressions were conducted. No justification needed
Software	100%	13.1	Describe the statistical platform used for inference (e.g., R)	Yes	R v4.5.1 was used for the analysis and is clearly stated in the Methods.
		13.2	Describe the packages used to run models	Yes	All packages and their versions are described in the Methods.
		13.3	Describe the functions used to run models	Yes	All important functions used for running the models are stated in the Methods.
		13.4	Describe any arguments that differed from the default settings	Yes	We have clearly stated all the important arguments that differed from the default settings. However, see data_analysis.Rmd for full description of the arguments for each model.
		13.5	Describe the version numbers of all software used	Yes	All packages and their versions are described in the Methods.
Non-independence	100%	14.1	Describe the types of non-independence encountered (e.g., phylogenetic, spatial, multiple measurements over time)	Yes	See data analysis for more details on how non-independence was handled.
		14.2	Describe how non-independence has been handled	Yes	See data analysis for more details on how non-independence was handled.
		14.3	Justify decisions made	Yes	See data analysis for more details on how non-independence was handled.
Meta-regression and model selection	100%	15.1	Provide a rationale for the inclusion of moderators (covariates) that were evaluated in meta-regression models	Yes	All the moderators that were evaluated were pre-registered with the rationale behind it clearly stated. See Data analysis for more details (but also see pre-registration).
		15.2	Justify the number of parameters estimated in models, in relation to the number of effect sizes and studies (e.g., interaction terms were not included due to insufficient sample sizes)	Yes	We described the random effects that were not included in the models in the Methods section
		15.3	Describe any process of model selection	Yes	No model selection was conducted
Publication bias and sensitivity analysis	100%	16.1	Describe assessments of the risk of bias due to missing results (e.g., publication, time-lag, and taxonomic biases)	Yes	Time-lag bias and small study bias are reported in the methods, results and discussion.
		16.2	Describe any steps taken to investigate the effects of such biases (if present)	Yes	We discussed the effects of potential biases and conducted a sensitivity analysis accounting these effects for the two main models.
		16.3	Describe any other analyses of robustness of the results, e.g., due to effect size choice, weighting or analytical model assumptions, inclusion or exclusion of subsets of the data, or the inclusion of alternative moderator variables in meta-regressions	Yes	All sensitivity analysis are mentioned in the methods, results and discussed where appropriate. All the results from the sensitivity analysis/robustness checks are reported as a table in the supplementary material.
Clarification of post hoc analyses	100%	17.1	When hypotheses were formulated after data analysis, this should be acknowledged.	Yes	All analysis are pre-registered and exploratory analysis are clearly stated under exploratory analysis both in the pre-registration and final article
Metadata, data, and code	100%	18.1	Share metadata (i.e. data descriptions)	Yes	See data and code availability statement for details.
		18.2	Share data required to reproduce the results presented in the manuscript	Yes	See data and code availability statement for details.

		18.3	Share additional data, including information that was not presented in the manuscript (e.g., raw data used to calculate effect sizes, descriptions of where data were located in papers)	Yes	See data and code availability statement for details.
		18.4	Share analysis scripts (or, if a software package with graphical user interface (GUI) was used, then describe full model specification and fully specify choices)	Yes	See data and code availability statement for details.
Results of study selection process	100%	19.1	Report the number of studies screened	Yes	Details in method and the PRISMA Flowchart in supplementary material
		19.2	Report the number of studies excluded at each stage of screening	Yes	Details in method and the PRISMA Flowchart in supplementary material
		19.3	Report brief reasons for exclusion from the full text stage	Yes	Details in the PRISMA Flowchart in supplementary material
		19.4	Present a Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)-like flowchart (www.prisma-statement.org).	Yes	Figure S1 in supplementary materials is a PRISMA Flowchart
Sample sizes and study characteristics	100%	20.1	Report the number of studies and effect sizes for data included in meta-analyses	Yes	Details in method and supplementary tables
		20.2	Report the number of studies and effect sizes for subsets of data included in meta-regressions	Yes	Details in method and supplementary tables
		20.3	Provide a summary of key characteristics for reported outcomes (either in text or figures; e.g., one quarter of effect sizes reported for vertebrates and the rest invertebrates)	Yes	All figures contain effect sizes (k) and unique articles (n) for different moderator levels. Summary of reported outcomes is also in the first paragraph of the Results.
		20.4	Provide a summary of limitations of included moderators (e.g., collinearity and overlap between moderators)	NA	All moderators tested were categorical and only publication year was a continuous moderator. Therefore, no collinearity or overlap is possible.
		20.5	Provide a summary of characteristics related to individual study quality (risk of bias)	Yes	This data is presented in dataset in columns blinding, missing_data, random_assignment
Meta-analysis	100%	21.1	Provide a quantitative synthesis of results across studies, including estimates for the mean effect size, with confidence/credible intervals	Yes	See results and supplementary material for detail of all the synthesis.
Heterogeneity	100%	22.1	Report indicators of heterogeneity in the estimated effect (e.g., I ² , tau ² and other variance components)	Yes	All estimates of heterogeneity are reported in the results and model table. Also see supplementary material for the results not reported in the main text.
Meta-regression	100%	23.1	Provide estimates of meta-regression slopes (i.e. regression coefficients) and confidence/credible intervals	Yes	See results and supplementary material for detail of all the synthesis.
		23.2	Include estimates and confidence/credible intervals for all moderator variables that were assessed (i.e. complete reporting)	Yes	See results and supplementary material for detail of all the synthesis.
		23.3	Report interactions, if they were included	NA	There were no interactions included
		23.4	Describe outcomes from model selection, if done (e.g., R ² and AIC)	NA	No model selection was conducted.
Outcomes of publication bias and sensitivity analysis	100%	24.1	Provide results for the assessments of the risks of bias (e.g., Egger's regression, funnel plots)	Yes	Risk of bias was assessed using meta-regression models with moderator levels for blinding, missing data, random assignment in experiments and reported in results (and see supplementary material for complete details). We also reported meta-regressions for publication bias (time-lag bias and small-study effects) in results (and see supplementary material for complete details)
		24.2	Provide results for the robustness of the review's results (e.g., subgroup analyses, meta-regression of study quality, results from alternative methods of analysis, and temporal trends)	Yes	All robustness checks/sensitivity analysis conducted are reported in full detail in the supplementary material and mentioned in the results clearly.
Discussion	100%	25.1	Summarise the main findings in terms of the magnitude of effect	Yes	
		25.2	Summarise the main findings in terms of the precision of effects (e.g., size of confidence intervals, statistical significance)	Yes	We use confidence intervals, prediction intervals and statistical significance to discuss the precision of the effects in the results and discussion section.
		25.3	Summarise the main findings in terms of their heterogeneity	Yes	We report and discuss the heterogeneity and explore some of the possible causes of it in our results and discussion.
		25.4	Summarise the main findings in terms of their biological/practical relevance	Yes	We have discussed all important findings in terms of their biological and practical implications
		25.5	Compare results with previous reviews on the topic, if available	Yes	We have discussed the results comparing them to other published literature (including two narrative reviews that summarise the field) but no quantitative review on the field exists to compare the results with.
		25.6	Consider limitations and their influence on the generality of conclusions, such as gaps in the available evidence (e.g., taxonomic and geographical research biases)	Yes	We discuss the limitations of the evidence.

Contributions and funding	100%	26.1	Provide names, affiliations, and funding sources of all co-authors	Yes	See section Fundings
		26.2	List the contributions of each co-author	Yes	See section Author Contributions
		26.3	Provide contact details for the corresponding author	Yes	See manuscript Author details: Shreya Dimri - shreya.dimri@uni-bielefeld.de
		26.4	Disclose any conflicts of interest	Yes	See section Conflict of interest
References	100%	27.1	Provide a reference list of all studies included in the systematic review or meta-analysis	Yes	See supplementary material for a list.
		27.2	List included studies as referenced sources (e.g., rather than listing them in a table or supplement)	Yes	We have included all the studies in our meta-analysis within the text of the article in appropriate places.




Figure S 1. PRISMA flowchart illustrating the steps and number of studies included in the systematic review and meta-analysis on green nest material and fitness outcomes.

Initial library used for search validation

Article Title	Authors	DOI	Journal	Year of Publication
Green incubation: avian offspring benefit from aromatic nest herbs through improved parental incubation behaviour	Gwinner, Helga and Capilla-Lasheras, Pablo and Cooper, Caren and Helm, Barbara	10.1098/rspb.2018.0376	Proceedings of the Royal Society B- Biological Sciences	2018
Telomere length and dynamics of spotless starling nestlings depend on nest-building materials used by parents	Soler, Juan J. and Ruiz-Castellano, Cristina and Figuerola, Jordi and Martin-Vivaldi, Manuel and Martinez-de la Puente, Josue and Ruiz-Rodriguez, Magdalena and Tomas, Gustavo	10.1016/j.anbehav.2017.01.018	Animal Behaviour	2017
Green plants in nests reduce offspring recruitment rates in the spotless starling	Polo, Vicente and Rubalcaba, Juan G. and Veiga, Jose P.	10.1093/beheco/arv056	Behavioural Ecology	2015
Nest size and aromatic plants in the nest as sexually selected female traits in blue tits	Tomas, Gustavo and Merino, Santiago and Martinez-de la Puente, Josue and Moreno, Juan and Morales, Judith and Rivero-de Aguilar, Juan	10.1093/beheco/art015	Behavioural Ecology	2013
Interacting effects of aromatic plants and female age on nest-dwelling ectoparasites and blood-sucking flies in avian nests	Tomas, G. and Merino, S. and Martinez-de la Puente, J. and Moreno, J. and Morales, J. and Lobato, E. and Rivero-de Aguilar, J. and del Cerro, S.	10.1016/j.beproc.2012.02.003	Behavioural Ecology	2012
Experimental Addition of Green Plants to the Nest Increases Testosterone Levels in Female Spotless Starlings	Polo, Vicente and Lopez-Rull, Isabel and Gil, Diego and Veiga, Jose P.	10.1111/j.1439-0310.2009.01724.x	Ethology	2010
Aromatic plants in nests of the blue tit <i>Cyanistes caeruleus</i> protect chicks from bacteria	Mennerat, Adele and Mirleau, Pascal and Blondel, Jacques and Perret, Philippe and Lambrechts, Marcel M. and Heeb, Philipp	10.1007/s00442-009-1418-6	Oecologia	2009
Do female spotless starlings <i>Sturnus unicolor</i> adjust maternal investment according to male attractiveness?	Lopez-Rull, Isabel and Gil, Diego	10.1111/j.1600-048X.2009.04553.x	Journal of Avian Biology	2009
Aromatic plants in nests of blue tits: positive effects on nestlings	Mennerat, Adele and Perret, Philippe and Bourgault, Patrice and Blondel, Jacques and Gimenez, Olivier and Thomas, Don W. and Heeb, Philipp and Lambrechts, Marcel M.	10.1016/j.anbehav.2008.11.008	Animal Behaviour	2009
Blue tits (<i>Cyanistes caeruleus</i>) respond to an experimental change in the aromatic plant odour composition of their nest	Mennerat, A.	10.1016/j.beproc.2008.07.003	Behavioural Process	2008
Experimental addition of greenery reduces flea loads in nests of a non-greenery using species, the tree swallow <i>Tachycineta bicolor</i>	Shutler, Dave and Campbell, Adam A.	10.1111/j.2007.0908-8857.04015.x	Journal of Avian Biology	2007
European starlings: nestling condition, parasites and green nest material during the breeding season	Gwinner, H and Berger, S	10.1007/s10336-005-0012-x	Journal of Ornithology	2005
Does fresh vegetation protect avian nests from ectoparasites? An experiment with tree swallows	Dawson, RD	10.1139/Z04-076	Canadian Journal of Zoology- Revue Canadienne De Zoologie	2004
Ectoparasitism and the role of green nest material in the European Starling	Fauth, P.T and Kremenz, D.G and Hines, J.E	10.1007/BF00328399	Oecologia	1991
Aromatic plants in blue tit <i>Cyanistes caeruleus</i> nests: no negative effect on blood-sucking <i>Protophila</i> blow fly larvae	Mennerat, Adele and Perret, Philippe and Caro, Samuel P. and Heeb, Philipp and Lambrechts, Marcel M.	10.1111/j.0908-8857.2008.04400.x	Journal of Avian Biology	2008

Figure S 2. Validation of the search using our own initial library of 15 relevant articles: We validated our search strategy by testing whether it retrieved a pre-compiled set of 15 known studies on green nest material. Searches conducted successfully recovered all 15 articles (100% retrieval), confirming the effectiveness of our search query

Decision Tree




Figure S 3. Decision tree used for screening studies during title–abstract and full-text screening process:

The diagram illustrates the stepwise inclusion and exclusion criteria applied to each article during screenings. At each decision point, reviewers selected *include/uncertain* or *exclude*, following pre-registered criteria to ensure consistency.

Fitness proxies included in or excluded from the meta-analysis

Summary of all fitness proxies included in the meta-analysis: Each column represents a hypothesis tested in the dataset—Courtship Hypothesis, Parental Care Hypothesis, and Both Hypotheses (studies addressing both). Entries list each unique fitness proxy followed by the number of effect sizes (k, numbers without brackets) and unique articles (n, numbers within brackets) contributing data, formatted as Proxy: k(n). Proxies marked with an asterisk (*) were included only in sensitivity analyses and not in the main meta-analytic models. Summary of fitness proxies completely removed from the meta-analysis or used only in sensitivity analyses: Fitness proxies were excluded when they could not be assigned a consistent directional link to fitness or when the literature on the validity of a proxy for fitness is highly debated. “Completely excluded” proxies were omitted from all models, whereas “Sensitivity-analysis only” proxies were tested separately to evaluate the robustness of the meta-analytic results. Values denote the number of effect sizes (k) and number of studies (n) for each proxy.

Table S 2.

Courtship Hypothesis			
Fitness Proxy	Effect Size(studies)	Definition (as measures/provided by authors of primary study)	Expected Direction (as described or expected by authors of primary study)
Sex Ratio*	9(2)	Sex Ratio is the hatching broods sex ratio measures as the proportion of male offspring in a breeding attempt.	In these experiments, adding fresh GNM is treated as a courtship signal. Under sex-allocation as described by the authors, females paired to more attractive males (that add more GNM), are expected to bias investment towards the sex with higher potential fitness returns, here assumed to be sons. So the predicted effect is a more male-biased sex ratio in nests with GNM (Polo et al., 2004; Veiga et al., 2008). However, we only include them in a sensitivity analysis since the evidence for whether birds can change sex ratio is debated (see Krackow, S. (2002), Why Parental Sex Ratio Manipulation is Rare in Higher Vertebrates (Invited Article). <i>Ethology</i> , 108: 1041-1056. https://doi.org/10.1046/j.1439-0310.2002.00843.x).
Females Testosterone*	4(2)	Circulating (plasma) testosterone concentration of breeding females (during laying or early incubation), quantified from a blood sample collected during the breeding attempt and assayed using ELISA.	In these experiments author predicted higher female testosterone in GNM-added nests. Experimental GNM addition is treated as increasing the perceived attractiveness/quality of the male (i.e., a courtship signal). The authors predict (and test) that this should elevate female testosterone because (i) females may increase competitive/aggressive capacity to maintain access to preferred males or nest sites under heightened female–female competition, and/or (ii) female endocrine state may shift as part of differential allocation when paired to (perceived) attractive males. This is described as a mechanistic mediator that would show competitive state of the females, which could indirectly affect reproductive outcomes (e.g., maintaining access to higher quality males thus benefiting the offspring or altering maternal investment strategies). (López-Rull & Gil, 2009; Polo et al., 2010). However, we only include them in a sensitivity analysis because in these studies, only one measure of testosterone was taken per female and the values of testosterone can considerably change within the time of the day or phase of breeding, for consistency with other physiological measures like blood glucose, which we only test in a sensitivity analysis, we also excluded it. Furthermore, an increase in female testosterone does not have clear fitness benefits associated with it.
Courtship Time	3(1)	Interval between first appearance of greenery to first egg.	Under the courtship hypothesis (greenery as a sexually selected ornament/mate-attraction cue), the expected relationship is that GNM should facilitate mate attraction and progression to laying, so addition would be expected to shorten courtship time (i.e., faster progression to laying). Early laying of eggs can then have several benefits for the offspring. (Brouwer & Komdeur, 2004)
Male Provisioning Rate	2(1)	Male provisioning rate is the male's feeding visit rate to nestlings, measured as the number of visits per hour to the nest box while provisioning chicks.	A higher male provisioning rates in nests supplied with aromatic plants (GNM) is expected. In this experimental context, the amount of aromatic plant material in the nest is treated as a female sexual signal that could elicit differential allocation by males. If males interpret aromatic plant addition as indicating higher female quality or higher expected reproductive value of the brood, it would predict increased male parental investment, measured here as increased provisioning rate. (Tomás et al., 2013)
Male Visit	2(1)	A participation or attendance measure indicating whether the male was present and entered the nest-box to provision nestlings.	A higher male visit in nests supplied with aromatic plants (GNM) is expected. In this experimental context, the amount of aromatic plant material in the nest is treated as a female sexual signal that could elicit differential allocation by males. If males interpret aromatic plant addition as indicating higher female quality or higher expected reproductive value of the brood, it would predict increased male parental investment, measured here as increased male visit to the nest. (Tomás et al., 2013)
Yolk Hormones*	2(1)	Yolk androgen concentrations measured from eggs (e.g., testosterone (T), androstenedione (A4))	A higher yolk androgen deposition is expected in experimental nests with added green plant material (GNM). Under the courtship hypothesis, females paired to more attractive males (who bring more GNM) are predicted to increase pre-laying maternal investment, including transferring higher androgen levels to eggs. (López-Rull & Gil, 2009). However, we only include them in a sensitivity analysis because, according to a meta-analysis on yolk hormone and fitness, there is no strong evidence for such a relationship. (Mentesana, L., Hau, M., D'Amelio, P.B., Adreani, N.M. and Sánchez-Tójar, A. (2025), Do Egg Hormones Have Fitness Consequences in Wild Birds? A Systematic Review and Meta-Analysis. <i>Ecology Letters</i> , 28: e70100. https://doi.org/10.1111/ele.70100)
Male Risk Taking	1(1)	Measured as a composite index of male parental risk-taking/investment, constructed as the mean of many binomial behavioural components spanning the nestling period. Higher scores indicate greater willingness to incur risk to attend the nest under disturbance or predation cues.	The authors interpret entering the nest-box under potential danger as risky and treat male risk-taking as an index of male parental investment. Under the courtship hypothesis, aromatic plant supplementation (a female trait/signal in this case) is expected to elicit greater male reproductive investment, expressed here as increased risk-taking. (Tomás et al., 2013)
Parental Care	Hypothesis		
Fitness Proxy	Effect Size(studies)	Definition (as measures/provided by authors of primary study)	Expected Direction (as described or expected by authors of primary study)
Ectoparasite Abundance/Load	Mites: 22(7) Blowflies number: 8(6) Fleas number: 6(6) Carnusflies load: 5(2)	These are the different measures of the amount of ectoparasites found in the nests or on the nestlings. Many different ectoparasites have been measured in different studies.	It is expected that the addition of GNM would reduce the amount/abundance/load of ectoparasites in the nests, thereby increasing the fitness of the nestlings. These are considered the most direct measures of the nest protection hypothesis.

	Blackflies number: 2(1) Midges number: 2(1) Beetles: 1(1) Blowfly Parasitoids Number: 1(1) Booklice And Barklice: 1(1) Butterflies And Moths: 1(1) Clown Beetles: 1(1) Flies: 1(1) Lice Score: 1(1) Rove Beetles: 1(1) Skin Beetles: 1(1) Springtails: 1(1) Tick Load: 1(1) Ticks: 1(1) True Bugs: 1(1) Wasps, Bees, And Ants: 1(1)		
Microorganism Abundance/ Load/Community metrics	Bacterial Load: 20(4) Bacterial Increase: 2(1) Bacterial Richness: 2(1) Leucocytozoon Prevalence: 3(1)	Different measures of microbial load, including increase or richness, are present in the nests or on the nestlings.	It is expected that the addition of GNM would reduce the diversity/abundance/ load of microbes in the nests, thereby increasing the fitness of the nestlings. These are considered the most direct measures of the nest protection hypothesis.
Hemoglobin Measure	9(6)	Hemoglobin concentration in blood is a measure of oxygen-carrying capacity and the general physiological condition of nestlings.	It is expected that the addition of GNM would increase the Hemoglobin Measure. Increased hemoglobin (improved physiological condition) under the parental care hypothesis, because of reduced parasitism/pathogen pressure and/or pharmacological benefits, would support a better physiological state.
Hematocrit %	1(1)	The percentage of blood volume composed of red blood cells is another indicator of oxygen transport and the physiological condition of the nestlings.	Increased hematocrit (improved physiological condition) under the parental care hypothesis, because of reduced parasitism/pathogen pressure and/or pharmacological benefits, would support a better physiological state.
Scab Score	5(3)	Score quantifying skin lesions/scabs on nestlings, typically interpreted as damage from ectoparasites.	With increased GNM we expect reduced scab score because of reduced ectoparasite load or reduced inflammation from secondary infection.
Telomere Measure	4(1)	Measure of telomere length (but in one case, telomere dynamics).	We expect a positive relation between telomere length and GNM. Telomere length shortens as the individual ages and the cell dies once there is not enough protection provided by the telomere cap. Therefore, the shorter the telomere, the worse they are in their fitness. We expect a negative relationship between telomere dynamics and GNM. Telomere dynamics are calculated as the rank difference in telomere length.
Nestling Immunity	2(2)	A direct measure of immune function (varies by study; e.g., PHA swelling response) intended to reflect immune performance/competence.	Increased immune response when GNM is added through the direct pharmacological effect of GNM, thus enabling a better immune response.
Glucose Measure*	2(2)	Blood glucose concentration, a metabolic indicator reflecting energetic state and/or physiological stress.	Authors of the primary studies expect the addition of GNM to decrease the value of Glucose concentration in the blood, i.e., it is closer to the baseline level, showing less stress. However, we only include them in a sensitivity analysis, as Glucose Measure is highly context-dependent. We do not consider glucose concentration to be a meaningful fitness proxy because (1) the blood samples of the birds were collected at random time of the day without any strict protocol for between sampling differences. Blood glucose levels could change very rapidly depending on the consumption of food and there could be great between individual variation in the population. Circadian rhythm could also influence the blood sugar levels. (2) Authors themselves call this a less robust proxy - "The level of blood glucose is considered a reverse (lower concentration of glucose is related to higher body condition) and a less robust (more variable) indicator of body condition than haemoglobin but is also useful in studies of ecophysiology"
For data from	Tachycineta bicolor (Because addition)	of GNM is not shown by this species,	these proxies were treated as parental-care proxies only)
Nestling Mass	1(1)	Body mass of nestlings measured at a different ages in Tachycineta bicolor.	Increased nestling mass would be expected with addition of GNM in Tachycineta bicolor. Since this bird does not show this behaviour in wild, the effect of GNM cannot be attributed to courtship hypothesis.
Nestling Size	1(1)	A composite size index measured in nestlings using other structural size metrics like tarsus length, wing length.	Increased nestling size would be expected with addition of GNM in Tachycineta bicolor. Since this bird does not show this behaviour in wild, the effect of GNM cannot be attributed to courtship hypothesis.
Number of Fledglings	1(1)	Count of young that fledge per nesting attempt.	If GNM reduces parasite/pathogen pressure and improves nestling condition, we should see an increased survival to fledgeling with GNM addition.
Number of Nestlings	1(1)	Count of nestlings present/survive to a certain age.	If GNM reduces early mortality, we expect more chicks to remain alive during the nestling period.
Reproductive Success	3(1)	Percentage of nests with complete reproductive success in Tachycineta bicolor.	With the addition of GNM, we should see increased reproductive success through reduced mortality.
Clutch Size	1(1)	Number of eggs laid in a clutch in Tachycineta bicolor.	With the addition of GNM, we should see an increased clutch size.

Both Hypothesis			
Fitness Proxy	Effect Size(studies)	Definition (as measures/provided by authors of primary study)	Expected Direction (as described or expected by authors of primary study)
Growth, size or condition of offspring	Nestling Mass: 20(9) Chick Mass: 16(4) Nestling Tarsus Length: 13(5) Fledgeling Mass: 7(2) Nestling Wing Length: 2(2) Fledgeling Tarsus-Metatarsus Lengths (Females): 1(1) Fledgeling Tarsus-Metatarsus Lengths (Males): 1(1) Body Condition Estimate: 3(1) Fat Score: 1 (1) Mouth Colouration: 3 (1) Chick Feather Development: 2 (1)	These proxies capture various measurements of offspring growth, structural size and development condition by measuring body mass, skeletal measures (e.g., tarsus/wing length), integrated body condition indices, fat reserves, and feather development, all of which reflect the developmental quality of the nestling at various measurement ages.	We expect GNM to have a positive effect on growth, size and body condition of the offspring. This could be because of reduced pathogens or drug effect, but also because of investment of parents through various mechanisms. Therefore, it is not possible to categorise this into a clear mechanism.
Reproductive Output/Success	Clutch Size: 16(8) Number Of Nestling: 7(5) Fledgling Success: 4 (4) Hatching Success: 4 (3) Local Recruitment Probability: 4 (1) Survival To Fledgeling: 3(1) Nest Mortality: 1 (1) Percent Survived: 1(1) Survival Per Brood: 1(1) Survival Rate Nestling: 1(1)	These proxies quantify the reproductive output and survival (hatching, fledgling, nest failure and recruitment) across different stages of breeding.	GNM is predicted to increase productivity and survival and decrease mortality. These proxies of fitness could improve via increased parental investment, better genetic makeup or through lower parasites in the nests. Therefore, it is difficult to categorise these proxies into a particular mechanism.
Laying Date	3(3)	The date of the clutch initiation (first egg laid), expressed as a relative count from the reference date.	Since earlier laying is often associated with higher reproductive output, we expected GNM would lead to advancement in laying date (i.e., smaller values of laying date proxy).
Hatching Date	1(1)	The date of the first egg hatched, expressed as a relative count from the reference date.	Earlier hatching can have several benefits due to the availability of better quality food; we expected GNM would lead to advancement in hatching date (i.e., smaller values of hatching date proxy).
Clutch Development Score	2(1)	It is a measure of how far embryonic development has progressed (egg, wet hatchling, dry hatchling), at a standardized time late in incubation.	Earlier development on a fixed day is interpreted as a benefit due to reduced exposure time to risks during incubation and improved postnatal condition. Therefore, it is expected that GNM would increase the clutch development score.
Egg Size	1(1)	As a measure of egg volume, derived from the length and width of the egg calculated as $0.45 \times L \times W^2$.	It is expected that GNM would lead to females laying larger eggs.
Feeding Rate*	1(1)	Measure of parental provisioning, quantified from nest-box video as the total number of feeding events standardised per hour during the recording period.	We only test this in a sensitivity analysis, because the relation of this to fitness can be argued. A higher feeding rate can indicate better care and resource collection, which might suggest higher fitness of the parents or brood. However, if feeding is high, it could also imply poor quality of food availability or compensation for the poor health of the nestling.
Time Spent Begging By Broods*	1(1)	Total begging duration, summed across nestlings, divided by brood size, and expressed as seconds of begging per hour of recording (s/h).	If GNM reduces parasite/pathogen burden and/or improves nestling condition, nestlings are expected to show reduced need, expressed as less begging time. However, we only test this in a sensitivity analysis because it is difficult and contentious to interpret. Begging is generally a signal that chicks are hungry and in need of care. More begging might indicate a greater demand for resources and therefore potentially less fitness in terms of immediate brood condition, but studies also suggest that begging intensity is an evolved behaviour to maximise parental care and could imply it is not related to fitness but rather a strategy to elicit care.
Fitness proxies	excluded from the	meta-analysis	
Fitness Proxy	Effect Size(studies)		Reason for exclusion
Male Mass	3(1)		Variables such as adult male or female mass as a fitness proxy are tricky. On one hand, adults in better condition could provide better parental care for their

			offspring and be more likely to survive to reproduce the following season, thereby increasing their overall fitness. However, whether adult mass is directly related to fitness remains unclear.
Female Body Condition	2(1)		The authors used female body condition simply to control for its effect. We are excluding it, as the relationship between adult body condition and GNM is rather tricky to interpret and difficult to describe in terms of a true direction of effect.
Female Provisioning Rate	2(1)		We will exclude this variable from all our analyses. Since females add the GNM as noted by the authors, it is the male that should adjust his investment accordingly. One could argue that female investment will be an indirect reflection of the male's investment; however, this relationship is very indirect, and the direction of this proxy in relation to the GNM addition can be argued both ways. That is, if females use GNM as an honest signal to increase male investment, their own investment would not be directly related to fitness. Alternatively, they might increase investment to compensate for a lack of male investment, or it could be negatively related to fitness because they invested both in bringing in the GNM and in higher provisioning. Therefore, we excluded it.
Measures of WBC (White Blood Cells)	Number of Basophils: 2(1) Number of Lymphocytes: 2(1) Eosinophils of Nestlings: 1(1) Heterophils of Nestlings: 1(1) Nestling Leukocytes: 1(1) WBC measures of Nestlings: 1(1)		It is very difficult to assign a direction of the relationship. Does an increase in WBC imply better or worse fitness? We consulted several researchers in the field and there was no clear consensus on this. Therefore, we decided to exclude it. For further details, see the discussion provided in the project materials.

Studies included in the meta-analysis

Summary of all studies included in the meta-analysis: For each study, the table reports the unique paper identifier used in this meta-analysis (**Paper ID**), the full article title (**Title**), the study authors (**Authors**), the journal in which the study was published (**Journal**), the publication year (**Year**), and the digital object identifier (**DOI**). The table includes both published studies and unpublished unpublished dataset and thesis.

Table S 3.

Paper ID	Title	Authors	Journal	Year	DOI
GNM_001	Consequences of experimental addition of fresh, aromatic plants into nests of blue tits (<i>Cyanistes caeruleus</i>) on the physiological condition of nestlings	Gładalski M; Kaliński A; Wawrzyniak J; Bańbura M; Markowski M; Skwarska J; Bańbura J	Behavioral Ecology and Sociobiology	2020	10.1007/s00265-020-2812-7
GNM_002	Development of nestling blue tits (<i>Cyanistes caeruleus</i>) is affected by experimental addition of aromatic plants	Pires BA; Belo ADF; Diamantino F; Rabaça JE; Merino S	Avian Biology Research	2020	10.1177/1758155920921075
GNM_010	'Green incubation': avian offspring benefit from aromatic nest herbs through improved parental incubation behaviour	Gwinner H; Capilla-Lasheras P; Cooper C; Helm B	Proceedings of the Royal Society B: Biological Sciences	2018	10.1098/rspb.2018.0376
GNM_016	Green plants in nests reduce offspring recruitment rates in the spotless starling	Polo V; Rubalcaba JG; Veiga JP	Behavioral Ecology	2015	10.1093/beheco/arv056
GNM_018	Nest size and aromatic plants in the nest as sexually selected female traits in blue tits	Tomás G; Merino S; Martínez-de la Puente J; Moreno J; Morales J; Rivero-de Aguilar J	Behavioral Ecology	2013	10.1093/beheco/art015
GNM_023	Experimental Addition of Green Plants to the Nest Increases Testosterone Levels in Female Spotless Starlings	Polo V; López-Rull I; Gil D; Veiga JP	Ethology	2010	10.1111/j.1439-0310.2009.01724.x
GNM_024	Do female spotless starlings (<i>Sturnus unicolor</i>) adjust maternal investment according to male attractiveness?	López-Rull I; Gil D	Journal of Avian Biology	2009	10.1111/j.1600-048X.2009.04553.x
GNM_028	Age dependent sex allocation in the polygynous spotless starling	Veiga JP; Polo V; Cordero PJ	Evolutionary Ecology	2008	10.1007/s10682-007-9166-8
GNM_032	Female starlings adjust primary sex ratio in response to aromatic plants in the nest	Polo V; Veiga JP; Cordero PJ; Viñuela J; Monaghan P	Proceedings of the Royal Society of London. Series B: Biological Sciences	2004	10.1098/rspb.2004.2801
GNM_069	Telomere length and dynamics of spotless starling nestlings depend on nest-building materials used by parents	Soler JJ; Ruiz-Castellano C; Figuerola J; Martín-Vidal M; Martínez-de la Puente J; Ruiz-Rodríguez M; Tomás G	Animal Behaviour	2017	10.1016/j.anbehav.2017.01.018
GNM_101	Interacting effects of aromatic plants and female age on nest-dwelling ectoparasites and blood-sucking flies in avian nests	Tomás G; Merino S; Martínez-de la Puente J; Moreno J; Morales J; Lobato E; Rivero-de Aguilar J; del Cerro S	Behavioural Processes	2012	10.1016/j.beproc.2012.02.003
GNM_121	Aromatic plants in nests of the blue tit <i>Cyanistes caeruleus</i> protect chicks from bacteria	Mennerat A; Mirléau P; Blondel J; Perret P; Lambrechts MM; Heeb P	Oecologia	2009	10.1007/s00442-009-1418-6
GNM_123	Aromatic plants in nests of blue tits: positive effects on nestlings	Mennerat A; Perret P; Bourgault P; Blondel J; Gimenez O; Thomas DW; Heeb P; Lambrechts MM	Animal Behaviour	2009	10.1016/j.anbehav.2008.11.008

Paper ID	Title	Authors	Journal	Year	DOI
GNM_128	Aromatic plants in blue tit <i>Cyanistes caeruleus</i> nests: no negative effect on blood-sucking <i>Protocalliphora</i> blow fly larvae	Mennerat A; Perret P; Caro SP; Heeb P; Lambrechts MM	Journal of Avian Biology	2008	10.1111/j.0908-8857.2008.04400.x
GNM_137	Experimental addition of greenery reduces flea loads in nests of a non-greenery using species, the tree swallow <i>Tachycineta bicolor</i>	Shutler D; Campbell AA	Journal of Avian Biology	2007	10.1111/j.2007.0908-8857.04015.x
GNM_143	European starlings: nestling condition, parasites and green nest material during the breeding season	Gwinner H; Berger S	Journal of Ornithology	2005	10.1007/s10336-005-0012-x
GNM_147	Does fresh vegetation protect avian nests from ectoparasites? An experiment with tree swallows	Dawson RD	Canadian Journal of Zoology	2004	10.1139/Z04-076
GNM_152	Green nesting material has a function in mate attraction in the European starling	Brouwer L; Komdeur J	Animal Behaviour	2004	10.1016/j.anbehav.2003.07.005
GNM_185	Effects of nest-box environment on fledgling success rate and pathogen load	Scott-Baumann JF; Morgan ER; Cogan TA	Parasitology	2022	10.1017/S0031182022000695
GNM_237	Nest Material Shapes Eggs Bacterial Environment	Ruiz-Castellano C; Tomás G; Ruiz-Rodríguez M; Martín-Gálvez D; Soler JJ	PLOS ONE	2016	10.1371/journal.pone.0148894
GNM_349	Ectoparasitism and the role of green nesting material in the European starling	Fauth PT; Krementz DG; Hines JE	Oecologia	1991	10.1007/BF00328399
GNM_355	Effect of biologically active plants used as nest material and the derived benefit to starling nestlings	Clark L; Mason JR	Oecologia	1988	10.1007/BF00379183
GNM_001_add	Is there a role for aromatic plants in blue tit (<i>Cyanistes caeruleus</i>) nests? Results from a correlational and an experimental study	Garrido-Bautista J; Ramos JA; Arce SI; Melero-Romero P; Ferreira R; Santos-Baena C; Guimaro HR; Martín-Villegas C; Moreno-Rueda G; Norte AC	Behavioral Ecology and Sociobiology	2023	10.1007/s00265-023-03393-9
GNM_001_rep	Effects of experimental nest treatment with herbs on ectoparasites and body condition of nestlings	Gładalski M; Norte AC; Bartos M; Demeško I; Kaliński A; Markowski M; Skwarska J; Wawrzyniak J; Zieliński P; Bańbura J	Behavioral Ecology	2025	10.1093/beheco/arae103
GNM_002_add	Green plants in starling nests: effects on nestlings	Gwinner H; Oltrogge M; Trost L; Nienaber U	Animal Behaviour	2000	10.1006/anbe.1999.1306
GNM_003_add	Sparrows use a medicinal herb to defend against parasites and increase offspring condition	Yang C; Ye P; Huo J; Møller AP; Liang W; Feeney WE	Current Biology	2020	10.1016/j.cub.2020.10.021
GNM_1_Unpublished	Greenery experiment – Common buzzard	Meinolf Ottensmann	Unpublished Data (Not peer-reviewed)	NA	NA
GNM_2_Unpublished	Aromatic plants in nests of blue tits (<i>Cyanistes caeruleus</i>)	Anna Pantazi	Master Thesis (Not peer-reviewed)	2024	https://hdl.handle.net/11250/3172874

Standardized Email Template




Figure S 4. **Standardized email template sent to all the authors whose article was included in the meta-analysis:** In cases where we needed the incomplete information, the fitness proxies of interest were mentioned. In all cases, we requested for any unpublished experimental data that could be used for the purpose of our meta-analysis

Description of Unpublished studies

Figure S 5.

We included two unpublished studies in our meta-analysis:

GNM_1_Unpublished:. Greenery manipulation experiment was conducted over two breeding seasons in 2018 and 2019. There were four treatment groups: Control, Spruce, Plastic (Branches of an artificial Christmas tree) and Greenery removal. Nests assigned to the spruce and plastic treatments, were lined with approximately ten twigs in such a way that the entire nest platform was covered. At nests assigned to the greenery removal treatment, all fresh greenery was carefully removed while the nest structure was left untouched. Greenery removal treatment was conditioned on the existence of fresh greenery at the first sampling occasion. Control nests were handled similarly but the greenery composition was not altered. For the purpose of the meta-analysis we did not include the plastic treatment, only, addition or removal of green material and control group. Sampling interval was 5 to 19 days (mean: 7.72 ± 0.54 , $n = 42$ nests).

Sample sizes 42 nests in total: 18 nests allocated to the spruce treatment, 9 greenery removal, 10 plastic and 5 controls

Fitness proxies included in the meta-analysis:

1. Body condition of the nestling
2. Ectoparasite load (Carnus hemapterus): Semi-quantitative measure with five levels, ranging from 0 (no infestation on any of the four locations) to 4 (C. hemapterus larvae or fresh biting marks present on all four examined body locations)
3. Endoparasite load (Leucocytozoon): Five levels 0 – 4

Data used for the meta-analysis can be found data/02_data_extraction/data_extraction_MO_checkedSD.xlsx

GNM_2_Unpublished:. An experiment on Blue Tits conducted in Bergen (Norway), adding lavender in nests and comparing nestling growth & survival with controls. 5 fresh leaves of lavender were added / replaced every 2nd day in broods in the “aromatic” group, from day 5 to day 13 after hatching. In the control group nothing was added, but the broods were otherwise treated similarly.

The following variables were provided in the dataset:

1. Weight D5, D7, D9, D11, D13 - Weight of nestlings at 5, 7, 9, 11, 13 days after hatching of first egg
2. Weight D14-17 - Final weight measure on nestlings at days 14-17 post-hatching, seen as fledging weight
3. TarsusLength D14-17 - Tarsus length measured on nestlings at days 14-17 post-hatching, 0.1mm accuracy

4. Percent Survived - Percentage of all chicks initially hatched in each group that survived until fledging (therefore no SD)
 5. Survival Per Brood - Percentage of chicks in each brood that survived until fledging (the unit is the brood here)
- This data was part of a Master thesis “**Aromatic plants in nests of blue tits (*Cyanistes caeruleus*)**” and all the details can be found at <https://hdl.handle.net/11250/3172874>.

Data used for the meta-analysis can be found data/02_data_extraction/data_extraction_UnpublishedThesis.xlsx

Complete List of Variables extracted from each study

Complete list of variables extracted from each study: comprehensive overview of all variables recorded during data extraction.
Figure S 6.

- Authors Complete information of all authors. Each author name to be separated by ;. Example: Dave Shutler and Adam A. Campbell → “Shutler D; Campbell A A”.
- Year of Publication The year the article was published.
- Population Location Geographical location of the study population extracted as provided by the authors along with any coordinates provided. If multiple sites are studied in a location, extract them as PopulationName_SiteA.
- Experiment ID An identifier given to the estimates coming from the same sub-population within a location.
- Group ID An identifier given to the estimates coming from the same group of experimental units (e.g., males and females in a population, first clutch and second clutch).
- Repeated Trait ID An identifier given to the repeated measurement of the same trait within the same individuals (e.g., chick mass on day 7, 14, and 21).
- Bird Species Scientific name of the bird species studied.
- Treatment Group Plant Species Scientific or common name of the plant species used as the green nest material (treatment group).
- Control Group Plant Species Scientific name of the plant(s) species used as control treatment(s) in the study. If the scientific name is not given, extract it as reported by the authors. If there is no material used for control, note “blank”.
- Comparison Type Experimental design used to code the comparison between the two groups: 1 = non-aromatic (control) vs. aromatic (treatment) 2 = no material (control) vs. aromatic (treatment) 3 = no material (control) vs. non-aromatic (treatment)
- CH Estimates related to the courtship hypothesis — yes (1) or no (0).
- PCH Estimates related to the nest protection or drug hypothesis — yes (1) or no (0).
- Measure of Central Tendency (treatment group) Estimated value of central tendency for the samples of the treatment group.
- Type of Measure of Central Tendency (treatment group) Type of central tendency value reported for the treatment group (e.g., Mean, Median, Mode).
- Measure of Dispersal (treatment group) Value of the measure of variation for the treatment group.
- Type of Measure of Dispersal (treatment group) Type of variation of the data (e.g., SE, SD, CI95) reported for the treatment group.
- Sample Size (treatment group) Sample size for the treatment group.
- Measure of Central Tendency (control group) Estimated value of central tendency for the samples of the control group.
- Type of Measure of Central Tendency (control group) Type of central tendency value reported for the control group (e.g., Mean, Median, Mode).
- Measure of Dispersal (control group) Value of the measure of variation for the control group.
- Type of Measure of Dispersal (control group) Type of variation of the data (e.g., SE, SD, CI95) reported for the control group.
- Sample Size (control group) Sample size for the control group.
- Fitness Proxy Measured Variable name used in the study as a proxy of fitness. Extracted as presented in the article and later standardized across studies.
- Category/Type of Fitness Proxy Measured Type of trait (i.e., fitness proxy) studied: behaviour, morphology, parasite and pathogenic load, phenology, physiology, or reproduction.
- Fitness Proxy Sign Sign assigned in relation to fitness: +1 if fitness increases with proxy (e.g., survival rate, body mass). -1 if fitness decreases with proxy (e.g., mortality, parasite load).
- Statistics Type Type of statistics used to address the effect of GNM on fitness when primary data were not reported (e.g., t-test, ANOVA). Extracted only when central tendency was not reported.
- Test Statistics Type Type of test statistic used (e.g., t, F).
- Statistics Value Value of the test statistic used.

- P Value Reported p-value (e.g., $p = 0.003$).
- Sign Relationship Positive when treatment increases the trait value and negative when it decreases the trait value. Used when only the statistical value was provided.
- Total Sample Size Total sample size used to calculate the statistical value.
- Degree of Freedom Degrees of freedom for the statistics used.
- Data Location The part of the text or figure from which the data were extracted.
- Parasite Type Type of parasite when fitness proxy measured parasite/pathogen load (Arthropod or Micro-organism).
- Time of Green Nest Material Addition Time of green nest material addition by authors or birds [continuous (c), before (b), or after (a) egg hatching].
- Blinding Whether blinding was conducted – yes (y) or no (n).
- Random Assignment Whether nests were assigned to treatment or control randomly – yes (y) or no (n).
- Missing Data Whether data were only partially or not reported – yes (y) or no (n).
- Shared Treatment Group Used to identify non-independence between shared treatment groups. Value = 1 if individuals were compared only once as a treatment group.
- Shared Control Group Used to identify non-independence between shared control groups. Value = 1 if individuals were compared only once as a control group.

Random Effects




Figure S 7. Alluvial plot showing the overlap among random effects extracted for the meta-analytic dataset: Each vertical bar represents a random-effect term i.e. Paper ID (28 levels), Experiment ID (48), Group ID (52), Repeated Trait ID (225), Observation ID (273), Bird Species (7), and Population ID (20). Connections between bars indicate shared values across levels. The figure shows that Experiment ID and Group ID were nearly identical, and that Repeated Trait ID had many levels almost as many as the Observation ID, and Repeated Trait ID was excluded from the final models. In contrast, Population ID captured additional clustering across studies and was therefore added to the random effects (not pre-registered).

Sensitivity analysis of sampling correlation values (ρ) in variance–covariance structures

Sensitivity to sampling correlation (ρ)										
Results for lnRR and SMD(H) Meta-analytic models										
ρ	Overall effect	SE	p	95% CI		LogLik	AIC	BIC	AICc	Effect.size
				CI low	CI high					
0.2	0.0168	0.0131	0.198	−0.0088	0.0424	−15.5399	43.0798	63.8881	43.4450	lnRR
0.3	0.0186	0.0138	0.180	−0.0086	0.0457	−17.4753	46.9505	67.7589	47.3157	lnRR
0.5	0.0190	0.0156	0.222	−0.0115	0.0495	−26.0231	64.0462	84.8546	64.4114	lnRR
0.7	0.0147	0.0181	0.416	−0.0207	0.0501	−43.3232	98.6464	119.4548	99.0117	lnRR
0.8	0.0086	0.0203	0.670	−0.0311	0.0483	−58.3988	128.7976	149.6060	129.1628	lnRR
0.2	0.1330	0.0501	0.008	0.0349	0.2311	−297.6150	607.2301	628.4066	607.5729	SMD(H)
0.3	0.1479	0.0556	0.008	0.0390	0.2568	−304.1710	620.3420	641.5186	620.6849	SMD(H)
0.5	0.1793	0.0665	0.007	0.0489	0.3098	−324.4948	660.9896	682.1661	661.3324	SMD(H)
0.7	0.2560	0.1180	0.030	0.0248	0.4872	−355.0793	722.1585	743.3351	722.5014	SMD(H)
0.8	0.2936	0.1508	0.051	−0.0019	0.5891	−372.2800	756.5600	777.7365	756.9028	SMD(H)

Figure S 8. **Sensitivity of overall effect estimates to the assumed sampling correlation (ρ):** The table shows model estimates of the overall effect, standard errors (SE), p -values, 95% confidence intervals (CI), and fit statistics (log-likelihood, AIC, BIC, and AICc) across a range of assumed sampling correlations (ρ) used for the variance-covariance matrices.

All sensitivity-analysis/robustness check

The overall results and conclusion were robust to several sensitivity analyses including the bivariate multilevel meta-analytic model including both effect sizes as response variables (lnRR = −0.015, p -value = 0.624 and SMD(H) = 0.250 , p -value = 0.069).

Summary of Sensitivity Analyses for lnRR and SMD(H) Intercept-only models								
Effect Size	Data used in model	Estimate	95% CI	95% PI	P-value	k (No. of Effect Sizes)	n (No. of Studies)	
lnRR	Including published data only	0.020	[−0.013, 0.052]	[−0.290, 0.329]	0.230	221	24	
lnRR	Excluding Experimental Design Non-Aromatic Vs. No added material	0.027	[−0.009, 0.063]	[−0.203, 0.256]	0.147	228	26	
lnRR	Excluding Experimental Design Aromatic Vs. Non-Aromatic	0.032	[−0.024, 0.088]	[−0.379, 0.443]	0.258	131	15	
lnRR	Including flagged proxies	0.024	[−0.006, 0.053]	[−0.272, 0.319]	0.113	259	28	
lnRR	Without those cases where author reported no effect of the treatment	0.019	[−0.013, 0.050]	[−0.291, 0.328]	0.238	222	26	
lnRR	Including flagged proxies and without the cases with no effect of the treatment	0.024	[−0.006, 0.054]	[−0.280, 0.328]	0.121	243	28	
lnRR	Excluding effect sizes where lnRR failed Geary's Test	0.029	[−0.003, 0.060]	[−0.271, 0.328]	0.074	182	26	
SMD(H)	Including published data only	0.192	[0.056, 0.327]	[−0.849, 1.232]	0.006	235	24	
SMD(H)	Excluding Experimental Design Non-Aromatic Vs. No added material	0.148	[0.019, 0.278]	[−0.816, 1.112]	0.025	242	26	
SMD(H)	Excluding Experimental Design Aromatic Vs. Non-Aromatic	0.220	[0.020, 0.419]	[−0.972, 1.411]	0.031	136	15	
SMD(H)	Including flagged proxies	0.191	[0.067, 0.315]	[−0.798, 1.181]	0.003	274	28	
SMD(H)	Without those cases where author reported no effect of the treatment	0.186	[0.048, 0.324]	[−0.896, 1.268]	0.008	237	26	
SMD(H)	Including flagged proxies and without the cases with no effect of the treatment	0.199	[0.069, 0.329]	[−0.854, 1.251]	0.003	258	28	
SMDH	Without the cases where SMDH was calculated from inferential statistics	0.179	[0.044, 0.314]	[−0.856, 1.215]	0.009	245	26	

Table S 4. Results of sensitivity analyses assessing the robustness of the overall meta-analytic results for lnRR and SMD(H) intercept-only models. We tested whether the main conclusions were sensitive to alternative analytical decisions, including the use of published data only, inclusion of flagged proxies, exclusion of cases reported as having no treatment effect, exclusion of specific experimental-design categories, exclusion of effect sizes calculated from inferential statistics or failing Geary's test. Across analyses, the number of effect sizes ranged from 131 to 274, representing 15 to 28 studies. Overall, the results were qualitatively robust across these alternative model specifications.

Plots without cropping the axis (showing complete X-axis)




Figure S 9. Overall effect of green nest material on all fitness proxies combined. The meta-analytic mean (black circle) with 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along with the data points (coloured circles) scaled by precision (inverse of standard error). Results are shown as an orchard plot for the intercept-only model (A) using log response ratio (lnRR), and (B) standardised mean difference (SMDH). k : number of effect sizes (number of studies is shown in brackets).




Figure S 10. There is no clear evidence to favour either the courtship or the parental care hypothesis as drivers of potential fitness benefits of green nest material. Orchard plot of the multilevel meta-regression for (A) log response ratio (lnRR) and (B) standardised mean difference (SMDH). Mean estimates (black circles), 95% confidence intervals (thicker bars), and prediction intervals (thinner bars) are shown along with the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k : number of effect sizes (number of studies is shown in brackets).




Figure S 11. Experimental designs comparing nests with aromatic green nest material against nests without any green material, and those adding the material continuously, lead to the largest fitness estimates. Orchard plots of the multilevel meta-regressions showing the overall effect of green nest material depending on experimental design for (A: log response ratio (lnRR); B: standardised mean difference (SMD(H)), and time of green nest material addition (C: log response ratio (lnRR); D: standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars) and prediction intervals (thinner bars) are shown along the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k: number of effect sizes (number of studies is shown in brackets).




Figure S 12. Orchard plot of the multilevel meta-regression for (A) log response ratio (lnRR) and (B) standardised mean difference (SMD(H)). Mean estimates (black circles), 95% confidence intervals (thicker bars), and prediction intervals (thinner bars) are shown along with the individual data points (transparently coloured circles) scaled by precision (inverse of standard error). k: number of effect sizes (number of studies is shown in brackets).

Risk of Bias Test

The risk of bias categories we evaluated (i.e., blinding, random assignment to each treatment and partial reporting) were not statistically significant for lnRR (Table S 5). SMD(H) was positive and statistically significant for studies without blinding ($SMD(H)_{No\ Blinding} = 0.184$, 95% CI: [0.051, 0.316], 95% PI: [-0.828, 1.195], p-value = 0.007, k = 242, n = 25; Table S 5), with random assignment to treatment ($SMD(H)_{Random\ Assignment} = 0.205$, 95% CI: [0.049, 0.360], 95% PI: [-0.813, 1.223], p-value = 0.010, k = 225, n = 23; Table S 5) and where no data was missing ($SMD(H)_{No\ Partial\ Reporting} = 0.196$, 95% CI: [0.055, 0.337], 95% PI: [-0.817, 1.209], p-value = 0.007, k = 216, n = 22; Table S 5). Notably, other levels of each category included fewer than four studies, indicating that the observed significance likely reflects the higher statistical power in the larger sample level rather than any systematic bias. All the three categories for risk of bias explained very little heterogeneity ($R^2_{marginal(lnRR)} > 0.8\%$; $R^2_{marginal(SMD(H))} > 0.7\%$).

Multilevel Meta Regressions for Risk of Bias											
lnRR						SMDH					
Level	Estimate	P-val	95% CI	95% PI	k	n	Estimate	P-val	95% CI	95% PI	k n
1. Blinding											
No	0.022	0.168	[-0.009, 0.053]	[-0.278, 0.322]	227	25	0.184	0.007	[0.051, 0.316]	[-0.828, 1.195]	242 25
Yes	-0.042	0.555	[-0.182, 0.098]	[-0.372, 0.288]	11	1	0.016	0.970	[-0.816, 0.847]	[-1.287, 1.318]	11 1
2. Random assignment to treatment											
No	0.002	0.958	[-0.080, 0.084]	[-0.309, 0.313]	22	3	0.070	0.705	[-0.292, 0.431]	[-0.999, 1.138]	28 3
Yes	0.022	0.199	[-0.011, 0.055]	[-0.280, 0.323]	216	23	0.205	0.010	[0.049, 0.360]	[-0.813, 1.223]	225 23
3. Partial Reporting											
No	0.021	0.205	[-0.011, 0.053]	[-0.280, 0.322]	201	22	0.196	0.007	[0.055, 0.337]	[-0.817, 1.209]	216 22
Yes	-0.001	0.989	[-0.110, 0.109]	[-0.320, 0.318]	37	4	0.071	0.695	[-0.286, 0.429]	[-0.994, 1.136]	37 4

Table S 5.

Publication Bias

Since published effect sizes did not seem to change considerably over time, our analyses suggest no clear evidence of decline effects (i.e., time-lag bias) in published effect sizes ($\ln RR_{\text{intercept}} = 0.021$, 95% CI: $[-0.011; 0.053]$, $p\text{-value} = 0.196$, $\ln RR_{\text{slope}} = -0.002$, 95% CI: $[-0.005 \text{ to } 0.001]$, $p\text{-value} = 0.225$, $k = 221$, $n = 24$, $R^2_{\text{marginal}(\ln RR)} = 1.8\%$; $SMDH_{\text{intercept}} = 0.193$, 95% CI: $[0.058 \text{ to } 0.329]$, $p\text{-value} = 0.005$, $SMDH_{\text{slope}} = -0.008$, 95% CI: $[-0.023 \text{ to } 0.006]$, $p\text{-value} = 0.264$, $k = 235$, $n = 24$, $R^2_{\text{marginal}(SMD(H))} = 2.9\%$).




Figure S 13. Tests for publication bias in the effect of green nest material on fitness outcomes. Plots show relationships between effect sizes and (A–B) the inverse of effective sample size testing for small-study effects, and (C–D) publication year (mean-centered) testing for time-lag bias, for log response ratio (lnRR, A and C) and standardized mean difference (SMD(H), B and D). Points represent individual effect sizes, scaled by precision (1/SE). Solid lines show meta-regression slopes, with 95% confidence intervals (dashed lines) and 95% prediction intervals (dotted lines). k : number of effect sizes included in each analysis.

Systematic Search Information

Figure S 14.

The search was conducted twice, the first time on 07.09.2022 and then repeated on 12.08.2024 to ensure it was the most up to date and did not miss any additional articles on the topic that were published meanwhile. For both the searches, we did not use any time limitation i.e. it included all the published literature till that given date. We used **Advance Search** option for both Web of Science Core Collection Database and Scopus Database using the search query given below.

Search string for the Web of Science Core Collection

Databases used in the search included

- Science Citation Index Expanded (SCI-EXPANDED) [1900-present]
- Social Sciences Citation Index (SSCI) [1900-present]
- Arts & Humanities Citation Index (AHCI) [1975 - present]
- Conference Proceedings Citation Index - Science (CPCI-S) [1991 - present]
- Conference Proceedings Citation Index - Social Science & Humanities (CPCI-SSH) [1991 - present]
- Emerging Sources Citation Index (ESCI) [2017- present]

We used advance search option and Title (TI), Abstract (AB) and Author Keywords (AK) to conduct our systematic

search using the following string

<TI=((("green*" OR "herb*" OR "aromatic*") AND "nest*" AND ("bird*" OR "aves" OR "avian" OR "ornithol*" OR "passerine*" OR "passeriform*" OR "songbird*" OR "Accipiter" OR "Aegyptius" OR "Aquila" OR "Aviceda" OR "Busarellus" OR "Butastur" OR "Buteo" OR "Buteogallus" OR "Chondrohierax" OR "Circetus" OR "Circus" OR "Clanga" OR "Elanoides" OR "Elanus" OR "Gampsonyx" OR "Geranoaetus" OR "Geranoaspiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR "Haliastur" OR "Harpagus" OR "Harpia" OR "Hieraaetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco" OR "Leptodon" OR "Leucopternis" OR "Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus" OR "Necrosyrtes" OR "Neophron" OR "Nisaetus" OR "Parabuteo" OR "Pernis" OR "Pithecophaga" OR "Polemaetus" OR "Pseudastur" OR "Rostrhamus" OR "Rupornis" OR "Sarcogyps" OR "Spilornis" OR "Spizaetus" OR "Stephanoaetus" OR "Terathopius" OR "Torgos" OR "Trigonoceps" OR "Cathartes" OR "Coragyps" OR "Gymnogyps" OR "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix" OR "Alopochen" OR "Amazonetta" OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR "Bucephala" OR "Cairina" OR "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloephaga" OR "Clangula" OR "Coscoroba" OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR "Histrionicus" OR "Hymenolaimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca" OR "Marmaronetta" OR "Melanitta" OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neochen" OR "Netta" OR "Nettapus" OR "Nomonyx" OR "Oxyura" OR "Plectropterus" OR "Polysticta" OR "Pteronetta" OR "Sarkidiornis" OR "Sibirionetta" OR "Somateria" OR "Spatula" OR "Tachyeres" OR "Tadorna" OR "Thalassornis" OR "Anhima" OR "Chauna" OR "Anseranas" OR "Aegothales" OR "Aerodramus" OR "Aeronautes" OR "Apus" OR "Chaetura" OR "Collocalia" OR "Cypseloides" OR "Cypsiurus" OR "Hirundapus" OR "Panyptila" OR "Streptoprocne" OR "Tachornis" OR "Tachymarptis" OR "Hemiprocne" OR "Abeillia" OR "Adelomyia" OR "Aglaeactis" OR "Aglaiocercus" OR "Amazilia" OR "Androdon" OR "Anopetia" OR "Anthocephala" OR "Anthracothorax" OR "Archilochus" OR "Atthis" OR "Augastes" OR "Avocettula" OR "Basilinna" OR "Boissonneaua" OR "Calliphlox" OR "Calypte" OR "Campylopterus" OR "Chaetocercus" OR "Chalcostigma" OR "Chalybura" OR "Chlorestes" OR "Chlorostilbon" OR "Chrysolampis" OR "Chrysuronia" OR "Clytolaema" OR "Coeligena" OR "Colibri" OR "Cyanophaia" OR "Cynanthus" OR "Discosura" OR "Doricha" OR "Doryfera" OR "Elvira" OR "Ensifera" OR "Eriocnemis" OR "Eugenes" OR "Eulampis" OR "Eulidia" OR "Eupetomena" OR "Eupherusa" OR "Eutoxeres" OR "Florisuga" OR "Glaucis" OR "Goethalsia" OR "Goldmania" OR "Haplophaedia" OR "Heliactin" OR "Heliangelus" OR "Heliodoxa" OR "Heliomaster" OR "Heliosthryx" OR "Hylocharis" OR "Juliamyia" OR "Klais" OR "Lafresnaya" OR "Lampornis" OR "Lamprolaima" OR "Lepidopyga" OR "Lesbia" OR "Leucippus" OR "Leucochloris" OR "Lophornis" OR "Mellisuga" OR "Metallura" OR "Microchera" OR "Myrmia" OR "Myrtis" OR "Oreatus" OR "Opisthoprora" OR "Oreonympha" OR "Oreotrochilus" OR "Orthorhynchus" OR "Oxygogon" OR "Panterpe" OR "Patagona" OR "Phaeochroa" OR "Phaethornis" OR "Phlogophilus" OR "Polyonymus" OR "Polytmus" OR "Pterophanes" OR "Ramphodon" OR "Ramphomicron" OR "Rhodopsis" OR "Sappho" OR "Schistes" OR "Selasphorus" OR "Sephanoidea" OR "Stephanoxis" OR "Sternoclyta" OR "Taphrospilus" OR "Thalaurania" OR "Thaumastura" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR "Rhabdotorrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus" OR "Upupa" OR "Antristomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus" OR "Hydropsalis" OR "Lurocalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus" OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius" OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariama" OR "Casuarius" OR "Dromaius" OR "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cerorhinca" OR "Fratricula" OR "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus" OR "Charadrius" OR "Elseynoris" OR "Erythrogonyx" OR "Hoploxypterus" OR "Oreopholus" OR "Peltodytes" OR "Plegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelochelidon" OR "Gygis" OR "Hydrocoloeus" OR "Hydroprogne" OR "Ichthyaeetus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR "Pagophila" OR "Phaetusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula" OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR "Recurvirostra" OR "Nycticriophes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR "Calidris" OR "Coenocorypha" OR "Gallinago" OR "Limnodromus" OR "Limosa" OR "Lymnocyrtus" OR "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagis" OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chalcophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula" OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR "Gymnophaps" OR "Hemiphaga" OR "Henicophaps" OR "Leptotila" OR "Leptotrygon" OR "Leucosarcia" OR "Lopholaimus" OR "Macropygia" OR "Metriopelia" OR "Ocyphaps" OR "Oena" OR "Otidiphaps" OR "Patagioenas" OR "Petrophassa" OR "Pezophaps" OR "Phapitreron" OR "Phaps" OR "Ptilinopus" OR "Raphus" OR "Reinwardtoena" OR "Spilopelia" OR "Streptopelia" OR "Treron" OR "Trugon" OR "Turacoena" OR "Turtur" OR "Uropelia" OR "Zenaida" OR "Zentrygon" OR "Actenoides" OR "Alcedo" OR "Caridonax" OR "Ceryle" OR "Ceyx" OR "Chloroceryle" OR "Cittura" OR "Corythornis" OR "Dacelo" OR

"Halcyon" OR "Ispidina" OR "Lacedo" OR "Megaceryle" OR "Melidora" OR "Syma" OR "Tanysiptera" OR "Todiramphus" OR "Atelornis" OR "Brachypteracias" OR "Geobiastes" OR "Coracias" OR "Eurystomus" OR "Merops" OR "Nyctyornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes" OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR "Chrysococcyx" OR "Clamator" OR "Coccyua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus" OR "Dasylophus" OR "Dromococcyx" OR "Eudynamys" OR "Geococcyx" OR "Guira" OR "Hierococcyx" OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicophaeus" OR "Piaya" OR "Rhinortha" OR "Scythrops" OR "Surniculus" OR "Tapera" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR "Anomalopteryx" OR "Emeus" OR "Euryapteryx" OR "Eurypyga" OR "Rhynochetos" OR "Caracara" OR "Daptrius" OR "Falco" OR "Herpetotheres" OR "Ibycter" OR "Micrastur" OR "Microhierax" OR "Milvago" OR "Phalcoboenus" OR "Polihierax" OR "Spiziapteryx" OR "Aburria" OR "Chamaepetes" OR "Crax" OR "Mitu" OR "Nothocrax" OR "Oreophasis" OR "Ortalis" OR "Pauxi" OR "Penelope" OR "Penelopina" OR "Pipile" OR "Alectura" OR "Megapodius" OR "Acryllium" OR "Guttera" OR "Numida" OR "Callipepla" OR "Colinus" OR "Cyrtonyx" OR "Dendrortyx" OR "Odontophorus" OR "Oreortyx" OR "Philortyx" OR "Ptilopachus" OR "Rhynchortyx" OR "Alectoris" OR "Ammoperdix" OR "Arborophila" OR "Argusianus" OR "Bambusicola" OR "Bonasa" OR "Caloperdix" OR "Centrocerus" OR "Chrysolophus" OR "Coturnix" OR "Crossoptilon" OR "Dendragapus" OR "Excalfactoria" OR "Falcipennis" OR "Francolinus" OR "Gallus" OR "Haematortyx" OR "Ithaginis" OR "Lagopus" OR "Lerwa" OR "Lophophorus" OR "Lophura" OR "Lyrurus" OR "Meleagris" OR "Pavo" OR "Peliperdix" OR "Perdix" OR "Phasianus" OR "Polyplectron" OR "Pternistis" OR "Pucrasia" OR "Rhizothera" OR "Rollulus" OR "Scleroptila" OR "Syrmaticus" OR "Tetrao" OR "Tetraogallus" OR "Tetraophasis" OR "Tragopan" OR "Tympanuchus" OR "Gavia" OR "Aramus" OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amaurornis" OR "Anurolimnas" OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia" OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrio" OR "Porphyriops" OR "Porzana" OR "Rallacula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura" OR "Leptosomus" OR "Corythaëola" OR "Corythaixoides" OR "Crinifer" OR "Musophaga" OR "Tauraco" OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax" OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis" OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis" OR "Smicornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna" OR "Nesillas" OR "Aegithalos" OR "Leptopoeile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes" OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafr" OR "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhynchophanes" OR "Callaeas" OR "Heteralocha" OR "Philesturnus" OR "Calyptophilus" OR "Campephaga" OR "Ceblepyris" OR "Coracina" OR "Edolisoma" OR "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amaurospiza" OR "Cardinalis" OR "Caryothraustes" OR "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Paserina" OR "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus" OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus" OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia" OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus" OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus" OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias" OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza" OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis" OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga" OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calyptomena" OR "Cymbirhynchus" OR "Eurylaimus" OR "Psalisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chloris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra" OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorrhous" OR "Hemignathus" OR "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza" OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus" OR "Certhiasomus" OR "Certhiaxis" OR "Cincludes" OR "Clibanornis" OR "Coryphistera" OR

"Cranioleuca" OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geocerthia" OR "Geositta" OR "Glyphorhynchus" OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR "Limnortites" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor" OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus" OR "Spartonoica" OR "Sylviorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonimus" OR "Thripadectes" OR "Thripophaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes" OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hyllopezus" OR "Myrmothera" OR "Alopochelidon" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon" OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psalidoprocne" OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR "Tachycineta" OR "Hyltiota" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Amblycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives" OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus" OR "Lamprosar" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar" OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura" OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus" OR "Leiothrix" OR "Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochalopecton" OR "Turdoides" OR "Bradypterus" OR "Cincloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malaconotus" OR "Nilaus" OR "Rhodophoneus" OR "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia" OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomyzon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes" OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis" OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius" OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cincloerthia" OR "Dumetella" OR "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis" OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus" OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Camicoloides" OR "Cercotrichas" OR "Chamaetylas" OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia" OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiomela" OR "Myioparus" OR "Myophonus" OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola" OR "Sheppardia" OR "Sholicola" OR "Stiphornis" OR "Tarsiger" OR "Thamnodaea" OR "Vauriella" OR "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkinogramma" OR "Leptocoma" OR "Nectarinia" OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleadryas" OR "Oriolus" OR "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR "Cyanistes" OR "Lophophanes" OR "Macholophus" OR "Melaniparus" OR "Melanochlora" OR "Pardaliparus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus" OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros" OR "Leiothlypis" OR "Limnithlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis" OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisospiza" OR "Atlapetes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza" OR "Melozona" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR "Pezopetes" OR "Pipilo" OR "Poocetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizelloides" OR "Xenospiza" OR "Zonotrichia" OR "Carpospiza" OR "Gymnoris" OR "Hypocryptadius" OR "Montifringilla" OR "Onychostreuthus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR "Gampsorhynchus" OR "Illadopsis" OR "Jabouilleia" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR "Malacopteron" OR "Napothea" OR "Pellorneum" OR "Rimator" OR "Trichastoma" OR "Amalocichla" OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca" OR "Monachella" OR "Pachycephalopsis" OR "Peneoenanthe" OR "Peneothello" OR "Petroica" OR "Poecilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyrannetes" OR "Xenopipo" OR "Erythropitta" OR "Hydornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalornis" OR "Euplectes" OR "Malimbus" OR "Plocepasser"

OR "Ploceus" OR "Quelea" OR "Sporopipes" OR "Pnoepyga" OR "Microbates" OR "Poliopitila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR "Cinclosoma" OR "Ptilorrhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla" OR "Atimastillas" OR "Baepogon" OR "Bleda" OR "Chlorocichla" OR "Criniger" OR "Eurillas" OR "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus" OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus" OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pteroptochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynch" OR "Culicicapa" OR "Elminia" OR "Acridotheres" OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora" OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotornis" OR "Leucopsar" OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprositornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR "Drymophila" OR "Dysithamnus" OR "Epinecropphylla" OR "Euchrepomis" OR "Formicivora" OR "Frederickena" OR "Gymnocichla" OR "Gymnopathys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus" OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myrmoderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes" OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania" OR "Pygipitila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus" OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis" OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Chariotspiza" OR "Chlorochrysa" OR "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba" OR "Compsospiza" OR "Compsotrupis" OR "Conirostrum" OR "Conothraupis" OR "Coryphospingus" OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis" OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Inaspiza" OR "Iridophanes" OR "Iridosornis" OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes" OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocomma" OR "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha" OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatrhinus" OR "Spelaornis" OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius" OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyrhamphus" OR "Schiffornis" OR "Terenotriccus" OR "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia" OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Troglodytes" OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus" OR "Zoothraupis" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola" OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Casiempis" OR "Casiornis" OR "Cnemarchus" OR "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis" OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus" OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myiodynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiorticcus" OR "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes" OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis" OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis" OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis" OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR "Urocynchramus" OR "Artamella" OR "Callicolpus" OR "Cyanolanius" OR "Euryceros" OR "Falculea" OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris" OR "Anomalospiza"

OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR "Vireo" OR "Vireolanus" OR "Zele-
donia" OR "Apalopteron" OR "Dasycrotapha" OR "Heleia" OR "Lophozosterops" OR "Sternhoptilus" OR "Yuhina" OR
"Zosterops" OR "Zosterornis" OR "Agamia" OR "Ardea" OR "Ardeola" OR "Botaurus" OR "Bubulcus" OR "Butorides"
OR "Cochlearius" OR "Dupetor" OR "Egretta" OR "Gorsachius" OR "Ixobrychus" OR "Nyctanassa" OR "Nycticorax"
OR "Pilherodius" OR "Syrrigma" OR "Tigriornis" OR "Tigrisoma" OR "Zebrilus" OR "Balaeniceps" OR "Pelecanus" OR
"Scopus" OR "Bostrychia" OR "Eudocimus" OR "Geronticus" OR "Lophotibis" OR "Mesembrinibis" OR "Nipponia"
OR "Phimosus" OR "Platalea" OR "Plegadis" OR "Pseudibis" OR "Theristicus" OR "Threskiornis" OR "Phaethon" OR
"Phoeniconaias" OR "Phoenicoparrus" OR "Phoenicopterus" OR "Bucco" OR "Chelidoptera" OR "Hapaloptila" OR
"Hypnelus" OR "Malacoptila" OR "Micromonacha" OR "Monasa" OR "Nonnulla" OR "Notharchus" OR "Nystalus" OR
"Capito" OR "Eubucco" OR "Brachygalba" OR "Galbalcyrhynchus" OR "Galbula" OR "Jacamaralcyon" OR "Jacamerops"
OR "Indicator" OR "Melichneutes" OR "Meligonomon" OR "Prodotiscus" OR "Bucconodon" OR "Gymnobucco" OR
"Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus" OR "Psilopogon" OR "Blythipicus" OR "Campephilus"
OR "Campetheres" OR "Celeus" OR "Chloropicus" OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Den-
drocopos" OR "Dendrocopetes" OR "Dendropicos" OR "Dinopium" OR "Dryobates" OR "Dryocopus" OR "Gecinulus" OR
"Geocolaptes" OR "Jynx" OR "Leiopicus" OR "Leuconotopicus" OR "Meiglyptes" OR "Melanerpes" OR "Micropternus"
OR "Mulleripicus" OR "Nesocittes" OR "Picoides" OR "Piculus" OR "Picumnus" OR "Picus" OR "Reinwardtipicus" OR
"Sasia" OR "Sphyrapicus" OR "Veniliornis" OR "Yungipicus" OR "Andigena" OR "Aulacorhynchus" OR "Pteroglossus"
OR "Ramphastos" OR "Selenidera" OR "Semnornis" OR "Aechmophorus" OR "Podiceps" OR "Podilymbus" OR "Polio-
cephalus" OR "Rollandia" OR "Tachybaptus" OR "Diomedea" OR "Phoebeastria" OR "Phoebebria" OR "Thalassarche" OR
"Hydrobates" OR "Oceanodroma" OR "Fregetta" OR "Garrodia" OR "Nesofregetta" OR "Oceanites" OR "Pelagodroma"
OR "Aphrodroma" OR "Ardena" OR "Bulweria" OR "Calonectris" OR "Daption" OR "Fulmarus" OR "Halobaena" OR
"Macronectes" OR "Pachyptila" OR "Pagodroma" OR "Pelecanoides" OR "Procellaria" OR "Pseudobulweria" OR "Ptero-
droma" OR "Puffinus" OR "Thalassoica" OR "Cacatua" OR "Callocephalon" OR "Calyptorhynchus" OR "Eolophus" OR
"Lophochroa" OR "Nymphicus" OR "Probosciger" OR "Alipiopsitta" OR "Amazona" OR "Anodorhynchus" OR "Ara" OR
"Aratinga" OR "Bolborhynchus" OR "Brotogeris" OR "Conuropsis" OR "Cyanoliseus" OR "Cyanopsitta" OR "Deroytus"
OR "Diopsittaca" OR "Enicognathus" OR "Eupsittula" OR "Forpus" OR "Graydidascalus" OR "Guarouba" OR "Guaruba"
OR "Hapalopsittaca" OR "Leptosittaca" OR "Myiopsittaca" OR "Nannopsittaca" OR "Orthopsittaca" OR "Pionites" OR
"Pionopsitta" OR "Pionus" OR "Poicephalus" OR "Primolius" OR "Psilopsiagon" OR "Psittacara" OR "Psittacus" OR
"Pyralia" OR "Pyrrhura" OR "Rhynchopsitta" OR "Thectocercus" OR "Touit" OR "Triclaria" OR "Agapornis" OR "Alis-
terus" OR "Aprosmictus" OR "Barnardius" OR "Bolbopsittacus" OR "Chalcopsitta" OR "Charmosyna" OR "Coracopsis"
OR "Cyanoramphus" OR "Cyclopsitta" OR "Eclectus" OR "Eos" OR "Eunymphicus" OR "Geoffroyus" OR "Glossopsitta"
OR "Lathamus" OR "Loriculus" OR "Lorius" OR "Melopsittacus" OR "Micropsitta" OR "Neophema" OR "Neopsephotus"
OR "Neopsittacus" OR "Northiella" OR "Oreopsittacus" OR "Parvipsitta" OR "Pezoporus" OR "Phigys" OR "Platyercus"
OR "Polytelis" OR "Prioniturus" OR "Prosopieia" OR "Psephotellus" OR "Psephotus" OR "Pseudeos" OR "Psittacula"
OR "Psittacula" OR "Psittaculirostris" OR "Psitteuteles" OR "Psittinus" OR "Psittirichas" OR "Purpureicephalus" OR
"Tanygnathus" OR "Trichoglossus" OR "Vini" OR "Nestor" OR "Strigops" OR "Pterocles" OR "Synrhaphes" OR "Rhea"
OR "Aptenodytes" OR "Eudyptes" OR "Eudyptula" OR "Megadyptes" OR "Pygoscelis" OR "Spheniscus" OR "Aegolius"
OR "Asio" OR "Athene" OR "Bubo" OR "Glaucidium" OR "Ketupa" OR "Lophostrix" OR "Megascops" OR "Micrathene"
OR "Ninox" OR "Otus" OR "Pseudoscops" OR "Psiloscoops" OR "Pulsatrix" OR "Sceloglaux" OR "Scotopelia" OR "Strix"
OR "Surnia" OR "Xenoglaux" OR "Phodilus" OR "Tyto" OR "Aepyornis" OR "Mullerornis" OR "Struthio" OR "Anhinga"
OR "Fregata" OR "Leucocarbo" OR "Microcarbo" OR "Nannopterum" OR "Phalacrocorax" OR "Morus" OR "Sula" OR
"Crypturellus" OR "Eudromia" OR "Nothocercus" OR "Nothoprocta" OR "Nothura" OR "Rhynchotus" OR "Tinamotis"
OR "Tinamus" OR "Apalharpactes" OR "Apaloderma" OR "Harpactes" OR "Pharomachus" OR "Trogon" OR "Tetrastes"
OR "Hesperiphona") AND ("experiment*" OR "manipulat*")) OR AB= (((("green*" OR "herb*" OR "aromatic*") AND
"nest*" AND ("bird*" OR "aves" OR "avian" OR "ornithol*" OR "passerine*" OR "passeriform*" OR "songbird*" OR
"Accipiter" OR "Aegyptius" OR "Aquila" OR "Aviceda" OR "Busarellus" OR "Butastur" OR "Buteo" OR "Buteogallus" OR
"Chondrohierax" OR "Circaetus" OR "Circus" OR "Clanga" OR "Elanoides" OR "Elanus" OR "Gampsonyx" OR "Gera-
noaetus" OR "Geranoospiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR "Haliastur" OR "Harpagus"
OR "Harpia" OR "Hieraaetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco" OR "Leptodon" OR "Leucopternis" OR
"Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus" OR "Necrosyrtes" OR "Neophron" OR "Nisaetus"
OR "Parabuteo" OR "Pernis" OR "Pithecopaga" OR "Polemaetus" OR "Pseudastur" OR "Rostrhamus" OR "Rupornis" OR
"Sarcogyps" OR "Spilornis" OR "Spizaetus" OR "Stephanoaetus" OR "Terathopus" OR "Torgos" OR "Trionoceps" OR
"Cathartes" OR "Coragyps" OR "Gymnogyps" OR "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix"
OR "Alopochen" OR "Amazonetta" OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR "Bucephala"
OR "Cairina" OR "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloephaga" OR "Clangula" OR
"Coscoroba" OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR "Histriornis" OR "Hymeno-
laimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca" OR "Marmaronetta" OR "Melanitta"
OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neoche" OR "Netta" OR "Nettapus" OR "Nomonyx" OR "Oxyura"
OR "Plectropterus" OR "Polysticta" OR "Pteronetta" OR "Sarkidiornis" OR "Sibirionetta" OR "Somateria" OR "Spatula"
OR "Tachyeres" OR "Tadorna" OR "Thalassornis" OR "Anhimus" OR "Chauna" OR "Anseranas" OR "Aegotheles" OR

Aceramus” OR “Aeronautes” OR “Apus” OR “Chaetura” OR “Collocalia” OR “Cypseloides” OR “Cypsiurus” OR “Hirundapus” OR “Panyptila” OR “Streptoprocne” OR “Tachornis” OR “Tachymarptis” OR “Hemiprocne” OR “Abeillia” OR “Adelomyia” OR “Aglaeactis” OR “Agelaiocercus” OR “Amazilia” OR “Androdon” OR “Anopetia” OR “Anthocephala” OR “Anthracothorax” OR “Archilochus” OR “Atthis” OR “Augastes” OR “Avocettula” OR “Basilinna” OR “Boissonneaua” OR “Calliphlox” OR “Calypste” OR “Campylopterus” OR “Chaetocercus” OR “Chalcostigma” OR “Chalybura” OR “Chlorestes” OR “Chlorostilbon” OR “Chrysolampis” OR “Chrysuronia” OR “Clytolaema” OR “Coeligena” OR “Colibri” OR “Cyanophaia” OR “Cynanthus” OR “Discosura” OR “Doricha” OR “Doryfera” OR “Elvira” OR “Ensifera” OR “Eriocnemis” OR “Eugenes” OR “Eulampis” OR “Eulidia” OR “Eupetomena” OR “Eupherusa” OR “Eutoxeres” OR “Florisuga” OR “Glaucis” OR “Goethalsia” OR “Goldmania” OR “Haplophaedia” OR “Heliactin” OR “Heliangelus” OR “Heliodoxa” OR “Helioaster” OR “Heliothryx” OR “Hylocharis” OR “Juliamya” OR “Klais” OR “Lafresnaya” OR “Lampornis” OR “Lamprolaima” OR “Lepidopyga” OR “Lesbia” OR “Leucippus” OR “Leucochloris” OR “Lophornis” OR “Mellisuga” OR “Metallura” OR “Microchera” OR “Myrmia” OR “Myrtis” OR “Ocreatus” OR “Opisthoprora” OR “Oreonympha” OR “Oreotrochilus” OR “Orthorhyncus” OR “Oxygogon” OR “Panterpe” OR “Patagona” OR “Phaeochroa” OR “Phaethornis” OR “Phlogophilus” OR “Polyonymus” OR “Polymnus” OR “Pterophanes” OR “Ramphodon” OR “Ramphomicron” OR “Rhodopsis” OR “Sappho” OR “Schistes” OR “Selasphorus” OR “Sephanoides” OR “Stephanoxis” OR “Sternoclyta” OR “Taphrosphilus” OR “Thalaurania” OR “Thaumastura” OR “Threnetes” OR “Topaza” OR “Trochilus” OR “Urochroa” OR “Urosticte” OR “Apteryx” OR “Anthracoceros” OR “Buceros” OR “Bycanistes” OR “Lophoceros” OR “Penelopides” OR “Rhabdotorrhinus” OR “Rhyticeros” OR “Tockus” OR “Bucorvus” OR “Phoenixulus” OR “Rhinopomastus” OR “Upupa” OR “Antrostomus” OR “Caprimulgus” OR “Chordeiles” OR “Eleothreptes” OR “Eurostopodus” OR “Hydropsalis” OR “Lurocalis” OR “Lyncornis” OR “Nyctidromus” OR “Nyctiphrynus” OR “Nyctipolus” OR “Nyctiprogne” OR “Phalaenoptilus” OR “Setopagis” OR “Systellura” OR “Uropsalis” OR “Nyctibius” OR “Batrachostomus” OR “Podargus” OR “Steatornis” OR “Cariama” OR “Casuaris” OR “Dromaius” OR “Aethia” OR “Alca” OR “Alle” OR “Brachyrhamphus” OR “Cepphus” OR “Cerorhinca” OR “Fratricula” OR “Pinguinus” OR “Ptychoramphus” OR “Synthliboramphus” OR “Uria” OR “Burhinus” OR “Anarhynchus” OR “Charadrius” OR “Elseyornis” OR “Erythrogonys” OR “Hoploxypterus” OR “Oreopholus” OR “Peltohyas” OR “Phegornis” OR “Pluvialis” OR “Thinornis” OR “Vanellus” OR “Chionis” OR “Dromas” OR “Cursorius” OR “Glareola” OR “Rhinoptilus” OR “Stiltia” OR “Haematopus” OR “Actophilornis” OR “Hydrophasianus” OR “Irediparra” OR “Jacana” OR “Metopidius” OR “Microparra” OR “Anous” OR “Chlidonias” OR “Chroicocephalus” OR “Creagrus” OR “Gelochelidon” OR “Gygis” OR “Hydrocoleus” OR “Hydroprogne” OR “Ichthyaelus” OR “Larosterna” OR “Larus” OR “Leucophaeus” OR “Onychoprion” OR “Pagophila” OR “Phaetusa” OR “Rhodostethia” OR “Rissa” OR “Rynchops” OR “Sterna” OR “Sternula” OR “Thalasseus” OR “Xema” OR “Pedionomus” OR “Pluvianellus” OR “Pluvianus” OR “Himantopus” OR “Recurvirostra” OR “Nycticryptes” OR “Rostratula” OR “Actitis” OR “Arenaria” OR “Bartramia” OR “Calidris” OR “Coenocorypha” OR “Gallinago” OR “Limnodromus” OR “Limosa” OR “Lymnocryptes” OR “Numenius” OR “Phalaropus” OR “Scolopax” OR “Tringa” OR “Xenus” OR “Stereocorius” OR “Attagus” OR “Thinocorus” OR “Turnix” OR “Ciconia” OR “Ephippiorhynchus” OR “Jabiru” OR “Leptoptilos” OR “Mycteria” OR “Colius” OR “Urocolius” OR “Aleutroenas” OR “Alopecoenas” OR “Caloenas” OR “Chalcophaps” OR “Claravis” OR “Columba” OR “Columbina” OR “Didunculus” OR “Drepanoptila” OR “Ducula” OR “Ectopistes” OR “Gallicolumba” OR “Geopelia” OR “Geophaps” OR “Geotrygon” OR “Goura” OR “Gymnophaps” OR “Hemiphaga” OR “Henicophaps” OR “Leptotila” OR “Leptotrygon” OR “Leucosarcia” OR “Lopholaimus” OR “Macropygia” OR “Metriopelia” OR “Ocyphaps” OR “Oena” OR “Otidiophaps” OR “Patagioenas” OR “Petrophassa” OR “Pezophaps” OR “Phapitreron” OR “Phaps” OR “Ptilinopus” OR “Raphus” OR “Reinwardtoena” OR “Spilopelia” OR “Streptopelia” OR “Treron” OR “Trugon” OR “Turacoena” OR “Turtur” OR “Uropelia” OR “Zenaida” OR “Zentrygon” OR “Actenoides” OR “Alcedo” OR “Caridonax” OR “Ceryle” OR “Ceyx” OR “Chloroceryle” OR “Cittura” OR “Corythornis” OR “Dacelo” OR “Halcyon” OR “Ispidina” OR “Lacedo” OR “Megaceryle” OR “Melidora” OR “Syma” OR “Tanysiptera” OR “Todiramphus” OR “Atelornis” OR “Brachypteracias” OR “Geobiastes” OR “Coracias” OR “Eurystomus” OR “Merops” OR “Nyctyonis” OR “Baryphthengus” OR “Electron” OR “Eumomota” OR “Hylomanes” OR “Momotus” OR “Todus” OR “Cacomantis” OR “Carpococcyx” OR “Centropus” OR “Cercococcyx” OR “Chrysococcyx” OR “Clamator” OR “Coccyua” OR “Coccyzus” OR “Coua” OR “Crotophaga” OR “Cuculus” OR “Dasylophus” OR “Dromococcyx” OR “Eudynamys” OR “Geococcyx” OR “Guira” OR “Hierococcyx” OR “Morococcyx” OR “Neomorpus” OR “Pachycoccyx” OR “Phaenicophaeus” OR “Piaya” OR “Rhinortha” OR “Scythrops” OR “Surniculus” OR “Tapera” OR “Urodynamis” OR “Zanclostomus” OR “Dinornis” OR “Anomalopteryx” OR “Emeus” OR “Euryapteryx” OR “Eurypyga” OR “Rhynchotos” OR “Caracara” OR “Daptrius” OR “Falco” OR “Herpetotheres” OR “Ibycter” OR “Micrastur” OR “Microhierax” OR “Milvago” OR “Phalcoboenus” OR “Polihierax” OR “Spiziapteryx” OR “Aburria” OR “Chamaepetes” OR “Crax” OR “Mitu” OR “Nothocrax” OR “Oreophasis” OR “Ortalis” OR “Pauxi” OR “Penelope” OR “Penelopina” OR “Pipile” OR “Alectura” OR “Megapodius” OR “Acryllium” OR “Guttera” OR “Numida” OR “Callipepla” OR “Colinus” OR “Cyrtonyx” OR “Dendrortyx” OR “Odontophorus” OR “Oreortyx” OR “Philortyx” OR “Ptilopachus” OR “Rhynchortyx” OR “Alectoris” OR “Ammoperdix” OR “Arborophila” OR “Argusianus” OR “Bambusicola” OR “Bonasa” OR “Caloperdix” OR “Centrocercus” OR “Chrysolophus” OR “Coturnix” OR “Crossopylon” OR “Dendragapus” OR “Excalfactoria” OR “Falcipennis” OR “Francolinus” OR “Gallus” OR “Haemataortyx” OR “Ithaginis” OR “Lagopus” OR “Lerwa” OR “Lophophorus” OR “Lophura” OR “Lyururus” OR “Meleagris” OR “Pavo” OR “Peliperdix” OR “Perdix” OR “Phasianus” OR “Polyplectron” OR “Pternistis” OR “Pucrasia” OR “Rhizothera” OR “Rollulus” OR “Scleroptila” OR “Syrmaticus” OR “Tetrao” OR “Tetraogallus” OR “Tetraophasis” OR “Tragopan” OR

"Tympanuchus" OR "Gavia" OR "Aramus" OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amauornis" OR "Anurolimnas" OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia" OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrio" OR "Porphyriops" OR "Porzana" OR "Rallacula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura" OR "Leptosomus" OR "Corythaeola" OR "Corythaixoides" OR "Crinifer" OR "Musophaga" OR "Tauraco" OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax" OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis" OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis" OR "Smicrornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna" OR "Nesillas" OR "Aegithalos" OR "Leptopoecile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes" OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafr" OR "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhinophanes" OR "Callaeas" OR "Heteralocha" OR "Philesturnus" OR "Calyptrophilus" OR "Campephaga" OR "Ceblepyris" OR "Coracina" OR "Edolisoma" OR "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amaurospiza" OR "Cardinalis" OR "Caryothraustes" OR "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Paserina" OR "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus" OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus" OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia" OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus" OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus" OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias" OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza" OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis" OR "Nigrita" OR "Oryzospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga" OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calyptromena" OR "Cymbirhynchus" OR "Eurylaimus" OR "Psalisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chloris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra" OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorrhous" OR "Hemignathus" OR "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza" OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus" OR "Certhiasomus" OR "Certhiaxis" OR "Cinclodes" OR "Clibanornis" OR "Coryphistera" OR "Cranioleuca" OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geococcyx" OR "Geositta" OR "Glyphorhynchus" OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR "Limnortyx" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor" OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus" OR "Spartonoica" OR "Sylviorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonotus" OR "Thripadectes" OR "Thripophaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes" OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hyllopezus" OR "Myrmothera" OR "Alopochelidon" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon" OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psolidoprocne" OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR "Tachycineta" OR "Hyliota" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Amblycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives" OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus" OR "Lamprospiza" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar" OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura" OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus" OR "Leiostrix" OR

"Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochalopecteron" OR "Turdoides" OR "Bradypterus" OR "Cincloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malaconotus" OR "Nilaus" OR "Rhodophoneus" OR "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia" OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomyzon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes" OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis" OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius" OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cinclocerthia" OR "Dumetella" OR "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis" OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus" OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Campicoloides" OR "Cercotrichas" OR "Chamaetylas" OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia" OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiommela" OR "Myioparus" OR "Myophonus" OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola" OR "Sheppardia" OR "Sholicola" OR "Stiphornis" OR "Tarsiger" OR "Thamnodora" OR "Vauriella" OR "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkinogramma" OR "Leptocoma" OR "Nectarinia" OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleodryas" OR "Oriolus" OR "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR "Cyanistes" OR "Lophophanes" OR "Macholophus" OR "Melaniparus" OR "Melanochlora" OR "Pardaliparus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus" OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros" OR "Leiothlypis" OR "Limnithlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis" OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisospiza" OR "Atlapetes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza" OR "Melozone" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR "Pezopetes" OR "Pipilo" OR "Poocetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizelloides" OR "Xenospiza" OR "Zonotrichia" OR "Carospiza" OR "Gymnoris" OR "Hypocryptadius" OR "Montifringilla" OR "Onychostreuthus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR "Gampsorhynchus" OR "Illadopsis" OR "Jabouillea" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR "Malacopteron" OR "Napothea" OR "Pellorneum" OR "Rimtor" OR "Trichastoma" OR "Amalocichla" OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca" OR "Monachella" OR "Pachycephalopsis" OR "Peneoanthe" OR "Peneothello" OR "Petroica" OR "Poecilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyrannetes" OR "Xenopipo" OR "Erythropitta" OR "Hydornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalornis" OR "Euplectes" OR "Malimbus" OR "Ploceopasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR "Pnoepyga" OR "Microbates" OR "Poliopitila" OR "Ramphocaelus" OR "Garritornis" OR "Prunella" OR "Cinclosoma" OR "Ptilorhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla" OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Criniger" OR "Eurillas" OR "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus" OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus" OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pteroptochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynch" OR "Culicicapa" OR "Elminia" OR "Acridotheres" OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora" OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotornis" OR "Leucopsar" OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopar" OR "Streptocitta" OR "Sturnia" OR "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprositornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR "Drymophila" OR "Dysithamnus" OR "Epinecrophylia" OR "Euchrepomis" OR "Formicivora" OR "Frederickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR

"Megastictus" OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myrmoderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neoctantes" OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania" OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus" OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis" OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Chariospiza" OR "Chlorochrysa" OR "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba" OR "Compsospiza" OR "Compsotrhaupis" OR "Conirostrum" OR "Conotrhaupis" OR "Coryphospingus" OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delotrhaupis" OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Incaspiza" OR "Iridophanes" OR "Iridosornis" OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes" OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocomma" OR "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha" OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichotrhaupis" OR "Volatinia" OR "Wetmorethraupis" OR "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatorhinus" OR "Spelaornis" OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius" OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia" OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Troglodytes" OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus" OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola" OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Capsiempis" OR "Casiornis" OR "Cnemarchus" OR "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis" OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus" OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myiodynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiotriccus" OR "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes" OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis" OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis" OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis" OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR "Urocynchramus" OR "Artamella" OR "Calicalicus" OR "Cyanolanius" OR "Euryceros" OR "Falculea" OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris" OR "Anomalospiza" OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR "Vireo" OR "Vireolanus" OR "Zeledonina" OR "Apalopteron" OR "Dasycrotapha" OR "Heleia" OR "Lophozosterops" OR "Sterrhoptilus" OR "Yuhina" OR "Zosterops" OR "Zosterornis" OR "Agamia" OR "Ardea" OR "Ardeola" OR "Botaurus" OR "Bubulcus" OR "Butorides" OR "Cochlearius" OR "Dupetor" OR "Egretta" OR "Gorsachius" OR "Ixobrychus" OR "Nyctanassa" OR "Nycticorax" OR "Ptilerodius" OR "Syrigma" OR "Tigriornis" OR "Tigrisoma" OR "Zebulus" OR "Balaeniceps" OR "Pelecanus" OR "Scopus" OR "Bostrychia" OR "Eudocimus" OR "Geronticus" OR "Lophotibis" OR "Mesembrinibis" OR "Nipponia" OR "Phimosus" OR "Platalea" OR "Plegadis" OR "Pseudibis" OR "Theristicus" OR "Threskiornis" OR "Phaethon" OR "Phoeniconaias" OR "Phoenicoparrus" OR "Phoenicopterus" OR "Bucco" OR "Chelidoptera" OR "Hapalopectila" OR "Hypnelus" OR "Malacoptila" OR "Micromonacha" OR "Monasa" OR "Nonnula" OR "Notharchus" OR "Nystalus" OR "Capito" OR "Eubucco" OR "Brachygalba" OR "Galbacyrhynchus" OR "Galbula" OR "Jacamaralcyon" OR "Jacamerops" OR "Indicator" OR "Melichneutes" OR "Meligonomon" OR "Prodotiscus" OR "Buccanodon" OR "Gymnobucco" OR "Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus" OR "Psilopogon" OR "Blythipicus" OR "Campephilus" OR "Campethera" OR "Celeus" OR "Chloropicus" OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Dendrocopos" OR "Dendrocytes" OR "Dendropicus" OR "Dinopium" OR "Dryobates" OR "Dryocopus" OR "Gecinulus" OR "Geocolaptes" OR "Jynx" OR "Leiopicus" OR "Leuconotopicus" OR "Meiglyptes" OR "Melanerpes" OR "Micropternus" OR "Mulleripicus" OR "Nesocittes" OR "Picoides" OR "Piculus" OR "Picumnus" OR "Picus" OR "Reinwardtipicus" OR "Sasia" OR "Sphyrapicus" OR "Veniliornis" OR "Yungipicus" OR "Andigena" OR "Aulacorhynchus" OR "Pteroglossus" OR "Ramphastos" OR "Selenidera" OR "Semnornis" OR "Aechmophorus" OR "Podiceps" OR "Podilymbus" OR "Polio-

cephalus" OR "Rollandia" OR "Tachybaptus" OR "Diomedea" OR "Phoebastria" OR "Phoebetria" OR "Thalassarche" OR "Hydrobates" OR "Oceanodroma" OR "Fregetta" OR "Garrodia" OR "Nesofregetta" OR "Oceanites" OR "Pelagodroma" OR "Aphrodroma" OR "Ardena" OR "Bulweria" OR "Calonectris" OR "Daption" OR "Fulmarus" OR "Halobaena" OR "Macronectes" OR "Pachyptila" OR "Pagodroma" OR "Pelecanoides" OR "Procellaria" OR "Pseudobulweria" OR "Pterodroma" OR "Puffinus" OR "Thalassoica" OR "Cacatua" OR "Callocephalon" OR "Calyptorhynchus" OR "Eolophus" OR "Lophochroa" OR "Nymphicus" OR "Probosciger" OR "Alipiopsitta" OR "Amazona" OR "Anodorhynchus" OR "Ara" OR "Aratinga" OR "Bolborhynchus" OR "Brotogeris" OR "Conuropsis" OR "Cyanoliseus" OR "Cyanopsitta" OR "Derophtus" OR "Diopsittaca" OR "Enicognathus" OR "Eupsittula" OR "Forpus" OR "Graydidascalus" OR "Guarouba" OR "Guaruba" OR "Hapalopsittaca" OR "Leptosittaca" OR "Myiopsitta" OR "Nannopsittaca" OR "Orthopsittaca" OR "Pionites" OR "Pionopsitta" OR "Pionus" OR "Poicephalus" OR "Primolius" OR "Psilopsiagon" OR "Psittacara" OR "Psittacus" OR "Pyralia" OR "Pyrrhura" OR "Rhynchopsitta" OR "Thectocercus" OR "Touit" OR "Triclaria" OR "Agapornis" OR "Alisterus" OR "Aprosmictus" OR "Barnardius" OR "Bolbopsittacus" OR "Chalcopsitta" OR "Charmosyna" OR "Coracopsis" OR "Cyanoramphus" OR "Cyclopsitta" OR "Eclectus" OR "Eos" OR "Eunymphicus" OR "Geoffroyus" OR "Glossopsitta" OR "Lathamus" OR "Loriculus" OR "Lorius" OR "Melopsittacus" OR "Micropsitta" OR "Neophema" OR "Neopsephotus" OR "Neopsittacus" OR "Northiella" OR "Oreopsittacus" OR "Parvipsitta" OR "Pezoporus" OR "Phigys" OR "Platycercus" OR "Polytelis" OR "Prioniturus" OR "Prosopeia" OR "Psephotellus" OR "Psephotus" OR "Pseudeos" OR "Psittacella" OR "Psittacula" OR "Psittaculirostris" OR "Psitteuteles" OR "Psittinus" OR "Psittrichas" OR "Purpureicephalus" OR "Tanygnathus" OR "Trichoglossus" OR "Vini" OR "Nestor" OR "Strigops" OR "Pterocles" OR "Syrrhaptes" OR "Rhea" OR "Aptenodytes" OR "Eudypetes" OR "Eudyptula" OR "Megadyptes" OR "Pygoscelis" OR "Spheniscus" OR "Aegolius" OR "Asio" OR "Athene" OR "Bubo" OR "Glaucidium" OR "Ketupa" OR "Lophotrix" OR "Megascops" OR "Micrathene" OR "Ninox" OR "Otus" OR "Pseudoscops" OR "Psiloscops" OR "Pulsatrix" OR "Sceloglaux" OR "Scotopelia" OR "Strix" OR "Surnia" OR "Xenoglaux" OR "Phodilus" OR "Tyto" OR "Aepyornis" OR "Mullerornis" OR "Struthio" OR "Anhinga" OR "Fregata" OR "Leucocarbo" OR "Microcarbo" OR "Nannopterum" OR "Phalacrocorax" OR "Morus" OR "Sula" OR "Crypturellus" OR "Eudromia" OR "Nothocercus" OR "Nothoprocta" OR "Nothura" OR "Rhynchotus" OR "Tinamotis" OR "Tinamus" OR "Apalharactes" OR "Apaloderma" OR "Harpactes" OR "Pharomachrus" OR "Trogon" OR "Tetrastes" OR "Hesperiphona" AND ("experiment*" OR "manipulat*")) OR AK= (((("green*" OR "herb*" OR "aromatic*") AND "nest*" AND ("bird*" OR "aves" OR "avian" OR "ornithol*" OR "passerine*" OR "passeriform*" OR "songbird*" OR "Accipiter" OR "Aegyptius" OR "Aquila" OR "Aviceda" OR "Busarellus" OR "Butastur" OR "Buteo" OR "Buteogallus" OR "Chondrohierax" OR "Circaetus" OR "Circus" OR "Clanga" OR "Elanoides" OR "Elanus" OR "Gampsonyx" OR "Geranoaetus" OR "Geranospiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR "Haliastur" OR "Harpagus" OR "Harpia" OR "Hieraetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco" OR "Leptodon" OR "Leucopternis" OR "Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus" OR "Necrosyrtes" OR "Neophron" OR "Nisaetus" OR "Parabuteo" OR "Pernis" OR "Pithecopaga" OR "Polemaetus" OR "Pseudastur" OR "Rostrhamus" OR "Rupornis" OR "Sarcogyps" OR "Spilornis" OR "Spizaetus" OR "Stephanoaetus" OR "Terathopius" OR "Torgos" OR "Trigonoceps" OR "Cathartes" OR "Coragyps" OR "Gymnogyps" OR "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix" OR "Alopochen" OR "Amazonetta" OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR "Bucephala" OR "Cairina" OR "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloephaga" OR "Clangula" OR "Coscoroba" OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR "Histrionicus" OR "Hymenolaimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca" OR "Marmaronetta" OR "Melanitta" OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neochen" OR "Netta" OR "Nettapus" OR "Nomonyx" OR "Oxyura" OR "Plectropterus" OR "Polysticta" OR "Pteronetta" OR "Sarkidiornis" OR "Sibirionetta" OR "Somateria" OR "Spatula" OR "Tachyeres" OR "Tadorna" OR "Thalassornis" OR "Anhima" OR "Chauna" OR "Anseranas" OR "Aegothales" OR "Aerodramus" OR "Aeronautes" OR "Apus" OR "Chaetura" OR "Collocalia" OR "Cypseloides" OR "Cypsiurus" OR "Hirundapus" OR "Panyptila" OR "Streptoprocne" OR "Tachornis" OR "Tachymarptis" OR "Hemiprocne" OR "Abeilia" OR "Adelomyia" OR "Aglaeactis" OR "Aglaiocercus" OR "Amazilia" OR "Androdon" OR "Anopetia" OR "Anthocephala" OR "Anthracothorax" OR "Archilochus" OR "Atthis" OR "Augastes" OR "Avocettula" OR "Basilinna" OR "Boissonneaua" OR "Calliphlox" OR "Calypte" OR "Campylopterus" OR "Chaetocercus" OR "Chalcostigma" OR "Chalybura" OR "Chlorestes" OR "Chlorostilbon" OR "Chrysolampis" OR "Chrysurnia" OR "Clytolaema" OR "Coeligena" OR "Colibri" OR "Cyanophaia" OR "Cynanthus" OR "Discosura" OR "Doricha" OR "Doryfera" OR "Elvira" OR "Ensifera" OR "Eriocnemis" OR "Eugenes" OR "Eulampis" OR "Eulidia" OR "Eupetomena" OR "Eupherusa" OR "Eutoxeres" OR "Florisuga" OR "Glaucis" OR "Goethalsia" OR "Goldmania" OR "Haplophaedia" OR "Heliactin" OR "Heliangelus" OR "Heliodoxa" OR "Heliomaster" OR "Heliophryx" OR "Hylocharis" OR "Juliamyia" OR "Klais" OR "Lafresnaya" OR "Lampornis" OR "Lamprolaima" OR "Lepidopyga" OR "Lesbia" OR "Leucippus" OR "Leucochloris" OR "Lophornis" OR "Mellisuga" OR "Metallura" OR "Microchera" OR "Myrmia" OR "Myrtis" OR "Ocreatus" OR "Opisthoprora" OR "Oreonympha" OR "Oreotrochilus" OR "Orthorhynchus" OR "Oxyopogon" OR "Panterpe" OR "Patagona" OR "Phaeochroa" OR "Phaethornis" OR "Phlogophilus" OR "Polyonymus" OR "Polytmus" OR "Pterophanes" OR "Ramphodon" OR "Ramphomicron" OR "Rhodopsis" OR "Sappho" OR "Schistes" OR "Selasphorus" OR "Sephanioides" OR "Stephanoxis" OR "Sternoclyta" OR "Taphrospilus" OR "Thalurania" OR "Thaumastura" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR "Rhabdotorrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus" OR

"Upupa" OR "Antrastomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus" OR "Hydropsalis" OR "Lurocalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus" OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius" OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariamā" OR "Casuarius" OR "Dromaius" OR "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cerorhinca" OR "Fratrercula" OR "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus" OR "Charadrius" OR "Elseyornis" OR "Erythrogonys" OR "Hoploxypterus" OR "Oreopholus" OR "Peltohyas" OR "Phegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelochelidon" OR "Gygis" OR "Hydrocoloeus" OR "Hydroprogne" OR "Ichthyæetus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR "Pagophila" OR "Phætusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula" OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR "Recurvirostra" OR "Nycticryphes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR "Calidris" OR "Coenocorypha" OR "Gallinago" OR "Limnodromus" OR "Limosa" OR "Lymnocyrtus" OR "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagus" OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chalcophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula" OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR "Gymnophaps" OR "Hemiphaga" OR "Henicophaps" OR "Leptotila" OR "Leptotrygon" OR "Leucosarcia" OR "Lopholaimus" OR "Macropygia" OR "Metriopelia" OR "Ocyphaps" OR "Oena" OR "Otidiphaps" OR "Patagioenas" OR "Petrophassa" OR "Pezophaps" OR "Phapitreron" OR "Phaps" OR "Ptilinopus" OR "Raphus" OR "Reinwardtoena" OR "Spilopelia" OR "Streptopelia" OR "Treron" OR "Trugon" OR "Turacoena" OR "Turtur" OR "Uropelia" OR "Zenaida" OR "Zentrygon" OR "Actenoides" OR "Alcedo" OR "Caridonax" OR "Ceryle" OR "Ceyx" OR "Chloroceryle" OR "Cittura" OR "Corythornis" OR "Dacelo" OR "Halcyon" OR "Ispidina" OR "Lacedo" OR "Megaceryle" OR "Melidora" OR "Syma" OR "Tanysiptera" OR "Todiramphus" OR "Atelornis" OR "Brachypteracias" OR "Geobiastes" OR "Coracias" OR "Eurystomus" OR "Merops" OR "Nyctornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes" OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR "Chrysococcyx" OR "Clamator" OR "Coccyua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus" OR "Dasylophus" OR "Dromococcyx" OR "Eudynamys" OR "Geococcyx" OR "Guirā" OR "Hierococcyx" OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicophaeus" OR "Piaya" OR "Rhinortha" OR "Scythrops" OR "Surniculus" OR "Taperā" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR "Anomalopteryx" OR "Emeus" OR "Euryapteryx" OR "Eurypyga" OR "Rhynochetos" OR "Caracara" OR "Daptrius" OR "Falco" OR "Herpetotheres" OR "Ibycter" OR "Micrastur" OR "Microhierax" OR "Milvago" OR "Phalcoboenus" OR "Polihierax" OR "Spiziapteryx" OR "Aburria" OR "Chamaepetes" OR "Crax" OR "Mitu" OR "Nothocrax" OR "Oreophasis" OR "Ortalis" OR "Pauxi" OR "Penelope" OR "Penelopina" OR "Pipile" OR "Alectura" OR "Megapodius" OR "Acryllium" OR "Guttera" OR "Numida" OR "Callipepla" OR "Colinus" OR "Cyrtonyx" OR "Dendrortyx" OR "Odontophorus" OR "Oreortyx" OR "Philortyx" OR "Ptilopachus" OR "Rhynchortyx" OR "Alectoris" OR "Ammoperdix" OR "Arborophila" OR "Argusianus" OR "Bambusicola" OR "Bonasa" OR "Caloperdix" OR "Centrocerus" OR "Chrysolophus" OR "Coturnix" OR "Crossoptilon" OR "Dendragapus" OR "Excalfactoria" OR "Falcipennis" OR "Francolinus" OR "Gallus" OR "Haematortyx" OR "Ithaginis" OR "Lagopus" OR "Lerwa" OR "Lophophorus" OR "Lophura" OR "Lyrurus" OR "Meleagris" OR "Pavo" OR "Peliperdix" OR "Perdix" OR "Phasianus" OR "Polyplectron" OR "Pternistis" OR "Pucrasia" OR "Rhizothera" OR "Rollulus" OR "Scleroptila" OR "Syrmaticus" OR "Tetrao" OR "Tetraogallus" OR "Tetraophasis" OR "Tragopan" OR "Tympanuchus" OR "Gavia" OR "Aramus" OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amauornis" OR "Anurolimnas" OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia" OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrio" OR "Porphyriops" OR "Porzana" OR "Rallacula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura" OR "Leptosomus" OR "Corythaëola" OR "Corythaixoides" OR "Crinifer" OR "Musophaga" OR "Tauraco" OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax" OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis" OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis" OR "Smicromis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna" OR "Nesillas" OR "Aegithalos" OR "Leptopoecile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes" OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafrā" OR "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhynchophanes" OR "Callaëas" OR "Heteralocha" OR "Philesturnus" OR "Calyptophilus" OR "Campephaga" OR "Cebilepyris" OR "Coracina" OR "Edolisoma" OR "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amaurospiza" OR "Cardinalis" OR "Caryothraustes" OR "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Passe-

rina" OR "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus" OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus" OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia" OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus" OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus" OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrolaema" OR "Procnias" OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza" OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis" OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR "Pytilia" OR "Spermophaga" OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calypomena" OR "Cymbirhynchus" OR "Eurylaimus" OR "Psarisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chloris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra" OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorhous" OR "Hemignathus" OR "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR "Magnumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza" OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus" OR "Certhiasomus" OR "Certhiaxis" OR "Cinclodes" OR "Clibanornis" OR "Coryphistera" OR "Cranioleuca" OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geocerthia" OR "Geositta" OR "Glyphorhynchus" OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR "Limnortites" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor" OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus" OR "Spartonoica" OR "Sylviorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonimus" OR "Thripadectes" OR "Thriponphaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes" OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hyllopezus" OR "Myrmothera" OR "Alopochelidon" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon" OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psalidoprocne" OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR "Tachycineta" OR "Hylia" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Amblycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives" OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus" OR "Lamprospiza" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar" OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura" OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus" OR "Leiothrix" OR "Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochalopteron" OR "Turdoides" OR "Bradypterus" OR "Cincloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malacconotus" OR "Nilaus" OR "Rhodophoneus" OR "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia" OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomyzon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes" OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis" OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius" OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cinlocerthia" OR "Dumetella" OR "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis" OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus" OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Campicoloides" OR "Cercotrichas" OR "Chamaetylas" OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia" OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiomela" OR "Myioparus" OR "Myophonus" OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola" OR "Sheppardia" OR "Sholicola" OR "Stiphornis" OR "Tarsiger"

OR "Thamnolaia" OR "Vauriella" OR "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkinogramma" OR "Leptocoma" OR "Nectarinia" OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleadryas" OR "Oriolus" OR "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR "Cyanistes" OR "Lophophanes" OR "Machlolophus" OR "Melaniparus" OR "Melanochlora" OR "Pardaliparus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus" OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros" OR "Leiothlypis" OR "Limnithlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis" OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisospiza" OR "Atlapetes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza" OR "Melozona" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR "Pezopetes" OR "Pipilo" OR "Poocetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizelloides" OR "Xenospiza" OR "Zonotrichia" OR "Carpospiza" OR "Gymnoris" OR "Hypocryptadius" OR "Montifringilla" OR "Onychostreus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR "Gampsorhynchus" OR "Illadopsis" OR "Jabouillea" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR "Malacopteron" OR "Napothera" OR "Pellorneum" OR "Rimator" OR "Trichastoma" OR "Amalocichla" OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca" OR "Monachella" OR "Pachycephalopsis" OR "Peneoenanthe" OR "Peneothello" OR "Petroica" OR "Poecilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyrannetes" OR "Xenopipo" OR "Erythropitta" OR "Hydornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalornis" OR "Euplectes" OR "Malimbus" OR "Plocepasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR "Pnoepyga" OR "Microbates" OR "Polioptila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR "Cinclosoma" OR "Ptilorrhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla" OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Criniger" OR "Eurillas" OR "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus" OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus" OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pterotochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorhynchus" OR "Culicicapa" OR "Elminia" OR "Acridotheres" OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora" OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprolornis" OR "Leucopsar" OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprositornis" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR "Drymophila" OR "Dysithamnus" OR "Epinecrophylia" OR "Euchrepomis" OR "Formicivora" OR "Frederickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus" OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myrmoderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes" OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania" OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus" OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis" OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Chariospiza" OR "Chlorochrysa" OR "Chlorophanes" OR "Chlorornis" OR "Chrysithlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba" OR "Compsospiza" OR "Compsorthraupis" OR "Conirostrum" OR "Conothraupis" OR "Coryphospingus" OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis" OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Inaspiza" OR "Iridophanes" OR "Iridosornis" OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes" OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocomma" OR "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha" OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatrhinus" OR "Spelaornis" OR "Stachyridopsis" OR

"Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius" OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia" OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Troglodytes" OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus" OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola" OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Capsiempis" OR "Casiornis" OR "Cnemarchus" OR "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis" OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus" OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myiodynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiotriccus" OR "Myiozetetes" OR "Neopipo" OR "Neoxolmis" OR "Nephelomyias" OR "Ochthoeca" OR "Ochthornis" OR "Oncostoma" OR "Ornithion" OR "Phaeomyias" OR "Philohydor" OR "Phyllomyias" OR "Phylloscartes" OR "Piprites" OR "Pitangus" OR "Platyrinchus" OR "Poecilotriccus" OR "Pogonotriccus" OR "Polioxolmis" OR "Polystictus" OR "Pseudelaenia" OR "Pseudocolopteryx" OR "Pseudotriccus" OR "Pyrocephalus" OR "Pyrrhomyias" OR "Ramphotrigon" OR "Rhynchocyclus" OR "Rhytipterna" OR "Satrapa" OR "Sayornis" OR "Serpophaga" OR "Silvicultrix" OR "Sirystes" OR "Stigmatura" OR "Sublegatus" OR "Suiriri" OR "Tachuris" OR "Taeniotriccus" OR "Todiostrostrum" OR "Tolmomyias" OR "Tumbezia" OR "Tyrannopsis" OR "Tyrannulus" OR "Tyrannus" OR "Uromyias" OR "Xenotriccus" OR "Xolmis" OR "Zimmerius" OR "Urocynchramus" OR "Artamella" OR "Calicalicus" OR "Cyanolanius" OR "Euryceros" OR "Falculea" OR "Hypositta" OR "Leptopterus" OR "Mystacornis" OR "Newtonia" OR "Oriolia" OR "Philentoma" OR "Prionops" OR "Pseudobias" OR "Schetba" OR "Tephrodornis" OR "Tylas" OR "Vanga" OR "Xenopirostris" OR "Anomalospiza" OR "Vidua" OR "Cyclarhis" OR "Erpornis" OR "Hylophilus" OR "Pteruthius" OR "Vireo" OR "Vireolanius" OR "Zeledonia" OR "Apalopteron" OR "Dasycrotapha" OR "Heleia" OR "Lophozosterops" OR "Sterrhoptilus" OR "Yuhina" OR "Zosterops" OR "Zosterornis" OR "Agamia" OR "Ardea" OR "Ardeola" OR "Botaurus" OR "Bubulcus" OR "Butorides" OR "Cochlearius" OR "Dupetor" OR "Egretta" OR "Gorsachius" OR "Ixobrychus" OR "Nyctanassa" OR "Nycticorax" OR "Pilharodius" OR "Syrigma" OR "Tigriornis" OR "Tigrisoma" OR "Zebrilus" OR "Balaeniceps" OR "Pelecanus" OR "Scopus" OR "Bostrychia" OR "Eudocimus" OR "Geronticus" OR "Lophotibis" OR "Mesembrinibis" OR "Nipponia" OR "Phimosus" OR "Platalea" OR "Plegadis" OR "Pseudibis" OR "Theristicus" OR "Threskiornis" OR "Phaethon" OR "Phoeniconaias" OR "Phoenicoparrus" OR "Phoenicopterus" OR "Bucco" OR "Chelidoptera" OR "Haploptila" OR "Hypnelus" OR "Malacoptila" OR "Micromonacha" OR "Monasa" OR "Nonnula" OR "Notharchus" OR "Nystalus" OR "Capito" OR "Eubucco" OR "Brachygalba" OR "Galbalcyrrhynchus" OR "Galbula" OR "Jacamaralcyon" OR "Jacamerops" OR "Indicator" OR "Melichneutes" OR "Meligonomon" OR "Prodotiscus" OR "Buccanodon" OR "Gymnobucco" OR "Lybius" OR "Pogoniulus" OR "Stactolaema" OR "Trachyphonus" OR "Psilopogon" OR "Blythipicus" OR "Campephilus" OR "Campethera" OR "Celeus" OR "Chloropicus" OR "Chrysocolaptes" OR "Chrysophlegma" OR "Colaptes" OR "Dendrocopos" OR "Dendrocytes" OR "Dendropicus" OR "Dinopium" OR "Dryobates" OR "Dryocopus" OR "Gecinulus" OR "Geocolaptes" OR "Jynx" OR "Leiopicus" OR "Leuconotopicus" OR "Meiglyptes" OR "Melanerpes" OR "Micropternus" OR "Mulleripicus" OR "Nesocittes" OR "Picoides" OR "Piculus" OR "Picumnus" OR "Picus" OR "Reinwardtipicus" OR "Sasia" OR "Sphyrapicus" OR "Veniliornis" OR "Yungipicus" OR "Andigena" OR "Aulacorhynchus" OR "Pteroglossus" OR "Ramphastos" OR "Selenidera" OR "Semnornis" OR "Aechmophorus" OR "Podiceps" OR "Podilymbus" OR "Poliocephalus" OR "Rollandia" OR "Tachybaptus" OR "Diomedea" OR "Phoebastria" OR "Phoebetria" OR "Thalassarche" OR "Hydrobates" OR "Oceanodroma" OR "Fregetta" OR "Garrodia" OR "Nesofregetta" OR "Oceanites" OR "Pelagodroma" OR "Aphrodroma" OR "Ardenna" OR "Bulweria" OR "Calonectris" OR "Daption" OR "Fulmarus" OR "Halobaena" OR "Macronectes" OR "Pachyptila" OR "Pagodroma" OR "Pelecanoides" OR "Procellaria" OR "Pseudobulweria" OR "Pterodroma" OR "Puffinus" OR "Thalassoica" OR "Cacatua" OR "Callocephalon" OR "Calyptorhynchus" OR "Eolophus" OR "Lophochroa" OR "Nymphicus" OR "Probosciger" OR "Alipiopsitta" OR "Amazona" OR "Anodorhynchus" OR "Ara" OR "Aratinga" OR "Bolborhynchus" OR "Brotogeris" OR "Conuropsis" OR "Cyanoliseus" OR "Cyanopsitta" OR "Derophtus" OR "Diopsittaca" OR "Enicognathus" OR "Eupsittula" OR "Forpus" OR "Graydidascalus" OR "Guarouba" OR "Guaruba" OR "Hapalopsittaca" OR "Leptosittaca" OR "Myiopsitta" OR "Nannopsittaca" OR "Orthopsittaca" OR "Pionites" OR "Pionopsitta" OR "Pionus" OR "Poicephalus" OR "Primolius" OR "Psilopsiagon" OR "Psittacara" OR "Psittacus" OR "Pyralia" OR "Pyrrhura" OR "Rhynchopsitta" OR "Thectocercus" OR "Touit" OR "Triclaria" OR "Agapornis" OR "Alisterus" OR "Aprosmictus" OR "Barnardius" OR "Bolbopsittacus" OR "Chalcopsitta" OR "Charmosyna" OR "Coracopsis" OR "Cyanoramphus" OR "Cyclopsitta" OR "Eclectus" OR "Eos" OR "Eunymphicus" OR "Geoffroyus" OR "Glossopsitta" OR "Lathamus" OR "Loriculus" OR "Lorius" OR "Melopsittacus" OR "Micropsitta" OR "Neophema" OR "Neopsephotus" OR "Neopsittacus" OR "Northiella" OR "Oreopsittacus" OR "Parvipsitta" OR "Pezoporus" OR "Phigys" OR "Platycercus" OR "Polytelis" OR "Prioniturus" OR "Prosopieia" OR "Psephotellus" OR "Psephotus" OR "Pseudeos" OR "Psittacella" OR "Psittacula" OR "Psittaculirostris" OR "Psittuteles" OR "Psittinus" OR "Psittichas" OR "Purpureicephalus" OR "Tanygnathus" OR "Trichoglossus" OR "Vini" OR "Nestor" OR "Strigops" OR "Pterocles" OR "Syrrhaptes" OR "Rhea"

OR "Aptenodytes" OR "Eudytes" OR "Eudiptula" OR "Megadyptes" OR "Pygoscelis" OR "Spheniscus" OR "Aegolius" OR "Asio" OR "Athene" OR "Bubo" OR "Glaucidium" OR "Ketupa" OR "Lophostrix" OR "Megascops" OR "Micrathene" OR "Ninox" OR "Otus" OR "Pseudoscops" OR "Psilosops" OR "Pulsatrix" OR "Sceloglaux" OR "Scotopelia" OR "Strix" OR "Surnia" OR "Xenoglaux" OR "Phodilus" OR "Tyto" OR "Aepyornis" OR "Mullerornis" OR "Struthio" OR "Anhinga" OR "Fregata" OR "Leucocarbo" OR "Microcarbo" OR "Nannopterum" OR "Phalacrocorax" OR "Morus" OR "Sula" OR "Crypturellus" OR "Eudromia" OR "Nothocercus" OR "Nothoprocta" OR "Nothura" OR "Rhynchotus" OR "Tinamotis" OR "Tinamus" OR "Apalharpactes" OR "Apaloderma" OR "Harpactes" OR "Pharomachrus" OR "Trogon" OR "Tetrastes" OR "Hesperiphona") AND ("experiment" OR "manipulat*"))>

The search was also conducted on SCOPUS using Advance Search Option and Title Abstract and Keyword (TITLE-ABS-KEY) on the same dates using the string

<TITLE-ABS-KEY (((("green*" OR "herb*" OR "aromatic*") AND "nest*" AND ("bird*" OR "aves" OR "avian" OR "ornithol*" OR "passerine" OR "passeriform" OR "songbird*" OR "Accipiter" OR "Aegyptius" OR "Aquila" OR "Aviceda" OR "Busarellus" OR "Butastur" OR "Buteo" OR "Buteogallus" OR "Chondrohierax" OR "Circaetus" OR "Circus" OR "Clanga" OR "Elanoides" OR "Elanus" OR "Gampsonyx" OR "Geranoaetus" OR "Geranospiza" OR "Gypaetus" OR "Gypohierax" OR "Gyps" OR "Haliaeetus" OR "Haliastur" OR "Harpagus" OR "Harpia" OR "Hieraetus" OR "Ictinaetus" OR "Ictinia" OR "Kaupifalco" OR "Leptodon" OR "Leucopternis" OR "Lophotriorchis" OR "Milvus" OR "Morphnarchus" OR "Morphnus" OR "Necrosyrtes" OR "Neophron" OR "Nisaetus" OR "Parabuteo" OR "Pernis" OR "Pithecophaga" OR "Polemaetus" OR "Pseudastur" OR "Rostrhamus" OR "Rupornis" OR "Sarcogyps" OR "Spilornis" OR "Spizaetus" OR "Stephanoaetus" OR "Terathopius" OR "Torgos" OR "Trigonoceps" OR "Cathartes" OR "Coragyps" OR "Gymnogyps" OR "Sarcoramphus" OR "Vultur" OR "Pandion" OR "Sagittarius" OR "Aix" OR "Alopochen" OR "Amazonetta" OR "Anas" OR "Anser" OR "Asarcornis" OR "Aythya" OR "Branta" OR "Bucephala" OR "Cairina" OR "Callonetta" OR "Cereopsis" OR "Chen" OR "Chenonetta" OR "Chloephaga" OR "Clangula" OR "Coscoroba" OR "Cyanochen" OR "Cygnus" OR "Dendrocygna" OR "Heteronetta" OR "Histrionicus" OR "Hymenolaimus" OR "Lophodytes" OR "Lophonetta" OR "Malacorhynchus" OR "Mareca" OR "Marmaronetta" OR "Melanitta" OR "Merganetta" OR "Mergellus" OR "Mergus" OR "Neochen" OR "Netta" OR "Nettapus" OR "Nomonyx" OR "Oxyura" OR "Plectropterus" OR "Polysticta" OR "Pteronetta" OR "Sarkidiornis" OR "Sibirionetta" OR "Somateria" OR "Spatula" OR "Tachyeres" OR "Tadorna" OR "Thalassornis" OR "Anhima" OR "Chauna" OR "Anseranas" OR "Aegotheles" OR "Aerodramus" OR "Aeronautes" OR "Apus" OR "Chaetura" OR "Collocalia" OR "Cypseloides" OR "Cypsiurus" OR "Hirundapus" OR "Panyptila" OR "Streptoprocne" OR "Tachornis" OR "Tachymarpis" OR "Hemiprocne" OR "Abeillia" OR "Adelomyia" OR "Aglaeactis" OR "Aglaiocercus" OR "Amazilia" OR "Androdon" OR "Anopetia" OR "Anthocephala" OR "Anthracothorax" OR "Archilochus" OR "Atthis" OR "Augastes" OR "Avocettula" OR "Basilinna" OR "Boissonneaua" OR "Calliphlox" OR "Calypte" OR "Campylopterus" OR "Chaetocercus" OR "Chalcostigma" OR "Chalybura" OR "Chlorestes" OR "Chlorostilbon" OR "Chrysolampis" OR "Chrysuronia" OR "Clytolaema" OR "Coeligena" OR "Colibri" OR "Cyanophaia" OR "Cynanthus" OR "Discosura" OR "Doricha" OR "Doryfera" OR "Elvira" OR "Ensifera" OR "Eriocnemis" OR "Eugenes" OR "Eulampis" OR "Eulidia" OR "Eupetomena" OR "Eupherusa" OR "Eutoxeres" OR "Florisuga" OR "Glaucis" OR "Goethalsia" OR "Goldmania" OR "Haplophaedia" OR "Heliactin" OR "Heliangelus" OR "Heliodoxa" OR "Heliomaster" OR "Heliothryx" OR "Hylocharis" OR "Juliamyia" OR "Klais" OR "Lafresnaya" OR "Lampornis" OR "Lamprolaima" OR "Lepidopyga" OR "Lesbia" OR "Leucippus" OR "Leucochloris" OR "Lophornis" OR "Mellisuga" OR "Metallura" OR "Microchera" OR "Myrmia" OR "Myrtis" OR "Ocreatus" OR "Opisthoprora" OR "Oreonympha" OR "Oreotrochilus" OR "Orthorhynchus" OR "Oxygogon" OR "Panterpe" OR "Patagona" OR "Phaeochroa" OR "Phaethornis" OR "Phlogophilus" OR "Polyonymus" OR "Polytmus" OR "Pterophanes" OR "Ramphodon" OR "Ramphomicron" OR "Rhodopis" OR "Sappho" OR "Schistes" OR "Selasphorus" OR "Sephanooides" OR "Stephanoxis" OR "Sternoclyta" OR "Taphrospilus" OR "Thalurania" OR "Thaumastura" OR "Threnetes" OR "Topaza" OR "Trochilus" OR "Urochroa" OR "Urosticte" OR "Apteryx" OR "Anthracoceros" OR "Buceros" OR "Bycanistes" OR "Lophoceros" OR "Penelopides" OR "Rhabdotorrhinus" OR "Rhyticeros" OR "Tockus" OR "Bucorvus" OR "Phoeniculus" OR "Rhinopomastus" OR "Upupa" OR "Antrostomus" OR "Caprimulgus" OR "Chordeiles" OR "Eleothreptus" OR "Eurostopodus" OR "Hydropsalis" OR "Lurolalis" OR "Lyncornis" OR "Nyctidromus" OR "Nyctiphrynus" OR "Nyctipolus" OR "Nyctiprogne" OR "Phalaenoptilus" OR "Setopagis" OR "Systellura" OR "Uropsalis" OR "Nyctibius" OR "Batrachostomus" OR "Podargus" OR "Steatornis" OR "Cariama" OR "Casuarius" OR "Dromaius" OR "Aethia" OR "Alca" OR "Alle" OR "Brachyramphus" OR "Cepphus" OR "Cerorhinca" OR "Fratrercula" OR "Pinguinus" OR "Ptychoramphus" OR "Synthliboramphus" OR "Uria" OR "Burhinus" OR "Anarhynchus" OR "Charadrius" OR "Elseynornis" OR "Erythrogonys" OR "Hoploxypterus" OR "Oreopholus" OR "Peltodytes" OR "Phegornis" OR "Pluvialis" OR "Thinornis" OR "Vanellus" OR "Chionis" OR "Dromas" OR "Cursorius" OR "Glareola" OR "Rhinoptilus" OR "Stiltia" OR "Haematopus" OR "Actophilornis" OR "Hydrophasianus" OR "Irediparra" OR "Jacana" OR "Metopidius" OR "Microparra" OR "Anous" OR "Chlidonias" OR "Chroicocephalus" OR "Creagrus" OR "Gelochelidon" OR "Gygis" OR "Hydrocoloeus" OR "Hydroprogne" OR "Ichthyiaetus" OR "Larosterna" OR "Larus" OR "Leucophaeus" OR "Onychoprion" OR "Pagophila" OR "Phaetusa" OR "Rhodostethia" OR "Rissa" OR "Rynchops" OR "Sterna" OR "Sternula" OR "Thalasseus" OR "Xema" OR "Pedionomus" OR "Pluvianellus" OR "Pluvianus" OR "Himantopus" OR "Recurvirostra" OR "Nycticryphes" OR "Rostratula" OR "Actitis" OR "Arenaria" OR "Bartramia" OR "Calidris" OR "Coenocorypha" OR "Gallinago" OR

"Limnodromus" OR "Limosa" OR "Lymnocryptes" OR "Numenius" OR "Phalaropus" OR "Scolopax" OR "Tringa" OR "Xenus" OR "Stercorarius" OR "Attagis" OR "Thinocorus" OR "Turnix" OR "Ciconia" OR "Ephippiorhynchus" OR "Jabiru" OR "Leptoptilos" OR "Mycteria" OR "Colius" OR "Urocolius" OR "Alectroenas" OR "Alopecoenas" OR "Caloenas" OR "Chalcophaps" OR "Claravis" OR "Columba" OR "Columbina" OR "Didunculus" OR "Drepanoptila" OR "Ducula" OR "Ectopistes" OR "Gallicolumba" OR "Geopelia" OR "Geophaps" OR "Geotrygon" OR "Goura" OR "Gymnophaps" OR "Hemiphaga" OR "Henicophaps" OR "Leptotila" OR "Leptotrygon" OR "Leucosarcia" OR "Lopholaimus" OR "Macropygia" OR "Metriopelia" OR "Ocyphaps" OR "Oena" OR "Otidiophaps" OR "Patagioenas" OR "Petrophassa" OR "Pezophaps" OR "Phapitreron" OR "Phaps" OR "Ptilinopus" OR "Raphus" OR "Reinwardtoena" OR "Spilopelia" OR "Streptopelia" OR "Treron" OR "Trugon" OR "Turacoena" OR "Turtur" OR "Uropelia" OR "Zenaida" OR "Zentrygon" OR "Actenoides" OR "Alcedo" OR "Caridonax" OR "Ceryle" OR "Ceyx" OR "Chloroceryle" OR "Cittura" OR "Corythornis" OR "Dacelo" OR "Halcyon" OR "Ispidina" OR "Lacedo" OR "Megaceryle" OR "Melidora" OR "Syma" OR "Tanysiptera" OR "Todiramphus" OR "Atelornis" OR "Brachypteracias" OR "Geobiastes" OR "Coracias" OR "Eurystomus" OR "Merops" OR "Nyctornis" OR "Baryphthengus" OR "Electron" OR "Eumomota" OR "Hylomanes" OR "Momotus" OR "Todus" OR "Cacomantis" OR "Carpococcyx" OR "Centropus" OR "Cercococcyx" OR "Chrysococcyx" OR "Clamator" OR "Coccyua" OR "Coccyzus" OR "Coua" OR "Crotophaga" OR "Cuculus" OR "Dasylophus" OR "Dromococcyx" OR "Eudynamys" OR "Geococcyx" OR "Guira" OR "Hierococcyx" OR "Morococcyx" OR "Neomorphus" OR "Pachycoccyx" OR "Phaenicophaeus" OR "Piaya" OR "Rhinortha" OR "Scythrops" OR "Surniculus" OR "Tapera" OR "Urodynamis" OR "Zanclostomus" OR "Dinornis" OR "Anomalopteryx" OR "Emeus" OR "Euryapteryx" OR "Eurypyga" OR "Rhynochetos" OR "Caracara" OR "Daptrius" OR "Falco" OR "Herpetotheres" OR "Ibycter" OR "Micrastur" OR "Microhierax" OR "Milvago" OR "Phalcoboenus" OR "Polihierax" OR "Spiziapteryx" OR "Aburria" OR "Chamaepetes" OR "Crax" OR "Mitu" OR "Nothocrax" OR "Oreophasis" OR "Ortalis" OR "Pauxi" OR "Penelope" OR "Penelopina" OR "Pipile" OR "Alectura" OR "Megapodius" OR "Acryllium" OR "Guttera" OR "Numida" OR "Callipepla" OR "Colinus" OR "Cyrtonyx" OR "Dendrortyx" OR "Odontophorus" OR "Oreortyx" OR "Philortyx" OR "Ptilopachus" OR "Rhynchortyx" OR "Alectoris" OR "Ammoperdix" OR "Arborophila" OR "Argusianus" OR "Bambusicola" OR "Bonasa" OR "Caloperdix" OR "Centrocerus" OR "Chrysolophus" OR "Coturnix" OR "Crossoptilon" OR "Dendragapus" OR "Excalfactoria" OR "Falcipennis" OR "Francolinus" OR "Gallus" OR "Haematortyx" OR "Ithaginis" OR "Lagopus" OR "Lerwa" OR "Lophophorus" OR "Lophura" OR "Lyrurus" OR "Meleagris" OR "Pavo" OR "Peliperdix" OR "Perdix" OR "Phasianus" OR "Polyplectron" OR "Pternistis" OR "Pucrasia" OR "Rhizothera" OR "Rollulus" OR "Scleroptila" OR "Syrmaticus" OR "Tetrao" OR "Tetraogallus" OR "Tetraophasis" OR "Tragopan" OR "Tympanuchus" OR "Gavia" OR "Aramus" OR "Anthropoides" OR "Antigone" OR "Balearica" OR "Bugeranus" OR "Grus" OR "Leucogeranus" OR "Heliornis" OR "Psophia" OR "Aenigmatolimnas" OR "Amaurolimnas" OR "Amauornis" OR "Anurolimnas" OR "Aramides" OR "Atlantisia" OR "Coturnicops" OR "Crex" OR "Dryolimnas" OR "Eulabeornis" OR "Fulica" OR "Gallicrex" OR "Gallinula" OR "Gallirallus" OR "Laterallus" OR "Lewinia" OR "Micropygia" OR "Neocrex" OR "Nesoclopeus" OR "Paragallinula" OR "Pardirallus" OR "Porphyrion" OR "Porphyrion" OR "Porzana" OR "Rallula" OR "Rallina" OR "Rallus" OR "Tribonyx" OR "Canirallus" OR "Sarothrura" OR "Leptosomus" OR "Corythaedra" OR "Corythaixoides" OR "Crinifer" OR "Musophaga" OR "Tauraco" OR "Opisthocomus" OR "Afrotis" OR "Ardeotis" OR "Chlamydotis" OR "Neotis" OR "Otis" OR "Tetrax" OR "Acanthisitta" OR "Pachyplichas" OR "Traversia" OR "Xenicus" OR "Acanthiza" OR "Acanthornis" OR "Aethomyias" OR "Aphelocephala" OR "Calamanthus" OR "Gerygone" OR "Hylacola" OR "Neosericornis" OR "Oreoscopus" OR "Origma" OR "Pachycare" OR "Pycnoptilus" OR "Pyrrholaemus" OR "Sericornis" OR "Smicrornis" OR "Acrocephalus" OR "Arundinax" OR "Calamonastides" OR "Hippolais" OR "Iduna" OR "Nesillas" OR "Aegithalos" OR "Leptopoecile" OR "Psaltiriparus" OR "Aegithina" OR "Alaemon" OR "Alauda" OR "Alaudala" OR "Ammomanes" OR "Calandrella" OR "Calendulauda" OR "Chersomanes" OR "Eremophila" OR "Eremopterix" OR "Galerida" OR "Lullula" OR "Melanocorypha" OR "Mirafr" OR "Pinarocorys" OR "Spizocorys" OR "Artamus" OR "Gymnorhina" OR "Melloria" OR "Bombycilla" OR "Buphagus" OR "Calcarius" OR "Plectrophenax" OR "Rhynchophanes" OR "Callaenas" OR "Heteralocha" OR "Philesturnus" OR "Calyptrophilus" OR "Campephaga" OR "Ceblepyris" OR "Coracina" OR "Edolisoma" OR "Lalage" OR "Malindangia" OR "Pericrocotus" OR "Amauropsiza" OR "Cardinalis" OR "Caryothraustes" OR "Chlorothraupis" OR "Cyanocompsa" OR "Cyanoloxia" OR "Granatellus" OR "Habia" OR "Paserina" OR "Periporphyrus" OR "Pheucticus" OR "Piranga" OR "Spiza" OR "Certhia" OR "Salpornis" OR "Abroscopus" OR "Cettia" OR "Horornis" OR "Phyllergates" OR "Tesia" OR "Urosphena" OR "Chloropsis" OR "Cinclus" OR "Apalis" OR "Artisornis" OR "Camaroptera" OR "Cisticola" OR "Eremomela" OR "Heliolais" OR "Hypergerus" OR "Neomixis" OR "Oreolais" OR "Orthotomus" OR "Phragmacia" OR "Poliolais" OR "Prinia" OR "Schistolais" OR "Urolais" OR "Urorhipis" OR "Climacteris" OR "Cormobates" OR "Conopophaga" OR "Struthidea" OR "Aphelocoma" OR "Calocitta" OR "Cissa" OR "Coloeus" OR "Corvus" OR "Cyanocitta" OR "Cyanocorax" OR "Cyanolyca" OR "Cyanopica" OR "Dendrocitta" OR "Garrulus" OR "Gymnorhinus" OR "Nucifraga" OR "Perisoreus" OR "Pica" OR "Platylophus" OR "Podoces" OR "Psilorhinus" OR "Ptilostomus" OR "Pyrrhocorax" OR "Urocissa" OR "Ampelioides" OR "Ampelion" OR "Carpornis" OR "Cephalopterus" OR "Conioptilon" OR "Cotinga" OR "Doliornis" OR "Gymnoderus" OR "Haematoderus" OR "Lipaugus" OR "Perissocephalus" OR "Phoenicircus" OR "Phytotoma" OR "Pipreola" OR "Porphyrulaema" OR "Procnias" OR "Pyroderus" OR "Querula" OR "Rupicola" OR "Snowornis" OR "Xipholena" OR "Zaratornis" OR "Dasyornis" OR "Dicaeum" OR "Prionochilus" OR "Dicrurus" OR "Donacobius" OR "Dulus" OR "Emberiza" OR "Amadina" OR "Amandava" OR "Cryptospiza" OR "Erythrura" OR "Estrilda" OR "Euodice" OR "Euschistospiza" OR "Lagonosticta" OR "Lonchura" OR "Mandingoa" OR "Neochmia" OR "Nesocharis" OR "Nigrita" OR "Ortygospiza" OR "Parmoptila" OR "Pyrenestes" OR

"Pytilia" OR "Spermophaga" OR "Stagonopleura" OR "Taeniopygia" OR "Uraeginthus" OR "Calyptomena" OR "Cymbirhynchus" OR "Eurylaimus" OR "Psarisomus" OR "Serilophus" OR "Smithornis" OR "Chamaeza" OR "Formicarius" OR "Acanthis" OR "Agraphospiza" OR "Akialoa" OR "Bucanetes" OR "Carduelis" OR "Carpodacus" OR "Chloris" OR "Chlorodrepanis" OR "Chlorophonia" OR "Chrysocorythus" OR "Coccothraustes" OR "Crithagra" OR "Drepanis" OR "Eophona" OR "Euphonia" OR "Fringilla" OR "Haemorhous" OR "Hemignathus" OR "Himatione" OR "Leucosticte" OR "Linaria" OR "Linurgus" OR "Loxia" OR "Loxioides" OR "Loxops" OR "Magumma" OR "Melamprosops" OR "Oreomystis" OR "Paroreomyza" OR "Pinicola" OR "Procarduelis" OR "Pseudonestor" OR "Psittirostra" OR "Pyrrhoplectes" OR "Pyrrhula" OR "Rhodopechys" OR "Rhodospiza" OR "Serinus" OR "Spinus" OR "Telespiza" OR "Anabacerthia" OR "Anabazenops" OR "Ancistrops" OR "Anumbius" OR "Aphrastura" OR "Asthenes" OR "Automolus" OR "Berlepschia" OR "Campylorhamphus" OR "Certhiasomus" OR "Certhiaxis" OR "Cinclodes" OR "Clibanornis" OR "Coryphistera" OR "Cranioleuca" OR "Deconychura" OR "Dendrexetastes" OR "Dendrocincla" OR "Dendrocolaptes" OR "Dendroplex" OR "Drymornis" OR "Drymotoxeres" OR "Furnarius" OR "Geocerthia" OR "Geositta" OR "Glyphorhynchus" OR "Heliobletus" OR "Hellmayrea" OR "Hylexetastes" OR "Lepidocolaptes" OR "Leptasthenura" OR "Limnortyx" OR "Limnornis" OR "Lochmias" OR "Margarornis" OR "Mazaria" OR "Megaxenops" OR "Metopothrix" OR "Microxenops" OR "Nasica" OR "Ochetorhynchus" OR "Phacellodomus" OR "Philydor" OR "Phleocryptes" OR "Premnoplex" OR "Premnornis" OR "Pseudasthenes" OR "Pseudocolaptes" OR "Pseudoseisura" OR "Pygarrhichas" OR "Roraimia" OR "Schoeniophylax" OR "Sclerurus" OR "Sittasomus" OR "Spartonoica" OR "Sylviorthorhynchus" OR "Synallaxis" OR "Syndactyla" OR "Tarphonomus" OR "Thripadectes" OR "Thripophaga" OR "Upucerthia" OR "Xenerpestes" OR "Xenops" OR "Xiphocolaptes" OR "Xiphorhynchus" OR "Grallaria" OR "Grallaricula" OR "Hyllopyz" OR "Myrmothera" OR "Alopochelidon" OR "Atticora" OR "Cecropis" OR "Delichon" OR "Haplochelidon" OR "Hirundo" OR "Neochelidon" OR "Notiochelidon" OR "Orochelidon" OR "Petrochelidon" OR "Phedina" OR "Progne" OR "Psolidoprocne" OR "Pseudhirundo" OR "Pseudochelidon" OR "Ptyonoprogne" OR "Riparia" OR "Stelgidopteryx" OR "Tachycineta" OR "Hylia" OR "Hypocolius" OR "Agelaioides" OR "Agelaius" OR "Agelasticus" OR "Amblycercus" OR "Amblyramphus" OR "Anumara" OR "Cacicus" OR "Chrysomus" OR "Curaeus" OR "Dives" OR "Dolichonyx" OR "Euphagus" OR "Gnorimopsar" OR "Gymnomystax" OR "Hypopyrrhus" OR "Icterus" OR "Lamprospiza" OR "Leistes" OR "Macroagelaius" OR "Molothrus" OR "Nesopsar" OR "Oreopsar" OR "Psarocolius" OR "Pseudoleistes" OR "Quiscalus" OR "Sturnella" OR "Xanthocephalus" OR "Xanthopsar" OR "Icteria" OR "Ifrita" OR "Irena" OR "Corvinella" OR "Eurocephalus" OR "Lanius" OR "Actinodura" OR "Argya" OR "Garrulax" OR "Grammatoptila" OR "Heterophasia" OR "Ianthocincla" OR "Laniellus" OR "Leiothrix" OR "Liocichla" OR "Minla" OR "Pterorhinus" OR "Trochaloxypterus" OR "Troglodytes" OR "Bradypterus" OR "Cinloramphus" OR "Helopsaltes" OR "Locustella" OR "Megalurus" OR "Poodytes" OR "Machaerirhynchus" OR "Graueria" OR "Hylia" OR "Macrosphenus" OR "Melocichla" OR "Sylvietta" OR "Chlorophoneus" OR "Dryoscopus" OR "Laniarius" OR "Malaconotus" OR "Nilaus" OR "Rhodophoneus" OR "Tchagra" OR "Telophorus" OR "Amytornis" OR "Chenorhamphus" OR "Clytomyias" OR "Malurus" OR "Stipiturus" OR "Melampitta" OR "Melanocharis" OR "Oedistoma" OR "Toxorhamphus" OR "Melanopareia" OR "Acanthagenys" OR "Acanthorhynchus" OR "Anthochaera" OR "Anthornis" OR "Caligavis" OR "Entomyzon" OR "Epthianura" OR "Glycifolia" OR "Gymnomyza" OR "Lichmera" OR "Manorina" OR "Melidectes" OR "Melilestes" OR "Meliphaga" OR "Melipotes" OR "Melithreptus" OR "Myzomela" OR "Nesoptilotis" OR "Philemon" OR "Phylidonyris" OR "Prothemadera" OR "Ptiloprora" OR "Ptilotula" OR "Pycnopygius" OR "Timeliopsis" OR "Xanthotis" OR "Menura" OR "Allenia" OR "Cinclocerthia" OR "Dumetella" OR "Margarops" OR "Melanoptila" OR "Melanotis" OR "Mimus" OR "Oreoscoptes" OR "Ramphocinclus" OR "Toxostoma" OR "Lamprospiza" OR "Mitrospingus" OR "Moho" OR "Mohoua" OR "Arses" OR "Carterornis" OR "Clytorhynchus" OR "Grallina" OR "Hypothymis" OR "Monarcha" OR "Myiagra" OR "Symposiachrus" OR "Terpsiphone" OR "Trochocercus" OR "Anthus" OR "Macronyx" OR "Motacilla" OR "Alethe" OR "Anthipes" OR "Brachypteryx" OR "Calliope" OR "Campicoloides" OR "Cercotrichas" OR "Chamaetylas" OR "Copsychus" OR "Cossypha" OR "Cyanoptila" OR "Cyornis" OR "Enicurus" OR "Erithacus" OR "Eumyias" OR "Ficedula" OR "Fraseria" OR "Irania" OR "Larvivora" OR "Leonardina" OR "Luscinia" OR "Melaenornis" OR "Monticola" OR "Muscicapa" OR "Myiomela" OR "Myiopus" OR "Myophonus" OR "Myrmecocichla" OR "Niltava" OR "Oenanthe" OR "Phoenicurus" OR "Pogonocichla" OR "Saxicola" OR "Sheppardia" OR "Sholicola" OR "Stiphodon" OR "Tarsiger" OR "Thamnolaea" OR "Vauriella" OR "Aethopyga" OR "Anabathmis" OR "Anthreptes" OR "Arachnothera" OR "Chalcomitra" OR "Cinnyris" OR "Cyanomitra" OR "Deleornis" OR "Hedydipna" OR "Kurochkegama" OR "Leptocoma" OR "Nectarinia" OR "Daphoenositta" OR "Nesospingus" OR "Nicator" OR "Notiomystis" OR "Aleodryas" OR "Oriolus" OR "Pitohui" OR "Sphecotheres" OR "Turnagra" OR "Colluricincla" OR "Falcunculus" OR "Melanorectes" OR "Pachycephala" OR "Pseudorectes" OR "Panurus" OR "Cicinnurus" OR "Diphyllodes" OR "Manucodia" OR "Paradisaea" OR "Ptiloris" OR "Paramythia" OR "Pardalotus" OR "Baeolophus" OR "Cephalopyrus" OR "Cyanistes" OR "Lophophanes" OR "Macholophus" OR "Melaniparus" OR "Melanochlora" OR "Pardaliparus" OR "Parus" OR "Periparus" OR "Poecile" OR "Pseudopodoces" OR "Sittiparus" OR "Sylviparus" OR "Basileuterus" OR "Cardellina" OR "Catharopeza" OR "Dendroica" OR "Geothlypis" OR "Helmitheros" OR "Leiothlypis" OR "Limnithlypis" OR "Mniotilta" OR "Myioborus" OR "Myiothlypis" OR "Oporornis" OR "Oreothlypis" OR "Parkesia" OR "Protonotaria" OR "Seiurus" OR "Setophaga" OR "Vermivora" OR "Aimophila" OR "Ammodramus" OR "Amphispiza" OR "Arremon" OR "Arremonops" OR "Artemisospiza" OR "Atlantes" OR "Calamospiza" OR "Chlorospingus" OR "Chondestes" OR "Junco" OR "Melospiza" OR "Melozone" OR "Oreothraupis" OR "Oriturus" OR "Passerculus" OR "Passerella" OR "Peucaea" OR "Pezopetes" OR "Pipilo" OR "Poocetes" OR "Pselliophorus" OR "Rhynchospiza" OR "Spizella" OR "Spizelloides" OR "Xenospiza" OR

"Zonotrichia" OR "Carospiza" OR "Gymnoris" OR "Hypocryptadius" OR "Montifringilla" OR "Onychostruthus" OR "Passer" OR "Petronia" OR "Pyrgilauda" OR "Alcippe" OR "Gampsorhynchus" OR "Illadopsis" OR "Jabouilleia" OR "Kenopia" OR "Laticilla" OR "Malacocincla" OR "Malacopteron" OR "Napothea" OR "Pellorneum" OR "Rimator" OR "Trichastoma" OR "Amalocichla" OR "Drymodes" OR "Eopsaltria" OR "Eugerygone" OR "Heteromyias" OR "Melanodryas" OR "Microeca" OR "Monachella" OR "Pachycephalopsis" OR "Peneoanthe" OR "Peneothello" OR "Petroica" OR "Poecilodryas" OR "Tregellasia" OR "Peucedramus" OR "Microligea" OR "Phaenicophilus" OR "Xenoligea" OR "Phylloscopus" OR "Picathartes" OR "Antilophia" OR "Ceratopipra" OR "Chiroxiphia" OR "Chloropipo" OR "Corapipo" OR "Cryptopipo" OR "Heterocercus" OR "Ilicura" OR "Lepidothrix" OR "Machaeropterus" OR "Manacus" OR "Masius" OR "Neopelma" OR "Pipra" OR "Pseudopipra" OR "Tyranneutes" OR "Xenopipo" OR "Erythropitta" OR "Hydroornis" OR "Pitta" OR "Batis" OR "Platysteira" OR "Anaplectes" OR "Bubalornis" OR "Euplectes" OR "Malimbus" OR "Plocepasser" OR "Ploceus" OR "Quelea" OR "Sporopipes" OR "Pnoepyga" OR "Microbates" OR "Polioptila" OR "Ramphocaenus" OR "Garritornis" OR "Prunella" OR "Cinclosoma" OR "Ptilorhoa" OR "Phainopepla" OR "Phainoptila" OR "Ptiliogonys" OR "Ailuroedus" OR "Amblyornis" OR "Ptilonorhynchus" OR "Sericulus" OR "Alophoixus" OR "Andropadus" OR "Arizelocichla" OR "Atimastillas" OR "Baeopogon" OR "Bleda" OR "Chlorocichla" OR "Criniger" OR "Eurillas" OR "Hemixos" OR "Hypsipetes" OR "Iole" OR "Ixos" OR "Neolestes" OR "Phyllastrephus" OR "Pycnonotus" OR "Spizixos" OR "Stelgidillas" OR "Tricholestes" OR "Regulus" OR "Anthoscopus" OR "Auriparus" OR "Remiz" OR "Rhagologus" OR "Eleoscytalopus" OR "Liosceles" OR "Psilorhamphus" OR "Pteroptochos" OR "Scelorchilus" OR "Scytalopus" OR "Teledromas" OR "Rhipidura" OR "Rhodinocichla" OR "Sapayoa" OR "Sitta" OR "Spindalis" OR "Chelidorphynx" OR "Culicicapa" OR "Elminia" OR "Acridotheres" OR "Agropsar" OR "Ampeliceps" OR "Aplonis" OR "Basilornis" OR "Cinnyricinclus" OR "Creatophora" OR "Gracula" OR "Gracupica" OR "Hartlaubius" OR "Hylopsar" OR "Lamprotornis" OR "Leucopsar" OR "Mino" OR "Notopholia" OR "Onychognathus" OR "Pastor" OR "Poeoptera" OR "Rhabdornis" OR "Sarcops" OR "Scissirostrum" OR "Speculipastor" OR "Spodiopsar" OR "Streptocitta" OR "Sturnia" OR "Sturnus" OR "Chamaea" OR "Chleuasicus" OR "Chloropeta" OR "Chrysomma" OR "Conostoma" OR "Fulvetta" OR "Lioparus" OR "Paradoxornis" OR "Pseudoalcippe" OR "Psittiparus" OR "Rhopophilus" OR "Sinosuthora" OR "Suthora" OR "Sylvia" OR "Teretistris" OR "Akletos" OR "Ampelornis" OR "Aprositoris" OR "Batara" OR "Cercomacra" OR "Cercomacroides" OR "Cymbilaimus" OR "Dichrozona" OR "Drymophila" OR "Dysithamnus" OR "Epinecrophylia" OR "Euchrepomis" OR "Formicivora" OR "Frederickena" OR "Gymnocichla" OR "Gymnopithys" OR "Hafferia" OR "Herpsilochmus" OR "Hylophylax" OR "Hypocnemis" OR "Hypocnemoides" OR "Hypoedaleus" OR "Isleria" OR "Mackenziaena" OR "Megastictus" OR "Microrhopias" OR "Myrmeciza" OR "Myrmelastes" OR "Myrmoborus" OR "Myrmochanes" OR "Myrmoderus" OR "Myrmophylax" OR "Myrmorchilus" OR "Myrmornis" OR "Myrmotherula" OR "Neotantes" OR "Oneillornis" OR "Percnostola" OR "Phaenostictus" OR "Phlegopsis" OR "Pithys" OR "Poliocrania" OR "Pygiptila" OR "Pyriglena" OR "Rhegmatorhina" OR "Rhopias" OR "Rhopornis" OR "Sakesphorus" OR "Sciaphylax" OR "Sclateria" OR "Sipia" OR "Taraba" OR "Thamnistes" OR "Thamnomanes" OR "Thamnophilus" OR "Willisornis" OR "Xenornis" OR "Anisognathus" OR "Bangsia" OR "Buthraupis" OR "Calochaetes" OR "Catamblyrhynchus" OR "Catamenia" OR "Charitospiza" OR "Chlorochrysa" OR "Chlorophanes" OR "Chlorornis" OR "Chrysothlypis" OR "Cissopis" OR "Cnemoscopus" OR "Coereba" OR "Compsospiza" OR "Compsotrappis" OR "Conirostrum" OR "Conotrappis" OR "Coryphospingus" OR "Creurgops" OR "Cyanerpes" OR "Cyanicterus" OR "Cypsnagra" OR "Dacnis" OR "Delothraupis" OR "Diglossa" OR "Diuca" OR "Dolospingus" OR "Donacospiza" OR "Dubusia" OR "Emberizoides" OR "Embernagra" OR "Eucometis" OR "Euneornis" OR "Geospiza" OR "Gubernatrix" OR "Haplospiza" OR "Hemispingus" OR "Hemithraupis" OR "Heterospingus" OR "Inaspiza" OR "Iridophanes" OR "Iridosornis" OR "Lanio" OR "Lophospingus" OR "Loxigilla" OR "Loxipasser" OR "Melanodera" OR "Melanospiza" OR "Melopyrrha" OR "Nemosia" OR "Nephelornis" OR "Oreomanes" OR "Oryzoborus" OR "Parkerthraustes" OR "Paroaria" OR "Phrygilus" OR "Piezorina" OR "Pipraeidea" OR "Poospiza" OR "Pyrrhocoma" OR "Ramphocelus" OR "Rhodospingus" OR "Saltator" OR "Saltatricula" OR "Schistochlamys" OR "Sericossypha" OR "Sicalis" OR "Sporophila" OR "Stephanophorus" OR "Tachyphonus" OR "Tangara" OR "Tersina" OR "Thlypopsis" OR "Thraupis" OR "Tiaris" OR "Trichothraupis" OR "Volatinia" OR "Wetmorethraupis" OR "Xenodacnis" OR "Xenospingus" OR "Tichodroma" OR "Macronus" OR "Pomatiorhinus" OR "Spelaornis" OR "Stachyridopsis" OR "Stachyris" OR "Iodopleura" OR "Laniisoma" OR "Laniocera" OR "Myiobius" OR "Onychorhynchus" OR "Oxyruncus" OR "Pachyramphus" OR "Schiffornis" OR "Terenotriccus" OR "Tityra" OR "Xenopsaris" OR "Campylorhynchus" OR "Cantorchilus" OR "Catherpes" OR "Cinnycerthia" OR "Cistothorus" OR "Cyphorhinus" OR "Henicorhina" OR "Microcerculus" OR "Odontorchilus" OR "Pheugopedius" OR "Salpinctes" OR "Thryomanes" OR "Thryophilus" OR "Thryothorus" OR "Troglodytes" OR "Uropsila" OR "Catharus" OR "Cichlopsis" OR "Entomodestes" OR "Geokichla" OR "Hylocichla" OR "Ixoreus" OR "Myadestes" OR "Neocossyphus" OR "Ridgwayia" OR "Sialia" OR "Stizorhina" OR "Turdus" OR "Zoothera" OR "Agriornis" OR "Alectrurus" OR "Anairetes" OR "Aphanotriccus" OR "Arundinicola" OR "Atalotriccus" OR "Attila" OR "Camptostoma" OR "Casiempis" OR "Casiornis" OR "Cnemarchus" OR "Cnemotriccus" OR "Cnipodectes" OR "Colonia" OR "Colorhamphus" OR "Conopias" OR "Contopus" OR "Corythopsis" OR "Culicivora" OR "Deltarhynchus" OR "Elaenia" OR "Empidonax" OR "Empidonomus" OR "Euscarthmus" OR "Fluvicola" OR "Griseotyrannus" OR "Gubernetes" OR "Hemitriccus" OR "Heteroxolmis" OR "Hirundinea" OR "Hymenops" OR "Inezia" OR "Knipolegus" OR "Lathrotriccus" OR "Legatus" OR "Leptopogon" OR "Lessonia" OR "Lophotriccus" OR "Machetornis" OR "Mecocerculus" OR "Megarynchus" OR "Mionectes" OR "Mitrephanes" OR "Muscigralla" OR "Muscisaxicola" OR "Myiarchus" OR "Myiodynastes" OR "Myiopagis" OR "Myiophobus" OR "Myiornis" OR "Myiotheretes" OR "Myiortic-

OR “Myiozetetes” OR “Neopipo” OR “Neoxolmis” OR “Nephemomyias” OR “Ochthoeca” OR “Ochthornis” OR “Oncostoma” OR “Ornithion” OR “Phaeomyias” OR “Philohydor” OR “Phyllosomyias” OR “Phylloscartes” OR “Piprites” OR “Pitangus” OR “Platyrinchus” OR “Poecilotriccus” OR “Pogonotriccus” OR “Polioxolmis” OR “Polystictus” OR “Pseudelaenia” OR “Pseudocolopteryx” OR “Pseudotriccus” OR “Pyrocephalus” OR “Pyrrhomyias” OR “Ramphotrigon” OR “Rhynchocyclus” OR “Rhytipterna” OR “Satrapa” OR “Sayornis” OR “Serpophaga” OR “Silvicultrix” OR “Sirystes” OR “Stigmatura” OR “Sublegatus” OR “Suiriri” OR “Tachuris” OR “Taeniotriccus” OR “Todiostrostrum” OR “Tolmomyias” OR “Tumbezia” OR “Tyrannopsis” OR “Tyrannulus” OR “Tyrannus” OR “Uromyias” OR “Xenotriccus” OR “Xolmis” OR “Zimmerius” OR “Urocynchramus” OR “Artamella” OR “Calicalicus” OR “Cyanolanius” OR “Euryceros” OR “Falculea” OR “Hypositta” OR “Leptopterus” OR “Mystacornis” OR “Newtonia” OR “Oriolia” OR “Philentoma” OR “Prionops” OR “Pseudobias” OR “Schetba” OR “Tephrodornis” OR “Tylas” OR “Vanga” OR “Xenopirostris” OR “Anomalospiza” OR “Vidua” OR “Cyclarhis” OR “Erpornis” OR “Hylophilus” OR “Pteruthius” OR “Vireo” OR “Vireolanius” OR “Zeledonia” OR “Apalopteron” OR “Dasycrotapha” OR “Heleia” OR “Lophozosterops” OR “Sternhoptilus” OR “Yuhina” OR “Zosterops” OR “Zosterornis” OR “Agamia” OR “Ardea” OR “Ardeola” OR “Botaurus” OR “Bubulcus” OR “Butorides” OR “Cochlearius” OR “Dupetor” OR “Egretta” OR “Gorsachius” OR “Ixobrychus” OR “Nyctanassa” OR “Nycticorax” OR “Ptilherodius” OR “Syrigma” OR “Tigriornis” OR “Tigrisoma” OR “Zebrilus” OR “Balaeniceps” OR “Pelecanus” OR “Scopus” OR “Bostrychia” OR “Eudocimus” OR “Geronticus” OR “Lophotibis” OR “Mesembrinibis” OR “Nipponia” OR “Phimosus” OR “Platalea” OR “Plegadis” OR “Pseudibis” OR “Theristicus” OR “Threskiornis” OR “Phaethon” OR “Phoeniconaias” OR “Phoenicoparrus” OR “Phoenicopterus” OR “Bucco” OR “Chelidoptera” OR “Hapaloptila” OR “Hypnelus” OR “Malacoptila” OR “Micromonacha” OR “Monasa” OR “Nonnula” OR “Notharchus” OR “Nystalus” OR “Capito” OR “Eubucco” OR “Brachygalba” OR “Galbalcyrhynchus” OR “Galbula” OR “Jacamaralcyon” OR “Jacamerops” OR “Indicator” OR “Melichneutes” OR “Meligonomon” OR “Prodotiscus” OR “Buccanodon” OR “Gymnobucco” OR “Lybius” OR “Pogoniulus” OR “Stactolaema” OR “Trachyphonus” OR “Psilopogon” OR “Blythipicus” OR “Campephilus” OR “Campethera” OR “Celeus” OR “Chloropicus” OR “Chrysocolaptes” OR “Chrysophlegma” OR “Colaptes” OR “Dendrocopos” OR “Dendrocopetes” OR “Dendropicos” OR “Dinopium” OR “Dryobates” OR “Dryocopus” OR “Gecinulus” OR “Geocolaptes” OR “Jynx” OR “Leipicus” OR “Leuconotopicus” OR “Meiglyptes” OR “Melanerpes” OR “Micropternus” OR “Mulleripicus” OR “Nesocittes” OR “Picoides” OR “Piculus” OR “Picumnus” OR “Picus” OR “Reinwardtipicus” OR “Sasia” OR “Sphyrapicus” OR “Veniliornis” OR “Yungipicus” OR “Andigena” OR “Aulacorhynchus” OR “Pteroglossus” OR “Ramphastos” OR “Selenidera” OR “Semnornis” OR “Aechmophorus” OR “Podiceps” OR “Podilymbus” OR “Poliocephalus” OR “Rollandia” OR “Tachybaptus” OR “Diomedea” OR “Phoebastria” OR “Phoebetria” OR “Thalassarche” OR “Hydrobates” OR “Oceanodroma” OR “Fregatta” OR “Garrodia” OR “Nesofregatta” OR “Oceanites” OR “Pelagodroma” OR “Aphrodroma” OR “Ardena” OR “Bulweria” OR “Calonectris” OR “Daption” OR “Fulmarus” OR “Halobaena” OR “Macronectes” OR “Pachyptila” OR “Pagodroma” OR “Pelecanoides” OR “Procellaria” OR “Pseudobulweria” OR “Pterodroma” OR “Puffinus” OR “Thalassoica” OR “Cacatua” OR “Callocephalon” OR “Calyptorhynchus” OR “Eolophus” OR “Lophochroa” OR “Nymphicus” OR “Probosciger” OR “Alipiopsitta” OR “Amazona” OR “Anodorhynchus” OR “Ara” OR “Aratinga” OR “Bolborhynchus” OR “Brotogeris” OR “Conuropsis” OR “Cyanoliseus” OR “Cyanopsitta” OR “Deroptyus” OR “Diopsittaca” OR “Enicognathus” OR “Eupsittula” OR “Forpus” OR “Graydidascalus” OR “Guarouba” OR “Guaruba” OR “Hapalopsittaca” OR “Leptosittaca” OR “Myiopsitta” OR “Nannopsittaca” OR “Orthopsittaca” OR “Pionites” OR “Pionopsitta” OR “Pionus” OR “Poicephalus” OR “Primolius” OR “Psilopsiagon” OR “Psittacara” OR “Psittacus” OR “Pyrilia” OR “Pyrrhura” OR “Rhynchopsitta” OR “Thectocercus” OR “Touit” OR “Triclaria” OR “Agapornis” OR “Alisterus” OR “Aprosmictus” OR “Barnardius” OR “Bolbopsittacus” OR “Chalcopsitta” OR “Charmosyna” OR “Coracopsis” OR “Cyanoramphus” OR “Cyclopsitta” OR “Eclectus” OR “Eos” OR “Eunymphicus” OR “Geoffroyus” OR “Glossopsitta” OR “Lathamus” OR “Loriculus” OR “Lorius” OR “Melopsittacus” OR “Micropsitta” OR “Neophema” OR “Neopsephotus” OR “Neopsittacus” OR “Northiella” OR “Oreopsittacus” OR “Parvipsitta” OR “Pezoporus” OR “Phigys” OR “Platycercus” OR “Polytelis” OR “Prioniturus” OR “Prosopieia” OR “Psephotellus” OR “Psephotus” OR “Pseudeos” OR “Psittacella” OR “Psittacula” OR “Psittaculirostris” OR “Psitteuteles” OR “Psittinus” OR “Psittichas” OR “Purpureicephalus” OR “Tanygnathus” OR “Trichoglossus” OR “Vini” OR “Nestor” OR “Strigops” OR “Pterocles” OR “Syrhaptes” OR “Rhea” OR “Aptenodytes” OR “Eudypetes” OR “Eudyptula” OR “Megadyptes” OR “Pygoscelis” OR “Spheniscus” OR “Aegolius” OR “Asio” OR “Athene” OR “Bubo” OR “Glaucidium” OR “Ketupa” OR “Lophostrix” OR “Megascops” OR “Micrathene” OR “Ninox” OR “Otus” OR “Pseudoscops” OR “Psiloscops” OR “Pulsatrix” OR “Sceloglaux” OR “Scotopelia” OR “Strix” OR “Surnia” OR “Xenoglaux” OR “Phodilus” OR “Tyto” OR “Aepyornis” OR “Mullerornis” OR “Struthio” OR “Anhinga” OR “Fregata” OR “Leucocarbo” OR “Microcarbo” OR “Nannopterum” OR “Phalacrocorax” OR “Morus” OR “Sula” OR “Crypturellus” OR “Eudromia” OR “Nothocercus” OR “Nothoprocta” OR “Nothura” OR “Rhynchotus” OR “Tinamotis” OR “Tinamus” OR “Apalharpactes” OR “Apaloderma” OR “Harpactes” OR “Pharomachrus” OR “Trogon” OR “Tetrastes” OR “Hesperiphona”) AND (“experiment*” OR “manipulat*”)))>