arXiv:2401.00018v1 [hep-lat] 27 Dec 2023

Combining Bayesian reconstruction entropy with maximum entropy method for analytic continuations of matrix-valued Green's functions

Songlin Yang,[1] Liang Du,[1,*] and Li Huang[1,†]

[1]*College of Physics and Technology, Guangxi Normal University, Guilin, Guangxi 541004, China*

(Dated: January 2, 2024)

The Bayesian reconstruction entropy is considered an alternative to the Shannon-Jaynes entropy, as it does not exhibit the asymptotic flatness characteristic of the Shannon-Jaynes entropy and obeys the scale invariance. It is commonly utilized in conjunction with the maximum entropy method to derive spectral functions from Euclidean time correlators produced by lattice QCD simulations. This study expands the application of the Bayesian reconstruction entropy to the reconstruction of spectral functions for Matsubara or imaginary-time Green's functions in quantum many-body physics. Furthermore, it extends the Bayesian reconstruction entropy to implement the positive-negative entropy algorithm, enabling the analytic continuations of matrix-valued Green's functions on an element-wise manner. Both the diagonal and off-diagonal components of the matrix-valued Green's functions are treated equally. Benchmark results for the analytic continuations of synthetic Green's functions indicate that the Bayesian reconstruction entropy, when combined with the preblur trick, demonstrates comparable performance to the Shannon-Jaynes entropy. Notably, it exhibits greater resilience to noises in the input data, particularly when the noise level is moderate.

I. INTRODUCTION

In quantum many-body physics, single-particle Green's function holds significant importance [1, 2]. It is typically computed using numerical methods such as quantum Monte Carlo [3–5] and functional renormalization group [6, 7], or many-body perturbative techniques like random phase approximation [8, 9] and GW approximation [10–13], within the framework of imaginary time or Matsubara frequency formulation. This quantity provides valuable insights into the dynamic response of the system. However, it cannot be directly compared to real-frequency experimental observables. Therefore, it is necessary to convert the calculated values to real frequencies, which presents the fundamental challenge of the analytic continuation problem. In theory, the retarded Green's function $G^R(\omega)$ and the spectral function $A(\omega)$ can be derived from the imaginary-time Green's function $G(\tau)$ or Matsubara Green's function $G(i\omega_n)$ by performing an inverse Laplace transformation [1, 2]. Due to the ill-conditioned nature of the transformation kernel, solving it directly is nearly impossible. In fact, the output for this inverse problem is highly sensitive to the input. Even minor fluctuations or noises in $G(\tau)$ or $G(i\omega_n)$ can result in nonsensical changes for $A(\omega)$ or $G^R(\omega)$, making analytic continuations extremely unstable.

In recent decades, numerous techniques have been developed to address the challenges associated with analytic continuation problems. These methods include Padé approximation (PA) [14–17], maximum entropy method (MaxEnt) [18–20], stochastic analytic continuation (SAC) [21–30], stochastic optimization method (SOM) [31–35], stochastic pole expansion (SPX) [36, 37], sparse modeling (SpM) [38, 39], Nevanlinna analytical continuation (NAC) [40, 41], causal projections [42], Prony fits [43–45], and machine learning assistant approaches [46–52], and so on. Each method has its own advantages and disadvantages. There is no doubt that the MaxEnt method is particularly popular due to its ability to maintain a balance between computational efficiency and accuracy [18–20]. Several open source toolkits, such as `ΩMAXENT` [53], `QMEM` [54], `MAXENT` by Kraberger *et al.* [55], `MAXENT` by Levy *et al.* [56], `ALF` [57], `ACFlow` [58], `eDMFT` [59], and `ana_cont` [60], have been developed to implement this method.

The MaxEnt method is rooted in Bayesian inference [61]. Initially, the spectral function $A(\omega)$ is constrained to be non-negative and is interpreted as a probability distribution. Subsequently, the MaxEnt method endeavors to identify the most probable spectrum by minimizing a functional $Q[A]$, which comprises the misfit functional $L[A]$ and the entropic term $S[A]$ [see Eq. (10)]. Here, $L[A]$ quantifies the difference between the input and reconstructed Green's function, while $S[A]$ is employed to regulate the spectrum [18]. The MaxEnt method is well-established for the diagonal components of matrix-valued Green's functions, as the corresponding spectral functions are non-negative (for fermionic systems). However, the non-negativity of the spectral functions cannot be assured for the off-diagonal components of matrix-valued Green's functions. As a result, conventional MaxEnt method fails in this situation.

Currently, there is a growing emphasis on obtaining the full matrix representation of the Green's function (or the corresponding self-energy function) on the real axis [54, 62]. This is necessary for calculating lattice quantities of interest in Dyson's equation [59]. Therefore, there is a strong demand for a reliable method to perform analytic continuation of the entire Green's function matrix [41–43]. To meet this requirement, several novel approaches have been suggested in recent years as enhancements to the conventional MaxEnt method. The three most significant advancements are as follows: (1) *Auxiliary Green's function algorithm*. It is possible to create an auxiliary Green's function by combining the off-diagonal and diagonal elements of the matrix-valued Green's function to ensure the positivity of the auxiliary spectrum [63–68]. The analytic continuation of the auxiliary Green's function can

* liangdu@gxnu.edu.cn
† lihuang.dmft@gmail.com

then be carried out using the standard MaxEnt method. (2) *Positive-negative entropy algorithm.* Kraberger *et al.* proposed that the off-diagonal spectral functions can be seen as a subtraction of two artificial positive functions. They extended the entropic term to relax the non-negativity constraint presented in the standard MaxEnt method [55, 60]. This enables simultaneous treatment of both the diagonal and off-diagonal elements, thereby restoring crucial constraints on the mathematical properties of the resulting spectral functions, including positive semi-definiteness and Hermiticity. (3) *Maximum quantum entropy algorithm.* Recently, Sim and Han generalized the MaxEnt method by reformulating it with quantum relative entropy, maintaining the Bayesian probabilistic interpretation [18, 61]. The matrix-valued Green's function is directly continued as a single object without any further approximation or ad-hoc treatment [54].

Although various algorithms have been proposed to enhance the usefulness of the MaxEnt method [53–55], further improvements in algorithms and implementations are always beneficial. As mentioned above, the essence of the MaxEnt method lies in the definition of the entropic term $S[A]$. Typically, it takes the form of Shannon-Jaynes entropy (dubbed $S_{\rm SJ}$) in the realm of quantum many-body physics [69]. However, alternative forms such as the Tikhonov regulator [70], the positive-negative entropy [55], and the quantum relative entropy [54] are also acceptable. It is worth noting that in the context of lattice QCD simulations, the extraction of spectral functions from Euclidean time correlators is of particular important. The MaxEnt method is also the primary tool for the analytic continuation of lattice QCD data [71]. Burnier and Rothkopf introduced an enhanced dimensionless prior distribution, known as the Bayesian reconstruction entropy (dubbed $S_{\rm BR}$) [72, 73], which demonstrates superior asymptotic behavior compared to $S_{\rm SJ}$ [69]. They discovered that by using $S_{\rm BR}$ in conjunction with the MaxEnt method, a significant improvement in the reconstruction of spectral functions for Euclidean time correlators can be achieved. They latter extended $S_{\rm BR}$ to support Bayesian inference of non-positive spectral functions in quantum field theory [74]. But to the best of our knowledge, the application of $S_{\rm BR}$ has been limited to post-processing for lattice QCD data thus far. Hence, some questions naturally arise. How effective is $S_{\rm BR}$ for solving analytic continuation problems in quantum many-body simulations? Is it truly superior to $S_{\rm SJ}$? Keeping these questions in mind, the primary objective of this study is to broaden the application scope of $S_{\rm BR}$. We at first verify whether $S_{\rm BR}$ can handle Matsubara (or imaginary-time) Green's functions. Then, we generalize $S_{\rm BR}$ to implement the positive-negative entropy algorithm [55] for analytic continuations of matrix-valued Green's functions. The simulated results indicate that $S_{\rm BR}$ works quite well in transforming imaginary-time or Matsubara Green's function data to real frequency, no matter whether the Green's function is a matrix or not. We find that $S_{\rm BR}$ has a tendency to sharpen the peaks in spectral functions, but this shortcoming can be largely overcame by the preblur trick [55, 60]. Overall, the performance of $S_{\rm BR}$ is comparable to that of $S_{\rm SJ}$ [69] for the examples involved.

The rest of this paper is organized as follows. Section II introduces the spectral representation of the single-particle Green's function. Furthermore, it explains the fundamental formalisms of the MaxEnt method, the Bayesian reconstruction entropy, and the principle of the positive-negative entropy algorithm. Section III is dedicated to various benchmark examples, including analytic continuation results for synthetic single-band Green's functions and multi-orbital matrix-valued Green's functions. The spectral functions obtained by $S_{\rm SJ}$ and $S_{\rm BR}$ are compared with the exact spectra. In Section IV, we further examine the preblur trick and the auxiliary Green's function algorithm for $S_{\rm SJ}$ and $S_{\rm BR}$. The robustness of $S_{\rm BR}$ with respect to noisy Matsubara data is discussed and compared with that of $S_{\rm SJ}$. A concise summary is presented in Section V. Finally, Appendix A introduces the goodness-of-fit functional. The detailed mathematical derivations for $S_{\rm SJ}$ and $S_{\rm BR}$ are provided in Appendices B and C, respectively.

II. METHOD

A. Spectral representation of single-particle Green's function

It is well known that the single-particle imaginary-time Green's function $G(\tau)$ can be defined as follows:

$$G(\tau) = \langle \mathcal{T}_\tau d(\tau) d^\dagger(0) \rangle, \tag{1}$$

where τ is imaginary time, $\mathcal{T}_\tau$ means the time-ordered operator, d ($d^\dagger$) denotes the annihilation (creation) operator [1, 2]. Given $G(\tau)$, Matsubara Green's function $G(i\omega_n)$ can be constructed via direct Fourier transformation:

$$G(i\omega_n) = \int_0^\beta d\tau\, e^{-i\omega_n \tau} G(\tau), \tag{2}$$

where β is the inverse temperature of the system ($\beta \equiv 1/T$), and ω_n is the Matsubara frequency. Note that ω_n is equal to $(2n+1)\pi/\beta$ for fermions and $2n\pi/\beta$ for bosons (n is an integer).

Let us assume that the spectral function of the single-particle Green's function is $A(\omega)$, then the spectral representations of $G(\tau)$ and $G(i\omega_n)$ read:

$$G(\tau) = \int_{-\infty}^{+\infty} K(\tau, \omega) A(\omega) d\omega, \tag{3}$$

and

$$G(i\omega_n) = \int_{-\infty}^{+\infty} K(i\omega_n, \omega) A(\omega) d\omega, \tag{4}$$

respectively. Here, $K(\tau, \omega)$ and $K(i\omega_n, \omega)$ are called the kernel functions. Their definitions are as follows:

$$K(\tau, \omega) = \frac{e^{-\tau\omega}}{1 \pm e^{-\beta\omega}}, \tag{5}$$

and

$$K(i\omega_n, \omega) = \frac{1}{i\omega_n - \omega}. \tag{6}$$

In the right-hand side of Eq. (5), + is for fermionic correlators and - is for bosonic correlators [18]. In the subsequent discussion, our focus will be on fermionic correlators.

We observe that Eqs. (3) and (4) are classified as Fredholm integral equations of the first kind. When $A(\omega)$ is given, it is relatively straightforward to compute the corresponding $G(\tau)$ and $G(i\omega_n)$ by numerical integration methods. However, the inverse problem of deducing $A(\omega)$ from $G(\tau)$ or $G(i\omega_n)$ is challenging due to the ill-conditioned nature of the kernel function K. To be more specific, the condition number of K is very large because of the exponential decay of K with ω and τ. Consequently, direct inversion of K becomes impractical from a numerical standpoint [19]. Furthermore, the $G(\tau)$ or $G(i\omega_n)$ data obtained from finite temperature quantum Monte Carlo simulations is not free from errors [3–5], further complicating the solution of Eqs. (3) and (4).

B. Maximum entropy method

The cornerstone of the MaxEnt method is Bayes' theorem. Let us treat the input Green's function G and the spectrum A as two events. According to Bayes' theorem, the posterior probability $P[A|G]$ can be calculated by:

$$P[A|G] = \frac{P[G|A]P[A]}{P[G]}, \tag{7}$$

where $P[G|A]$ is the likelihood function, $P[A]$ is the prior probability, and $P[G]$ is the evidence [61]. $P[G|A]$ is assumed to be in direct proportion to $e^{-L[A]}$, where the misfit functional $L[A]$ reads:

$$L[A] = \frac{1}{2}\chi^2[A] = \frac{1}{2}\sum_{i=1}^{N}\left(\frac{G_i - \tilde{G}_i[A]}{\sigma_i}\right)^2. \tag{8}$$

Here, N is the number of input data points, σ denotes the standard deviation of G, $\tilde{G}[A]$ is the reconstructed Green's function via Eqs. (3) and (4), and $\chi^2[A]$ is called the goodness-of-fit functional (see Appendix A for more details). On the other hand, $P[A]$ is supposed to be in direct proportion to $e^{\alpha S[A]}$, where α is a regulation parameter and S is entropy. Since the evidence $P[G]$ can be viewed as a normalization constant, it is ignored in what follows. Thus,

$$P[A|G] \propto e^{Q}, \tag{9}$$

where Q is a functional of A [18]. It is defined as follows:

$$Q[A] = \alpha S[A] - L[A] = \alpha S[A] - \frac{1}{2}\chi^2[A]. \tag{10}$$

The basic idea of the MaxEnt method is to identify the optimal spectrum $\hat{A}$ that maximizes the posterior probability $P[A|G]$ (or equivalently $Q[A]$). In essence, our goal is to determine the most favorable $\hat{A}$ that satisfies the following equation:

$$\left.\frac{\partial Q}{\partial A}\right|_{A=\hat{A}} = 0. \tag{11}$$

Eq. (11) can be easily solved by the standard Newton's algorithm [75]. In Appendices A, B and C, all terms in the right-hand side of Eq. (10) are elaborated in detail. We also explain how to solve Eq. (11) efficiently.

C. Bayesian reconstruction entropy

The entropy S is also known as the Kullback-Leibler distance sometimes. In principle, there are multiple choices for it. Perhaps $S_{\rm SJ}$ is the most frequently used in quantum many-body physics and condensed matter physics [18, 69]. It reads:

$$S_{\rm SJ}[A] = \int d\omega \left[A(\omega) - D(\omega) - A(\omega)\log\left(\frac{A(\omega)}{D(\omega)}\right)\right]. \tag{12}$$

Here, $D(\omega)$ is called the default model, providing the essential features of spectra. Usually $D(\omega)$ is a constant or a Gaussian function, but it can be determined by making use of high-frequency behavior of input data as well [53]. If both $A(\omega)$ and $D(\omega)$ have the same normalization, Eq. (12) reduces to:

$$S_{\rm SJ}[A] = -\int d\omega\, A(\omega)\log\left(\frac{A(\omega)}{D(\omega)}\right). \tag{13}$$

The $S_{\rm BR}$ introduced by Burnier and Rothkopf [72, 73] is dominant in high-energy physics and particle physics. It reads:

$$S_{\rm BR}[A] = \int d\omega \left[1 - \frac{A(\omega)}{D(\omega)} + \log\left(\frac{A(\omega)}{D(\omega)}\right)\right]. \tag{14}$$

Note that $S_{\rm SJ}$ is constructed from four axioms [18]. While for $S_{\rm BR}$, two of these axioms are replaced [72]. First of all, the scale invariance is incorporated in $S_{\rm BR}$. It means that $S_{\rm BR}$ only depends on the ratio between A and D. The integrand in Eq. (14) is dimensionless, such that the choice of units for A and D will not change the result of the spectral reconstruction. Second, $S_{\rm BR}$ favors choosing smooth spectrum. The spectra that deviate between two adjacent frequencies, ω_1 and ω_2, should be penalized.

D. Positive-negative entropy

The formulas discussed above are only correct for positive definite spectra with a finite norm, i.e.,

$$\int d\omega\, A(\omega) > 0. \tag{15}$$

However, the norm could be zero for off-diagonal elements of matrix-valued Green's functions due to the fermionic anti-commutation relation [1, 42]:

$$\int d\omega\, A(\omega) = 0. \tag{16}$$

This suggests that the spectral function is not positive definite any more. The equations for $S_{\rm SJ}$ and $S_{\rm BR}$ [i.e., Eqs. (12) and (14)] should be adapted to this new circumstance. The positive-negative entropy algorithm proposed by Kraberger *et al.* is a graceful solution to this problem [55, 60]. They rewrite the off-diagonal spectral function $A(\omega)$ as the subtraction of two positive definite spectra:

$$A(\omega) = A^{+}(\omega) - A^{-}(\omega). \tag{17}$$