

Quicksort and Selection

Abhiram Ranade

January 18, 2016

Randomized Quicksort

Randomized Quicksort

Quicksort(X){

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .

Randomized Quicksort

Quicksort(X) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

We will estimate expected time, expectation over pivot choices.

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

We will estimate expected time, expectation over pivot choices.

Possible to write recurrence.

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

We will estimate expected time, expectation over pivot choices.

Possible to write recurrence. Instead we will use a more global method.

Randomized Quicksort

Quicksort(X) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

We will estimate expected time, expectation over pivot choices.

Possible to write recurrence. Instead we will use a more global method.

we show that over the entire execution the expected number of comparisons is $O(n \log n)$, where n is the number of keys.

Randomized Quicksort

Quicksort(X){

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. Return Quicksort(S) || M || Quicksort(L)

}

We will estimate expected time, expectation over pivot choices.

Possible to write recurrence. Instead we will use a more global method.

we show that over the entire execution the expected number of comparisons is $O(n \log n)$, where n is the number of keys.

Time = $O(\text{number of comparisons}) \Rightarrow \text{Time} = O(n \log n)$

Analysis

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesnt matter where they are in the given list.

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked,
OR

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesnt matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked,
OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked,
OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked,
OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked,
OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

$p(x_{ij} = 1) = 2/(j - i + 1)$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked, OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

$p(x_{ij} = 1) = 2/(j - i + 1)$

$$\sum_{i < j} E[x_{ij}] =$$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked, OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

$p(x_{ij} = 1) = 2/(j - i + 1)$

$$\sum_{i < j} E[x_{ij}] = \sum_{j=2}^{j=n} \sum_{i=1}^{i=j-1} \frac{2}{j - i + 1}$$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked, OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

$p(x_{ij} = 1) = 2/(j - i + 1)$

$$\sum_{i < j} E[x_{ij}] = \sum_{j=2}^{j=n} \sum_{i=1}^{i=j-1} \frac{2}{j - i + 1} \leq 2 \sum_{j=2}^{j=n} \frac{1}{j} = 2 \ln n$$

Analysis

Random variable x_{ij} = whether i th smallest key is compared with the j th smallest.

Doesn't matter where they are in the given list.

Expected number of comparisons: $\sum_{i < j} x_{ij}$

x_{ij} is a 0-1 variable $\Rightarrow E[x_{ij}] = p(x_{ij} = 1)$

i th smallest, j th smallest compared if

1. i th is picked before elements with rank $i + 1, \dots, j$ are picked, OR
2. j th is picked before elements with rank $i, \dots, j - 1$ are picked

Probability of 1: $1/(j - i + 1)$

all ranks i, \dots, j equally likely to be picked before one another

$p(x_{ij} = 1) = 2/(j - i + 1)$

$$\sum_{i < j} E[x_{ij}] = \sum_{j=2}^{j=n} \sum_{i=1}^{i=j-1} \frac{2}{j-i+1} \leq 2 \sum_{j=2}^n j = n \ln n \leq 2n \ln n$$

Remarks

Remarks

- ▶ In actual implementations, quicksort is done "in-place": S list is built from the beginning of the array, and L from the end.

Remarks

- ▶ In actual implementations, quicksort is done "in-place": S list is built from the beginning of the array, and L from the end.
- ▶ Quicksort is among the fastest sorting algorithms.

Remarks

- ▶ In actual implementations, quicksort is done "in-place": S list is built from the beginning of the array, and L from the end.
- ▶ Quicksort is among the fastest sorting algorithms.
- ▶ Make sure you understand what we proved: Expected time for any instance is $O(n \log n)$. Hence also the expected time for the worst instance.

Selection

Selection

Input: Keys x_1, \dots, x_n , Integer r .

Selection

Input: Keys x_1, \dots, x_n , Integer r .

Output: r th smallest key.

Selection

Input: Keys x_1, \dots, x_n , Integer r .

Output: r th smallest key.

"Easy" algorithm: Sort x_i . Return r th from resulting order.

Selection

Input: Keys x_1, \dots, x_n , Integer r .

Output: r th smallest key.

"Easy" algorithm: Sort x_i . Return r th from resulting order.

Time: $O(n \log n)$

Selection

Input: Keys x_1, \dots, x_n , Integer r .

Output: r th smallest key.

"Easy" algorithm: Sort x_i . Return r th from resulting order.

Time: $O(n \log n)$

Can be done in $O(n)$ time.

Next.

Idea: Adapt Quicksort

Idea: Adapt Quicksort

```
Quickselect(X,r){
```

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
 Else if $r \leq |S| + |M|$ return p .

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
 Else if $r \leq |S| + |M|$ return p .
 Else return Quickselect($L, r - |S| - |M|$).

}

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
 Else if $r \leq |S| + |M|$ return p .
 Else return Quickselect($L, r - |S| - |M|$).

}

Adapt Quicksort analysis to give $O(n)$ time

Homework

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
 Else if $r \leq |S| + |M|$ return p .
 Else return Quickselect($L, r - |S| - |M|$).

}

Adapt Quicksort analysis to give $O(n)$ time

Homework

Today:

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Quickselect($L, r - |S| - |M|$).

}

Adapt Quicksort analysis to give $O(n)$ time

Homework

Today:

Deterministic way to ensure $p =$ approximate median.

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Quickselect($L, r - |S| - |M|$).

}

Adapt Quicksort analysis to give $O(n)$ time

Homework

Today:

Deterministic way to ensure $p =$ approximate median.

\Rightarrow Problem nearly halves in each recursion.

Idea: Adapt Quicksort

Quickselect(X, r) {

1. Pick pivot p at random from X .
2. Distribute elements of X into 3 lists:
 S : smaller than p . M : equal to p . L : larger than p .
3. If $r \leq |S|$ return Quickselect(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Quickselect($L, r - |S| - |M|$).

}

Adapt Quicksort analysis to give $O(n)$ time

Homework

Today:

Deterministic way to ensure p = approximate median.

\Rightarrow Problem nearly halves in each recursion.

\Rightarrow Total time = $O(n)$.

Deterministic selection algorithm

Deterministic selection algorithm

Select (X, r) {

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.
3. Distribute X into lists:

Recurse!

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

3. Distribute X into lists:

S : smaller than p , M : equal to p , L : larger than p .

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.
3. Distribute X into lists:
S : smaller than p , M: equal to p , L: larger than p .
4. If $r \leq |S|$ return Select(S, r).

Recurse!

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.
3. Distribute X into lists:
S : smaller than p , M: equal to p , L: larger than p .
4. If $r \leq |S|$ return Select(S, r).
Else if $r \leq |S| + |M|$ return p .

Recurse!

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

3. Distribute X into lists:

S : smaller than p , M : equal to p , L : larger than p .

4. If $r \leq |S|$ return Select(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Select($L, r - |S| - |M|$).

}

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

3. Distribute X into lists:

S : smaller than p , M : equal to p , L : larger than p .

4. If $r \leq |S|$ return Select(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Select($L, r - |S| - |M|$).

}

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

Proof soon.

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

3. Distribute X into lists:

S : smaller than p , M : equal to p , L : larger than p .

4. If $r \leq |S|$ return Select(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Select($L, r - |S| - |M|$).

}

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

Proof soon.

Lemma: $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Obvious

Deterministic selection algorithm

Select (X, r) {

1. If $n = |X| < 50$, sort and return r th.
2. Choose pivot p as follows
 - ▶ Create $\lfloor n/5 \rfloor$ 5 tuples from X . Some elements remain.
 - ▶ Find $m_i = \text{median of } i\text{th 5-tuple}$.
 - ▶ $p = \text{median of all } m_i$.

Recurse!

3. Distribute X into lists:

S : smaller than p , M : equal to p , L : larger than p .

4. If $r \leq |S|$ return Select(S, r).
Else if $r \leq |S| + |M|$ return p .
Else return Select($L, r - |S| - |M|$).

}

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

Proof soon.

Lemma: $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Obvious

Lemma: Using recursion tree $T(n) = O(n)$

Proof soon.

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor$

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Proof of equality: Let $n = 10p + q \dots$

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Proof of equality: Let $n = 10p + q \dots$

Each median is less than or equal to 3 elements in its 5 tuple.

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Proof of equality: Let $n = 10p + q \dots$

Each median is less than or equal to 3 elements in its 5 tuple.

Number of elements less than or equal to p : $\geq 3 \lfloor n/10 \rfloor$.

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Proof of equality: Let $n = 10p + q \dots$

Each median is less than or equal to 3 elements in its 5 tuple.

Number of elements less than or equal to p : $\geq 3 \lfloor n/10 \rfloor$.

Thus $|L| < n - 3 \lfloor n/10 \rfloor$.

Lemma: $|L|, |S| \leq n - 3 \lfloor n/10 \rfloor$

p = median of $\lfloor n/5 \rfloor$ medians of 5 tuples.

Number of medians less than or equal to p : $\lfloor \lfloor n/5 \rfloor / 2 \rfloor = \lfloor n/10 \rfloor$

Proof of equality: Let $n = 10p + q \dots$

Each median is less than or equal to 3 elements in its 5 tuple.

Number of elements less than or equal to p : $\geq 3 \lfloor n/10 \rfloor$.

Thus $|L| < n - 3 \lfloor n/10 \rfloor$.

Similarly $|S|$.

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level})$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq n/10 - 1$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50$.

$$\text{Solving } T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\text{Thus subproblem size sum } \leq n/5 + (n - 3 \lfloor n/10 \rfloor)$$

$$\text{Solving } T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 \end{aligned}$$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50) \end{aligned}$$

$$\text{Solving } T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50 - 3 \times 4) \end{aligned}$$

$$\text{Solving } T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25} \end{aligned}$$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25} \end{aligned}$$

Thus constant fraction reduction even with floors!

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50$.

Thus subproblem size sum $\leq n/5 + (n - 3 \lfloor n/10 \rfloor)$
 $\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25}$

Thus constant fraction reduction even with floors!

Time for leaf level:

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25} \end{aligned}$$

Thus constant fraction reduction even with floors!

Time for leaf level: Time for 50 element sequence = c'

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

$$\text{Thus } \lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50.$$

$$\begin{aligned} \text{Thus subproblem size sum} &\leq n/5 + (n - 3 \lfloor n/10 \rfloor) \\ &\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25} \end{aligned}$$

Thus constant fraction reduction even with floors!

Time for leaf level: Time for 50 element sequence = c'

Total leaf level time $\leq c'n$

Solving $T(n) \leq T(\lfloor n/5 \rfloor) + T(n - 3 \lfloor n/10 \rfloor) + cn$

Ignoring floors: Subproblem sizes add up to $9n/10$.

\Rightarrow Work reduces by factor $9/10$ each level of recursion tree.

Work decreases geometrically by levels, so work = $O(\text{work at top level}) = O(n)$.

Analysis accounting for floor:

We stop recursion when $n < 50 \Rightarrow$ In the recurrence $n \geq 50$

Thus $\lfloor n/10 \rfloor \geq n/10 - 1 \geq n/10 - n/50 \geq 4n/50$.

Thus subproblem size sum $\leq n/5 + (n - 3 \lfloor n/10 \rfloor)$
 $\leq \frac{n}{50}(10 + 50 - 3 \times 4) = n \frac{24}{25}$

Thus constant fraction reduction even with floors!

Time for leaf level: Time for 50 element sequence = c'

Total leaf level time $\leq c'n$

Overall: $O(n)$

Remarks

Remarks

- ▶ Clever recursive use of selection itself to find pivot.

Remarks

- ▶ Clever recursive use of selection itself to find pivot.
- ▶ Not quite "divide" and conquer: Two recursive calls overlap in elements.

Remarks

- ▶ Clever recursive use of selection itself to find pivot.
- ▶ Not quite "divide" and conquer: Two recursive calls overlap in elements.
- ▶ Need to be careful to ensure that the two subproblems have total size $\leq n$.

Remarks

- ▶ Clever recursive use of selection itself to find pivot.
- ▶ Not quite "divide" and conquer: Two recursive calls overlap in elements.
- ▶ Need to be careful to ensure that the two subproblems have total size $\leq n$.
- ▶ Exercise: Will the algorithm work if we took medians of 3-tuples or of 7-tuples?

Remarks

- ▶ Clever recursive use of selection itself to find pivot.
- ▶ Not quite "divide" and conquer: Two recursive calls overlap in elements.
- ▶ Need to be careful to ensure that the two subproblems have total size $\leq n$.
- ▶ Exercise: Will the algorithm work if we took medians of 3-tuples or of 7-tuples?
- ▶ We can improve constants, e.g. 24/25. Key emphasis in this course: do not worry about the constant, but get a simple clean argument.

A detour: some more notation

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<$, $>$ "

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<$, $>$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

Limits and asymptotic comparisons:

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

Limits and asymptotic comparisons:

► $\lim_{n \rightarrow \infty} f(n)/g(n) = 0 : f = o(g)$

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

Limits and asymptotic comparisons:

- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n) = 0 : f = o(g)$
- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n)$ is finite and positive: $f = \theta(g)$

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

Limits and asymptotic comparisons:

- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n) = 0 : f = o(g)$
- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n)$ is finite and positive: $f = \theta(g)$
- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n) = \infty : f = \omega(g)$

A detour: some more notation

We said " θ means $=$ ", " O means \leq ", " Ω means \geq "

Asymptotically, and ignoring constants

Classes Little o, Little omega: "mean $<, >$ "

$o(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = 0\}$ $f = o(g)$ means " $f < g$ "

$\omega(g) = \{f \mid \lim_{n \rightarrow \infty} f(n)/g(n) = \infty\}$ $f = \omega(g)$ means " $f > g$ "

Example: $n = o(n \log n)$. $n^3 = \omega(n^2)$.

"Is there an $o(n \log n)$ time sorting algorithm?"

Limits and asymptotic comparisons:

- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n) = 0 : f = o(g)$
- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n)$ is finite and positive: $f = \theta(g)$
- ▶ $\lim_{n \rightarrow \infty} f(n)/g(n) = \infty : f = \omega(g)$

$$o(g) \subset O(g), \quad \omega(g) \subset \Omega(g)$$