

Pre-exposure Prophylaxis (PrEP) & Post-exposure Prophylaxis (PEP) – Summary

Key Symptoms: See HIV summary sheet

Eligibility

- **PrEP:** daily or episodic ART in those at high risk of HIV acquisition
 - **Gay, Bisexual and Men who have Sex with Men (GBMSM)**
 - Syphilis, rectal STI in last 12 months
 - Recurrent PEP use
 - Ongoing relationship with HIV+ partner with risk of transmission
 - High-incidence risk index ≥ 11
 - **Heterosexual**
 - Ongoing relationship with HIV+ partner with risk of transmission (viral load is NOT undetectable)
 - **Persons Who Inject Drugs (PWID)**
 - Consider in those who share needles with HIV+ people whose viral loads are NOT undetectable (= non-negligible risk)
- **nPEP:** administration of ART after potential event has occurred
 - **Recommend** if HIV-negative person presents within 72 hours of mod or high-risk exposure with a person with substantial risk for HIV transmission
 - **Consider** if HIV-negative person presents within 72 hours of mod to high-risk exposure with low but non-negligible risk for HIV transmission
- **Occupational PEP:** Administration of ART after potential event has occurred, usually w/in healthcare environment involving percutaneous exposure
 - **High Risk:** Percutaneous exposure to bodily fluids
 - **Mod Risk:** Mucous membrane exposure, bite wounds, exposure of broken skin to body fluids
 - **Low Risk:** Intact skin, exposure to low-risk fluids (urine, feces, etc.)

Non-Pharmacological Treatment: Safe sex practices, patient education, needle exchange program

Pharmacological Treatment

- **PrEP**
 - **Preferred:** Tenofovir DF/Emtricitabine **OR** Tenofovir AF/Emtricitabine **OR** Cabotegravir IM
 - Evidence/drug preference varies by population (see slide 8)
 - **Alternative regimen:** “PRN regimen” Tenofovir DF/Emtricitabine 2 tabs PO once within 2-24 hours prior to exposure followed by 1 tablet daily until 48 hours after the last exposure
- **nPEP/occupation PEP** (see lecture for dose amounts as it may vary between preferred and alternative regimens)
 - **Preferred:** Tenofovir AF/Emtricitabine/Biktegravir PO daily **OR** Tenofovir DF/Emtricitabine + Dolutegravir PO daily

Monitoring Parameters: see slide 13 for nPEP and slide 21 for PEP

Additional Key Points

- Do not delay treatment (aim <72 hours) = better outcomes
- See key points under HIV & Opportunistic Infections summary page