

COPD– Summary

Classification of COPD Severity (E>B>A):

- **Group A:** 0-1 exacerbations in past year (not leading to hospital admission) AND mMRC 0-1 OR CAT <10
- **Group B:** 0-1 exacerbations in past year (not leading to hospital admission) AND mMRC ≥ 2 OR CAT ≥ 10
- **Group E:** ≥ 2 exacerbations in past year (not leading to hospital admission) OR ≥ 1 exacerbation(s) in past year leading to hospital admission

Recommend: smoking cessation, exercise/active lifestyle, vaccinations (yearly flu, pneumococcal, COVID-19, Tdap, RSV)

Pharmacological Treatment

- **Group A:** SABA PRN OR SAMA PRN OR LABA OR LAMA
- **Group B:** LAMA + LABA
- **Group E:** LAMA + LABA (consider LABA+LAMA+ICS if serum eos>300)

Stepping Up Pharmacological Treatment

- **If symptoms not optimally controlled AND...**
 - **Already on LAMA or LABA:** switch to LAMA+LABA
 - **Already on LAMA+LABA:** consider switching molecule/inhaler device
- **If exacerbations are not optimally minimized/controlled AND...**
 - **Already on LAMA or LABA:** switch to LAMA+LABA (if eos <300) OR LAMA+LABA+ICS (if eos >300)
 - **Already on LAMA+LABA:** roflumilast (if eos <100) OR LAMA+LABA+ICS (if eos >100)
 - **Already on LAMA+LABA+ICS:** azithromycin

Acute exacerbation can be treated with antibiotics if the patient has 2-3 of the following symptoms:

increased dyspnea, increased sputum volume, increased sputum purulence* (*must be present)

- **Empiric antibiotic selection if <4 exacerbations/year:** amoxicillin x5 days OR doxycycline x5 days
- **Empiric antibiotic selection if ≥ 4 exacerbations in the past year OR had failed amoxicillin, doxy, or SMZ/TMP OR has had antibiotics in the past 3 months:** amox/clav x5days OR cefuroxime x5 days OR levofloxacin x5 days OR moxifloxacin x5days
 - Alternatives: azithromycin x3-5 days OR clarithromycin x5 days OR ciprofloxacin x5 days (reserved for risk of Pseudomonas)