

# HOMOGENEOUS LINEAR RECURRENCES

# Linear Recurrences

If  $A = (a_0, a_1, a_2, \dots, a_n, \dots)$  is a sequence, we say that  $A$  satisfies a linear recurrence relation with constant coefficients if for all  $n \geq m$ ,  $a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_m a_{n-m} + g(n)$ , where  $c_1, c_2, \dots, c_m$  is some set of constants and  $g(n)$  is a function depending on  $n$ . If  $c_m \neq 0$ , we say that the recurrence relation has order  $m$ , and in case  $g(n) = 0$ , we say the recurrence is homogeneous.

---

In this section we study methods for solving homogeneous recurrence relations, which usually means that we want to find a solution  $a_n = f(n)$  of the recurrence that satisfies the initial conditions  $a_0 = b_0, a_1 = b_1, \dots, a_{m-1} = b_{m-1}$ , for a given set of constants  $b_0, b_1, b_2, \dots, b_{m-1}$ .

Solve  $a_n = r a_{n-1}$  with  $a_0 = k$ .

$$a_n = k r^n$$

In this section we show how to solve recurrence relations of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_r a_{n-r} \quad (1)$$

The general solution to (1) will involve a sum of individual solutions of the form  $a_n = \alpha^n$ .

Solve the recurrence relation  $a_n = 2a_{n-1} + 3a_{n-2}$  with  $a_0 = a_1 = 1$ .

Find a formula for the number of ways for the elf in Example 2 of Section 7.1 to climb the  $n$  stairs.

The recurrence relation obtained in Example 2 of Section 7.1 was the Fibonacci relation  $a_n = a_{n-1} + a_{n-2}$ , with the initial conditions  $a_1 = 1$ ,  $a_2 = 2$ , or equivalently,  $a_0 = a_1 = 1$ . The associated characteristic equation is obtained by setting  $a_n = \alpha^n$ :

# The Case of Repeated Roots

In the first part of this section, we saw that when the characteristic equation of a homogeneous linear recurrence relation had distinct roots, a general solution could be obtained by looking at linear combinations of **m fundamental solutions**, one for each characteristic root.

In this case, for each characteristic root, there was an associated fundamental solution  $a_n = r_i^n$ .

However, when the characteristic equation has **repeated roots, two or more fundamental solutions coincide**. Therefore we have **fewer** than m fundamental solutions that are exponential functions. Thus the linear combinations of exponential functions cannot form a general solution because there are not enough fundamental solutions.

It is still possible, though, to find a general solution for a homogeneous linear recurrence even when the characteristic equation has repeated roots by looking at a slightly larger class of fundamental solutions.



Solve the recurrence relation  $a_n = 6a_{n-1} - 9a_{n-2}$  with initial conditions  $a_0 = 1$  and  $a_1 = 4$ .



Find a formula for  $a_n$  satisfying the relation  $a_n = -2a_{n-2} - a_{n-4}$  with  $a_0 = 0$ ,  $a_1 = 1$ ,  $a_2 = 2$ , and  $a_3 = 3$ .

