

Journal of Fluid Mechanics

Date of delivery: November 11, 2015

Journal and vol/article ref: flm 1500642

Number of pages (not including this page): 12

page 1 of 2

This proof is sent to you on behalf of Cambridge University Press.

Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.

Please return the marked proofs **within 7 days** of receipt to: **JFMproduction@cambridge.org**IMPORTANT. Please ensure that you return your completed **OFFPRINT ORDER FORM** at the same time as returning your proofs

PLEASE NOTE THIS PDF IS FOR PROOF CHECKING PURPOSES ONLY. IT SHOULD NOT BE DISTRIBUTED TO THIRD PARTIES AND MAY NOT REPRESENT THE FINAL VERSION.

IMPORTANT INFORMATION ABOUT CORRECTING AND RETURNING YOUR PROOFS – PLEASE READ

Your PDF proofs now show changes made to the MAIN TEXT by the copy-editor. Inserted text is shown in red. Locations of deleted text are indicated with a yellow rectangle/grey arrow. When viewed using the freely available Adobe Reader software, the deleted text becomes visible when you hover with the pointer over the yellow rectangle. (This does not happen with other PDF viewers.)

There may be typesetting changes to mathematical formulae that are not marked in this way; therefore please **proofread all mathematical expressions carefully**. Also **ALL figures are relabelled, please check all figure labels carefully against your originals**.

HOW TO RETURN YOUR PROOFS

You can mark up proofs either on screen using the enabled electronic editing tools or by hand on the hardcopy.

Marking up electronically. All proofs are enabled to allow electronic annotation in the freely available Adobe Reader software. Using your cursor select the text for correction, right click and use the most appropriate single tool (i.e. 'Replace', 'Cross out' or 'Add note to text'), to insert place the cursor then go to Tools/Comment&mark up/Text edits/Insert text at cursor. Please do not use the 'Sticky note' function as its placement is not precise enough. 'Show comments' allows all marks to be clearly identified, please do not emphasise your marks in any way. Please return the file as an attachment via email.

Marking up by hand. Please use the proof correction symbols indicated keeping marks clear and concise. Scanned proofs can be returned via email.

As only the marked proofs are sent on for correction they must be self-contained, i.e. ALL CORRECTIONS AND ANY NOTES MUST BE MARKED CLEARLY AND SUCCINCTLY ON THE PROOFS. Please do not send lists or explanations in the email or as separate documents, these will not be included in the correction process.

Journal of Fluid Mechanics

If you have no corrections to make please inform the Editorial Assistant by email.

Postal address – to be used only where other options are not possible

Journal of Fluid Mechanics – Mrs A. R. James

Cambridge University Press

Edinburgh Building

Shaftesbury Road

Cambridge CB2 8RU, UK Tel: +44 (0)1223 326061

For further information about Journals Production please consult our FAQs at http://journals.cambridge.org/production_faqs

Please note:

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Please answer carefully any author queries.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.
- If you have problems with the file please contact

JFMproduction@cambridge.org























Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.



















Important: you must return any forms included with your proof.

Please do not reply to this email

NOTE - for further information about **Journals Production** please consult our **FAQs** at http://journals.cambridge.org/production_faqs

Proof correction symbols

Correction required	Marginal mark	Corresponding mark in text
Insert new material	Inserted material, followed by caret, 	
Delete		(Cross out what is not wanted)
Substitute new material for something typeset	Write new material followed by 	(Cross out material to be replaced)
Leave as printed	 under material to remain
Insert space		 where space is required
Take out space (close up)		 around material to be closed up
Insert hyphen		 where hyphen is required
Change letter(s) to capital(s)		 under letter(s) required
Change letter(s) to small capital(s)		 under letter(s) required
Change to small letter(s)		(Encircle required letter(s))
Change to bold type		 under letter(s) to be altered
Change to italic type		 under letter(s) to be altered
Begin a new paragraph		 before first word of new paragraph

Correction required	Marginal mark	Corresponding mark in text
No new paragraph		 between paragraphs
Inferior (subscript) letter, e.g. x		 if to be inserted; cross out incorrect letter if a replacement
Superior (superscript) letter, e.g. p		 if to be inserted; cross out incorrect letter if a replacement
Transpose letters or words		 between letters or words
Indicate letters set upside-down		(Encircle letters to be altered)
Broken letter; replace by similar but undamaged character		(Encircle letters to be altered)
Insert parentheses		 where parentheses required
Insert square brackets		 where square brackets required
Insert apostrophe		 where apostrophe required
Insert quotation marks		 where quotation marks required

Offprint order form



CAMBRIDGE
UNIVERSITY PRESS

PLEASE COMPLETE AND RETURN THIS FORM WITH THE PROOF TO THE JOURNAL OF FLUID MECHANICS EDITORIAL ASSISTANT. WE WILL BE UNABLE TO SEND OFFPRINTS UNLESS A RETURN ADDRESS AND ARTICLE DETAILS ARE PROVIDED.

VAT REG NO. GB 823 8476 09

Journal of Fluid Mechanics (FLM)

(Office use only)

Volume:

no:

Offprints

Offprints may be purchased at the price shown below. The total order will be sent by surface mail within one month of publication to the address given below.

Number of offprints required:

Article ref. no. (FLM no. from front page of proofs): **FLM**

Email:

Offprints to be sent to (print in **BLOCK CAPITALS**):

Post/Zip Code:

Telephone:

Date (dd/mm/yy):

Author(s):

Article Title:

All enquiries about offprints should be addressed to **the publisher**: Journals Production Department, Cambridge University Press, The Edinburgh Building, Shaftesbury Road, Cambridge CB2 8RU, UK.

Charges for offprints (excluding VAT) Please circle the appropriate charge:

Number of copies	25	50	100	150	200	per 50 extra
1-4 pages	£68	£109	£174	£239	£309	£68
5-8 pages	£109	£163	£239	£321	£399	£109
9-16 pages	£120	£181	£285	£381	£494	£120
17-24 pages	£131	£201	£331	£451	£599	£131
Each Additional 1-8 pages	£20	£31	£50	£70	£104	£20

Methods of payment

If you live in Belgium, France, Germany, Ireland, Italy, Portugal, Spain or Sweden and are not registered for VAT we are required to charge VAT at the rate applicable in your country of residence. If you live in any other country in the EU and are not registered for VAT you will be charged VAT at the UK rate.

If registered, please quote your VAT number, or the VAT number of any agency paying on your behalf if it is registered.

VAT Number:

Payment **must** be included with your order, please tick which method you are using:

- ☐ Cheques should be made out to Cambridge University Press.
- ☐ Payment by someone else. Please enclose the official order when returning this form and ensure that when the order is sent it mentions the name of the journal and the article title.
- ☐ Payment may be made by any credit card bearing the Interbank Symbol.

Card Number:

Expiry Date (mm/yy):

/

Card Verification Number:

The card verification number is a 3 digit number printed on the **back** of your **Visa** or **Master card**, it appears after and to the right of your card number. For **American Express** the verification number is 4 digits, and printed on the **front** of your card, after and to the right of your card number.

Signature of
card holder:

Amount

(Including VAT

if appropriate): £

Please advise if address registered with card company is different from above



Linear stability analysis for *monami* in a submerged seagrass bed

Q1 Ravi Singh¹, M. M. Bandi², Amala Mahadevan³ and Shreyas Mandre^{4,†}

Q2 ¹Department of Physics, Brown University, Providence, RI 02912, USA

²OIST Graduate University, Okinawa 904-0495, Japan

³Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

⁴School of Engineering, Brown University, Providence, RI 02912, USA

(Received 3 September 2015; revised 28 October 2015; accepted 28 October 2015)

The onset of *monami* – the synchronous waving of seagrass beds driven by a steady flow – is modelled as a linear instability of the flow. Unlike previous works, our model considers the drag exerted by the grass in establishing the steady flow profile, and in damping out perturbations to it. We find two distinct modes of instability, which we label modes 1 and 2. Mode 1 is closely related to Kelvin–Helmholtz instability modified by vegetation drag, whereas mode 2 is unrelated to Kelvin–Helmholtz instability and arises from an interaction between the flow in the vegetated and unvegetated layers. The vegetation damping, according to our model, leads to a finite threshold flow for both of these modes. Experimental observations for the onset and frequency of waving compare well with model predictions for the instability onset criteria and the imaginary part of the complex growth rate respectively, but experiments lie in a parameter regime where the two modes can not be distinguished.

Key words: coastal engineering, geophysical and geological flows, instability

1. Introduction

Seagrasses exhibit a rich set of dynamical behaviours due to their collective interaction with fluid flows. The hydrodynamic processes resulting from this behaviour influence many environmental processes such as sedimentation, transport of dissolved oxygen, plant growth and biomass production (Fonseca & Kenworthy 1987; Grizzle *et al.* 1996; Nepf 1999, 2012). One response of the submerged grass beds to steady currents is the formation of coherent large-amplitude oscillations, known as *monami*.

† Email address for correspondence: shreyasmandre@gmail.com

(Ackerman & Okubo 1993). In this article, we present a linear hydrodynamic instability underlying the onset of these coherent oscillations.

Current explanations of *monami* (Ikeda & Kanazawa 1996; Raupach, Finnigan & Brunet 1996; Ghisalberti & Nepf 2002) invoke the existence of a shear layer at the top of the grass bed (henceforth called grass top) due to vegetation drag. Its instability, through a mechanism similar to the Kelvin–Helmholtz (KH) instability, is thought to lead to coherent eddies over the grass bed. The grass responds to these eddies by deforming, which leads to large-amplitude synchronous oscillations of the grass blades. This picture can be used to derive a simple scaling dependence of the *monami* frequency on the flow speed and the shear layer thickness, and to understand transport in the seagrass bed (Nepf & Vivoni 2000; Ghisalberti & Nepf 2002, 2004; Okamoto, Nezu & Ikeda 2012).

However, several aspects of this explanation remain unsatisfactory. The instability is modelled using the inviscid Rayleigh equation, and despite the role of drag in producing the shear layer, its role in damping the perturbations is ignored (Raupach *et al.* 1996). Furthermore, the shear layer is assumed not to be influenced by the top and bottom boundaries of the domain, although the experimentally measured thickness of the shear layer is in many cases comparable to the unvegetated water depth. The velocity profile of the free shear layer is assumed *ad hoc* to be piecewise linear (Py, De Langre & Moulia 2006) or hyperbolic tangent (Raupach *et al.* 1996; Ghisalberti & Nepf 2002), with parameters fitted using the experimental observations. The origin of these profiles, the values of the fitted parameters, and their effect on *monami* remain unexplained. Finally, no existing theory explains the threshold flow speed, observed in the laboratory (Ghisalberti & Nepf 2002) and in the field (Grizzle *et al.* 1996), below which *monami* is not observed.

Here, we present a mathematical model for the linear instability that accounts for these effects. Although *monami* is manifested in the motion of the grass, the drag exerted by the vegetation on the flow is central to the hypothesized instability. The instability and the resulting flow structures persist in laboratory experiments even when flexible grass mimics are replaced by rigid dowels (Ghisalberti & Nepf 2002, 2006). Therefore, to develop the essential mathematical model, we assume the grass blades to be rigid and oriented vertically (along the y direction) on average. The vegetation is modelled as a continuum drag on the fluid acting perpendicular to the blade orientation and proportional to the grass density. In the limit of dense vegetation, the steady profile established in the presence of the drag exhibits a localized region of enhanced shear gradient near the grass top and drives a flow instability. While some features of the instability are similar to KH instability, we also find significant differences. This comparison is the focus of this article.

2. Mathematical model

The vegetation is assumed to be sufficiently dense so that the drag exerted by it may be modelled by a continuous body force \mathbf{f} entering the equation governing the flow as

$$\nabla \cdot \mathbf{u} = 0, \quad \rho(\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u}) = -\nabla p + \mu \nabla^2 \mathbf{u} + \mathbf{f}, \quad (2.1a,b)$$

where ρ , $\mathbf{u} = (u, v)$ and p are the fluid density, velocity, and the dynamic pressure respectively, and μ is the (dynamic) eddy viscosity. The Reynolds number of the flow based on the scale of the grass blade is $O(10^2\text{--}10^3)$; therefore, neglecting skin friction, we model the form drag on the vegetation as $\mathbf{f} = -N_g C_N \rho \mathbf{u} |\mathbf{u}| \hat{\mathbf{x}}$ (Nepf 1999; Nepf & Vivoni 2000; Ghisalberti & Nepf 2004), where N_g is the number of grass blades per




FIGURE 1. (a) Schematic set-up and comparison of our steady flow profile with that from the experiments in Ghisalberti & Nepf (2004) (case I from table 1) and its approximation with $U_0 = 7.28 \text{ cm s}^{-1}$ and $\delta = 5.02 \text{ cm}$ in our model. The grass extends up to $y = h_g$ in the water column of depth $2H$. The steady velocity profile can be decomposed into a parabolic profile in the unvegetated region, a uniform profile deep within the vegetation and a boundary layer of thickness δ near the grass top. (b) The dependence of the boundary layer thickness (estimated as $|U/U_y|$ at $y = h_g$ from the numerical solution of (3.1)) on the vegetation density parameter $R\tilde{N}_g$.

unit horizontal area, C_N is the drag coefficient for the flow normal to the grass, \hat{x} is the unit vector in the horizontal direction and d is the average blade width projected perpendicular to the flow. In the interest of simplicity, we model the turbulence using an eddy viscosity; see § 5.3 for a detailed discussion. In the field, C_N , N_g and μ vary with position, but we do not expect these variations to be central to the instability mechanism, and therefore take them to be constants.

3. Linear stability analysis

We first calculate the fully developed steady solution $\mathbf{u} = U(y)\hat{x}$ of (2.1) driven by a constant pressure gradient dP/dx in a water column of depth $2H$ and vegetated depth h_g , and use it to non-dimensionalize the mathematical model. The momentum balance (2.1) for $U(y)$ simplifies to

$$-\frac{dP}{dx} + \mu U''(y) - S(y)\rho C_N d N_g U|U| = 0, \quad (3.1)$$

where $S(y) = 1$ for $0 < y < h_g$ and $S(y) = 0$ for $h_g < y < 2H$. Equation (3.1) is solved subject to no shear at the boundaries, i.e. $U'(0) = U'(2H) = 0$. The lower boundary condition is appropriate for dense vegetation because the shear stress exerted by the bottom surface is expected to be negligible compared with the vegetation drag (Nepf & Vivoni 2000), whereas the upper boundary condition models the free interface. A comparison of the steady flow profile from the solution of (3.1) with experimental measurements by Ghisalberti & Nepf (2004) is shown in figure 1. The profile $U(y)$ has three distinct regions, as determined by a matched asymptotic analysis (Hinch 1991). Within the vegetation, it is approximately uniform with $U(y) \approx U_g = \sqrt{(-dP/dx)/(\rho C_N d N_g)}$, arising from a balance of the drag with the pressure gradient. Outside the vegetation, the velocity has a simple parabolic profile.

At the grass top, continuity of shear stresses results in a boundary layer of thickness δ . Denoting U_{bl} to be the velocity scale in the boundary layer and $U_0 = (-dP/dx) H^2/\mu$ to be the velocity scale in the unvegetated region, the balance between viscous forces and vegetation drag implies $\mu U_{bl}/\delta^2 \sim \rho C_N d N_g U_{bl}^2$, and the continuity of shear stress across the grass top implies $U_{bl}/\delta \sim U_0/H$. Solving for δ and U_{bl} yields $\delta/H \sim U_{bl}/U_0 \sim (R\tilde{N}_g)^{-1/3}$, where $\tilde{N}_g = (C_N d H N_g)$ is the vegetation frontal area per bed area and $R = \rho U_0 H/\mu$ is the Reynolds number of the flow. A numerical estimate of δ (estimated as U/U_y at $y = h_g$) is compared with this prediction in figure 1 (inset). We identify the boundary layer to be analogous to the shear layer invoked by Ghisalberti & Nepf (2002, 2004) in the previous explanation of *monami*. This dependence of δ on N_g gives us a way to systematically investigate the effect of the shear layer thickness on the instability mechanism. Figure 1 also shows that the asymptotic regime of a thin boundary layer is expected to hold for $R\tilde{N}_g \gtrsim 100$. In this notation, $U_g/U_0 = (R\tilde{N}_g)^{-1/2}$ (used later in deriving (5.3)).

Next, we substitute $\mathbf{u} = (U + \tilde{u}, \tilde{v})$, $p = P + \tilde{p}$ in (2.1) and expand to linear order to investigate the evolution of small perturbations (\tilde{u}, \tilde{v}) , which obey

$$\rho(u_t + Uu_x + vU_y) = -p_x + \mu \nabla^2 u - 2S\rho C_N d N_g Uu, \quad (3.2)$$

$$\rho(v_t + Uv_x) = -p_y + \mu \nabla^2 v, \quad \nabla \cdot \mathbf{u} = 0, \quad (3.3a,b)$$

where the tildes are dropped. These equations are non-dimensionalized using the half-channel height H , velocity U_0 and time H/U_0 , leading to three non-dimensional parameters, namely R , \tilde{N}_g and the vegetation submergence ratio h_g/H . We also use δ/H in lieu of \tilde{N}_g to parametrize the vegetation density and help to elucidate the instability mechanism. Using a stream function ψ with $u = \psi_y$, $v = -\psi_x$ to satisfy mass balance, we seek a solution of the form $(u, v, \psi) = (\hat{u}(y), \hat{v}(y), \phi(y))e^{ikx + \sigma t}$ to obtain a modified Orr–Sommerfeld equation (Drazin & Reid 1981; Chu, Wu & Khayat 1991; Chen & Jirka 1997),

$$R^{-1}(D^2 - k^2)^2 \phi = [(\sigma + ikU)(D^2 - k^2) - ikU_{yy}]\phi + \tilde{N}_g D(2SUD\phi), \quad (3.4)$$

where $D = d/dy$, and subject to the boundary conditions $\phi = D^2\phi = 0$ at $y = 0$ and $y = 2$. The growth rate σ for a given wavenumber k appears as an eigenvalue that allows a non-trivial solution ϕ of (3.4). We solve (3.4) numerically for σ and ϕ .

4. Results – unstable modes and critical parameters

A threshold in R , above which the flow is unstable ($\text{Re}(\sigma) > 0$) for at least one k , emerges from the solution of (3.4). The dependence of this threshold R , and the corresponding marginally stable wavenumber k , on δ/H and h_g/H is shown in figure 2, and is found to compare well with experimental observations of Ghisalberti & Nepf (2002). The threshold R increases with the vegetation density, indicating a competition between the destabilizing shear in the flow and the stabilizing effect of damping due to vegetation drag. The frequency ($\text{Im}(\sigma)$) of the fastest growing mode also agrees well with the observed behaviour – frequency of *monami*, maxima in the velocity spectra and frequency of vortex passage in laboratory scale experiments (Ghisalberti & Nepf 2002) – for cases where the vegetation was sufficiently dense to be modelled by a continuum drag field, as shown in figure 3. The measured values of the eddy viscosity vary with depth in the water column (Ghisalberti & Nepf 2004); we use $\mu = 0.1$ Pa s as a representative value from this range for comparison with




FIGURE 2. Critical Reynolds number, threshold Reynolds number for modes 1 and 2 (a) and the corresponding marginally stable wavenumber (b) for different submergence ratios as a function of vegetation density parametrized by the boundary layer thickness. Parameters from experiments reported by Ghisalberti & Nepf (2002) to exhibit or suppress synchronous waving are also included in (b).




FIGURE 3. Comparison of the experimentally measured dominant frequency f_o (in Hz) with the predictions $f_p = \text{Im}(\sigma)$ from the solution of (3.4). The experimental data are obtained from Ghisalberti & Nepf (2002) and Vivoni (1998).

experiments. Figure 2 also shows that the predicted dimensionless wavenumber of the dominant mode for parameters corresponding to the laboratory scale experiments ($4 < H/\delta < 10$) is very close to unity, and therefore the dimensional wavelength is approximately $2\pi H$.

To better understand the instability mechanism, we consider the dependence of the fastest growing wavenumber on δ . The fastest growing wavenumber first increases proportional to H/δ , but at a critical δ discontinuously jumps and remains $O(1)$ (see figure 2). To aid in explaining this behaviour, we show heat maps of $\text{Re}(\sigma)$ as a function of R and k for different values of h_g/H and \tilde{N}_g in figure 4. The smallest R on the neutral curve ($\text{Re}(\sigma) = 0$) sets the threshold. We observe that as \tilde{N}_g increases, the unstable region splits into two; we refer to the region with the higher k as ‘mode 1’ and the one with the lower k as ‘mode 2’. For $h_g/H \lesssim 0.9$, the unstable region for mode 1 recedes to higher R , and for $h_g/H \gtrsim 0.9$, the region shrinks to zero size. In either case, due to such behaviour the most unstable mode transitions discontinuously from mode 1 to mode 2. All experimental data we have found correspond to a




FIGURE 4. Heat maps of $\text{Re}(\sigma)$ and the neutral curve ($\text{Re}(\sigma) = 0$) as a function of wavenumber and R for (a) $h_g/H = 0.8$, $\tilde{N}_g = 500$; (b) $h_g/H = 0.8$, $\tilde{N}_g = 850$; (c) $h_g/H = 0.8$, $\tilde{N}_g = 925$; (d) $h_g/H = 1$, $\tilde{N}_g = 900$; (e) $h_g/H = 1$, $\tilde{N}_g = 1400$; (f) $h_g/H = 1$, $\tilde{N}_g = 1700$. As \tilde{N}_g increases, the unstable region splits into two regions labelled as ‘mode 1’ and ‘mode 2’. For \tilde{N}_g below (above) a critical value, mode 1 (mode 2) sets the threshold R .

vegetation density for which the unstable region in the R – k space has not split into two, so we are unable to determine whether the flow instability in the laboratory scale experiments (Ghisalberti & Nepf 2002) is due to mode 1 or mode 2.

5. Discussion – comparison of unstable modes with the KH instability mechanism

The distinct asymptotic behaviour of the two modes as $\tilde{N}_g \gg 1$ distinguishes them from each other and facilitate comparison with the KH instability mechanism.

5.1. Mode 1

We numerically observe that the threshold Reynolds number for mode 1 instability scales as $R \sim (H/\delta)^2$ (or $R \propto \tilde{N}_g^2$). Our calculations also show that mode 1 asymptotically localizes to the boundary layer near the grass top, and exhibits highest growth for a perturbation of $k \sim H/\delta$ (see figure 2). The behaviour of this critical Reynolds number can be understood by considering the limit $R \gg 1$ and $\tilde{N}_g \gg 1$. Estimating the size of various terms of (3.4) within the boundary layer in this limit helps us to understand the behaviour of the critical R . Using $D \sim H/\delta$, $\sigma \sim O(1)$ and $U = U_{bl} \sim \delta/H$ in the boundary layer, the magnitude of the advection term is $(H/\delta)^2$ (or $R^{2/3}\tilde{N}_g^{2/3}$), and the viscous and vegetation drag terms are $(1/R)(\delta/H)^{-4}$ (or $(R^{1/3}\tilde{N}_g)^{4/3}$). The advection term, viscous term and vegetation drag terms balance when $R \sim (H/\delta)^2$ (or $R \sim \tilde{N}_g^2$).

To further understand the mechanism of mode 1, we rescale (3.4) near the grass top using the boundary layer scalings $\eta = y/(\delta/H)$, $U(y) = (\delta/H)\bar{U}(\eta)$ and $k = (H/\delta)\bar{k}$. With these scalings (3.4) simplifies to

$$(\bar{D}^2 - \bar{k}^2)^2 \phi = (R/\tilde{N}_g^2)^{1/3} [(\sigma + i\bar{k}\bar{U})(\bar{D}^2 - \bar{k}^2) - i\bar{k}\bar{U}_{\eta\eta}] \phi + \bar{D}(2S\bar{U}\bar{D}\phi), \quad (5.1)$$

in a region of thickness $O(\delta)$ near $y = h_g$, where $\bar{D} = d/d\eta$. Since (R/\tilde{N}_g^2) is the only remaining parameter in (5.1), the mode shape and solution are expected to converge




FIGURE 5. Plot of the neutral mode 1 (solid) and mode 2 (dashed) shapes $|\phi|$ in the limit of small δ/H for $h_g/H = 0.2$. The approach of the mode shapes to each other for these small values of δ/H indicates that the dense vegetation asymptote is reached. Mode 1 shapes appear to be self-similar in shape as $\delta \rightarrow 0$. The inset shows that the rescaled $|\phi|$ for mode 1 as a function of $(y - h_g)/\delta$ approaches a universal shape, indicating that an asymptotic limit has been reached.

	KH	Mode 1	Mode 2
Base velocity profile	$U(y) = U_0 \tanh(y/\delta)$		Equation (3.1)
Domain	$-\infty < y < \infty$		$-1 < y < 1$
Inflection point	Exists at $y = 0$	$U''(y)$ discontinuous at $y = h_g$	
Shear layer thickness	δ		$\delta \sim H(R\tilde{N}_g)^{-1/3}$
Linearized dynamics	Equation (5.2)		Equation (3.4)
Dense grass limit	No grass included	Equation (5.1)	Equation (5.3)
Critical parameters	None	$R \propto \tilde{N}_g^2$	$R \propto \tilde{N}_g$
Most unstable k as $\delta \rightarrow 0$	$\propto H/\delta$	$\propto H/\delta$	$O(1)$
Mode localized?	Yes, near $y = 0$	Yes, near $y = h_g$	No, spans water column

TABLE 1. Comparison between the KH instability and the two unstable modes resulting from solution of (3.4).

in the limit $R \gg 1$, $\tilde{N}_g \gg 1$, but R/\tilde{N}_g^2 fixed. Our numerical findings confirm this expectation; the critical R scales as $(H/\delta)^2$, as shown in figure 2, and the mode shapes are self-similar with length scale δ , as shown in figure 5.

Mode 1 shares many characteristics with the KH instability (see table 1). The fastest growing wavenumber at the critical R scales as $k \propto (H/\delta)$, Similarly to the KH instability. The extent of the unstable mode is also localized to the boundary layer region. The porous nature of the vegetation implies that a weak flow of magnitude $U_{bl} = U_0\delta/H$ penetrates a thin boundary layer region δ , and therefore the shear gradient $U_{yy} \sim U_0/\delta H$ is largest in this region. The strong shear gradient U_{yy} in the boundary layer plays a central role in destabilizing the flow and localizing the instability to that region.

Our detailed description of mode 1, given by (5.1), also highlights key differences from formulations of the KH instability. The KH instability is usually described using the inviscid Rayleigh equation,

$$(\sigma + ikU)(D^2 - k^2)\phi = ikU_{yy}\phi, \quad (5.2)$$

and is therefore not parametrized by the Reynolds number. Describing the instability using the Orr–Sommerfeld equation introduces the Reynolds number as a parameter, but shear flows with \tanh profiles are unstable for all values of the parameter (Drazin & Reid 1981). Therefore, based on inviscid formulations of the KH instability, the origin of the threshold flow conditions observed in experiments and in the field is unclear.

In our model, (turbulent eddy) viscosity sets the scale of the boundary layer, and therefore for mode 1. However, the boundary layer is established only in the vegetated region; the velocity profile does not saturate on the scale of δ in the unvegetated region. The threshold flow condition arises from a competition between the destabilizing role of fluid inertia, which is very similar to the role played in the KH instability, and the vegetation drag. The vegetation drag may not be neglected within this boundary layer, and therefore plays a central role in the mode 1 instability mechanism.

5.2. Mode 2

The threshold condition for mode 2 is numerically observed to be $R \propto (\delta/H)^{-3/2}$ (or $R \propto \tilde{N}_g$) for $k \sim O(1)$, as shown in figure 2, which can be understood by assuming $R \gg 1$ but fixed $R/\tilde{N}_g \sim O(1)$. In this limit, the non-dimensional flow in the grass bed is $U_g/U_0 \sim (R\tilde{N}_g)^{-1/2} \ll 1$, and therefore $ikU \ll \sigma$ may be neglected in comparison to σ . Furthermore, U_{yy} decays to zero within the grass outside the boundary layer. Outside the grass, the turbulent viscous stress term is negligible compared with the inertial term because $R \gg 1$. Thus, (3.4) simplifies to

$$\sigma(D^2 - k^2)\phi = -2(\tilde{N}_g/R)^{1/2}D^2\phi, \quad \text{for } y < h_g, \quad (5.3a)$$

$$(\sigma + ikU)(D^2 - k^2)\phi = ikU_{yy}\phi, \quad \text{for } y > h_g. \quad (5.3b)$$

The only remaining parameter in (5.3) is R/\tilde{N}_g . For fixed R/\tilde{N}_g , the mode shape converges in the aforementioned limit, in agreement with our numerical results shown in figure 5. This convergence indicates that we have identified the correct asymptotic limit to investigate mode 2. The parameter R/\tilde{N}_g therefore sets the threshold, justifying the numerically observed asymptotic behaviour $R \propto \tilde{N}_g$ (or $R \sim (\delta/H)^{-3/2}$; see figure 2 for comparison with numerical results).

The structure of this mode in the aforementioned limit is such that ϕ is continuous at $y = h_g$, but $D\phi$ undergoes a rapid transition there, on the scale of the boundary layer thickness δ . The eigenvalues and the mode shape are otherwise independent of δ . Therefore, we conclude that the boundary layer only plays a secondary role of regularizing the discontinuity in the tangential velocity arising at $y = h_g$ in this instability mechanism. The enhanced shear in the boundary layer plays no role for this mode of instability.

Mode 2 has characteristics distinct from the KH instability. Outside the grass, the unstable mode shape is governed by the inviscid Rayleigh equation (5.2). An inflection point in $U(y)$ is a necessary condition for instability arising from (5.2) according to Rayleigh's criteria (Rayleigh 1879). However, for our $U(y)$ profiles, $U_{yy}(y) = -1$ above the grass and therefore does not change sign for $y > h_g$. Instead, the dynamics is coupled with the flow in the grass bed described by (5.3a) in $y < h_g$. The absence of U_{yy} in (5.3a) indicates that U_{yy} is approximated to be zero in $y < h_g$, and therefore the positive values of U_{yy} that occur in the boundary layer do not affect this mode of

instability to leading order. Furthermore, the presence of the critical parameter R/\tilde{N}_g in (5.3a) indicates the presence of alternative destabilizing dynamics, involving the interaction of flow in the unvegetated region governed by (5.2) with the flow in the vegetated region incorporating the drag. Therefore, we conclude that **mode 2** is distinct from the KH instability, and owes its existence to vegetation drag.

5.3. Role of the turbulence model

We have modelled the turbulence using a constant eddy viscosity. The simplicity of this model allows us to make progress and capture the essential features of the instability. However, this simplicity in some cases only provides a qualitatively accurate description of the flow. In this subsection, we present an account of the advantages and shortcomings of assuming a constant eddy viscosity to model the turbulence.

As a consequence of the constant eddy viscosity, the boundary layer thickness δ scales as $H(R\tilde{N}_g)^{-1/3}$. Experimental observations show that the boundary layer thickness scales instead as \tilde{N}_g^{-1} (Nepf, White & Murphy 2007). While the precise boundary layer thickness is governed by the details of the turbulence model, the existence of this boundary layer for dense vegetation is independent of the turbulence model. We have captured one possible realization of this feature using a constant eddy viscosity. Experiments have also shown that a model based on a mixing length l better approximates the turbulent characteristics of the flow with $l \sim \delta$; i.e. the boundary layer itself establishes eddies to transport momentum. The eddy viscosity corresponding to this model is $\mu \sim \rho U \delta$, and the leading-order balance between turbulent momentum transport and vegetation drag is $\mu U / \delta^2 \sim \rho C_N d N_g U^2$. Substituting μ yields $\delta/H \sim \tilde{N}_g^{-1}$, in agreement with the experimental observations.

Within our framework, the mixing length model implies a scale for the eddy viscosity $\mu \sim \rho U_{bl} \delta$ at the grass top, which corresponds to an effective $R \sim U_0 H / U_{bl} \delta$. Furthermore, matching the slope of the velocity profile from the boundary layer to the unvegetated flow implies $U_{bl} / U_0 \sim \delta / H$, and therefore $R \sim (H/\delta)^2$. Substituting this relation in $\delta/H \sim (R\tilde{N}_g)^{-1/3}$ and solving for δ yields $\delta/H \sim \tilde{N}_g^{-1}$. This simple scaling analysis shows that the boundary layer thickness depends on the turbulence model, and indicates that turbulence models based on mixing lengths will yield more realistic scalings for the boundary layer thickness. At the same time, the qualitative features of the instability are represented by our analysis.

The **mode 1** instability is driven by the intense shear on the scale of the boundary layer. The driving mechanism for this instability is similar to that of the KH instability, and relies only on the presence of this shear as presented in the $\tilde{U}_{\eta\eta}$ term in (5.1). Therefore, we expect **mode 1** instability to be exhibited independent of the turbulence model. We further expect the fastest growing wavenumber to be proportional to $1/\delta$, and the mode to be localized to the boundary layer because these results have a basis in dimensional analysis. The threshold parameters for **mode 1**, however, may depend on the precise turbulence model used.

For the **mode 2** instability, the turbulent momentum transport is found to be irrelevant to leading order. In the asymptotic limit of dense grass, (5.3) shows that the instability is driven by the interaction of the unvegetated flow with the vegetation drag. The influence of the turbulence model is limited to the regularization of the sharp transition in tangential velocity across the grass top. Therefore, we expect **mode 2** and its features to be preserved even if a different turbulence model is used.

5.4. Comparison with previous models

A modified version of the Orr–Sommerfeld equation was analysed previously by Chu *et al.* (1991), Chen & Jirka (1997) and White & Nepf (2007) in the context of instabilities in depth-averaged shallow water flows, where bottom friction replaces or augments vegetation drag. They assumed the steady profile to be a hyperbolic tangent, the drag to be isotropic and the flow domain to be infinite in y . White & Nepf (2007) also neglected the eddy viscosity in their stability analysis. While a detailed investigation needed to compare the consequences of the different assumptions is outside the scope of this paper, we discuss similarities and differences between their results and ours. These investigations only found one unstable mode. This is most likely so because the calculations were restricted to a parameter regime where the two modes have not yet been separated from each other, as is the case shown in figure 4 for the lowest \tilde{N}_g . These investigations also found that increasing the drag could further destabilize the flow, which is consistent with our interpretation of the mode 2 instability mechanism.

The analogous oscillation of terrestrial canopies in wind, known as *honami* (Inoue 1956; Raupach *et al.* 1996), is different because the atmospheric boundary layer is much larger than the vegetation height. In the framework of our model, the limit of $h_g/H \ll 1$ while $\delta/h_g = \text{constant}$ can be used to represent the hydrodynamic instability for the terrestrial case. We find that in this case, the transition from mode 1 to mode 2 occurs at such a large vegetation density that mode 2 is irrelevant. Hence, only the KH-like characteristics are observed in the terrestrial case.

While predictions of the threshold R for the onset of the instability and the frequency of oscillations are comparable to experimental observations, the deviation of our model predictions from the observations may be attributed to the various simplifications in our model. In real meadows, the grass is flexible, and the drag coefficient is known to vary from the bottom to the tip of the grass blades due to variation in vegetation characteristics (Vivoni 1998; Nepf & Vivoni 2000). The turbulence model for the flow through the meadow can also be improved from one with constant eddy viscosity (Ghisalberti & Nepf 2002, 2004). Although these model improvements might lead to a better agreement between the observed and predicted quantities, the insight furnished by (5.1) and (5.3), and therefore our main conclusions, remain useful.

We now test the assumption of an undeformable grass bed due to the dominant restoring force of buoyancy, using the criterion that the buoyancy time scale be much shorter than the hydrodynamic time scale H/U_0 . For a common seagrass, *Zostera marina*, the relative density difference is $\Delta\rho/\rho \approx 0.25$, the volume fraction is $V_f \approx 0.1$ and $H = 1$ m (Fonseca 1998), yielding the buoyancy time scale to be $\sqrt{\rho H/V_f \Delta\rho g} \approx 2$ s. The hydrodynamic time scale assuming $U_0 \approx 0.1$ m s⁻¹ is 10 s, and is therefore longer than the buoyancy time scale. We have neither accounted for the pre-factors appearing in the scaling argument nor considered cases when the time scale separation is not so evident. Accounting for these factors can lead to further interesting behaviour (Py *et al.* 2006; Gosselin & De Langre 2009), and motivates further investigation.

6. Conclusion

In conclusion, we have shown that the hydrodynamic instability underlying *monami* differs from the traditional KH instability due to the presence of the vegetation drag. The threshold flow condition observed in the field and in laboratory experiments arises due to the presence of this drag. Furthermore, our linear stability

analysis reveals two modes, namely **mode 1** with a mechanism similar to **the KH instability** and **mode 2** with characteristics distinctly different from **the KH instability**. We are unable to determine based on observations, and therefore have refrained from identifying, which mode is observed in experiments, **and this** still remains a subtle question and subject of future investigation. Since the two modes merge for the experimental parameters, **the KH instability** may not be assumed to underlie *monami*. The spatial structure of the instability modes has direct implications for transport in the grass bed. Our analysis also informs flow structure formation in many other related scenarios, such as flow over coral reefs **and** permeable sediments, and flow through urban environments, **and therefore is** expected to have a wider impact.

Acknowledgement

M.M.B. was supported by the Collective Interactions Unit, OIST Graduate University, while visiting Brown University. We thank H. Nepf, M. Ghisalberti, M. Luhar **and** L. Mahadevan for helpful discussions.

~~Supplementary movies/material~~

~~Supplementary movies/material is available at <http://dx.doi.org/10.1017/jfm.2015.642>.~~

References

- ACKERMAN, J. D. & OKUBO, A. 1993 Reduced mixing in a marine macrophyte canopy. *Funct. Ecol.* **7**, 305–309.
- CHEN, D. & JIRKA, G. H. 1997 Absolute and convective instabilities of plane turbulent wakes in a shallow water layer. *J. Fluid Mech.* **338**, 157–172.
- CHU, V. H., WU, J.-H. & KHAYAT, R. E. 1991 Stability of transverse shear flows in shallow open channels. *J. Hydraul. Engng* **117** (10), 1370–1388.
- DRAZIN, P. G. & REID, W. H. 1981 *Hydrodynamic Stability*. Cambridge University Press.
- FONSECA, M. 1998 Exploring the basic pattern expression in seagrass landscape. PhD thesis, University of California, Berkeley.
- FONSECA, M. S. & KENWORTHY, W. J. 1987 Effects of current on photosynthesis and distribution of seagrasses. *Aquat. Bot.* **27** (1), 59–78; ~~environmental impacts on seagrasses.~~
- GHISALBERTI, M. & NEPF, H. M. 2002 Mixing layer and coherent structures in vegetated aquatic flows. *J. Geophys. Res.* **107**.
- GHISALBERTI, M. & NEPF, H. M. 2004 The limited growth of vegetated shear layers. *Water Resour. Res.* **40** (7).
- GHISALBERTI, M. & NEPF, H. M. 2006 The structures of shear layer in flows over rigid and flexible canopies. *Environ. Fluid Mech.* 277–301.
- GOSSELIN, F. & DE LANGRE, E. 2009 Destabilising effects of plant flexibility in air and aquatic vegetation canopy flows. *Eur. J. Mech. (B/Fluids)* **28** (2), 271–282.
- GRIZZLE, R. E., SHORT, F. T., NEWELL, C. R., HOVEN, H. & KINDBLUM, L. 1996 Hydrodynamically induced synchronous waving of seagrasses: ‘monami’ and its possible effects on larval mussel settlement. *J. Expl Mar. Biol. Ecol.* **206** (1–2), 165–177.
- HINCH, E. J. 1991 *Perturbation Methods*. Cambridge University Press.
- IKEDA, S. & KANAZAWA, M. 1996 Three-dimensional organized vortices above flexible water plants. *J. Hydraul. Engng* **122** (11), 634–640.
- INOUE, E. 1956 Studies of the phenomena of waving plants caused by wind. *J. Agric. Meteorol.* **11** (4), 147–151.
- NEPF, H. M. 1999 Drag, turbulence, and diffusion in flow through emergent vegetation. *Water Resour. Res.* **35** (2), 479–489.

- NEPF, H. M. 2012 Flow and transport in regions with aquatic vegetation. *Annu. Rev. Fluid Mech.* **44**, 123–142.
- NEPF, H. M. & VIVONI, E. R. 2000 Flow structure in depth-limited vegetated flow. *J. Geophys. Res.* **105**, 28547–28557.
- NEPF, H. M., WHITE, B. & MURPHY, E. 2007 Retention time and dispersion associated with submerged aquatic canopies. *Water Resour. Res.* **43**.
- OKAMOTO, T., NEZU, I. & IKEDA, H. 2012 Vertical mass and momentum transport in open-channel flows with submerged vegetations. *J. Hydro-Environ. Res.* **6** (4), 287–297.
- PY, C., DE LANGRE, E. & MOULIA, B. 2006 A frequency lock-in mechanism in the interaction between wind and crop canopies. *J. Fluid Mech.* **568**, 425–449.
- RAUPACH, M. R., FINNIGAN, J. J. & BRUNET, Y. 1996 Coherent eddies and turbulence in vegetation canopies: the mixing layer analogy. *Boundary-Layer Meteorol.* **78**, 351–382.
- RAYLEIGH, L. 1879 On the stability, or instability, of certain fluid motions. *Proc. Lond. Math. Soc.* **1** (1), 57–72.
- VIVONI, E. R. 1998 Turbulence structure of a model seagrass meadow. Master’s thesis, MIT.
- WHITE, B. L. & NEPF, H. M. 2007 Shear instability and coherent structures in shallow flow adjacent to a porous layer. *J. Fluid Mech.* **593**, 1–32.

Please do not answer on this page but find the appropriate point in the text and make your annotation there.

Author Queries

Journal: FLM

Article id: 00642

Author: R. Singh, M. M. Bandi, A. Mahadevan and S. Mandre

Short title: Linear stability analysis for *monami* in a submerged seagrass bed

Q1 (Page 1)

Au: The distinction between surnames can be ambiguous, therefore to ensure accurate tagging for indexing purposes online (eg for PubMed entries), please check that the highlighted surnames have been correctly identified, that all names are in the correct order and spelt correctly.

Q2 (Page 1)

Au: Please check that the affiliations are correct for all authors. Also please ensure that department, institution, town, postal code and country are included if missing.

Q3 (Page 11)

Au: Supplementary movies in the article are not cited in the text. Please cite it in appropriate place(s).

Q4 (Page 11)

Au: Fonseca & Kenworthy (1987): Please check the reference given here is OK.

Q5 (Page 11)

Au: Ghisalberti & Nepf (2002, 2004), Nepf (2007): Please provide page/article number, if available.

Q6 (Page 11)

Au: Ghisalberti & Nepf (2006): Please provide volume number, if available.