






(a)
175
150
125
100
75
50
25
0
MSE($\widehat{u}_t$) ((mm/s)^2)
0
10
20
30
t (observations)
(b)
Regularized projector
$\overline{P}_t^{\text{iso}}$
$P_{\widehat{x}_t}$
$\overline{P}_t^{+}$
9
10
11
12
13
MSE($\widehat{u}$) ((mm/s)^2)
$\overline{P}_t^{+}$
$P_{\widehat{x}_t}$
Regularized projector
GREAT



(a)
Regularized projector
$\overline{P}_t^{\text{iso}}$
$P_{\widehat{\mathcal{X}}_t}$
0.0
0.5
1.0
1.5
2.0
$\text{MSE}_{\text{ref}} - \text{MSE}_{\text{B}}$ ($(\text{mm/s})^2$)



(a)
10^{-10}
10^{-12}
10^{-15}
Aggregation discrepancy
parameter
mean
mode
0
10
20
30
Observations per block
(b)
Delayed action
Common endpoint
Block likelihood
Response MSE ((mm/s)^2)
5
10
15
20
25
30
35
0
10
20
30
Observations per block
(c)
Action delay (s)
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
1.6
0
10
20
30
Observations per block



Interval contains zero
$P_{\widehat{\mathcal{X}}_t}$
$\overline{P}_t^{\text{iso}}$
GREAT
GREAT EW
Regularized projector
1
4
16
32
Number of observations
100
50
0
−50
−100
$\text{MSE}_{\text{ref}} - \text{MSE}_{\text{B}}$ $((\text{mm/s})^2)$


Interval contains zero
$P_{\widehat{x}_t}$
$\overline{P}_t^{\text{iso}}$
GREAT
GREAT EW
Regularized projector
1
4
16
32
Number of observations
100
50
0
−50
−100
$\text{MSE}_{\text{ref}} - \text{MSE}_{\text{B}}$ $((\text{mm/s})^2)$


Interval contains zero
$P_{\widehat{\mathcal{X}}_t}$
$\overline{P}_t^{\text{iso}}$
GREAT
GREAT EW
Regularized projector
$\overline{P}_t^{+}, G_t = I$
$P_{\widehat{\mathcal{X}}_t}, G_t = I$
4
16
32
64
128
192
Number of observations
100
50
0
−50
−100
$\text{MSE}_{\text{ref}} - \text{MSE}_{\text{B}}$ $((\text{mm/s})^2)$


Interval contains zero
$P_{\widehat{\mathcal{X}}_t}$
$\overline{P}_t^{\text{iso}}$
GREAT
GREAT EW
Regularized projector
$\overline{P}_t^{+}, G_t = I$
$P_{\widehat{\mathcal{X}}_t}, G_t = I$
4
16
32
64
128
192
Number of observations
100
50
0
−50
−100
$\text{MSE}_{\text{ref}} - \text{MSE}_{\text{B}}$ $((\text{mm/s})^2)$


(a)
(b)
(c)
n_1
n_2
n_3
$q([n])$
10^1
10^0
10^{-2}
$v^\top(\overline{P} - \overline{P}^2)v/\|v\|^2$
θ (rad)
$\overline{P}_t^+$
$\overline{P}_t^{\text{iso}}$
0.09
0.10
0.11
0.12
0.13
0.14
0.15
0
$\pi/2$
π
$3\pi/2$
2π


(a)
(b)
(c)
(d)
n_1
n_2
n_3
$q([n])$
10^1
10^0
10^{-2}
Cumulative probability
Chordal distance
$q_{A^{(1)}}$
$q_{A^{(2)}}$
$v^\top(\overline{P} - \overline{P}^2)v$
θ (rad)