



# Structure Preserving Discretizations using Retraction Maps

**Shreyas N B**  
210010061

**A project report submitted in partial fulfillment of  
the requirements for the degree of Bachelor of Technology,  
Specialisation Area of Systems and Controls Engineering**

**Indian Institute of Technology, Bombay  
Department of Aerospace Engineering**

**Advisor: Prof. Ravi Banavar  
Co-Advisor: Prof. Krishnendu Haldar**

November 1, 2024



# Statement of Integrity

I declare that this written submission represents my ideas in my own words and where others' ideas or words have been included, I have adequately cited and referenced the original sources.

I also declare that I have adhered to all principles of academic honesty and integrity and have not misrepresented, fabricated or falsified any idea/data/fact/source in my submission.

I understand that any violation of the above will be cause for disciplinary action by the Institute and can also evoke penal action from the sources which have thus not been properly cited or from whom proper permission has not been taken when needed.

IIT Bombay, November 1, 2024

Shreyas N B  
210010061



# Abstract

Mechanical systems are most often described by a set of continuous-time, nonlinear, second-order differential equations (SODEs) of a particular structure governed by the covariant derivative. The digital implementation of controllers for such systems requires a discrete model of the system and hence requires numerical discretization schemes. Feedback linearizability of such sampled systems, however, depends on the discretization scheme employed.

In this thesis, we utilize retraction maps and their lifts to construct feedback linearizable discretizations for SODEs which can be applied to many mechanical systems.

**Keywords:** geometric integrators, retraction maps, discrete systems, feedback linearization, mechanical systems



# Acknowledgement

I would like to thank Prof. David M. Diego, Insituto de Ciencias Matemáticas (ICMAT), for his guidance in the development of the theory in this document.

I would also like to acknowledge the support of my family, friends, and colleagues, who have supported me throughout the development of this work.





# Contents

<b>List of Figures</b>	<b>xi</b>
<b>List of Tables</b>	<b>xiii</b>
<b>List of Symbols</b>	<b>xv</b>
<b>1 Retraction Maps</b>	<b>1</b>
1.1 Introduction . . . . .	1
1.2 Discretization maps . . . . .	2
1.2.1 Examples . . . . .	2
1.3 Lifts of discretization maps . . . . .	2
1.3.1 Tangent Lifts of Discretization Maps . . . . .	3
<b>Appendix A Appendix Title Here</b>	<b>5</b>



# List of Figures

1.1	Retraction maps: A visualization . . . . .	1
3.1	$\mathcal{D}^{TM}$ and $\mathcal{D}^{TN}$ commute as shown . . . . .	3



# List of Tables

2.1	Examples of discretization maps . . . . .	2
-----	---	---



# List of Symbols

$\mathcal{R}$  retraction map  
 $\mathcal{D}$  discretization map





# Chapter 1

## Retraction Maps

### 1.1 Introduction

The notion of a retraction map is fundamental in research areas like optimization theory, machine learning, numerical analysis, and in this context, geometric integrators.

Many mechanical systems usually evolve on manifolds, which naturally requires some method of discretely approximating the dynamics on the manifold (i.e., the geodesic).

In Riemannian geometry, this idea is given by the exponential map. On a Riemannian manifold  $(M, g)$ , we define  $\exp : T_x M \rightarrow M$  as the exponential map at the point  $x$ . For instance, if  $\gamma : [0, 1] \rightarrow M$  is a unique geodesic on  $M$ , and  $\gamma(0) = x$ , then  $\exp_x(v) = \gamma(1)$ , where  $v \in T_x M$  is the initial velocity of the geodesic at  $x$  such that  $\dot{\gamma}(0) = v$ .




Figure 1.1: Retraction maps: A visualization

Let  $M$  be an  $n$  dimensional manifold, and  $TM$  be its tangent bundle.

**Definition 1.1.1.** We define a **retraction map** on a manifold  $M$  as a smooth map  $\mathcal{R} : TM \rightarrow M$ , such that if  $\mathcal{R}_x$  be the restriction of  $\mathcal{R}$  to  $T_x M$ , then the following properties are satisfied:

1.  $\mathcal{R}_x(0_x) = x$  where  $0_x$  is the zero element of  $T_x M$ .
2.  $D\mathcal{R}_x(0_x) = T_{0_x} \mathcal{R}_x = Id_{T_x M}$ , where  $Id_{T_x M}$  is the identity mapping on  $T_x M$ .

Here, the first property is trivial, whereas the second property is known as the **local rigidity condition** since, given  $v \in T_x M$ , the curve  $\gamma_v(t) = \mathcal{R}_x(tv)$  has initial velocity  $v$  at  $x$ . Hence,

$$\dot{\gamma}_v(t) = \langle D\mathcal{R}_x(tv), v \rangle \implies \dot{\gamma}_v(0) = Id_{T_x M}(v) = v$$

## 1.2 Discretization maps

**Definition 1.2.1.** A map  $\mathcal{D} : U \subset TM \longrightarrow M \times M$  given by

$$\mathcal{D}(x, v) = (\mathcal{R}_x^1(v), \mathcal{R}_x^2(v))$$

where  $U$  is the open neighborhood of the zero section  $0_x \in TM$ , is called a **discretization map** on  $M$ , if the following properties are satisfied:

1.  $\mathcal{D}(x, 0_x) = (x, x)$
2.  $T_{0_x}\mathcal{R}_x^2 - T_{0_x}\mathcal{R}_x^1 = Id_{T_xM}$ , which is the identity map on  $T_xM$  for any  $x \in M$ .

Using this definition, one can prove (not included here) that the discretization map  $\mathcal{D}$  is a local diffeomorphism around the zero section  $0_x \in TM$ . This is a crucial property for the construction of geometric integrators, since we need to be able to define  $\mathcal{D}^{-1}(x_k, x_{k+1})$ .

Thus, given a vector field  $X \in \mathfrak{X}(M)$  on  $M$ , i.e.,  $X : M \longrightarrow TM$  such that  $\tau_M \circ X = Id_M$ , where  $\tau_M : TM \longrightarrow M$  is the canonical projection on the tangent bundle, we can approximate the integral curve by the following first-order discrete equation:

$$hX(\tau_M(\mathcal{D}^{-1}(x_k, x_{k+1}))) = \mathcal{D}^{-1}(x_k, x_{k+1})$$

Hence, given an initial condition  $x_0$ , we may be able to solve the discrete equation iteratively to obtain the sequence  $\{x_k\}$  which is indeed an approximation of  $\{x(kh)\}$ , where  $x(t)$  is the integral curve of  $X$  with initial condition  $x_0$  and time-step  $h$ .

### 1.2.1 Examples

We consider a few examples of discretization maps on  $\mathbb{R}^n$ :

Discretization map $\mathcal{D}$	Scheme	Order
$\mathcal{D}(x, v) = (x, x + v)$	Forward Euler $x_{k+1} = x_k + hX(x_k)$	$\mathcal{O}(h)$
$\mathcal{D}(x, v) = (x - v, x)$	Backward Euler $x_k = x_{k+1} - hX(x_{k+1})$	$\mathcal{O}(h)$
$\mathcal{D}(x, v) = \left(x - \frac{v}{2}, x + \frac{v}{2}\right)$	Symmetric Euler $x_{k+1} = x_k + hX\left(\frac{x_k + x_{k+1}}{2}\right)$	$\mathcal{O}(h^2)$

Table 2.1: Examples of discretization maps

## 1.3 Lifts of discretization maps

As mentioned before, discretization maps are diffeomorphisms around the zero section  $0_x \in TM$ . This is useful because typically when studying mechanical systems, we would like to define the discretization map on the tangent bundle  $TM$  (for Lagrangian frameworks) or the cotangent bundle  $T^*M$  (for Hamiltonian frameworks), in order to generate geometric integrators on the manifold.

Thus, since discretization maps can be defined on different manifolds, we denote  $\mathcal{D}^{TM} : TM \longrightarrow M \times M$  as a discretization map on  $M$ .

### 1.3.1 Tangent Lifts of Discretization Maps

Given a smooth map  $\varphi : M \rightarrow N$  between two  $n$ -dimensional manifolds  $M$  and  $N$ , we can define the **tangent lift** of  $\varphi$  as the map  $T\varphi : TM \rightarrow TN$  such that

$$T\varphi(v_x) = T_x\varphi(v_x) \in T_{\varphi(x)}N$$

where  $v_x \in T_xM$  and  $T_x\varphi$  is the tangent map of  $\varphi$ , whose matrix is the Jacobian at  $x \in M$ , in a local chart.

**Proposition 1.3.1.** *Let  $M$  and  $N$  be two  $n$ -dimensional manifolds, and  $\varphi : M \rightarrow N$  be a smooth map (diffeomorphism). For a given discretization map  $\mathcal{D}^{TM}$  on  $M$ , the map  $\mathcal{D}_\varphi := (\varphi \times \varphi) \circ \mathcal{D}^{TM} \circ T\varphi^{-1}$  is a discretization map on  $N$  i.e.,  $\mathcal{D}_\varphi \equiv \mathcal{D}^{TN} : TN \rightarrow N \times N$ .*

*Proof.* For any given  $y \in N$ , we have that

$$\begin{aligned} \mathcal{D}_\varphi(y, 0_y) &= ((\varphi \times \varphi) \circ \mathcal{D}^{TM} \circ T\varphi^{-1})(y, 0_y) \\ &= ((\varphi \times \varphi) \circ \mathcal{D}^{TM} \circ T\varphi^{-1})(\varphi(x), 0_{\varphi(x)}) \\ &= (\varphi \times \varphi) \circ \mathcal{D}^{TM}(x, 0_x) \\ &= (\varphi \times \varphi)(x, x) = (y, y) \end{aligned}$$

which proves the first condition. For the second condition, let  $v_y \in T_yN$ , be a given vector.

$$\begin{aligned} (T_{0_x}\mathcal{R}_{x,\varphi}^2 - T_{0_x}\mathcal{R}_{x,\varphi}^1)(y, u_y) &= \left. \frac{d}{ds} \right|_{s=0} (\mathcal{R}_{x,\varphi}^2(y, su_y) - \mathcal{R}_{x,\varphi}^1(y, su_y)) \\ &= \left. \frac{d}{ds} \right|_{s=0} (\varphi \circ \mathcal{R}_x^1 \circ T\varphi^{-1}(y, su_y)) - (\varphi \circ \mathcal{R}_x^2 \circ T\varphi^{-1}(y, su_y)) \\ &= T_y\varphi \left( \left. \frac{d}{ds} \right|_{s=0} [\mathcal{R}_x^1(t(T\varphi^{-1}(y, u_y)))] - [\mathcal{R}_x^2(t(T\varphi^{-1}(y, u_y)))] \right) \\ &= T_y\varphi(T_y\varphi^{-1}(y, u_y)) = (y, u_y) \end{aligned}$$

Thus, both the conditions from Definition 1.1.1 are satisfied. □

The above proposition can be visualized as shown below in Figure 3.1.

$$\begin{array}{ccc} TM & \xrightarrow{T\varphi} & TN \\ \mathcal{D}^{TM} \downarrow & & \downarrow \mathcal{D}^{TN} \\ M \times M & \xrightarrow{\varphi \times \varphi} & N \times N \end{array}$$

Figure 3.1:  $\mathcal{D}^{TM}$  and  $\mathcal{D}^{TN}$  commute as shown



## **Appendix A**

### **Appendix Title Here**

Write your Appendix content here.