

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

working Rule:-

Step I) Given $f(x)=0$ — ①

find initial root x_0 such that $f(x_0) \neq 0$ (i.e. x_0 is near to root of ①).

Step-II

find $f(x_0)$ and $f'(x_0)$
derivative of $f(x)$

Step-III

first approximate root by N-R method

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

find $f(x_1)$ and $f'(x_1)$

Step-IV - 2nd approximator $a_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$

like that 3rd approximate root

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)}$$

∴ General formula

$$x_n = x_{n-1} - \frac{f(x_{n-1})}{f'(x_{n-1})}$$

Repeat the Step-II upto get correct accurate root.

Q ⇒ Find the real root of eqⁿ $x^3 - 3x + 1 = 0$ by N/w Raphson method correct upto 4 decimal places.

Solⁿ given

Let $f(x) = 0$

$$x^3 - 3x + 1 = 0 \quad \text{--- ①}$$

To find x_0

$$f(0) = (0)^3 - 3(0) + 1 = 1 > 0$$

$$f(1) = 1 - 3(1) + 1 = -1 < 0$$

$$f(0.5) = (0.5)^3 - 3(0.5) + 1 = -0.375 < 0$$

$$f(0.4) = (0.4)^3 - 3(0.4) + 1 = -0.136 < 0$$

$$f(0.3) = (0.3)^3 - 3(0.3) + 1 = 0.127 > 0$$

choosing $x_0 = 0.3$

$$f(x_0) = 0.127$$

$$f'(x) = 3x^2 - 3 = 3(0.3)^2 - 3 = -2.73$$

1st approximate root by Newton-Raphson method

$$x_1 = x_0 - \left[\frac{f(x_0)}{f'(x_0)} \right]$$

$$= 0.3 - \left[\frac{0.127}{-2.73} \right] = 0.3 + \left[\frac{127}{273} \right] = 0.346520$$

$$f(x_1) = x^3 - 3x + 1 = 0.002048$$

$$f'(x_1) = 3x^2 - 3 = -2.639771$$

2nd approximate root by N-R method

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = 0.34652 - \frac{(0.002048)}{(-2.639771)}$$

$$x_2 = 0.347295$$

$$f(x_2) = f(0.347295) = x^3 - 3x + 1 = 0.000035$$

$$f'(x_2) = f'(0.347295) = 3x^2 - 3 = 3(0.347295)^2 - 3 = -2.638158$$

3rd approximate root by N-R method

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} = 0.347295 - \frac{0.000035}{(-2.638158)}$$

$$= 0.347295 - 0.000013266$$

$$= 0.3472$$

hence the approximate root correct upto 4 decimal places

$$\boxed{x = 0.3472} - \text{Ans}$$

Q.2 find the real root of eqⁿ cube of 48 by N-R method correct upto 4-decimal places.
solⁿ given

$$x^3 = 48$$

$$x^3 - 48 = 0$$

Let $\therefore f(x) = x^3 - 48 = 0$

Find x_0 $f(0) = -48$ $f(1) = -47$ $f(2) = -40 < 0$
 $f(3) = 27 - 48 = -21 < 0$
 $f(4) = 64 - 48 = 16 > 0$

\therefore Find $f(3.5) = (3.5)^3 - 48 = -5.125 < 0$
 $f(3.6) = (3.6)^3 - 48 = -1.344 < 0$
 $f(3.7) = (3.7)^3 - 48 = 2.6553 > 0$

by choosing $x_0 = 3.6$ $f(x_0) = -1.344$

$$f'(x) = 3x^2 = f'(3.6) = 3(3.6)^2 = 38.88$$

1st approximate root by N-R Method.

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$= 3.6 - \frac{(-1.344)}{(38.88)} = 3.634567$$

$$f(x_1) = x^3 - 48 = 0.012910$$

$$f'(x_1) = 3x^2 = 39.630231$$

2nd approximate root by N-R method.

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = 3.634567 - \frac{0.012910}{(39.630231)}$$

$$= 3.634241$$

$$f(x_2) = x^3 - 48 = 0.00007$$

$$f'(x_2) = 3x^2 = 3(3.634241)^2 = 39.623144$$

3rd approximate root by N-R method.

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} = 3.634241 - \frac{0.000007}{39.623144}$$

$$= \underline{\underline{3.634240}}$$

hence, the approximate root correct upto 4 decimal places.

$$x = 3.634240.$$