

# Assignment 5

## Probability and random Variables

Shreyas Wankhede

Indian Institute of Technology Hyderabad

May 19, 2022

# Outline

- 1 figure
- 2 Example
- 3 Independence of events
- 4 statements
- 5 explanation
- 6 Generalization

figure




Figure: fig-1

# Example

## Papoulis book example 2.19

The below example is an illustration that emphasizes that three events can be independent in pairs but not independent

Suppose that the events A,B,C of Fig-1 have the same probability.

$$P(A) = P(B) = P(C) = \frac{1}{5}$$

and the intersections AB,AC,BC,ABC also have the same probability

$$p = P(AB) = P(BC) = P(AC)$$

# Independence of events

## Independence of three events:

The events  $A_1, A_2$  and  $A_3$  are called (mutually) independent if they are independent in pairs:

$$P(A_i A_j) = P(A_i) \times P(A_j) \quad (1)$$

and

$$P(A_1 A_2 A_3) = P(A_1) \times P(A_2) \times P(A_3) \quad (2)$$

# statements

if  $p = \frac{1}{25}$

then these events are independent in pairs but they are not independent because

$$P(ABC) \neq P(A) \times P(B) \times P(C)$$

if  $p = \frac{1}{125}$

then

$$P(ABC) = P(A) \times P(B) \times P(C)$$

but the events are not independent because

$$P(AB) \neq P(A) \times P(B)$$

# Explanation

From the independence of the events  $A$ ,  $B$ , and  $C$  it follows that:

- 1 Anyone of them is independent of the intersection of the other two. Indeed, from (1) and (2) it follows that

$$P(A_1 A_2 A_3) = P(A_1) \times P(A_2) \times P(A_3) = P(A_1) \times P(A_2 A_3) \quad (3)$$

Hence the events  $A_1$  and  $A_2 A_3$  are independent

- 2 If we replace one or more of these events with their complements, the resulting events are also independent. Indeed since,

$$A_1 A_2 = A_1 A_2 A_3 \cup A_1 A_2 \bar{A}_3; \quad P(\bar{A}_3) = 1 - P(A_3)$$

we conclude with (3) that

$$P(A_1 A_2 \bar{A}_3) = P(A_1 A_2) - P(A_1 A_2) \times P(A_3) = P(A_1) \times P(A_2) \times P(\bar{A}_3)$$

- 3 Anyone of them is independent of the union of the other two.

To show that the events  $A_1$  and  $A_2 \cup A_3$  are independent, it suffices to show that the events  $A_1$  and  $\overline{A_2 \cup A_3} = \overline{A_2} \overline{A_3}$  are independent. This follows from 1 and 2



# Generalization

Generalization: The independence of  $n$  events can be defined inductively: Suppose that we have defined independence of  $k$  events for every  $k < n$ . We then say that the events  $A_1, \dots, A_n$  are independent if any  $k < n$  of them are independent and

$$P(A_1 \dots A_n) = P(A_1) \dots P(A_n) \quad (4)$$

This completes the definition for any  $n$  because we have already defined independence for  $n = 2$