

# Pingala Series

Shreyas wankhede

## CONTENTS

1	JEE 2019	1
2	Pingala Series	2
3	Power of the Z transform	3

**Abstract**—This manual provides a simple introduction to Transforms

## 1 JEE 2019

Let

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \geq 1 \quad (1.1)$$

$$b_n = a_{n-1} + a_{n+1}, \quad n \geq 2, \quad b_1 = 1 \quad (1.2)$$

Verify the following using a python code.

1.1

$$\sum_{k=1}^n a_k = a_{n+2} - 1, \quad n \geq 1 \quad (1.3)$$

**Solution:** : Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/1.1.py
```




Fig. 1.1

1.2

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{10}{89} \quad (1.4)$$

**Solution:** : Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/1.1.py
```




Fig. 1.2

1.3

$$b_n = \alpha^n + \beta^n, \quad n \geq 1 \quad (1.5)$$

**Solution:** : Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/1.1.py
```

1.4

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{8}{89} \quad (1.6)$$

**Solution:** : Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/1.4.py
```




Fig. 1.3




Fig. 1.4

From the figure we can see that  $b(n)$  doesn't converge to  $\frac{8}{89}$ , it converges to  $\frac{12}{89}$

## 2 PINGALA SERIES

2.1 The *one sided* Z-transform of  $x(n)$  is defined as

$$X^+(z) = \sum_{n=0}^{\infty} x(n)z^{-n}, \quad z \in \mathbb{C} \quad (2.1)$$

2.2 The *Pingala* series is generated using the difference equation

$$x(n+2) = x(n+1) + x(n) \quad (2.2)$$

$$x(0) = x(1) = 1, n \geq 0 \quad (2.3)$$

Generate a stem plot for  $x(n)$ .

**Solution:** Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/2.1.py
```

Use the following command in the terminal to run the code




Fig. 2.2

2.3 Find  $X^+(z)$ .

**Solution:** Applying positive Z-transform (linear operator) on both sides of equation 2.3

$$\sum_{k=0}^{\infty} x(k+2)z^{-k} = \sum_{k=0}^{\infty} x(k+1)z^{-k} + \sum_{k=0}^{\infty} x(k)z^{-k} \quad (2.4)$$

$$z^2(X^+(z) - x(0) - z^{-1}x(1)) = z(X^+(z) - x(0)) + X^+(z) \quad (2.5)$$

$$\Rightarrow X^+(z) = \frac{z^2}{z^2 - z - 1} \quad (2.6)$$

$$\Rightarrow X^+(z) = \frac{1}{1 - z^{-1} - z^{-2}} \quad (2.7)$$

2.4 Find  $x(n)$ .

**Solution:**

$$\frac{1}{1 - z^{-1} - z^{-2}} = \frac{1}{(1 - \alpha z^{-1})(1 - \beta z^{-1})} \quad (2.8)$$

where  $\alpha, \beta$  are the roots of equation

$$z^2 - z - 1 = 0 \quad (2.9)$$

Hence,

$$X^+(z) = \frac{1}{(1 - \alpha z^{-1})(1 - \beta z^{-1})} \quad (2.10)$$

$$\Rightarrow X^+(z) = \frac{1}{\alpha - \beta} \left( \frac{\alpha}{1 - \alpha z^{-1}} - \frac{\beta}{1 - \beta z^{-1}} \right) \quad (2.11)$$

As

$$a^n u(n) \stackrel{Z}{=} \frac{1}{1 - az^{-1}} \quad |z| > |a| \quad (2.12)$$

$$\Rightarrow a^{n+1} u(n) \stackrel{Z}{=} \frac{a}{1 - az^{-1}} \quad |z| > |a| \quad (2.13)$$

$$\therefore x(n) = \frac{1}{\alpha - \beta} (\alpha^{n+1} - \beta^{n+1}) \quad (2.14)$$

2.5 Sketch

$$y(n) = x(n-1) + x(n+1), \quad n \geq 0 \quad (2.15)$$

**Solution:** Run the below python code

```
https://github.com/
shreyaswankhede12/EE3900/blob
/master/Assignment%201/pingala
/2.2.py
```




Fig. 2.5

2.6 Find  $Y^+(z)$ .

**Solution:** Taking +ve Z-transform on both sides of the equation 2.15, we get

$$\sum_{k=0}^{\infty} y(k) z^{-k} = \sum_{k=0}^{\infty} x(k+1) z^{-k} + \sum_{k=0}^{\infty} x(k-1) z^{-k} \quad (2.16)$$

$$Y^+(z) = z(X^+(z) - x(0)) + z^{-1} X^+(z) \quad (2.17)$$

$$\therefore x(-1) = 0$$

$$Y^+(z) = \frac{z + z^{-1}}{1 - z^{-1} - z^{-2}} - z \quad (2.18)$$

$$\therefore Y^+(z) = \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (2.19)$$

2.7 Find  $y(n)$ .

**Solution:**  $y(n)$  is coefficient of  $z^{-n}$  in  $Y^+(z)$

$$Y^+(z) = \frac{1}{1 - z^{-1} - z^{-2}} + \frac{2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (2.20)$$

Applying inverse Z-transform to 2.20, we get

$$y(k) = \frac{\alpha^{k+1} - \beta^{k+1}}{\alpha - \beta} + 2 \frac{\alpha^k - \beta^k}{\alpha - \beta} \quad (2.21)$$

$$= \frac{\alpha^{k+2} + \alpha^k - \beta^{k+2} - \beta^k}{\alpha - \beta} \quad (2.22)$$

$$= \frac{\alpha^{k+2} - \beta^{k+2} + \alpha\beta^{k+1} - \beta\alpha^{k+1}}{\alpha - \beta} [\because \alpha\beta = -1] \quad (2.23)$$

$$\therefore y(k) = \alpha^{k+1} + \beta^{k+1}$$

### 3 POWER OF THE Z TRANSFORM

3.1 Show that

$$\sum_{k=1}^n a_k = \sum_{k=0}^{n-1} x(k) = x(n) * u(n-1) \quad (3.1)$$

**Solution:**

$$x(k) = a(k+1) \quad (3.2)$$

$$\Rightarrow \sum_{k=0}^{n-1} x(k) = \sum_{k=0}^{n-1} a(k+1) \quad (3.3)$$

$$\Rightarrow \sum_{k=1}^n a_k = \sum_{k=0}^{n-1} x(k) \quad (3.4)$$

$$x(n) * u(n-1) = \sum_{k=-\infty}^{\infty} x(k) u(n-1-k) \quad (3.5)$$

$$u(n-1-k) = \begin{cases} 0 & k > n-1 \\ 1 & k \leq n-1 \end{cases} \quad (3.6)$$

$$x(k) = 0 \quad \forall k < 0 \quad (3.7)$$

$$\Rightarrow x(n) * u(n-1) = \sum_{k=0}^{n-1} x(k) \quad (3.8)$$

3.2 Show that

$$a_{n+2} - 1, \quad n \geq 1 \quad (3.9)$$

can be expressed as

$$[x(n+1) - 1] u(n) \quad (3.10)$$

**Solution:** :

$$x(k) = a(k+1) \quad (3.11)$$

$$\Rightarrow x(k+1) = a(k+2) \quad (3.12)$$

$$a(k+2) - 1 = x(k+1) - 1 \quad (3.13)$$

$$\therefore a_{n+2} - 1 = [x(k+1) - 1]u(k) \quad [\forall n \geq 1] \quad (3.14)$$

3.3 Show that

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = \frac{1}{10} X^+ (10) \quad (3.15)$$

**Solution:** :

$$X^+(z) = \sum_{k=0}^{\infty} x(k)z^{-k} = z \sum_{k=1}^{\infty} a(k)z^{-k} \quad (3.16)$$

$$z = 10 \quad (3.17)$$

$$\Rightarrow 10 \sum_{k=1}^{\infty} \frac{a_k}{10^k} = \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = X^+ (10) \quad (3.18)$$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = \frac{1}{10} X^+ (10) \quad (3.19)$$

3.4 Show that

$$\alpha^n + \beta^n, \quad n \geq 1 \quad (3.20)$$

can be expressed as

$$w(n) = (\alpha^{n+1} + \beta^{n+1})u(n) \quad (3.21)$$

and find  $W(z)$ .

**Solution:** : Applying Z-transform on 3.21

$$W(z) = \sum_{n=-\infty}^{\infty} (\alpha^{n+1} + \beta^{n+1})u(n)z^{-n} \quad (3.22)$$

$$= \sum_{n=0}^{\infty} (\alpha^{n+1} + \beta^{n+1})z^{-n} \quad (3.23)$$

$$= \alpha \sum_{n=0}^{\infty} (\alpha z^{-1})^n + \beta \sum_{n=0}^{\infty} (\beta z^{-1})^n \quad (3.24)$$

$$\text{ROC: } |z| > \max(\alpha, \beta) \quad (3.25)$$

$$= \frac{\alpha}{1 - \alpha z^{-1}} + \frac{\beta}{1 - \beta z^{-1}} \quad (3.26)$$

$$= \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (3.27)$$

3.5 Show that

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} = \frac{1}{10} Y^+ (10) \quad (3.28)$$

**Solution:** :

$$y(k) = b(k+1) \quad (3.29)$$

$$\sum_{k=0}^{\infty} y(k)z^{-k} = \sum_{k=0}^{\infty} b(k+1)z^{-k} = Y^+(z) \quad (3.30)$$

$$\sum_{k=0}^{\infty} y(k)z^{-k} = z \sum_{k=1}^{\infty} b(k)z^{-k} = Y^+(z) \quad (3.31)$$

$$z = 10 \quad (3.32)$$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} = \frac{1}{10} Y^+ (10) \quad (3.33)$$

3.6 Solve the JEE 2019 problem.

**Solution:** :

$$X^+(z) = z \sum_{k=1}^{\infty} a(k)z^{-k} = \frac{1}{1 - z^{-1} - z^{-2}} \quad (3.34)$$

$$z = 10 \quad (3.35)$$

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10 \left(1 - \frac{1}{10} - \frac{1}{100}\right)} \quad (3.36)$$

$$= \frac{10}{89} \quad (3.37)$$

$$y(k) = \alpha^{k+1} + \beta^{k+1} \quad (3.38)$$

$$y(k) = b(k+1) \quad (3.39)$$

$$\Rightarrow b(k) = \alpha^k + \beta^k \quad (3.40)$$

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} Y^+ (10) \quad (3.41)$$

$$= \frac{1}{10} \left[ \frac{1 + \frac{2}{10}}{1 - \frac{1}{10} - \frac{1}{100}} \right] \quad (3.42)$$

$$\therefore Y^+(z) = \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (3.43)$$

$$= \frac{12}{89} \quad (3.44)$$

Run the following code to get the expressions of  $x(n)$  and  $y(n)$

<https://github.com/shreyaswankhede12/EE3900/blob/master/Assignment%201/pingala>

/2.2.py