

Machine Learning HW1

Shreyas Shukla

September 16, 2024

Question 1:

Find the transpose of the following matrices and vectors (10 pts):

1.

Matrix:

$$\begin{bmatrix} 7 & 4 & 5 & 6 \\ 2 & 8 & 9 & 7 \\ 1 & 0 & 2 & 3 \end{bmatrix}$$

Transpose:

$$\begin{bmatrix} 7 & 2 & 1 \\ 4 & 8 & 0 \\ 5 & 9 & 2 \\ 6 & 7 & 3 \end{bmatrix}$$

2.

Matrix:

$$\begin{bmatrix} a & b \\ c & d \\ e & f \end{bmatrix}$$

Transpose:

$$\begin{bmatrix} a & c & e \\ b & d & f \end{bmatrix}$$

3.

Matrix:

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 2 & 0 & 0 & 8 \end{bmatrix}$$

Transpose:

$$\begin{bmatrix} 1 & 5 & 2 \\ 2 & 6 & 0 \\ 3 & 7 & 0 \\ 4 & 8 & 8 \end{bmatrix}$$

4.

Matrix:

$$\begin{bmatrix} 2 & 2 & 8 & 10 \end{bmatrix}$$

Transpose:

$$\begin{bmatrix} 2 \\ 2 \\ 8 \\ 10 \end{bmatrix}$$

5.

Matrix:

$$\begin{bmatrix} 1 \\ 2 \\ 2 \\ 8 \end{bmatrix}$$

Transpose:

$$\begin{bmatrix} 1 & 2 & 2 & 8 \end{bmatrix}$$

Question 2:

Perform the following operations. If the dimension doesn't match, just indicate "dimension mismatch". (10 pts):

1.

$$\begin{bmatrix} -4 \\ -7 \end{bmatrix} - \begin{bmatrix} 7 \\ 3 \end{bmatrix} = \begin{bmatrix} -4 - 7 \\ -7 - 3 \end{bmatrix} = \begin{bmatrix} -11 \\ -10 \end{bmatrix}$$

2.

$$\begin{bmatrix} -2 \\ 0 \\ 3 \end{bmatrix} + \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 + 2 \\ 0 + 1 \\ 3 + 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 4 \end{bmatrix}$$

3.

$$\begin{bmatrix} 1 & 2 & 5 \end{bmatrix} + \begin{bmatrix} -7 \\ 8 \\ 3 \end{bmatrix} = \text{"dimension mismatch"}$$

4.

$$\begin{bmatrix} 1 & 2 & 5 \end{bmatrix} + \begin{bmatrix} -7 & 8 & 3 \end{bmatrix} = \begin{bmatrix} 1-7 & 2+8 & 5+3 \end{bmatrix} = \begin{bmatrix} -6 & 10 & 8 \end{bmatrix}$$

5.

$$\begin{bmatrix} 1 \\ 2 \\ 5 \\ 0 \end{bmatrix} + \begin{bmatrix} -1 \\ 1 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 1-1 \\ 2+1 \\ 5+3 \\ 0+3 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 8 \\ 3 \end{bmatrix}$$

Question 3:

Given

$$x = \begin{bmatrix} 2 \\ 2 \\ 6 \end{bmatrix}, \quad X = \begin{bmatrix} -4 & 7 & 5 \\ 1 & -9 & 2 \\ 8 & 3 & 6 \end{bmatrix}, \quad Y = \begin{bmatrix} 2 & 4 & 6 \\ 1 & 0 & 13 \\ -4 & 5 & 5 \end{bmatrix}, \quad Z = \begin{bmatrix} 4 & -1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

Calculate the following (10 pts):

1. $x^\top XY$

$$\begin{aligned} x^\top &= [2 \quad 2 \quad 6] \\ x^\top X &= [2 \quad 2 \quad 6] \begin{bmatrix} -4 & 7 & 5 \\ 1 & -9 & 2 \\ 8 & 3 & 6 \end{bmatrix} = [44 \quad 4 \quad 52] \\ [44 \quad 4 \quad 52] \begin{bmatrix} 2 & 4 & 6 \\ 1 & 0 & 13 \\ -4 & 5 & 5 \end{bmatrix} &= [(44 \cdot 2 + 4 \cdot 1 + 52 \cdot -4) \quad (44 \cdot 4 + 4 \cdot 0 + 52 \cdot 5) \quad (44 \cdot 6 + 4 \cdot 13 + 52 \cdot 5)] \\ &= [88 + 4 - 208 \quad 176 + 0 + 260 \quad 264 + 52 + 260] \\ &= [-116 \quad 436 \quad 576] \end{aligned}$$

2. YX

$$\begin{aligned} YX &= \begin{bmatrix} 2 & 4 & 6 \\ 1 & 0 & 13 \\ -4 & 5 & 5 \end{bmatrix} \begin{bmatrix} -4 & 7 & 5 \\ 1 & -9 & 2 \\ 8 & 3 & 6 \end{bmatrix} \\ &= \begin{bmatrix} (-8 + 4 + 48) & (14 - 36 + 18) & (10 + 8 + 36) \\ (-4 + 104) & (7 + 39) & (5 + 78) \\ (16 + 5 + 40) & (-28 - 45 + 15) & (-20 + 10 + 30) \end{bmatrix} \\ &= \begin{bmatrix} 44 & -4 & 54 \\ 100 & 46 & 83 \\ 61 & -58 & 20 \end{bmatrix} \end{aligned}$$

3. $2X + 2Y + X^\top Y$

$$2X = \begin{bmatrix} -8 & 14 & 10 \\ 2 & -18 & 4 \\ 16 & 6 & 12 \end{bmatrix}, \quad 2Y = \begin{bmatrix} 4 & 8 & 12 \\ 2 & 0 & 26 \\ -8 & 10 & 10 \end{bmatrix}$$

$$2X + 2Y = \begin{bmatrix} -8+4 & 14+8 & 10+12 \\ 2+2 & -18+0 & 4+26 \\ 16+-8 & 6+10 & 12+10 \end{bmatrix} = \begin{bmatrix} -4 & 22 & 22 \\ 4 & -18 & 30 \\ 8 & 16 & 22 \end{bmatrix}$$

$$X^\top = \begin{bmatrix} -4 & 1 & 8 \\ 7 & -9 & 3 \\ 5 & 2 & 6 \end{bmatrix}$$

$$X^\top Y = \begin{bmatrix} -4 & 1 & 8 \\ 7 & -9 & 3 \\ 5 & 2 & 6 \end{bmatrix} \begin{bmatrix} 2 & 4 & 6 \\ 1 & 0 & 13 \\ -4 & 5 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} (-8+1-32) & (-16+0+40) & (-24+13+40) \\ (14-9-12) & (28+0+15) & (42-117+15) \\ (10+2-24) & (20+0+30) & (30+26+30) \end{bmatrix}$$

$$= \begin{bmatrix} -39 & 24 & 29 \\ -7 & 43 & -60 \\ -12 & 50 & 86 \end{bmatrix}$$

$$2X + 2Y + X^\top Y = \begin{bmatrix} -4 & 22 & 22 \\ 4 & -18 & 30 \\ 8 & 16 & 22 \end{bmatrix} + \begin{bmatrix} -39 & 24 & 29 \\ -7 & 43 & -60 \\ -12 & 50 & 86 \end{bmatrix} = \begin{bmatrix} -43 & 46 & 51 \\ -3 & 25 & -30 \\ -4 & 66 & 108 \end{bmatrix}$$

4. YZ^\top

$$Z^\top = \begin{bmatrix} 4 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$YZ^\top = \begin{bmatrix} 2 & 4 & 6 \\ 1 & 0 & 13 \\ -4 & 5 & 5 \end{bmatrix} \begin{bmatrix} 4 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} (2 \cdot 4 + 4 \cdot -1 + 6 \cdot 0) & (2 \cdot 1 + 4 \cdot 1 + 6 \cdot 1) \\ (1 \cdot 4 + 0 \cdot -1 + 13 \cdot 0) & (1 \cdot 1 + 0 \cdot 1 + 13 \cdot 1) \\ (-4 \cdot 4 + 5 \cdot -1 + 5 \cdot 0) & (-4 \cdot 1 + 5 \cdot 1 + 5 \cdot 1) \end{bmatrix}$$

$$= \begin{bmatrix} (8-4+0) & (2+4+6) \\ (4+0+0) & (1+0+13) \\ (-16-5+0) & (-4+5+5) \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 12 \\ 4 & 14 \\ -21 & 6 \end{bmatrix}$$

5. $Z^\top Zx$

$$Z^\top Z = \begin{bmatrix} 4 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 4 & -1 & 0 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} (4 \cdot 4 + 1 \cdot 1) & (4 \cdot -1 + 1 \cdot 1) & (4 \cdot 0 + 1 \cdot 1) \\ (-1 \cdot 4 + 1 \cdot 1) & (-1 \cdot -1 + 1 \cdot 1) & (-1 \cdot 0 + 1 \cdot 1) \\ (0 \cdot 4 + 1 \cdot 1) & (0 \cdot -1 + 1 \cdot 1) & (0 \cdot 0 + 1 \cdot 1) \end{bmatrix}$$

$$= \begin{bmatrix} 17 & -3 & 1 \\ -3 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 17 & -3 & 1 \\ -3 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 6 \end{bmatrix} = \begin{bmatrix} 17 \cdot 2 + -3 \cdot 2 + 1 \cdot 6 \\ -3 \cdot 2 + 2 \cdot 2 + 1 \cdot 6 \\ 1 \cdot 2 + 1 \cdot 2 + 1 \cdot 6 \end{bmatrix} = \begin{bmatrix} 34 - 6 + 6 \\ -6 + 4 + 6 \\ 2 + 2 + 6 \end{bmatrix} = \begin{bmatrix} 34 \\ 4 \\ 10 \end{bmatrix}$$

Question 4:

Given

$$X = \begin{bmatrix} 1 & 7 \\ -8 & 6 \\ 3 & 1 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 4 & 6 & -8 \\ 1 & -3 & 5 & 7 \\ 11 & 13 & 9 & 9 \\ -1 & 0 & 8 & 7 \end{bmatrix}, \quad Z = \begin{bmatrix} 0 & 4 & 8 & 1 \\ -3 & 5 & 17 & 11 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 17 \\ 2 & 14 \end{bmatrix}, \quad b = [19 \quad -11 \quad 13], \quad c = \begin{bmatrix} 12 \\ -11 \\ 10 \end{bmatrix}, \quad x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad y = [3 \quad 3 \quad 1]$$

Calculate the following both by hand and by numpy (20 pts):

1. $xyy^\top x^\top$

$$xyy^\top x^\top = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [3 \quad 3 \quad 1] \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix} [1 \quad 2 \quad 3] = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [19] = 19 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 19 \\ 38 \\ 57 \end{bmatrix}$$

2. $3c - 9$

$$3c - 9 = 3 \begin{bmatrix} 12 \\ -11 \\ 10 \end{bmatrix} - 9 = \begin{bmatrix} 36 \\ -33 \\ 30 \end{bmatrix} - \begin{bmatrix} 9 \\ 9 \\ 9 \end{bmatrix} = \begin{bmatrix} 27 \\ -42 \\ 21 \end{bmatrix}$$

3. $\langle A, A \rangle \cdot A$

$$\langle A, A \rangle = \text{trace}(A^\top A) = \text{trace} \left(\begin{bmatrix} 1 & 2 \\ 17 & 14 \end{bmatrix} \begin{bmatrix} 1 & 17 \\ 2 & 14 \end{bmatrix} \right) = \text{trace} \left(\begin{bmatrix} 5 & 45 \\ 45 & 485 \end{bmatrix} \right) = 5 + 485 = 490$$

$$\langle A, A \rangle \cdot A = 490 \cdot \begin{bmatrix} 1 & 17 \\ 2 & 14 \end{bmatrix} = \begin{bmatrix} 490 & 8330 \\ 980 & 6860 \end{bmatrix}$$

4. $2XZ$

$$2XZ = 2 \begin{bmatrix} 1 & 7 \\ -8 & 6 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 0 & 4 & 8 & 1 \\ -3 & 5 & 17 & 11 \end{bmatrix} = 2 \begin{bmatrix} -21 & 39 & 127 & 78 \\ 24 & -2 & -88 & -58 \\ -3 & 17 & 41 & 14 \end{bmatrix} = \begin{bmatrix} -42 & 78 & 254 & 156 \\ 48 & -4 & -176 & -116 \\ -6 & 34 & 82 & 28 \end{bmatrix}$$

5. $(bc)A$

$$bc = \begin{bmatrix} 19 & -11 & 13 \end{bmatrix} \begin{bmatrix} 12 \\ -11 \\ 10 \end{bmatrix} = 19 \cdot 12 + (-11) \cdot (-11) + 13 \cdot 10 = 228 + 121 + 130 = 479$$

$$(bc)A = 479 \begin{bmatrix} 1 & 17 \\ 2 & 14 \end{bmatrix} = \begin{bmatrix} 479 & 8143 \\ 958 & 6706 \end{bmatrix}$$

6. bXA

$$\begin{aligned} bXA &= \begin{bmatrix} 19 & -11 & 13 \end{bmatrix} \begin{bmatrix} 1 & 7 \\ -8 & 6 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 17 \\ 2 & 14 \end{bmatrix} \\ &= \begin{bmatrix} 19 & -11 & 13 \end{bmatrix} \begin{bmatrix} 15 & 115 \\ -10 & -94 \\ 5 & 65 \end{bmatrix} = \begin{bmatrix} 460 & 4064 \end{bmatrix} \end{aligned}$$

7. $c \odot x$

$$c \odot x = \begin{bmatrix} 12 \\ -11 \\ 10 \end{bmatrix} \odot \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 12 \\ -22 \\ 30 \end{bmatrix}$$

8. $b \odot y$

$$b \odot y = \begin{bmatrix} 19 & -11 & 13 \end{bmatrix} \odot \begin{bmatrix} 3 & 3 & 1 \end{bmatrix} = \begin{bmatrix} 57 & -33 & 13 \end{bmatrix}$$

9. $c \otimes x$

$$c \otimes x = \begin{bmatrix} 12 \\ -11 \\ 10 \end{bmatrix} \otimes \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 12 & 24 & 36 \\ -11 & -22 & -33 \\ 10 & 20 & 30 \end{bmatrix}$$

10. YZ^\top

$$\begin{aligned} YZ^\top &= \begin{bmatrix} 0 & 4 & 6 & -8 \\ 1 & -3 & 5 & 7 \\ 11 & 13 & 9 & 9 \\ -1 & 0 & 8 & 7 \end{bmatrix} \begin{bmatrix} 0 & -3 \\ 4 & 5 \\ 8 & 17 \\ 1 & 11 \end{bmatrix} \\ &= \begin{bmatrix} 39 & 127 \\ -10 & -94 \\ 73 & 241 \\ 24 & 115 \end{bmatrix} \end{aligned}$$

Question 5:

Write the code that performs the following operations using numpy. Make sure you print the results. (20 pts):

- Create two matrices and two vectors:

$$X = \begin{bmatrix} 2 & 3 \\ 9 & 1 \end{bmatrix}, Y = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}, x = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, y = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

- Write the code that extracts the 1st column of X .
- Write the code that extracts the 2nd row of Y .
- Write the code that extracts the bottom right value of X .
- Write the code that adds $X + Y$.
- Write the code that performs $X + YX$.
- Write the code that performs the Hadamard product $x \odot y$.
- Write the code that performs the outer product $x \otimes y$.
- Write the code that performs the dot product XY .
- Write the code that performs the inner product $\langle X, Y \rangle$.

Question 6:

Given vectors

$$v = \begin{bmatrix} 5 \\ 0 \\ 0 \end{bmatrix}, w = \begin{bmatrix} 8 \\ 1 \\ 1 \end{bmatrix}, x = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}, y = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 3 \end{bmatrix}, z = \begin{bmatrix} 2 \\ 5 \end{bmatrix}$$

1. L1-norm:

$$\text{L1-norm}(v) = |5| + |0| + |0| = 5$$

$$\text{L1-norm}(w) = |8| + |1| + |1| = 10$$

$$\text{L1-norm}(x) = |3| + |1| + |2| = 6$$

$$\text{L1-norm}(y) = |2| + |1| + |0| + |3| = 6$$

$$\text{L1-norm}(z) = |2| + |5| = 7$$

2. L2-norm:

$$\text{L2-norm}(v) = \sqrt{5^2 + 0^2 + 0^2} = \sqrt{25} = 5$$

$$\text{L2-norm}(w) = \sqrt{8^2 + 1^2 + 1^2} = \sqrt{64 + 1 + 1} = \sqrt{66} \approx 8.12$$

$$\text{L2-norm}(x) = \sqrt{3^2 + 1^2 + 2^2} = \sqrt{9 + 1 + 4} = \sqrt{14} \approx 3.74$$

$$\text{L2-norm}(y) = \sqrt{2^2 + 1^2 + 0^2 + 3^2} = \sqrt{4 + 1 + 0 + 9} = \sqrt{14} \approx 3.74$$

$$\text{L2-norm}(z) = \sqrt{2^2 + 5^2} = \sqrt{4 + 25} = \sqrt{29} \approx 5.39$$

3. L-infinity norm:

$$\text{L-infinity norm}(v) = \max(|5|, |0|, |0|) = 5$$

$$\text{L-infinity norm}(w) = \max(|8|, |1|, |1|) = 8$$

$$\text{L-infinity norm}(x) = \max(|3|, |1|, |2|) = 3$$

$$\text{L-infinity norm}(y) = \max(|2|, |1|, |0|, |3|) = 3$$

$$\text{L-infinity norm}(z) = \max(|2|, |5|) = 5$$

Question 7:

Use Numpy to write the code that loads the hw1.csv file and prints out the data matrix. Please show the code as well as the output. (10 pts)

All coding questions are in .ipynb file