

Limits for functions of one variable




Examples



Examples


Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.



Examples

Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.

Consider the function $f(x) = x^2$. Then we can rewrite the above statement as $f(a_n) \rightarrow f(a)$ whenever $a_n \rightarrow a$.



Examples

Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.

Consider the function $f(x) = x^2$. Then we can rewrite the above statement as $f(a_n) \rightarrow f(a)$ whenever $a_n \rightarrow a$.

In contrast, consider the floor function $g(x) = \lfloor x \rfloor$.

Examples

Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.

Consider the function $f(x) = x^2$. Then we can rewrite the above statement as $f(a_n) \rightarrow f(a)$ whenever $a_n \rightarrow a$.

In contrast, consider the floor function $g(x) = \lfloor x \rfloor$.

If one takes a sequence a_n decreasing to 2, then indeed $g(a_n) \rightarrow g(2) = 2$.

Examples

Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.

Consider the function $f(x) = x^2$. Then we can rewrite the above statement as $f(a_n) \rightarrow f(a)$ whenever $a_n \rightarrow a$.

In contrast, consider the floor function $g(x) = \lfloor x \rfloor$.

If one takes a sequence a_n decreasing to 2, then indeed $g(a_n) \rightarrow g(2) = 2$.

However, if one takes a sequence a_n increasing to 2, then $g(a_n) \rightarrow 1$.

Examples

Recall that for any convergent sequence $a_n \rightarrow a$, we obtain that $a_n^2 \rightarrow a^2$.

Consider the function $f(x) = x^2$. Then we can rewrite the above statement as $f(a_n) \rightarrow f(a)$ whenever $a_n \rightarrow a$.

In contrast, consider the floor function $g(x) = \lfloor x \rfloor$.


If one takes a sequence a_n decreasing to 2, then indeed $g(a_n) \rightarrow g(2) = 2$.

However, if one takes a sequence a_n increasing to 2, then $g(a_n) \rightarrow 1$.

Note also that for $g(x)$ this happens at each integer value and if a is a non-integer value, then indeed $g(a_n) \rightarrow g(a)$ whenever $a_n \rightarrow a$.

Another example


Consider the function $f(x) = \begin{cases} 1 & \text{if } x \text{ is a rational number} \\ 0 & \text{if } x \text{ is not a rational number} \end{cases}$.



Another example

Consider the function $f(x) = \begin{cases} 1 & \text{if } x \text{ is a rational number} \\ 0 & \text{if } x \text{ is not a rational number} \end{cases}$.

Recall that $\sqrt{2}$ is an irrational number.



Another example

Consider the function $f(x) = \begin{cases} 1 & \text{if } x \text{ is a rational number} \\ 0 & \text{if } x \text{ is not a rational number} \end{cases}$.

Recall that $\sqrt{2}$ is an irrational number. We can construct a sequence of rational numbers which tends to $\sqrt{2}$.

Another example

Consider the function $f(x) = \begin{cases} 1 & \text{if } x \text{ is a rational number} \\ 0 & \text{if } x \text{ is not a rational number} \end{cases}$.

Recall that $\sqrt{2}$ is an irrational number. We can construct a sequence of rational numbers which tends to $\sqrt{2}$.

Then $f(a_n) = 1 \forall n$ whereas $f(\sqrt{2}) = 0$. Thus even though $a_n \rightarrow a$, $f(a_n) \not\rightarrow f(\sqrt{2})$.

$$\sqrt{2} = 1.414\dots$$

$a_1 = 1.4$, $a_2 = 1.41$, $a_3 = 1.414$

a_n = the number obtained by taking the first n terms after the decimal point in $\sqrt{2}$.

Another example

Consider the function $f(x) = \begin{cases} 1 & \text{if } x \text{ is a rational number} \\ 0 & \text{if } x \text{ is not a rational number} \end{cases}$.

Recall that $\sqrt{2}$ is an irrational number. We can construct a sequence of rational numbers which tends to $\sqrt{2}$.


Then $f(a_n) = 1 \forall n$ whereas $f(\sqrt{2}) = 0$. Thus even though $a_n \rightarrow a$,
 $f(a_n) \not\rightarrow f(\sqrt{2})$.

Limit of a function at a point from the left



Limit of a function at a point from the left


Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .



Limit of a function at a point from the left

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .

If there is a real number L such that $f(a_n) \rightarrow L$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n < a$, then we say that **the limit of f at a from the left exists and equals L** .



Limit of a function at a point from the left

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .

If there is a real number L such that $f(a_n) \rightarrow L$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n < a$, then we say that **the limit of f at a from the left exists and equals L** .

We denote this by $\lim_{x \rightarrow a^-} f(x) = L$.

Limit of a function at a point from the left

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .


If there is a real number L such that $f(a_n) \rightarrow L$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n < a$, then we say that **the limit of f at a from the left exists and equals L** .

We denote this by $\lim_{x \rightarrow a^-} f(x) = L$.

If there is no such number L then we say that the limit of f at a from the left does not exist.


An equivalent way of thinking of $\lim_{x \rightarrow a^-} f(x) = L$ is that as x comes closer and closer to a from the left, $f(x)$ eventually comes closer and closer to L .

Limit of a function at a point from the right



Limit of a function at a point from the right


Similarly, if there is a real number R such that $f(a_n) \rightarrow R$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n > a$, then we say that **the limit of f at a from the right exists and equals R .**



Limit of a function at a point from the right

Similarly, if there is a real number R such that $f(a_n) \rightarrow R$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n > a$, then we say that **the limit of f at a from the right exists and equals R .**

We denote this by $\lim_{x \rightarrow a^+} f(x) = L$.



Limit of a function at a point from the right

Similarly, if there is a real number R such that $f(a_n) \rightarrow R$ for all sequences a_n such that $a_n \rightarrow a$ and $a_n > a$, then we say that **the limit of f at a from the right exists and equals R .**

We denote this by $\lim_{x \rightarrow a^+} f(x) = L$.

If there is no such number R then we say that the limit of f at a from the right does not exist.

An equivalent way of thinking of $\lim_{x \rightarrow a^+} f(x) = R$ is that as x comes closer and closer to a from the left, $f(x)$ eventually comes closer and closer to R .

Limit of a function at a point



Limit of a function at a point


Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .



Limit of a function at a point

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .

Suppose the limit of f at a from both sides (i.e. left and right) exist and are equal i.e. $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$. Then we say that the limit of f at a exists and equals the left (and right) limit.



Limit of a function at a point

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n belongs to the domain of definition of f .

Suppose the limit of f at a from both sides (i.e. left and right) exist and are equal i.e. $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$. Then we say that the limit of f at a exists and equals the left (and right) limit.

We denote it by $\lim_{x \rightarrow a} f(x)$ or $f(a)$

$$\lim_{x \rightarrow a^-} x^2 = a^2$$

$$\lim_{x \rightarrow a^+} x^2 = a^2$$

$$g(x) = \lfloor x \rfloor : x \rightarrow a$$

$$\lim_{x \rightarrow a^-} \lfloor x \rfloor = 1$$

$$\lim_{x \rightarrow a^+} \lfloor x \rfloor = 2$$

$$f(x) = \begin{cases} 0 & \text{if } x \in \mathbb{Q} \\ 1 & \text{if } x \notin \mathbb{Q} \end{cases}$$

$$\lim_{x \rightarrow a^-} f(x) = 0$$

$$\lim_{x \rightarrow a^+} f(x) = 1$$

$$\lim_{x \rightarrow a} f(x) \text{ DNE}$$

$$\lim_{x \rightarrow a} x^2 : f(x) = x^2$$

$$\lim_{x \rightarrow a^-} \lfloor x \rfloor = \lfloor a \rfloor = g(a); f(a) \neq g(a)$$

$$\lim_{x \rightarrow a^+} \lfloor x \rfloor = 2 : \lim_{x \rightarrow 2} g(x) = 2$$

$$\lim_{x \rightarrow a} f(x) \text{ DNE}$$

$$\lim_{x \rightarrow a} f(x) \text{ DNE}$$


$$\lim_{x \rightarrow a} f(x) \text{ DNE}$$

The limit as $x \rightarrow (\pm)\infty$



The limit as $x \rightarrow (\pm)\infty$


Let f be a function such that there is an M such that it is defined for all $x > M$.



The limit as $x \rightarrow (\pm)\infty$

Let f be a function such that there is an M such that it is defined for all $x > M$.

Suppose for all sequences x_n diverging to ∞ , there exists L such that $f(x_n)$ converges to L (i.e. as x becomes larger and larger, $f(x)$ eventually gets closer and closer to L).




The limit as $x \rightarrow (\pm)\infty$

Let f be a function such that there is an M such that it is defined for all $x > M$.

Suppose for all sequences x_n diverging to ∞ , there exists L such that $f(x_n)$ converges to L (i.e. as x becomes larger and larger, $f(x)$ eventually gets closer and closer to L). Then we say that

$\lim_{x \rightarrow \infty} f(x)$ exists and equals L .



The limit as $x \rightarrow (\pm)\infty$

Let f be a function such that there is an M such that it is defined for all $x > M$.

Suppose for all sequences x_n diverging to ∞ , there exists L such that $f(x_n)$ converges to L (i.e. as x becomes larger and larger, $f(x)$ eventually gets closer and closer to L). Then we say that
$$\lim_{x \rightarrow \infty} f(x) \text{ exists and equals } L.$$

Similarly, let f be a function such that there is an N such that it is defined for all $x < N$.

The limit as $x \rightarrow (\pm)\infty$

Let f be a function such that there is an M such that it is defined for all $x > M$.

Suppose for all sequences x_n diverging to ∞ , there exists L such that $f(x_n)$ converges to L (i.e. as x becomes larger and larger, $f(x)$ eventually gets closer and closer to L). Then we say that
$$\lim_{x \rightarrow \infty} f(x) \text{ exists and equals } L.$$

Similarly, let f be a function such that there is an N such that it is defined for all $x < N$.

Suppose for all sequences x_n diverging to $-\infty$, there exists L such that $f(x_n)$ converges to L (i.e. as x becomes smaller and smaller, $f(x)$ eventually gets closer and closer to L).

The limit as $x \rightarrow (\pm)\infty$

Let f be a function such that there is an M such that it is defined for all $x > M$.

Suppose for all sequences x_n diverging to ∞ , there exists L such that $f(x_n)$ converges to L (i.e. as x becomes larger and larger, $f(x)$ eventually gets closer and closer to L). Then we say that


$$\lim_{x \rightarrow \infty} f(x) \text{ exists and equals } L.$$

Similarly, let f be a function such that there is an N such that it is defined for all $x < N$.


Suppose for all sequences x_n diverging to $-\infty$, there exists L such that $f(x_n)$ converges to L (i.e. as x becomes smaller and smaller, $f(x)$ eventually gets closer and closer to L). Then we say that

$$\lim_{x \rightarrow -\infty} f(x) \text{ exists and equals } L.$$

The function $\frac{1}{x}$




The function $\frac{1}{x}$



$$\lim_{x \rightarrow \infty} \frac{1}{x} = \lim_{x \rightarrow -\infty} \frac{1}{x} = 0.$$

The function $\frac{1}{x}$



$\lim_{x \rightarrow \infty} \frac{1}{x} = \lim_{x \rightarrow -\infty} \frac{1}{x} = 0$. Both $\lim_{x \rightarrow 0^-} \frac{1}{x}$ and $\lim_{x \rightarrow 0^+} \frac{1}{x}$ do not exist.

Some basic examples

$$1. \lim_{x \rightarrow a} x^k; k \geq 0$$

$$= a^k$$

$$3. \lim_{x \rightarrow a} e^x$$

$$= e^a$$

$$5. \lim_{x \rightarrow a} \sin(x)$$

$$= \sin(a)$$

$$2. \lim_{x \rightarrow a} x^k; k < 0, a \neq 0$$

$$= a^k$$


$$4. \lim_{x \rightarrow a} \log_e(x); a > 0$$

$$= \log_e(a)$$

$$6. \lim_{x \rightarrow a} \tan(x); a \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$= \tan(a)$$

Finding limits by substitution : beware




Finding limits by substitution : beware

Suppose we want to find the value of the limit of a function $f(x)$ at the point a i.e. $\lim_{x \rightarrow a} f(x)$.



Finding limits by substitution : beware

Suppose we want to find the value of the limit of a function $f(x)$ at the point a i.e. $\lim_{x \rightarrow a} f(x)$. Often we can **substitute** the value of a in the expression for $f(x)$ and obtain the limit.



Finding limits by substitution : beware

Suppose we want to find the value of the limit of a function $f(x)$ at the point a i.e. $\lim_{x \rightarrow a} f(x)$. Often we can **substitute** the value of a in the expression for $f(x)$ and obtain the limit.

Unfortunately, this does not work when the function gets slightly complicated or the point a does not belong to the domain of definition of $f(x)$.

Finding limits by substitution : beware

Suppose we want to find the value of the limit of a function $f(x)$ at the point a i.e. $\lim_{x \rightarrow a} f(x)$. Often we can **substitute** the value of a in the expression for $f(x)$ and obtain the limit.

Unfortunately, this does not work when the function gets slightly complicated or the point a does not belong to the domain of definition of $f(x)$.

Examples :

$$\lim_{x \rightarrow 2} \frac{x^2 - 5x + 6}{x - 2}.$$

Finding limits by substitution : beware

Suppose we want to find the value of the limit of a function $f(x)$ at the point a i.e. $\lim_{x \rightarrow a} f(x)$. Often we can **substitute** the value of a in the expression for $f(x)$ and obtain the limit.

Unfortunately, this does not work when the function gets slightly complicated or the point a does not belong to the domain of definition of $f(x)$.

Examples :

$$\lim_{x \rightarrow 2} \frac{x^2 - 5x + 6}{x - 2}$$

$$\begin{aligned} &= \frac{(x-2)(x-3)}{(x-2)} \\ &= x-3. \end{aligned}$$

$$\frac{2^2 - 5 \times 2 + 6}{2 - 2} = \frac{0}{0}$$


$$\lim_{x \rightarrow 0} \frac{1}{x}$$

DNE

cannot substitute

Some known limits


$$1. \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1.$$



Some known limits


$$1. \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1.$$

$$2. \lim_{x \rightarrow 0} \frac{1 + \log_e(x)}{x} = 1.$$



Some known limits

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1.$$



$$\frac{\sin(1^\circ)}{1^\circ} \leq 1.$$

$$2. \lim_{x \rightarrow 0} \frac{1 + \log_e(x)}{x}$$



$$3. \lim_{x \rightarrow \infty} \frac{a + be^x}{c + de^x} = \frac{b}{d}.$$




$$\begin{aligned} &= \lim_{x \rightarrow \infty} \frac{a + be^x}{c + de^x} \\ &\quad \cdot \frac{e^{-x} + d^{-x}}{e^{-x} + d^{-x}} \\ &= \frac{a \cdot 0 + b}{c \cdot 0 + d} = b/d. \end{aligned}$$

Continuity of a function at a point



Continuity of a function at a point

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n and a belong to the domain of f .



Continuity of a function at a point

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n and a belong to the domain of f .

Then the function f is said to be continuous at the point a if
 $\lim_{x \rightarrow a} f(x) = f(a)$.



Continuity of a function at a point

Let f be a function and a be a point such that $a_n \rightarrow a$ where a_n and a belong to the domain of f .

Then the function f is said to be continuous at the point a if
 $\lim_{x \rightarrow a} f(x) = f(a)$

i.e. continuity means "the limit at a can be obtained by evaluating the function at a ".

$$f(x) = \begin{cases} \frac{\sin(x)}{x} & x \neq 0 \\ 1 & x = 0 \end{cases}$$

Thank you

