


Critical points : local maxima and minima




Local maxima/minima



Local maxima/minima


Recall the following notions from maths 1 for a function $f(x)$:



Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:


An **interval of increase (respectively decrease)** is an interval on which the function $f(x)$ is increasing (respectively decreasing)



Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:

An **interval of increase (respectively decrease)** is an interval on which the function $f(x)$ is increasing (respectively decreasing) i.e. if $x_1 < x_2$ are points belonging to that interval, then $f(x_1) < f(x_2)$ (respectively $f(x_1) > f(x_2)$).



Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:

An **interval of increase** (respectively decrease) is an interval on which the function $f(x)$ is increasing (respectively decreasing) i.e. if $x_1 < x_2$ are points belonging to that interval, then $f(x_1) < f(x_2)$ (respectively $f(x_1) > f(x_2)$).

A point x is a **turning point** of $f(x)$ where one of the following phenomena take place :

Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:

An **interval of increase (respectively decrease)** is an interval on which the function $f(x)$ is increasing (respectively decreasing) i.e. if $x_1 < x_2$ are points belonging to that interval, then $f(x_1) < f(x_2)$ (respectively $f(x_1) > f(x_2)$).

A point x is a **turning point** of $f(x)$ where one of the following phenomena take place :

1. either there is an interval of increase ending at x and an interval of decrease beginning at x

Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:

An **interval of increase (respectively decrease)** is an interval on which the function $f(x)$ is increasing (respectively decreasing) i.e. if $x_1 < x_2$ are points belonging to that interval, then $f(x_1) < f(x_2)$ (respectively $f(x_1) > f(x_2)$).

A point x is a **turning point** of $f(x)$ where one of the following phenomena take place :

1. either there is an interval of increase ending at x and an interval of decrease beginning at x
2. or there is an interval of decrease ending at x and an interval of increase beginning at x .

Local maxima/minima

Recall the following notions from maths 1 for a function $f(x)$:

An **interval of increase (respectively decrease)** is an interval on which the function $f(x)$ is increasing (respectively decreasing) i.e. if $x_1 < x_2$ are points belonging to that interval, then $f(x_1) < f(x_2)$ (respectively $f(x_1) > f(x_2)$).

A point x is a **turning point** of $f(x)$ where one of the following phenomena take place :

1. either there is an interval of increase ending at x and an interval of decrease beginning at x
2. or there is an interval of decrease ending at x and an interval of increase beginning at x .

A turning point as in 1 above is called a **local maximum** and a turning point as in 2 above is called a **local minimum**.

Example




Figure: $\Gamma((x^2 - 4x + 3.8)(x + 2)x^3)$

Example





Figure: $\Gamma (x \cos(x))$

Tangents at local maxima/minima



Tangents at local maxima/minima

Observe that in both examples, tangents at the turning points are horizontal (i.e. flat or parallel to the X-axis).



Tangents at local maxima/minima

Observe that in both examples, tangents at the turning points are horizontal (i.e. flat or parallel to the X -axis).


This is a general phenomenon that is easy to visualize.




Tangents at local maxima/minima

Observe that in both examples, tangents at the turning points are horizontal (i.e. flat or parallel to the X -axis).

This is a general phenomenon that is easy to visualize.




Local maxima/minima and derivatives



Local maxima/minima and derivatives


How do we prove that the tangent (**if it exists**) is horizontal at a turning point?



Local maxima/minima and derivatives

How do we prove that the tangent (**if it exists**) is horizontal at a turning point?

Recall that the tangent to f at a exists (and is not vertical) is equivalent to f being differentiable at a .




Local maxima/minima and derivatives

How do we prove that the tangent (**if it exists**) is horizontal at a turning point?

Recall that the tangent to f at a exists (and is not vertical) is equivalent to f being differentiable at a . and its equation is then given by

$$y = f'(a)(x - a) + f(a).$$



Local maxima/minima and derivatives

How do we prove that the tangent (if it exists) is horizontal at a turning point?

Recall that the tangent to f at a exists (and is not vertical) is equivalent to f being differentiable at a . and its equation is then given by

$$y = f'(a)(x - a) + f(a).$$

Thus, proving that the tangent (if it exists) is horizontal at a turning point is equivalent to showing that $f'(a) = 0$.

a is a turning point & $f'(a)$ exists.


Suppose a is a local maximum. For some $(a - \epsilon, a + \epsilon)$

if $0 < h < \epsilon$ then $f(a-h) < f(a)$ & $f(a+h) < f(a)$.

$$\begin{aligned}f'(a) &= \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h} \leq 0 \quad \Rightarrow f'(a) = 0. \\&= \lim_{h \rightarrow 0^-} \frac{f(a+h) - f(a)}{h} \geq 0\end{aligned}$$

Critical points and saddle points


Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.



Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.




Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.



Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.

Thus, in order to find turning points (i.e. local maxima/minima), we will first find critical points.

Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.

Thus, in order to find turning points (i.e. local maxima/minima), we will first find critical points.

Suppose f is differentiable.

Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.

Thus, in order to find turning points (i.e. local maxima/minima), we will first find critical points.

Suppose f is differentiable. Is every critical point a turning point?

Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.

Thus, in order to find turning points (i.e. local maxima/minima), we will first find critical points.

Suppose f is differentiable. Is every critical point a turning point?

Unfortunately not e.g. 0 for $f(x) = (x^2 - 4x + 3.8)(x + 2)x^3$.

Critical points and saddle points

Thus, if f is differentiable at a turning point a , it satisfies $f'(a) = 0$.

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

Every turning point is a critical point.

Thus, in order to find turning points (i.e. local maxima/minima), we will first find critical points.

Suppose f is differentiable. Is every critical point a turning point?

Unfortunately not e.g. 0 for $f(x) = (x^2 - 4x + 3.8)(x + 2)x^3$.

A **saddle point** is a critical point which is not a local maximum or local minimum.

The second derivative test

Suppose f is differentiable.



The second derivative test


Suppose f is differentiable. How do we classify critical points?



The second derivative test

Suppose f is differentiable. How do we classify critical points?

Just like the first derivative f' checks for the monotonicity of f ,
the second derivative f'' checks for the monotonicity of f' .




The second derivative test

Suppose f is differentiable. How do we classify critical points?

Just like the first derivative f' checks for the monotonicity of f ,
the second derivative f'' checks for the monotonicity of f' .

So if f is twice differentiable, we check f'' at all the critical points.



The second derivative test

Suppose f is differentiable. How do we classify critical points?

Just like the first derivative f' checks for the monotonicity of f , the second derivative f'' checks for the monotonicity of f' .

So if f is twice differentiable, we check f'' at all the critical points.

1. If a is a critical point and $f'(a) > 0$, then a is a local minimum.

The second derivative test

Suppose f is differentiable. How do we classify critical points?

Just like the first derivative f' checks for the monotonicity of f , the second derivative f'' checks for the monotonicity of f' .

So if f is twice differentiable, we check f'' at all the critical points.

1. If a is a critical point and $f'(a) > 0$, then a is a local minimum.
2. If a is a critical point and $f'(a) < 0$, then a is a local maximum.

The second derivative test


Suppose f is differentiable. How do we classify critical points?

Just like the first derivative f' checks for the monotonicity of f , the second derivative f'' checks for the monotonicity of f' .

So if f is twice differentiable, we check f'' at all the critical points.

1. If a is a critical point and $f'(a) > 0$, then a is a local minimum.
2. If a is a critical point and $f'(a) < 0$, then a is a local maximum.
3. If a is a critical point and $f'(a) = 0$, then the test is **inconclusive**.

Examples



Examples

$$f(x) = x^3 - 12x$$

$$f'(x) = 3x^2 - 12 \quad \text{Setting it to 0,}$$

we obtain $3x^2 - 12 = 0 \Rightarrow x^2 = 4 \Rightarrow x = \pm 2.$

$$f''(x) = 6x \quad f''(-2) = -12 < 0$$

Critical points: ± 2
 $f''(2) = 12 > 0$
 $\therefore 2$ is a local minimum & -2 is a local maximum

$$f(x) = \cos(x)$$

$$f'(x) = -\sin(x).$$

$$f''(x) = -\cos(x).$$

$$f''(k\pi) = -\cos(k\pi) = \begin{cases} -1 & \text{if } k \text{ is an even integer} \\ 1 & \text{if } k \text{ is an odd integer} \end{cases}$$

$$f(x) = x^3 + x^2 - x + 5$$

$k\pi$; k is even are local maxima
 $k\pi$; k is odd are local minima.

$$f'(x) = 3x^2 + 2x - 1 \quad \text{Setting it to 0,}$$

we obtain $3x^2 + 2x - 1 = 0 \Rightarrow \text{Roots are } \frac{-2 \pm \sqrt{4+12}}{2 \times 3}$

Roots: $-1, \frac{1}{3} = \frac{-2 \pm \sqrt{16}}{6}$


$$f''(x) = 6x + 2.$$

$f''(-1) = -6 + 2 = -4$: local maximum

$f''(\frac{1}{3}) = 2 + 2 = 4$: local minimum

Local maxima/minima on closed intervals


Sometimes we want to find the local extrema of a function f on a closed interval $I = [a, b]$.



Local maxima/minima on closed intervals

Sometimes we want to find the local extrema of a function f on a closed interval $I = [a, b]$.

In that case, in addition to finding the extrema via the previously discussed method, it is possible that the end points could also be local extrema and so we have to consider them separately.



Local maxima/minima on closed intervals

Sometimes we want to find the local extrema of a function f on a closed interval $I = [a, b]$.

In that case, in addition to finding the extrema via the previously discussed method, it is possible that the end points could also be local extrema and so we have to consider them separately.

Example : $f(x) = x^2$ on the interval $[-1, 1]$.


$$f'(x) = 2x.$$

Set it to 0.

$$f''(x) = 2. \quad f''(0) = 2.$$


\therefore It is a local minimum.

-1 & 1 are also local maxima.



(Global) maximum/minimum


Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .



(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.



(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.

Fact : If the interval I is closed and bounded and f is continuous, the maximum and minimum must exist.

(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.

Fact : If the interval I is closed and bounded and f is continuous, the maximum and minimum must exist.

Note that the maximum and minimum are in particular local maxima or local minima

(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.

Fact : If the interval I is closed and bounded and f is continuous, the maximum and minimum must exist.

Note that the maximum and minimum are in particular local maxima or local minima unless they are on **boundary points**.

(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.

Fact : If the interval I is closed and bounded and f is continuous, the maximum and minimum must exist.

Note that the maximum and minimum are in particular local maxima or local minima unless they are on **boundary points**.

Thus to find the maximum and minimum, we find the critical points and the boundary points and check the value of f on all of them.

(Global) maximum/minimum

Sometimes we want to find the maximum or minimum value of a function f on a particular interval I .

In general this may not exist e.g. $f(x) = \frac{1}{x}$ on $I = (0, \infty)$.

Fact : If the interval I is closed and bounded and f is continuous, the maximum and minimum must exist.

Note that the maximum and minimum are in particular local maxima or local minima unless they are on **boundary points**.

Thus to find the maximum and minimum, we find the critical points and the boundary points and check the value of f on all of them.

We can in fact do this on any function which is defined **piecewise** continuously with finitely many pieces on a closed and bounded interval.

Example

$$f(x) = \begin{cases} x^3 + x^2 - x + 5 & \text{if } 0 \leq x \leq 100 \\ x^3 + 2x^2 + x - 5 & \text{if } -100 \leq x < 0 \end{cases}$$

$$f'(x) = \begin{cases} 3x^2 + 2x - 1 & \text{if } 0 < x < 100 \\ 3x^2 + 4x + 1 & \text{if } -100 \leq x < 0 \end{cases}$$

$$\therefore \frac{-2 \pm \sqrt{4+12}}{6}, \quad \frac{-2 \pm \sqrt{16-12}}{6}$$

Critical Pts.
 $\therefore -100, 0, 100.$

Bdry Pts.
 $\therefore \frac{-2 \pm 4}{6} = \left\{ \begin{array}{l} 1/3 \\ -1 \end{array} \right\}, \quad \frac{-2 \pm 2}{6} = \left\{ \begin{array}{l} 0 \\ -2/3 \end{array} \right\}$

$$f''(x) = \begin{cases} 6x + 2 & \text{loc. min.} \\ 6x + 4 & f''(-1/3) = 4 \\ & f''(0) = 4 \end{cases}$$

$$f''(-1) = -4 \quad \text{local max.}$$

$$f''(-2/3) = 0$$

$$f(-1) = -1 + 2 + (-1) - 5 = -5$$


$$f(0) = 5 \quad f(-2/3) = -\frac{8}{27} + \frac{8}{9} + \frac{(-2)}{3} - 5$$

$$f(1/3) =$$

Glob. max. $f(100) = 100^3 + 100^2 - 100 + 5$

Glob. min. $f(-100) = (-100)^3 + 2 \times 100^2 - 100 - 5$

Warning example : $f(x) = x^4$



$$f'(x) = 4x^3 \Rightarrow \text{critical point is } 0.$$

$$f''(x) = 12x^2 \Rightarrow f''(0) = 0. \text{ Inconclusive.}$$

Thank you

