NEERWATCH

Krishna Raja Sagara

Generated 2026-10-22 · satellite monitoring report

Chlorophyll-a Concentration Classification — US EPA Reference

Satellite-derived screening estimate; trophic categories are not water-safety or toxicity determinations.

Observation	Result
Earlier comparable acquisition	2026-10-09T05:22:41.024000Z: 25.15 µg/L
Latest processed acquisition	2026-10-12T05:17:39.024000Z: 24.72 µg/L
EPA trophic category	Eutrophic
Concentration change	-1.71%
Trend	Decreasing

EPA trophic reference: ≤2 oligotrophic; >2–7 mesotrophic; >7–30 eutrophic; >30 hypereutrophic (µg/L).
Source: US EPA National Lakes Assessment. https://nationallakesassessment.epa.gov/webreport/#trophic-state


Method and limitations

Satellite-derived chlorophyll-a estimates have not yet been validated against Karnataka field measurements.


Gilerson red/red-edge equation applied to Sentinel-2 L2A Sen2Cor surface reflectance as an experimental approximation to water-leaving reflectance. ACOLITE aquatic atmospheric correction has not been performed. Values outside the project screening range of 1–100 µg/L are excluded. No field accuracy score is claimed.

Acquisition: 2026-10-12T05:17:39.024000Z
Scene: S2B_MSIL2A_20261012T051739_R062_T43PFP_20261012T091927
Method: Gilerson 2010 two-band; Sentinel-2 L2A BOA screening adaptation v1

True-colour image



Chlorophyll-a map from the same acquisition



Dated Sentinel-2 estimates

2026-10-09T05:22:41.024000Z — 25.15 µg/L
2026-10-12T05:17:39.024000Z — 24.72 µg/L