



Semester 1 Assessment, 2021

School of Mathematics and Statistics

MAST30025 Linear Statistical Models Assignment 3

Submission deadline: **Friday May 28, 5pm**

This assignment consists of 3 pages (including this page)

Instructions to Students

Writing

- There are 5 questions with marks as shown. The total number of marks available is 48.
- This assignment is worth 7% of your total mark.
- You may choose to either typeset your assignment in L^AT_EX or handwrite and scan it to produce an electronic version.
- You may use R for this assignment, including the `lm` function unless specified. If you do, include your R commands and output.
- Write your answers on A4 paper. Page 1 should only have your student number, the subject code and the subject name. Write on one side of each sheet only. Each question should be on a new page. The question number must be written at the top of the page.

Scanning

- Put the pages in question order and all the same way up. Use a scanning app to scan all pages to PDF. Scan directly from above. Crop pages to A4. Check PDF is readable.

Submitting

- Go to the Gradescope window. Choose the Canvas assignment for this assignment. Submit your file as a single PDF document only. Get Gradescope confirmation on email.
- It is your responsibility to ensure that your assignments are submitted correctly and on time, and problems with online submissions are not a valid excuse for submitting a late or incorrect version of an assignment.

Question 1 (7 marks)

Let A be an $n \times p$ matrix with $n \geq p$.

- (a) Show directly that $r(A^c A) = r(A)$.
- (b) Show directly that $A^c A$ is idempotent.
- (c) Show directly that $A(A^T A)^c A^T$ is unique (invariant to the choice of conditional inverse).

Question 2 (11 marks)

We are interested in examining the lifetime of three different types of bulb (in 1000 hours). A study is conducted and the following data obtained:

| Bulb type | | |
|-----------|----|----|
| 1 | 2 | 3 |
| 22 | 16 | 28 |
| 23 | 18 | 27 |
| 24 | 19 | 29 |
| 22 | | 29 |
| 26 | | |

We fit the model

$$y_{ij} = \mu + \tau_i + \varepsilon_{ij},$$

where μ is the overall mean and τ_i is the effect of the i th type of bulb.

For this question, you may NOT use the `lm` function in R.

- (a) Find a conditional inverse for $X^T X$, using the algorithm given in Theorem 6.2.
- (b) Find s^2 .
- (c) Is $\mu + 2\tau_1 + \tau_2$ estimable?
- (d) Find a 90% confidence interval for the lifetime of the 2nd type of bulb.
- (e) Test the hypothesis that there is no difference in lifetime between 1st and 3rd types of bulb, at the 5% significance level.

Question 3 (5 marks)

Consider a linear model with only categorical predictors, written in matrix form as $\mathbf{y} = X_1 \boldsymbol{\beta}_1 + \boldsymbol{\varepsilon}_1$. Suppose we add some continuous predictors, resulting in an expanded model $\mathbf{y} = X \boldsymbol{\beta} + \boldsymbol{\varepsilon}$. Now consider a quantity $\mathbf{t}^T \boldsymbol{\beta}$, where $\mathbf{t}^T = [\mathbf{t}_1^T | \mathbf{t}_2^T]$ is partitioned according to the categorical and continuous predictors. Show that if $\mathbf{t}_1^T \boldsymbol{\beta}_1$ is estimable in the first model, then $\mathbf{t}^T \boldsymbol{\beta}$ is estimable in the second model.

If you write $X = [X_1 | X_2]$, you may assume that $r(X) = r(X_1) + r(X_2)$.

Hint: Use Theorems 6.9 and 6.3. For any vectors \mathbf{z}_1 and \mathbf{z}_2 , you can write

$$\left[\begin{array}{cc|c} X_1^T X_1 & X_1^T X_2 & X_1^T X_1 \mathbf{z}_1 \\ X_2^T X_1 & X_2^T X_2 & X_2^T X_2 \mathbf{z}_2 \end{array} \right] = \left[\begin{array}{cc} X_1^T & \mathbf{0} \\ \mathbf{0} & X_2^T \end{array} \right] \left[\begin{array}{cc|c} X_1 & X_2 & X_1 \mathbf{z}_1 \\ X_1 & X_2 & X_2 \mathbf{z}_2 \end{array} \right].$$

Question 4 (18 marks)

Data was collected on the world record times (in seconds) for the one-mile run. For males, the records are from the period 1861–1999, and for females, from the period 1967–1996. The data is given in the file `mile.csv`, available on the LMS.

- (a) Plot the data, using different colours and/or symbols for male and female records. Without drawing diagnostic plots, do you think that this data satisfies the assumptions of the linear model? Why or why not?
- (b) Test the hypothesis that there is no interaction between the two predictor variables. Interpret the result in the context of the study.
- (c) Write down the final fitted models for the male and female records. Add lines corresponding to these models to your plot from part (a).
- (d) Calculate a point estimate for the year when the female world record will equal the male world record. Do you expect this estimate to be accurate? Why or why not?
- (e) Is the year when the female world record will equal the male world record an estimable quantity? Is your answer consistent with part (d)?
- (f) Calculate a 95% confidence interval for the amount by which the gap between the male and female world records narrow every year.
- (g) Test the hypothesis that the male world record decreases by 0.4 seconds each year.

Question 5 (7 marks)

You wish to perform a study to compare 2 medical treatments (and a placebo) for a disease. Treatment 1 is an experimental new treatment, and costs \$5000 per person. Treatment 2 is a standard treatment, and costs \$2000 per person. Treatment 3 is a placebo, and costs \$1000 per person. You are given \$100,000 to complete the study. You wish to test if the treatments are effective, i.e., $H_0 : \tau_1 = \tau_2 = \tau_3$.

- (a) Determine the optimal allocation of the number of units to assign to each treatment.
- (b) Perform the random allocation. You must use R for randomisation and include your R commands and output.

End of Assignment — Total Available Marks = 48