




THE CANADIAN CHEMISTRY CONTEST 2020
PART A – MULTIPLE CHOICE QUESTIONS (60 minutes)

All contestants should attempt this part of the contest before proceeding to Part B and/or Part C.



The only reference material allowed is the CIC/CCO Periodic Table provided. You must complete answers on the Scantron Sheet provided. Students may use a scientific calculator. No phones or communication devices are allowed.

- 1) 15 mL of 0.25 mol L⁻¹ aqueous silver nitrate reacts with 0.10 g of zinc as follows:




What mass of silver metal should the reaction produce?

- A) 0.16 g B) 0.20 g **C) 0.33 g** D) 0.40 g E) 0.81 g
- 2) Based on VSEPR theory, which of the following has a seesaw molecular geometry?
- A) H₂O B) CH₄ C) XeF₄ D) BF₃ **E) SeF₄**
- 3) A student combines 100 mL of two clear, equimolar, colourless solutions and observes a solid white precipitate form at the bottom of the beaker. A representation of the resultant particles in the beaker is in the diagram below. Which combination of reactants would best align with the information provided and diagram of resultant particles in the beaker?
- A) lead (II) nitrate (aq) + potassium chloride (aq)
B) zinc nitrate (aq) + sodium phosphate (aq)
C) nickel (II) nitrate (aq) + lithium bromide (aq)
D) copper (II) nitrate (aq) + sodium hydroxide (aq)
E) silver nitrate (aq) + magnesium sulfate (aq)




- 4) A student drew structural diagrams for some of the isomers of octane and wrote down each isomer's boiling point (b.p.). There is an error in the student's work. For which structure did the student **incorrectly** record the boiling point?




- 5) In which of the following substances does sulfur have the lowest oxidation state?
- A) S₈ B) Na₂S₂O₃ C) Na₂SO₄ **D) H₂S** E) SO₂
- 6) Given the graph of the successive ionization energies for two unknown atoms, which combination provides the most likely identity of X and Y?

Unknown:	X	Y
A)	Al	Si
B)	Kr	Rb
C)	Cl	Ar
D)	Se	Br
E)	Rb	Sr



- 7) Bufotenin (structure below) is a substance found in the skin of certain species of toads. In large quantities, it can have a psychoactive effect on humans, changing brain function and altering behaviour. What is the correct molecular formula of bufotenin?



- A) **C₁₂H₁₆N₂O** B) C₁₀H₁₂N₂O C) C₁₁H₁₄N₂O
 C) C₁₂H₁₈NO E) C₁₂H₁₄N₂O
- 8) How many constitutional isomers and how many stereoisomers does the organic molecule with the formula C₃H₉N have?

	constitutional isomers	stereoisomers		constitutional isomers	stereoisomers
A)	2	2	D)	4	0
B)	3	2	E)	3	0
C)	4	2			


- 9) Consider the following equilibrium in a sealed container



Which of the following changes would most likely shift the equilibrium position towards reactants?

- A) **Increasing the temperature** D) Adding nitrogen monoxide gas
 B) Adding argon gas E) Removing ammonia
 C) Decreasing the volume of the sealed container
- 10) When a sample of an unknown compound with a formula of C_xH_y combusts with an excess amount of oxygen, the reaction produces 132.001 g of carbon dioxide and 72.064 g of water. Based on the information given, this unknown compound is:
- A) methane B) ethane **C) propane** D) butane E) octane

- 11) Hydrogen peroxide decomposes to form oxygen gas and water via the following chemical equation:



The density of 30% hydrogen peroxide is 1.11 g mL⁻¹. If 50.0 mL of a 30% by mass hydrogen peroxide solution decomposes, what is the accompanying change in energy?


- A) **96 kJ released** B) 117 kJ absorbed C) 173 kJ released
 D) 192 kJ released E) 392 kJ absorbed
- 12) A mixture of ethanol and nitric acid, called nital, is an industrial etching agent. A student prepared a 20.0 mL solution of nital using 0.70 mL of nitric acid of unknown concentration and 19.3 mL of 98% ethanol. The student determined by titration that the final concentration of nitric acid in the nital etching agent was 4.0 % by mass. The density of 98% ethanol was 0.79 g mL⁻¹ and the density of the unknown concentration of aqueous HNO₃ was 1.4 g mL⁻¹. What was the **original** concentration of HNO₃, in w/w, in the 0.70 mL reagent the student used for the nital solution preparation?
- A) 63% **B) 66%** C) 70% D) 73% E) 93%

- 13) A student placed 3 identical metal cubes into an oven and heated them to the same final temperature. The student entirely submerged each of the 3 cubes in separate beakers as shown in the diagram below. All solutions had approximately the same density and all were at room temperature before adding the metal cubes. The maximum temperature compared as follows:


$$T_{\text{max}} \text{ beaker B} > T_{\text{max}} \text{ beaker A} > T_{\text{max}} \text{ beaker C.}$$

Which of the following is true about the relationship of the specific heat capacities of solutions A, B and C?


- A) C_A > C_C > C_B
 B) C_C > C_B > C_A
C) C_C > C_A > C_B
 D) C_B > C_A > C_C
 E) C_B > C_C > C_A



- 14) The organic compound orthocaine is a local anesthetic developed over one hundred years ago to treat pain. The chemical name of orthocaine is methyl 3-amino-4-hydroxybenzoate. Which of the following structures represents a molecule of orthocaine?



- 15) Given the following standard reduction potentials.



Which of the following statement(s) is/are true?

- I. Na is the most easily oxidized
- II. Cu is the best reducing agent.
- III. Cu^{2+} cannot be spontaneously reduced by Al
- IV. Na^{+} has a greater attraction for electrons than Al^{3+}

- A) I only B) I and II only C) I, II and III only
D) II and IV only E) II, III, and IV only

- 16) Which of the following should have the same electron arrangement around the central atom as BF_4^{-} ?

- A) IF_4^{-} B) XeCl_4 C) ClF_4^{+} D) SF_4 E) CCl_4





- 17) A student adds two moles of liquid water initially at 273 K to three moles of liquid water at 363 K in a perfectly insulated container. The total volume of water remains constant. Assume that the molar heat capacity of liquid water is constant and independent of temperature. What is the final equilibrium temperature of the water?

- A) 298 K B) 309 K C) 318 K D) 327 K E) 358 K


- 18) One mole of uranium-238 decays slowly. If uranium-238 decays according to first order kinetics, after how many half-lives is there likely only one atom of uranium-238 left?

- A) 85 B) 10 C) 238 D) 41 E) 79

- 19) Consider the structures and associated pKa values in the table below. Which statement below the table best explains the pKa values observed?

			
acetic acid	chloroacetic acid	dichloroacetic acid	trichloroacetic acid
pKa = 4.76	pKa = 2.85	pKa = 1.35	pKa = 0.66

- A) Electron density is removed from the ionizable proton by induction, which increases the strength of the acid
B) The presence of electron-donating alkyl groups increases the stability of the conjugate base, increasing the strength of the acid
C) Resonance stabilization of the conjugate base increases the strength of the acid
D) The presence of electronegative atoms decreases the stability of the conjugate base, decreasing the strength of the acid
E) Resonance stabilization of the conjugate base decreases the strength of the acid




Using the equilibria above, determine which of the following combinations of solutions will generate a buffer with a pH of approximately 5?

- A) 50. mL of 0.1 mol L⁻¹ Na₂HPO₄ + 100. mL 0.1 mol L⁻¹ NaH₂PO₄
 B) 100. mL of 0.1 mol L⁻¹ NaCH₃COO + 100. mL of 0.1 mol L⁻¹ NaOH
 C) 100. mL of 0.1 mol L⁻¹ NaH₂PO₄ + 100. mL of 0.1 mol L⁻¹ HCl
 D) 100. mL of 0.1 mol L⁻¹ NaH₂PO₄ + 100. mL of 0.1 mol L⁻¹ NaOH
 E) 100. mL of 0.1 mol L⁻¹ CH₃COOH + 50. mL of 0.1 mol L⁻¹ NaOH

- 21) The *reaction mass efficiency* (RME) helps a student determine how much reactant material a product contains at the end of a chemical reaction. RME is calculated as follows:

$$\text{reaction mass efficiency} = \frac{\text{mass of desired product}}{(\text{total input mass} - \text{mass of recycled material})} \times 100\%$$

0.205 moles of phenylboronic acid (C₆H₇BO₂) was reacted with 0.205 moles of 4-iodophenol (C₆H₅IO) and 0.615 moles of potassium carbonate to form 0.185 moles of 4-phenylphenol (shown below). 300 mg of catalytic palladium on carbon was used and completely recycled.




- A) 18.4 % B) 20.3 % C) 24.2 % D) 25.5 % E) 35.1 %

- 22) The crossover temperature for a reaction is the temperature at which a chemical reaction changes from occurring spontaneously to being non-spontaneous. For a particular chemical reaction, the entropy change is -157 J mol⁻¹·K⁻¹ and the enthalpy change is -82 kJ mol⁻¹. What is the crossover temperature for the reaction?

- A) 0 K B) 75K C) 239 K D) 522K E) There is no crossover temperature for this reaction


- 23) One source of oxygen in chemical oxygen generators for emergencies in submarines and aircraft is potassium perchlorate, which decomposes as follows:




Assume ideal gas behaviour for the reaction, a temperature of 25.0°C and a pressure of 1.00 bar. If 138.5 grams of potassium perchlorate decompose inside an oxygen generator, what volume of oxygen does the reaction generate if there is a 78.0% yield?

- A) 9.65 L B) 19.3 L C) 24.8 L D) 38.6 L E) 49.6 L

- 24) Metal sulfides have considerable applications in industry, including their use as components of semiconducting materials. Consider the following equilibrium, where M represents a transition metal:




S²⁻ ions liberated in solution readily react and form HS⁻ and OH⁻ according to the following equilibrium:



Which of the following will increase the solubility of the metal sulfide?

- A) Increasing the concentration of OH⁻ by adding a strong base
 B) Adding a strong acid
 C) Increasing the amount of the metal sulfide present
 D) Diluting the solution present
 E) Adding MCl₂

- 25) The ΔH_f° of liquid water is -285.8 kJ mol⁻¹. The enthalpy of combustion of 1 mol of acetic acid (CH₃COOH) is -874.1 kJ mol⁻¹. Given the reaction:



- A) -2232.7 B) -1553.4 C) -981.8 D) -484.5 E) -194.8

End of Part A of the contest
Go back and check your work