Sets and functions

Subject: Mathematics | Level: Class 9 | Curriculum: Sindh Board
Prepared by: SHS Virtual Academy

1. Topic Overview and Learning Objectives

Sets and functions form the backbone of modern mathematics and are essential topics in the Sindh Board Class 9 curriculum. A set is a well-defined collection of distinct objects, and understanding how these collections interact through operations like union, intersection, and difference is crucial for logical reasoning. These concepts provide the language used to describe relationships between different groups of data, which is a skill you will use throughout your academic career.

In board examinations, questions on sets often carry significant weight, frequently appearing in both the objective and subjective sections of the paper. Examiners look for your ability to represent sets using roster, set-builder, and descriptive forms, as well as your proficiency in constructing Venn diagrams. Mastering these basics is not just about passing the exam; it is about building the mathematical maturity required for higher-level algebra and calculus in later years.

The concept of a function, which is a special type of relation, is equally important. It helps us understand how one variable depends on another, a principle that is vital in physics, economics, and engineering. By learning how to identify domains, ranges, and the nature of mappings, you are developing the analytical skills necessary to solve complex problems. Focus on the definitions and the logical flow of these operations, and you will find that these topics are both manageable and rewarding.

2. Core Concepts and Detailed Revision Notes

2.1 Introduction to Sets and Representations

A set is defined as a collection of well-defined and distinct objects. In mathematics, we use capital letters like A, B, or C to denote sets, while the elements inside are usually represented by small letters. It is vital to understand that the order of elements does not matter, and each element must be unique. There are three primary ways to represent a set: the descriptive form, which uses a statement; the tabular or roster form, which lists elements within curly brackets; and the set-builder form, which uses a rule or property to define the members. For example, the set of natural numbers less than five can be written as {1, 2, 3, 4}. Understanding these representations is the first step in solving any set-related problem in your exams, as you will often be asked to convert between these forms.

Key Points to Remember:

1. A set must contain well-defined and distinct elements.
2. The descriptive form uses a sentence to describe the set.
3. The tabular form lists elements separated by commas.
4. The set-builder form uses a variable and a defining property.
5. The empty set is denoted by curly brackets or the symbol phi.

2.2 Operations on Sets

Once you understand what a set is, the next step is to perform operations on them. The union of two sets, A and B, is the set of all elements that are in A, in B, or in both, denoted as A U B. The intersection, A n B, includes only the elements that are common to both sets. The difference of sets, A - B, consists of elements that are in A but not in B. These operations are best visualized using Venn diagrams, which are geometric representations that show the relationship between sets. When solving problems involving these operations, always ensure you identify the universal set first, as it defines the boundary for all elements under consideration. Practicing these operations will help you solve complex problems involving multiple sets and their complements efficiently during your board exams.

Key Points to Remember:

1. Union combines all elements from both sets without repetition.
2. Intersection selects only the shared elements.
3. Difference removes elements of the second set from the first.
4. Complement of a set A contains elements in the universal set not in A.
5. Venn diagrams are essential for visual proof and understanding.

2.3 Relations and Ordered Pairs

A relation is a subset of the Cartesian product of two sets. To understand this, we first define an ordered pair (x, y), where the order is significant. The Cartesian product A x B is the set of all possible ordered pairs where the first element is from A and the second is from B. A relation is simply any subset of this product. For instance, if set A represents students and set B represents their roll numbers, a relation could map each student to their specific number. In your exams, you might be asked to find the Cartesian product of two sets or identify whether a given set of ordered pairs constitutes a valid relation. Remember that a relation is a broader concept than a function, and every function is a relation, but not every relation is a function.

Key Points to Remember:

1. An ordered pair (x, y) is distinct from (y, x).

2. The Cartesian product A x B contains all possible pairs.
3. A relation is any subset of the Cartesian product.
4. Domain is the set of all first elements in the ordered pairs.
5. Range is the set of all second elements in the ordered pairs.

2.4 Understanding Functions

A function is a special type of relation where every element of the domain is mapped to exactly one element of the range. Think of it like a machine: you put one input in, and you get exactly one output out. If an input leads to two different outputs, it is not a function. In mathematics, we often write this as f: A -> B. To verify if a relation is a function, check the first elements of the ordered pairs; if any first element repeats with a different second element, it is not a function. Functions can be one-to-one, onto, or into, depending on how the elements map between sets. Mastering the concept of domain and range for functions will prepare you for more advanced topics like algebra and coordinate geometry in later classes.

Key Points to Remember:

1. A function maps each domain element to exactly one range element.
2. No two ordered pairs in a function can have the same first element.
3. The domain is the set of all valid inputs.
4. The range is the set of all resulting outputs.
5. Vertical line test can identify functions on a graph.

3. Key Definitions, Formulas, and Rules

1. **Union of sets:** A U B = {x | x is in A or x is in B}
2. **Intersection of sets:** A n B = {x | x is in A and x is in B}
3. **Difference of sets:** A - B = {x | x is in A and x is not in B}
4. **Complement of a set:** A' = U - A, where U is the universal set
5. **Cartesian Product:** A x B = {(a, b) | a is in A and b is in B}
6. **Empty Set:** A set with no elements, denoted by {} or phi
7. **Domain of a relation:** The set of all first components of the ordered pairs
8. **Range of a relation:** The set of all second components of the ordered pairs
9. **Function definition:** A relation where each input has exactly one output

4. Step-by-Step Worked Examples

Example 1: Finding Union and Intersection

Problem: Given A = {1, 2, 3} and B = {3, 4, 5}, find A U B and A n B.

Solution Steps:

Step 1: To find the union, list all elements from both sets without repeating the common element 3.

Step 2: A U B = {1, 2, 3, 4, 5}.

Step 3: To find the intersection, identify the element present in both sets.

Step 4: The only common element is 3.

Step 5: A n B = {3}.

Final Answer: A U B = {1, 2, 3, 4, 5} and A n B = {3}

Example 2: Calculating Cartesian Product

Problem: Find the Cartesian product A x B for A = {a, b} and B = {1, 2}.

Solution Steps:

Step 1: Pair each element of A with each element of B.

Step 2: Pair 'a' with 1 and 2: (a, 1), (a, 2).

Step 3: Pair 'b' with 1 and 2: (b, 1), (b, 2).

Step 4: Collect all pairs into a set.

Final Answer: A x B = {(a, 1), (a, 2), (b, 1), (b, 2)}

Example 3: Verifying a Function

Problem: Determine if the relation R = {(1, 2), (2, 3), (3, 2)} is a function.

Solution Steps:

Step 1: Check the first elements of each ordered pair: 1, 2, 3.

Step 2: Since all first elements are unique, each input maps to exactly one output.

Step 3: A relation where no first element repeats is a function.

Final Answer: Yes, it is a function because every domain element maps to a unique range element.

Example 4: Finding the Complement

Problem: If U = {1, 2, 3, 4, 5} and A = {1, 3, 5}, find A'.

Solution Steps:

Step 1: The complement A' consists of elements in U that are not in A.

Step 2: Remove {1, 3, 5} from {1, 2, 3, 4, 5}.

Step 3: The remaining elements are 2 and 4.

Final Answer: A' = {2, 4}

5. Common Exam Mistakes to Avoid

1. Forgetting to list elements only once in a set; remember that sets do not contain duplicates.
2. Confusing union (U) with intersection (n); remember U is like a cup that holds everything, while n is the overlap.
3. Assuming every relation is a function; always check if the first element repeats.
4. Mixing up the order in Cartesian products; (a, b) is not the same as (b, a).
5. Neglecting to define the universal set when calculating the complement; the complement is always relative to U.
6. Writing the domain or range as a single number instead of a set; always use curly brackets.

6. Practice Questions and Model Answers

Question 1

If A = {x, y, z} and B = {z, w}, find A - B.

Answer 1: A - B = {x, y} because z is in both sets and is removed.

Question 2

Is the relation R = {(1, 2), (1, 3), (2, 4)} a function? Why?

Answer 2: No, it is not a function because the input 1 maps to two different outputs, 2 and 3.

Question 3

Define the domain of the relation R = {(5, 10), (6, 12), (7, 14)}.

Answer 3: The domain is the set of first elements: {5, 6, 7}.

Question 4

If U = {a, b, c, d} and A = {a, b}, find the complement of A.

Answer 4: A' = {c, d}.

Question 5

Find the intersection of A = {2, 4, 6} and B = {6, 8, 10}.

Answer 5: A n B = {6}.