# Heat and thermodynamics

Subject: Physics | Level: Class 10 | Curriculum: Sindh Board
Prepared by: SHS Virtual Academy

## 1. Topic Overview and Learning Objectives

Heat and thermodynamics form the backbone of physical science, explaining how energy moves and transforms within our universe. In your Class 10 curriculum, understanding these concepts is vital because they bridge the gap between simple mechanical motion and the microscopic behavior of atoms and molecules. By grasping how heat flows from hotter to colder bodies, you are learning the very laws that govern everything from the steam engine to the cooling systems in your own homes.

For the Sindh Board examinations, this chapter is a high-yield area where examiners frequently test your conceptual clarity through both numerical problems and descriptive theory. You will often find questions regarding thermal expansion, specific heat capacity, and the gas laws. Mastering these topics not only guarantees marks in the theory section but also builds a strong foundation for the numerical problems that appear in Section B and Section C of your board paper.

The fundamental intuition here is to view heat as a form of energy in transit. When you add heat to a substance, you are essentially increasing the kinetic energy of its constituent particles, which manifests as a rise in temperature or a change in state. This chapter teaches you to quantify these changes using precise mathematical relationships. As you study, try to visualize the invisible dance of molecules, as this will make the laws of thermodynamics feel intuitive rather than just a set of formulas to memorize.

## 2. Core Concepts and Detailed Revision Notes

### 2.1 Temperature and Thermal Expansion

Temperature is a measure of the average kinetic energy of the molecules in a substance, while heat is the total energy transferred between systems. When materials are heated, their molecules vibrate more vigorously, causing the substance to expand. This phenomenon is known as thermal expansion. In solids, we observe linear expansion, where the length of a rod increases proportionally to the change in temperature. This is crucial in engineering, such as leaving gaps in railway tracks or bridges to accommodate expansion during the hot summer months in Sindh. Understanding the coefficient of linear expansion allows us to predict these changes accurately. It is important to distinguish between temperature, which is an intensive property, and heat, which is an extensive property, as this distinction is a common point of confusion during board exams.

**Key Points to Remember:**

1. Temperature measures average kinetic energy of particles.
2. Thermal expansion occurs due to increased molecular vibration.
3. Linear expansion depends on the original length and temperature change.
4. Coefficient of linear expansion is unique to each material.
5. Expansion gaps are necessary in construction to prevent structural damage.

### 2.2 Specific Heat Capacity and Latent Heat

Specific heat capacity is defined as the amount of heat energy required to raise the temperature of one kilogram of a substance by one degree Celsius. Substances with high specific heat, like water, take longer to heat up and cool down, which is why coastal areas have moderate climates. Conversely, latent heat refers to the energy absorbed or released during a phase change, such as melting ice into water or boiling water into steam, without any change in temperature. During these phase transitions, the energy is used to break or form intermolecular bonds rather than increasing the kinetic energy of the particles. Mastering these concepts is essential for solving calorimetry problems where you must calculate the final equilibrium temperature of mixed substances, a frequent favorite for board examiners.

**Key Points to Remember:**

1. Specific heat capacity is energy per unit mass per degree change.
2. Water has a high specific heat capacity compared to metals.
3. Latent heat of fusion is required for solid to liquid transition.
4. Latent heat of vaporization is required for liquid to gas transition.
5. Temperature remains constant during a phase change.

### 2.3 Gas Laws and Kinetic Theory

The behavior of gases is described by the relationship between pressure, volume, and temperature. Boyle's Law states that for a fixed mass of gas at a constant temperature, the pressure is inversely proportional to the volume. Charles's Law tells us that at constant pressure, the volume of a gas is directly proportional to its absolute temperature. These laws, combined with the Pressure Law, form the Ideal Gas Equation. The kinetic theory of gases provides the microscopic explanation for these laws, assuming that gas molecules are in constant, random motion and undergo perfectly elastic collisions. By understanding these laws, you can explain how a bicycle tire pressure changes with temperature or why a gas cylinder must be handled with care, as these real-world applications are often the basis for conceptual questions in your exams.

**Key Points to Remember:**

1. Boyle's Law: P1V1 equals P2V2 at constant temperature.
2. Charles's Law: V1/T1 equals V2/T2 at constant pressure.
3. Absolute zero is defined as 0 Kelvin or -273.15 degrees Celsius.
4. Ideal gas molecules are assumed to have negligible volume.
5. Pressure results from collisions of molecules with container walls.

### 2.4 Modes of Heat Transfer

Heat energy is transferred through three primary mechanisms: conduction, convection, and radiation. Conduction occurs primarily in solids through the direct collision of vibrating atoms or the movement of free electrons. Convection is the transfer of heat in fluids (liquids and gases) through the actual movement of the heated material, creating convection currents. Radiation is the transfer of energy via electromagnetic waves, which does not require a medium and can travel through a vacuum, explaining how the sun's heat reaches the Earth. In your exams, you might be asked to compare these methods or explain their applications, such as why thermos flasks are designed to minimize all three types of heat transfer. Understanding these mechanisms is vital for both theoretical explanations and practical problem-solving in physics.

**Key Points to Remember:**

1. Conduction is the main heat transfer method in solids.
2. Convection currents drive weather patterns and ocean currents.
3. Radiation requires no medium and travels at the speed of light.
4. Thermos flasks use vacuum layers to stop conduction and convection.
5. Shiny surfaces are good reflectors and poor emitters of radiation.

## 3. Key Definitions, Formulas, and Rules

1. **Linear Expansion:** Delta L = alpha - L0 - Delta T
2. **Heat Equation:** Q = m - c - Delta T
3. **Latent Heat:** Q = m - L
4. **Boyle's Law:** P1 - V1 = P2 - V2
5. **Charles's Law:** V1 / T1 = V2 / T2
6. **Ideal Gas Law:** P - V = n - R - T
7. **Temperature Conversion:** T(K) = T(C) + 273
8. **Specific Heat Capacity:** c = Q / (m - Delta T)
9. **Pressure Law:** P1 / T1 = P2 / T2

## 4. Step-by-Step Worked Examples

### Example 1: Linear Expansion Calculation

**Problem:** A steel rod is 2.0 meters long at 20 degrees Celsius. If it is heated to 100 degrees Celsius, calculate the change in length. (Alpha for steel = 1.2 x 10^-5 per degree Celsius)

**Solution Steps:**

Step 1: Identify given values: L0 = 2.0 m, Delta T = 100 - 20 = 80 C, alpha = 1.2 x 10^-5.

Step 2: Use formula: Delta L = alpha - L0 - Delta T.

Step 3: Substitute values: Delta L = (1.2 x 10^-5) - 2.0 - 80.

Step 4: Calculate: Delta L = 0.00192 meters.

**Final Answer:** 0.00192 meters or 1.92 mm

### Example 2: Heat Energy Required

**Problem:** How much heat is required to raise the temperature of 0.5 kg of water from 20 degrees Celsius to 80 degrees Celsius? (Specific heat of water = 4200 J/kg C)

**Solution Steps:**

Step 1: Identify given values: m = 0.5 kg, c = 4200 J/kg C, Delta T = 80 - 20 = 60 C.

Step 2: Use formula: Q = m - c - Delta T.

Step 3: Substitute values: Q = 0.5 - 4200 - 60.

Step 4: Calculate: Q = 126,000 Joules.

**Final Answer:** 126,000 Joules

### Example 3: Boyle's Law Application

**Problem:** A gas occupies 2 liters at a pressure of 1 atm. If the pressure is increased to 2 atm at constant temperature, what is the new volume?

**Solution Steps:**

Step 1: Identify given values: P1 = 1 atm, V1 = 2 L, P2 = 2 atm.

Step 2: Use formula: P1 - V1 = P2 - V2.

Step 3: Rearrange for V2: V2 = (P1 - V1) / P2.

Step 4: Substitute values: V2 = (1 - 2) / 2 = 1 L.

**Final Answer:** 1 Liter

### Example 4: Temperature Conversion

**Problem:** Convert 30 degrees Celsius to the Kelvin scale.

**Solution Steps:**

Step 1: Identify given value: T(C) = 30.

Step 2: Use formula: T(K) = T(C) + 273.

Step 3: Substitute: T(K) = 30 + 273.

Step 4: Calculate: T(K) = 303 K.

**Final Answer:** 303 Kelvin

## 5. Common Exam Mistakes to Avoid

1. Forgetting to convert Celsius to Kelvin when using gas laws, which leads to incorrect results.
2. Confusing specific heat capacity with latent heat; remember that specific heat is for temperature change, latent is for phase change.
3. Neglecting to calculate the change in temperature (Delta T) correctly, often using the final temperature instead of the difference.
4. Mixing up units, such as using grams instead of kilograms in the heat equation (Q = mcDeltaT).
5. Assuming that heat transfer stops at equilibrium, when in fact it continues at a net rate of zero.
6. Failing to include the correct SI units in the final answer, which results in a deduction of marks in board exams.

## 6. Practice Questions and Model Answers

### Question 1

Define specific heat capacity and state its SI unit.

**Answer 1:** Specific heat capacity is the amount of heat required to raise the temperature of 1 kg of a substance by 1 Kelvin (or 1 degree Celsius). Its SI unit is Joules per kilogram per Kelvin (J/kg K).

### Question 2

Why do we use mercury in thermometers?

**Answer 2:** Mercury is used because it has a uniform coefficient of expansion, a high boiling point, a low freezing point, and it is a good conductor of heat, making it highly sensitive to temperature changes.

### Question 3

A gas at 300 K has a volume of 5 cubic meters. If the temperature is doubled at constant pressure, what is the new volume?

**Answer 3:** Using Charles's Law (V1/T1 = V2/T2): 5/300 = V2/600. V2 = (5 - 600) / 300 = 10 cubic meters.

### Question 4

Explain why a metal spoon feels colder than a wooden spoon at room temperature.

**Answer 4:** Metal is a good conductor of heat, so it quickly conducts heat away from your hand. Wood is an insulator, so it does not conduct heat away from your hand as quickly, making it feel warmer.

### Question 5

What is the difference between heat and temperature?

**Answer 5:** Heat is the total energy transferred between systems due to a temperature difference, while temperature is a measure of the average kinetic energy of the molecules in a substance.