Basic trigonometric ratios

Subject: Mathematics | Level: Class 10 | Curriculum: Federal Board
Prepared by: SHS Virtual Academy

1. Topic Overview and Learning Objectives

Trigonometry, derived from the Greek words for triangle measurement, is a cornerstone of your Class 10 Mathematics curriculum. It provides the essential tools to analyze the relationships between the angles and the sides of right-angled triangles. Understanding these ratios is not merely an academic exercise but a gateway to advanced mathematics, physics, and engineering, forming the basis for everything from architectural design to navigation.

In the context of the Federal Board examinations, this topic is highly significant because it frequently appears in both objective and subjective sections. Examiners often test your conceptual clarity through questions involving finding unknown sides or angles, as well as real-world application problems. Scoring well here requires a solid grasp of the definitions of sine, cosine, and tangent, along with their reciprocals, as these concepts are building blocks for later chapters like heights and distances.

The fundamental intuition behind trigonometry is the concept of similarity. Because all right-angled triangles with the same acute angle are similar, the ratios of their corresponding sides remain constant regardless of the size of the triangle. By memorizing the SOH-CAH-TOA mnemonic and understanding how to identify the opposite, adjacent, and hypotenuse sides relative to a specific angle, you will unlock the ability to solve complex geometric problems with ease and precision.

2. Core Concepts and Detailed Revision Notes

2.1 Defining the Right Triangle Sides

To master trigonometry, you must first be able to correctly identify the three sides of a right-angled triangle relative to a given acute angle. The hypotenuse is always the longest side, located directly opposite the 90-degree angle. The other two sides are labeled based on their position relative to the angle you are focusing on, which we usually denote by the Greek letter theta. The side directly across from theta is called the opposite side, while the side that helps form the angle alongside the hypotenuse is called the adjacent side. If you misidentify these sides, your entire ratio calculation will be incorrect. Always take a moment to draw the triangle and label the hypotenuse, opposite, and adjacent sides clearly before you begin your calculations to ensure accuracy in your exam work. Remember that the labels opposite and adjacent are strictly dependent on the specific angle you are currently evaluating within the triangle.

Key Points to Remember:

1. The hypotenuse is always opposite the right angle.
2. The opposite side is across from the reference angle.
3. The adjacent side is next to the reference angle.
4. Labels change if the reference angle changes.
5. Always identify sides before applying ratios.

2.2 The Primary Trigonometric Ratios

The three primary trigonometric ratios are sine, cosine, and tangent. These ratios compare the lengths of two sides of a right triangle for a specific angle. Sine is defined as the ratio of the opposite side to the hypotenuse. Cosine is the ratio of the adjacent side to the hypotenuse. Tangent is the ratio of the opposite side to the adjacent side. A very helpful way to remember these is the mnemonic SOH-CAH-TOA, where SOH stands for Sine equals Opposite over Hypotenuse, CAH stands for Cosine equals Adjacent over Hypotenuse, and TOA stands for Tangent equals Opposite over Adjacent. By keeping this simple phrase in mind, you can quickly recall the definitions during the pressure of an exam. Practice writing these definitions out until they become second nature to you, as they form the foundation for all subsequent trigonometric identities and problem-solving techniques.

Key Points to Remember:

1. Sine = Opposite / Hypotenuse.
2. Cosine = Adjacent / Hypotenuse.
3. Tangent = Opposite / Adjacent.
4. Use SOH-CAH-TOA to memorize ratios.
5. Ratios are dimensionless values.

2.3 Reciprocal Trigonometric Ratios

Beyond the primary ratios, we have three reciprocal ratios: cosecant, secant, and cotangent. These are simply the multiplicative inverses of the primary ratios. Cosecant is the reciprocal of sine, meaning it is the ratio of the hypotenuse to the opposite side. Secant is the reciprocal of cosine, defined as the ratio of the hypotenuse to the adjacent side. Finally, cotangent is the reciprocal of tangent, defined as the ratio of the adjacent side to the opposite side. While these might seem like extra formulas to memorize, they are logically connected to the primary ratios. If you know that sine is opposite over hypotenuse, you automatically know that cosecant is hypotenuse over opposite. Understanding these relationships helps you simplify complex trigonometric expressions and solve equations more efficiently during your board examinations, especially when dealing with trigonometric identities.

Key Points to Remember:

1. Cosecant = 1 / Sine.
2. Secant = 1 / Cosine.
3. Cotangent = 1 / Tangent.
4. Reciprocals are inverses of primary ratios.
5. These ratios are essential for identities.

2.4 Solving for Unknown Sides and Angles

The true power of trigonometry lies in its ability to find unknown measurements. If you are given one side and one acute angle of a right triangle, you can use the appropriate trigonometric ratio to find the length of any other side. For example, if you know the angle and the hypotenuse, you can use sine to find the opposite side or cosine to find the adjacent side. Conversely, if you know two sides, you can find the angle by using the inverse trigonometric functions, written as arcsin, arccos, or arctan on your calculator. This process requires careful algebraic manipulation. Always ensure your calculator is set to degree mode before performing these calculations, as radian mode will give you incorrect results that will lead to lost marks on your paper. Always verify your final side lengths using the Pythagorean theorem to ensure your results are consistent with the properties of a right-angled triangle.

Key Points to Remember:

1. Use ratios to find missing sides.
2. Use inverse functions to find angles.
3. Ensure calculator is in degree mode.
4. Check your work with Pythagoras theorem.
5. Algebraic rearrangement is often required.

3. Key Definitions, Formulas, and Rules

1. **sin(theta):** Opposite / Hypotenuse
2. **cos(theta):** Adjacent / Hypotenuse
3. **tan(theta):** Opposite / Adjacent
4. **csc(theta):** 1 / sin(theta) = Hypotenuse / Opposite
5. **sec(theta):** 1 / cos(theta) = Hypotenuse / Adjacent
6. **cot(theta):** 1 / tan(theta) = Adjacent / Opposite
7. **tan(theta):** sin(theta) / cos(theta)
8. **Pythagorean Theorem:** a^2 + b^2 = c^2
9. **cot(theta):** cos(theta) / sin(theta)

4. Step-by-Step Worked Examples

Example 1: Finding a missing side

Problem: In a right triangle, the angle theta is 30 degrees and the hypotenuse is 10 cm. Find the length of the opposite side.

Solution Steps:

Step 1: Identify the ratio: sin(theta) = Opposite / Hypotenuse.

Step 2: Substitute known values: sin(30) = Opposite / 10.

Step 3: Solve for Opposite: Opposite = 10 - sin(30).

Step 4: Calculate: Opposite = 10 - 0.5 = 5.

Final Answer: 5 cm

Example 2: Finding an angle

Problem: In a right triangle, the opposite side is 3 cm and the adjacent side is 4 cm. Find the angle theta.

Solution Steps:

Step 1: Identify the ratio: tan(theta) = Opposite / Adjacent.

Step 2: Substitute values: tan(theta) = 3 / 4 = 0.75.

Step 3: Use inverse tangent: theta = arctan(0.75).

Step 4: Calculate: theta = 36.87 degrees.

Final Answer: 36.87 degrees

Example 3: Using secant ratio

Problem: If cos(theta) = 0.8, find the value of sec(theta).

Solution Steps:

Step 1: Recall definition: sec(theta) = 1 / cos(theta).

Step 2: Substitute value: sec(theta) = 1 / 0.8.

Step 3: Calculate: 1 / 0.8 = 1.25.

Final Answer: 1.25

Example 4: Applying Pythagoras

Problem: In a right triangle, the opposite side is 5 and the adjacent side is 12. Find the hypotenuse.

Solution Steps:

Step 1: Use Pythagoras: c^2 = a^2 + b^2.

Step 2: Substitute: c^2 = 5^2 + 12^2.

Step 3: Calculate: c^2 = 25 + 144 = 169.

Step 4: Take square root: c = sqrt(169) = 13.

Final Answer: 13

5. Common Exam Mistakes to Avoid

1. Forgetting to set the calculator to degree mode, which leads to completely wrong numerical answers.
2. Confusing the opposite and adjacent sides by not paying attention to which angle is being referenced.
3. Mixing up the primary ratios (SOH-CAH-TOA) during the exam due to nervousness.
4. Rounding off intermediate steps too early, which causes significant errors in the final result; always keep precision until the end.
5. Assuming that the hypotenuse can be one of the legs of the right triangle; always remember it is the longest side.
6. Failing to write the units (like cm or degrees) in the final answer, which can lead to point deductions.

6. Practice Questions and Model Answers

Question 1

If sin(theta) = 1/2, what is the value of csc(theta)?

Answer 1: Since csc(theta) = 1 / sin(theta), csc(theta) = 1 / (1/2) = 2.

Question 2

Find the length of the adjacent side if the hypotenuse is 20 and cos(theta) = 0.6.

Answer 2: Using cos(theta) = Adjacent / Hypotenuse, we get 0.6 = Adjacent / 20. Thus, Adjacent = 0.6 - 20 = 12.

Question 3

What is the value of tan(45 degrees)?

Answer 3: In a 45-45-90 triangle, the opposite and adjacent sides are equal, so tan(45) = 1/1 = 1.

Question 4

If tan(theta) = 1, find theta.

Answer 4: theta = arctan(1) = 45 degrees.

Question 5

Express cot(theta) in terms of sin(theta) and cos(theta).

Answer 5: Since cot(theta) = 1 / tan(theta) and tan(theta) = sin(theta) / cos(theta), then cot(theta) = cos(theta) / sin(theta).