

1 Setup python session and notebook

```
[32]: import numpy as np
import pencil as pc
import matplotlib.pyplot as plt
```

2 Get simulation as object

We will use the sample in 'pencil-code/python/tutorials/sample_simulations/2d_streaming_instability' in this notebook. So let's start by reading that simulation in as a simulation object.

```
[33]: SIMns = pc.get_sim('no_source')
SIMrd = pc.get_sim('ramp_decay')
SIMsd = pc.get_sim('step_decay')
SIMnd = pc.get_sim('step_no_decay')
```

3 Read timeseries

```
[34]: tsns = pc.read.ts(file_name="time_series.dat",datadir=SIMns.datadir)
tsrd = pc.read.ts(file_name="time_series.dat",datadir=SIMrd.datadir)
tssd = pc.read.ts(file_name="time_series.dat",datadir=SIMsd.datadir)
tsnd = pc.read.ts(file_name="time_series.dat",datadir=SIMnd.datadir)

tsall = [tsns, tsrd, tssd, tsnd]
labels = ['no_source', 'ramp_decay', 'step_decay', 'step_no_decay']
```

Read 40 lines.

Read 40 lines.

Read 40 lines.


Read 40 lines.

```
[35]: fig, ax = plt.subplots(2, len(tsall), figsize=(20,10))
for id,sim in enumerate(tsall):
    ax[0,id].plot(sim.t, sim.EEGW, label=r'$EEGW$')
    ax[1,id].plot(sim.t, sim.EEM, label=r'$EEM$')
    ax[0,id].set_title(labels[id])
    ax[0,id].set_xlabel(r'$t$')
```

```

ax[0,id].set_ylabel(r'$EEGW$')
ax[1,id].set_xlabel(r'$t$')
ax[1,id].set_ylabel(r'$EEM$')
ax[0,id].legend()
ax[1,id].legend()
plt.tight_layout()
plt.show()

```



3.0.1 Plotinf everything together

```

[36]: # Create the figure
plt.figure(figsize=(10, 6))

# Plot each dataset with its own x and y values
plt.plot(tsns.t, tsns.EEGW, label=r'$ns$')
plt.plot(tsrd.t, tsrd.EEGW, label=r'$rd$')
plt.plot(tssd.t, tssd.EEGW, label=r'$sd$')
plt.plot(tsnd.t, tsnd.EEGW, label=r'$nd$')

# Add labels and legend
plt.xlabel(r'$t$')
plt.ylabel(r'$EEGW$')
plt.legend()


# limits
plt.xlim(1, 1.10)
plt.ylim(-1e-7, 1e-6)
# Optional: Add grid for better readability

```

```
plt.grid(True, alpha=0.3)
```

```
# Show the plot
```

```
plt.show()
```



4 Reading parameters from the corespontid simulations

Here I will show all the parameters used in the corresponding simulation

```
[37]: params_ns = pc.read.param(datadir=SIMns.datadir,param2=True)
      params_rd = pc.read.param(datadir=SIMrd.datadir,param2=True)
      params_sd = pc.read.param(datadir=SIMsd.datadir,param2=True)
      params_nd = pc.read.param(datadir=SIMnd.datadir,param2=True)
      paramskeys_ns = list(params_ns.keys)
      paramskeys_rd = list(params_rd.keys)
      paramskeys_sd = list(params_sd.keys)
      paramskeys_nd = list(params_nd.keys)

      params_all = [params_ns, params_rd, params_sd, params_nd]
      params_labels = ['no_source', 'ramp_decay', 'step_decay', 'step_no_decay']
```

```
[38]: #Sources parameters for all simulations
      for id,sim in enumerate(params_all):
          print("-----")
```

```

print(f"key: {params_labels[id]}")
print(f"source5: {params_all[id].source5:e}")
print(f"source5_tdep: {params_all[id].source5_tdep}")
print(f"source5_expt: {params_all[id].source5_expt}")

```

```

-----
key: no_source
source5: 0.000000e+00
source5_tdep: const
source5_expt: 0.0

-----
key: ramp_decay
source5: 1.000000e+09
source5_tdep: step_decay_linear_ramp
source5_expt: 0.5

-----
key: step_decay
source5: 1.000000e+09
source5_tdep: step
source5_expt: 0.0

-----
key: step_no_decay
source5: 1.000000e+09
source5_tdep: step
source5_expt: 0.0

```

4.1 Generating power spectrum

4.2 general test

```

[39]: # read powr spec
      pwns = pc.read.power(datadir=SIMns.datadir)
      print(pwns.krms)

```

```

power_GWh.dat
powerhel_GWs.dat
power_Tpq.dat
power_Str.dat
powerhel_kin.dat
power_GWs.dat
powerhel_mag.dat
power_sp.dat
power_krms.dat
power_SCL.dat
power_mag.dat
powerhel_Str.dat
power_VCT.dat
power_kin.dat


```

```
powerhel_GWh.dat
[ 0.    1.29  2.24  3.14  4.07  5.11  6.13  7.08  8.03  9.06 10.1  11.1
 12.   13.   14.1  15. ]
```

```
[40]: # Compare kinetic energy spectrum at different times
plt.figure(figsize=(10, 6))

# Early, middle, and late snapshots
times = [0, len(pwns.t)//2, -1]
for idx in times:
    plt.loglog(pwns.krms, pwns.kin[idx], 'o-', label=f't = {pwns.t[idx]:.2f}')

plt.xlabel('k')
plt.ylabel('Kinetic Energy')
plt.title('Evolution of Kinetic Energy Spectrum')
plt.legend()
plt.grid(True, alpha=0.3)
plt.show()
```



```
[41]: # Track the evolution of specific k modes
plt.figure(figsize=(10, 6))
k_indices = [0, 5, 10, 15]


for k_idx in k_indices:
```

```

plt.semilogy(pwns.t, pwns.kin[:, k_idx], '-', label=f'k = {pwns.krms[k_idx]:
↪.3f}')

plt.xlabel('Time')
plt.ylabel('Kinetic Energy')
plt.title('Time Evolution of Kinetic Energy at Select k Values')
plt.legend()
plt.grid(True, alpha=0.3)
plt.show()

```



```


[42]: # Compare different types of spectra at the same time
plt.figure(figsize=(10, 6))

# Check if these spectra exist before plotting
if hasattr(pwns, 'kin'):
    plt.loglog(pwns.krms, pwns.kin[-1], 'o-', label='Kinetic')
if hasattr(pwns, 'mag'):
    plt.loglog(pwns.krms, pwns.mag[-1], 's-', label='Magnetic')
if hasattr(pwns, 'hel_kin'):
    plt.loglog(pwns.krms, np.abs(pwns.hel_kin[-1]), '^-', label='Kinetic_
↪Helicity')

plt.xlabel('k')
plt.ylabel('Power')

```


```
plt.title(f'Different Power Spectra at t = {pwns.t[-1]:.2f}')
plt.legend()
plt.grid(True, alpha=0.3)
plt.show()
```



```
[43]: plt.figure(figsize=(10, 6))

# Plot the spectrum at different times
times = [0, len(pwns.t)//2, -1]
for idx in times:
    plt.loglog(pwns.krms, pwns.GWs[idx], label=f't = {pwns.t[idx]:.2f}')

plt.xlabel('k')
plt.ylabel('GW Energy Spectrum')
plt.title('Gravitational Wave Energy Spectrum')
plt.legend()
plt.grid(True, alpha=0.3)
plt.show()
```



4.2.1 Power for the simulations


```
[44]: pwns = pc.read.power(datadir=SIMns.datadir)
      pwrnd = pc.read.power(datadir=SIMrd.datadir)
      pwrnsd = pc.read.power(datadir=SIMsd.datadir)
      pwrnd = pc.read.power(datadir=SIMnd.datadir)


      powers_all = [pwns, pwrnd, pwrnsd, pwrnd]
      powers_labels = ['no_source', 'ramp_decay', 'step_decay', 'step_no_decay']
```


```
power_GWh.dat
powerhel_GWs.dat
power_Tpq.dat
power_Str.dat
powerhel_kin.dat
power_GWs.dat
powerhel_mag.dat
power_sp.dat
power_krms.dat
power_SCL.dat
power_mag.dat
powerhel_Str.dat
power_VCT.dat
power_kin.dat
powerhel_GWh.dat
```



power_GWh.dat
powerhel_GWs.dat
power_Tpq.dat
power_Str.dat
powerhel_kin.dat
power_GWs.dat
powerhel_mag.dat
power_sp.dat
power_krms.dat
power_SCL.dat
power_mag.dat
powerhel_Str.dat
power_VCT.dat
power_kin.dat
powerhel_GWh.dat
power_GWh.dat
powerhel_GWs.dat
power_Tpq.dat
power_Str.dat
powerhel_kin.dat
power_GWs.dat
powerhel_mag.dat
power_sp.dat
power_krms.dat
power_SCL.dat
power_mag.dat
powerhel_Str.dat
power_VCT.dat
power_kin.dat
powerhel_GWh.dat
power_GWh.dat
powerhel_GWs.dat
power_Tpq.dat
power_Str.dat
powerhel_kin.dat
power_GWs.dat
powerhel_mag.dat
power_sp.dat
power_krms.dat
power_SCL.dat
power_mag.dat
powerhel_Str.dat
power_VCT.dat
power_kin.dat
powerhel_GWh.dat

```
[45]: for id,sim in enumerate(powers_all):
    times = [i for i in range(0, len(sim.t))]
    for idx in times:
        plt.loglog(sim.krms, sim.GWs[idx], label=f't = {sim.t[idx]:.2f}')
    plt.xlabel('k')
    plt.ylabel('GW Energy Spectrum')
    plt.title(f'Gravitational Wave Energy Spectrum for {powers_labels[id]}')
    #plt.legend()
    plt.grid(True, alpha=0.3)
    plt.show()
```









```
[46]: for id,sim in enumerate(powers_all):
    times = [i for i in range(0, len(sim.t))]
    for idx in times:
        plt.loglog(sim.krms, sim.mag[idx], label=f't = {sim.t[idx]:.2f}')
    plt.xlabel('k')
    plt.ylabel('Magnetic Spectrum')
    plt.title(f'Magnetic Energy Spectrum for {powers_labels[id]}')
    #plt.legend()
    plt.grid(True, alpha=0.3)
    plt.show()
```

