

MATH 8170,
Fall 2016
HW 1
Shuai Wei

1.

$$\begin{aligned} P(N(t) = k) &= P(T_k \leq t < T_{k+1}) \\ &= P(T_k \leq t) - P(T_{k+1} \leq t) \end{aligned}$$

$\{T_1 = X_1\}$ may be interpreted as the rewarded recieved after conducted one bernoulli trail. $X_1 = 1$ if trial fails and $X_1 = 2$ if it succeeds.
 $\{T_n = X_n\}$ may be interpreted as the rewards n recieved after conducting k bernoulli trails, and then we have $n - k$ succeses and $k - (n - k) = 2k - n$ fails. Moreover, the rewards n range from k to $2k$.

$$P(T_k = n) = \binom{k}{n-k} 0.2^{n-k} 0.8^{2k-n}, n = k, k+1, \dots, 2k$$

So

$$\begin{aligned} P(N(t) = k) &= \sum_{n=k}^{\min(\lfloor t \rfloor, 2k)} P(T_k = n) - \sum_{n=k+1}^{\lfloor t \rfloor} P(T_{k+1} = n) \\ &= \sum_{n=k}^{\min(\lfloor t \rfloor, 2k)} \binom{k}{n-k} 0.2^{n-k} 0.8^{2k-n} - \sum_{n=k+1}^{\min(\lfloor t \rfloor, 2k+2)} \binom{k+1}{n-k-1} 0.2^{n-k-1} 0.8^{2k+2-n} \end{aligned}$$

2.

Assume $T_0 = 0$.

$$\begin{aligned} E(A(t)B(t)) &= \int_0^\infty E(A(t)B(t)|T_1 = s) dF(s) \\ &= \int_0^t E(A(t)B(t)|T_1 = s) dF(s) + \int_t^\infty E(A(t)B(t)|T_1 = s) dF(s) \\ &= \int_0^t E(A(t-s)B(t-s) dF(s) + E\left((t - T_{N(t)})(T_{N(t)+1} - t) | T_1 \geq t\right) \\ &= \int_0^t E(A(t-s)B(t-s) dF(s) + E(t(T_1 - t) | T_1 \geq t) \end{aligned}$$

Then $h(t) = E(t(T_1 - t)|T_1 \geq t)$, so h is continuous a.e on $[0, \infty)$.
so for $t \geq 1$

$$\begin{aligned} |h(t)| &= E(t(T_1 - t)|T_1 \geq t) \leq E(T_1/2(T_1 - T_1/2)|T_1 \geq t) \\ &= E(T_1^2/4|T_1 \geq t) \\ &= \frac{1}{4}E(T_1^2|T_1 \geq t) = b(t). \end{aligned}$$

It is obvious that $b(t)$ is directly Riemann integrable since F has finite second moment.

Besides, F is nonarithmetic.

So the key renewal theorem is applicable, and

$$\begin{aligned} \lim_{t \rightarrow \infty} E(A(t)B(t)) &= \frac{1}{E(X_1)} \int_0^\infty E(t(T_1 - t)|T_1 \geq t) dt \\ &= \frac{1}{E(X_1)} \int_0^\infty E(tT_1 - t^2|T_1 \geq t) dt \\ &= \frac{1}{E(X_1)} \int_0^\infty \int_t^\infty E(tT_1 - t^2|T_1 = s) dF(s) dt \\ &= \frac{1}{E(X_1)} \int_0^\infty \int_t^\infty E(ts - t^2) dF(s) dt \\ &= \frac{1}{E(X_1)} \int_0^\infty \int_t^\infty ts - t^2 dF(s) dt \\ &= \frac{1}{E(X_1)} \int_0^\infty \left(t \int_t^\infty s dF(s) - t^2 \int_t^\infty dF(s) \right) dt \\ &= \frac{1}{\int_0^\infty s dF(s)} \int_0^\infty \left(t \int_t^\infty s dF(s) - t^2 (1 - F(t)) \right) dt \end{aligned}$$

3.

$$\begin{aligned} P(N(t) \text{ is odd}) &= P(N(t) \text{ is odd}, T_1 > t) + P(N(t) \text{ is odd}, T_1 \leq t) \\ &= P(N(t) \text{ is odd}, T_1 \leq t) \\ &= \int_0^t P(N(t) \text{ is odd} | T_1 = s) dF(s) \\ &= \int_0^t P(N(t-s) \text{ is even}) dF(s) \\ &= \int_0^t (1 - P(N(t-s) \text{ is odd})) dF(s) \\ &= \int_0^t dF(s) - \int_0^t P(N(t-s) \text{ is odd}) dF(s) \\ &= F(t) - \int_0^t P(N(t-s) \text{ is odd}) dF(s) \end{aligned}$$

Let $H(t) = P(N(t) \text{ is odd})$, then $H(0) = 0$ and $F(t) = 1 - e^{-\lambda t}$ and $dF(s) = \lambda e^{-\lambda s} ds$.

$$\begin{aligned} H(t) &= F(t) - \int_0^t H(t-s) dF(s) \\ &= 1 - e^{-\lambda t} - \int_0^t H(t-s) \lambda e^{-\lambda s} ds \\ &= 1 - e^{-\lambda t} - \lambda e^{-\lambda t} \int_0^t H(x) e^{\lambda x} dx \end{aligned}$$

Take derivatives on both sides, we have

$$\begin{aligned} H'(t) &= \lambda e^{-\lambda t} + \lambda^2 e^{-\lambda t} \int_0^t H(x) e^{\lambda x} dx - \lambda e^{-\lambda t} H(t) e^{\lambda t} \\ &= \lambda e^{-\lambda t} + \lambda^2 e^{-\lambda t} \int_0^t H(x) e^{\lambda x} dx - \lambda H(t) \\ &= \lambda e^{-\lambda t} + \lambda(1 - e^{-\lambda t} - H(t)) - \lambda H(t) \\ &= \lambda - 2\lambda H(t) \end{aligned}$$

So

$$H'(t) + 2\lambda H(t) = \lambda$$

It can be found that $\frac{\lambda}{2}$ is the special solution and $Ce^{-2\lambda t}, C \neq 0$ are the general solutions.

So $H(t) = Ce^{-2\lambda t} + \frac{\lambda}{2}$ is the general form of the solution.

Since $H(0) = C + \frac{\lambda}{2} = 0$, we have $C = -\frac{\lambda}{2}$.

Thus,

$$H(t) = -\frac{\lambda}{2} e^{-2\lambda t} + \frac{\lambda}{2}.$$

4.

Proof. (a)

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{T_n^D}{n} &= \lim_{n \rightarrow \infty} \frac{X_1 + \sum_{i=2}^n X_i}{n} \\ &= \lim_{n \rightarrow \infty} \frac{\sum_{i=2}^n X_i}{n} \\ &= \lim_{n \rightarrow \infty} \frac{\sum_{i=2}^n X_i}{n-1} \frac{n-1}{n} \\ &= \lim_{n \rightarrow \infty} \frac{\sum_{i=2}^n X_i}{n-1} \lim_{n \rightarrow \infty} \frac{n-1}{n} \\ &= E[X_2] \quad w.p.1 \end{aligned}$$

by central limit theorem since $X_n, n \geq 2$ are *iid*.

Then there exists an event $A \subseteq \Omega$ with $P(A) = 1$ and $\lim_{t \rightarrow \infty} \frac{T_n^D(\omega)}{n} = \mu$ for each $\omega \in A$.

We observe that $T_n^D(\omega) < \infty$ for each n and $T_n^D(\omega) \rightarrow \infty$ as $n \rightarrow \infty$, which implies $\lim_{t \rightarrow \infty} N^D(t, \omega) = \infty$.
 Fix an $\omega \in A$. For each $t > 0$,

$$T_{N^D(t)}^D(\omega) \leq t < T_{N^D(t)+1}^D(\omega).$$

Let t large enough such that $N^D(t, \omega) \geq 1$, and divide the above equalities by $N^D(t, \omega)$, so we have

$$\frac{T_{N^D(t)}^D(\omega)}{N^D(t, \omega)} \leq \frac{t}{N^D(t, \omega)} < \frac{T_{N^D(t)}^D(\omega) + 1}{N^D(t, \omega)}$$

Namely,

$$\frac{T_{N^D(t)}^D(\omega)}{N^D(t, \omega)} \leq \frac{t}{N^D(t, \omega)} < \frac{T_{N^D(t)}^D(\omega) + 1}{N^D(t, \omega) + 1} \frac{N^D(t, \omega) + 1}{N^D(t, \omega)}$$

Let $t \rightarrow \infty$, then

$$\frac{T_{N^D(t)}^D(\omega)}{N^D(t, \omega)} \rightarrow E[X_2] \text{ and } \frac{T_{N^D(t)}^D(\omega) + 1}{N^D(t, \omega) + 1} \frac{N^D(t, \omega) + 1}{N^D(t, \omega)} \rightarrow E[X_2].$$

So

$$\lim_{t \rightarrow \infty} \frac{N_D(t)}{t} = \frac{1}{E[X_2]}$$

(b) Step1:

Note that the *r.v* $N^D(t) + 1$ is a stopping time *w.r.t* $\{X_n\}_{n=1}^\infty$.
 Since

$$1(N^D(t) + 1 \geq k) = 1 - 1(N^D(t) + 1 < k) = 1 - 1(N^D(t) + 1 \leq k - 1),$$

$1(N^D(t) + 1 \geq k)$ is a function of X_0, X_1, \dots, X_{k-1} , and $1(N^D(t) \geq k)$ is independent of X_k .

Therefore,

$$\begin{aligned}
E(T_{N^D(t)+1}^D) &= E\left(\sum_{k=1}^{N^D(t)+1} X_k\right) \\
&= E\left(\sum_{k=1}^{\infty} X_k 1(N^D(t) + 1 \geq k)\right) \\
&= \sum_{k=1}^{\infty} E\left(X_k 1(N^D(t) + 1 \geq k)\right) \\
&= \sum_{k=1}^{\infty} E(X_k) P(N^D(t) + 1 \geq k) \\
&= E(X_1) P(N^D(t) + 1 \geq 1) + \sum_{k=2}^{\infty} E(X_k) P(N^D(t) + 1 \geq k) \\
&= E(X_1) + \sum_{k=1}^{\infty} E(X_{k+1}) P(N^D(t) + 1 \geq k+1) \\
&= E(X_1) + \sum_{k=1}^{\infty} E(X_{k+1}) P(N^D(t) \geq k) \\
&= E(X_1) + E(X_2) \sum_{k=1}^{\infty} P(N^D(t) \geq k) \\
&= E(X_1) + E(X_2) E(N^D(t)).
\end{aligned}$$

So

$$t < E(T_{N^D(t)+1}^D) = E(X_1) + E(X_2) E(N^D(t)).$$

Then

$$\frac{E(N^D(t))}{t} > \frac{1}{E(X_2)} - \frac{E(X_1)}{E(X_2)} \frac{1}{t}.$$

Thus,

$$\lim_{t \rightarrow \infty} \frac{E(N^D(t))}{t} \geq \frac{1}{E(X_2)}.$$

Step2:

Fix $a > 0$, and define $X_n^{(a)} = \min\{X_n, a\}$.

Let $T_n^{D(a)} = \sum_{k=1}^n X_k^{(a)}$, and define a new renewal process $\{N^{D(a)}(t)\}$ whose points are given by $\{T_n^{D(a)}\}$.

Then

$$E\left(T_{N^{D(a)}(t)+1}^{D(a)}\right) = E(X_1^a) + E(X_2^{(a)}) E(N^{D(a)}(t)).$$

by following the similat process with Step 1.

Next we note that

$$T_{N^{D(a)}(t)+1}^{D(a)} = T_{N^{D(a)}(t)}^{D(a)} + X_{N^{D(a)}(t)+1}^{(a)} \leq t + a.$$

So

$$\begin{aligned}
E\left(T_{N^{D(a)}(t)+1}^{D(a)}\right) &\leq t + a \implies E(X_1^a) + E(X_2^{(a)})E(N^{D(a)}(t)) \leq t + a \\
&\implies \frac{E(N^{D(a)}(t))}{t} \leq \frac{1}{E(X_2^{(a)})} + \frac{a - E(X_1^a)}{E(X_2^{(a)})} \frac{1}{t} \\
&\implies \lim_{t \rightarrow \infty} \sup \frac{E(N^{D(a)}(t))}{t} \leq \frac{1}{E(X_2^{(a)})}
\end{aligned}$$

Since $N^D(t) < N^{D(a)}(t)$, we have

$$\lim_{t \rightarrow \infty} \sup \frac{E(N^D(t))}{t} \leq \frac{1}{E(X_2^{(a)})}.$$

Let $a \rightarrow \infty$, and we have

$$\lim_{t \rightarrow \infty} \sup \frac{E(N^D(t))}{t} \leq \frac{1}{E(X_2)}$$

According to the above two steps, we have

$$\lim_{t \rightarrow \infty} \frac{E(N^D(t))}{t} = \frac{1}{E(X_2)}$$

□

5.

- (a) Define X_i as how long it takes the miner to make the i -th journey.
Define the stopping time N as the first journey which takes the miner 2 days.
Verify:

$$1(N = n) = 1(X_1 = 4 \text{ or } 6 \text{ for } i = 1, 2, \dots, n-1, X_{n-1} = 2)$$

$$P(N = i) = \left(\frac{2}{3}\right)^{i-1} \frac{1}{3}, \quad i = 1, 2, \dots$$

So $N \sim Geo\left(\frac{1}{3}\right)$ and

$$E(N) = 3$$

Moreover,

$$P(X_i = j) = \frac{1}{3} \text{ for } j = 2, 4, 6.$$

So for $i = 1, 2, \dots$,

$$E(X_i) = \frac{1}{3}(2 + 4 + 6) = 4.$$

- (b) By Wald's equality,

$$E(T) = E(N)E(X_i) = 3 * 4 = 12$$

(c)

$$\begin{aligned} E\left(\sum_{i=1}^N X_i | N = n\right) &= E\left(\sum_{i=1}^n X_i\right) \\ &= \sum_{i=1}^n E(X_i) \\ &= 4n \end{aligned}$$

since n is a constant and X_i are *iid*.

(d) By part (c), we have

$$E(T|N) = E\left(\sum_{i=1}^N X_i | N\right) = 4N.$$

So

$$E(T) = E(E(T|N)) = E(4N) = 4E(N) = 4 * 3 = 12.$$

6.

Let μ_G denote the mean service time.

Then the mean time of a cycle is $\mu_G + \frac{1}{\lambda}$ by the memoryless property of the Poisson process.

So the proportion of time (on the average) the server is busy is $\frac{\mu_G}{\mu_G + \frac{1}{\lambda}}$.