

MATH 8170,

Fall 2016

HW 2

Shuai Wei

**1.**

Let  $X_A(t)$  denote the number of organisms in state A at time  $t$ .

Let  $X_B(t)$  denote the number of organisms in state B.

Then  $X = \{X_A, X_B\}$  is a CTMC with sojourn time rate  $v_{ij} = \alpha i + \beta j$  for each state  $(i, j) \in E$ , where the state space is

$$E = \{(i, j)\}_{i, j \geq 0, i+j \geq 1},$$

where  $i, j \geq 0$  and  $i + j \geq 1$ .

The sojourn time  $T_{ij}$  for the state  $(i, j)$  is exponentially distributed with rate  $\alpha i + \beta j$ .

The transition probability is :

$$P_{(i,j)(i+2,j-1)} = \frac{\beta j}{\alpha i + \beta j};$$

$$P_{(i,j)(i-1,j+1)} = \frac{\alpha i}{\alpha i + \beta j};$$

**2.**

Let  $X(t)$  denote the number of customers in the system at time  $t$ .

The time until the next entering/birth  $X \sim \exp(\lambda \alpha_i)$ , where  $i$  is the number of customers already in the system.

The time until the next service ending/death  $Y \sim \exp(\mu)$ .

The birth rate  $\lambda \alpha_i$ , the death rate is  $\mu$ .

Then  $X(t)$  is a CTMC with sojourn time rate  $v_i = \mu + \lambda \alpha_i$  for state  $i$ .

Since it can only change its state by increasing by one or decreasing by one, it is a birth and death process.

The state space is

$$E = \{0, 1, 2, 3, \dots\}.$$

The transition probability is :

$$P_{ij} = \begin{cases} P(X < Y), & j = i + 1, i \geq 1 \\ P(Y < X), & j = i - 1, i \geq 1 \end{cases} = \begin{cases} \frac{\lambda \alpha_i}{\mu + \lambda \alpha_i}, & j = i + 1, i \geq 1 \\ \frac{\mu}{\mu + \lambda \alpha_i}, & j = i - 1, i \geq 1 \end{cases}$$

When  $i = 0$ ,  $P_{01} = 1$ .

### 3.

Yes, it is a CTMC.

Assume the number of people who have a infection is  $k$  given  $1 \leq k \leq N$ , then the state space is

$$E = \{k, n + 1, \dots, N\}.$$

For  $1 \leq n \leq N$ , let  $\tau_n$  be the time when a new infection occurs.

When a contact occurs at time  $t$  with  $X(t) = i$ , the probability of one being infected is

$$\frac{N(N-i)}{\binom{N}{2}} = \frac{2i(N-i)}{N(N-1)}.$$

Since contacts between two members of this population occur in accordance with a Poisson process having rate  $\lambda$ ,

the sojourn time  $T_n = \tau_{n+1} - \tau_n$  for  $X(\tau_n) = i$  with  $k \leq i < N$  and  $1 \leq n \leq N$  satisfies

$$T_n \sim \exp\left(\frac{2i(N-i)}{N(N-1)}\lambda\right).$$

For  $k \leq i < N$ ,

$$\begin{aligned} P(X_{n+1} = i + 1, T_{n+1} > t \mid X_n = i, (X_j, T_j), j < n) &= P(X_{n+1} = i + 1, T_{n+1} > t \mid X_n = i) \\ &= P(X_1 = i + 1, T_1 > t \mid X_0 = i) \\ &= \exp\left(-\frac{2i(N-i)}{N(N-1)}\lambda t\right). \end{aligned}$$

If the system enters the state  $N$ , it will stay there forever.

4.

Define a CTMC with  $E = \{0, 1, 2, 3\}$ .

The state 0 means both machine 1 and 2 operate.

The state 1 means machine 2 operates but machine 1 is being repaired.

The state 2 means machine 1 operates but machine 2 is being repaired.

The state 3 means both machine 1 and 2 are being repaired.

The operate time

$$O_i \sim \exp(\lambda_i)$$

for  $i = 1, 2$ .

The repair time

$$R_i \sim \exp(\mu_i)$$

for  $i = 1, 2$ .

Then by the memoryless property of exponential distribution,

$$P_{01} = P(O_1 < O_2) = \frac{\lambda_1}{\lambda_1 + \lambda_2}, \quad P_{02} = P(O_2 < O_1) = \frac{\lambda_2}{\lambda_1 + \lambda_2},$$

$$P_{10} = P(R_1 < O_2) = \frac{\mu_1}{\mu_1 + \lambda_2}, \quad P_{13} = P(O_2 < R_1) = \frac{\lambda_2}{\mu_1 + \lambda_2},$$

$$P_{20} = P(R_2 < O_1) = \frac{\mu_2}{\lambda_1 + \mu_2}, \quad P_{23} = P(O_1 < R_2) = \frac{\lambda_1}{\lambda_1 + \mu_2},$$

$$P_{31} = P(R_2 < R_1) = \frac{\mu_2}{\mu_1 + \mu_2}, \quad P_{32} = P(R_1 < R_2) = \frac{\mu_1}{\mu_1 + \mu_2}.$$

Besides,

$$P_{03} = P_{30} = P_{12} = P_{21} = 0.$$

We know

$$v_0 = \lambda_1 + \lambda_2, \quad v_1 = \mu_1 + \lambda_2, \quad v_2 = \lambda_1 + \mu_2, \quad v_3 = \mu_1 + \mu_2.$$

So

$$q_{01} = v_0 P_{01} = \lambda_1, \quad q_{02} = v_0 P_{02} = \lambda_2,$$

$$q_{10} = v_1 P_{10} = \mu_1, \quad q_{13} = v_1 P_{13} = \lambda_2,$$

$$q_{20} = v_2 P_{20} = \mu_2, \quad q_{23} = v_2 P_{23} = \lambda_1,$$

$$q_{31} = v_3 P_{31} = \mu_2, \quad q_{32} = v_3 P_{32} = \mu_1,$$

and

$$q_{03} = q_{30} = q_{12} = q_{21} = 0.$$

Moreover,  $q_{ii} = v_i$  for  $i = 0, 1, 2, 3$ . Thus, the transition rate matrix is

$$Q = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} -(\lambda_1 + \lambda_2) & \lambda_1 & 0 & \lambda_2 \\ \mu_1 & -(\mu_1 + \lambda_2) & 0 & \lambda_2 \\ \mu_2 & 0 & -(\lambda_1 + \mu_2) & \lambda_1 \\ 0 & \mu_2 & \mu_1 & -(\mu_1 + \mu_2) \end{pmatrix} \end{matrix}$$

Next we compute the transition matrix.

Consider first the case that there is just the machine 1.

Definte a CTMC with state space  $E = \{o, r\}$ .

State  $o$  means it operates and  $r$  means it is being repaired.

Then  $v_o = \lambda_1$  and  $v_r = \mu_1$ .

So the transition rate matrix is

$$Q_1 = \begin{matrix} & \begin{matrix} o & r \end{matrix} \\ \begin{matrix} o \\ r \end{matrix} & \begin{pmatrix} -\lambda_1 & \lambda_1 \\ \mu_1 & -\mu_1 \end{pmatrix} \end{matrix}$$

According to the computational results from a similar example in class, we have

$$P_{oo}^1(t) = \frac{\mu_1}{\lambda_1 + \mu_1} + \frac{\lambda_1}{\lambda_1 + \mu_1} e^{-(\lambda_1 + \mu_1)t}, \quad P_{or}^1(t) = \frac{\lambda_1}{\lambda_1 + \mu_1} - \frac{\lambda_1}{\lambda_1 + \mu_1} e^{-(\lambda_1 + \mu_1)t},$$

$$P_{rr}^1(t) = \frac{\lambda_1}{\lambda_1 + \mu_1} - \frac{\mu_1}{\lambda_1 + \mu_1} e^{-(\lambda_1 + \mu_1)t}, \quad P_{ro}^1(t) = \frac{\mu_1}{\lambda_1 + \mu_1} - \frac{\mu_1}{\lambda_1 + \mu_1} e^{-(\lambda_1 + \mu_1)t}.$$

In addition, we have the similar result for the machine 2.

Since the machines act independently of each other,

