

Homework 4, MATH 8010

Shuai Wei

2.

(b) *Proof.* Let T_n be the n^{th} replacement for $n \geq 1$ and set $T_0 = 0$.

Then $\{T_n - T_{n-1}\}_{n=1}^{\infty}$ is an *iid* sequence of nonnegative random variables.

Let S_1 be the first in-use failure.

Let $N = \inf\{n \geq 1 : T_n - T_{n-1} < T\}$.

Since $\mathbb{1}(N \leq n) = 1 - \mathbb{1}(N > n) = 1 - \mathbb{1}(T_1 - T_0 \geq T, T_2 - T_1 > T, \dots, T_n - T_{n-1} \geq T)$, N is a stopping time w.r.t. $\{T_n - T_{n-1}\}_{n=0}^{\infty}$.

Then

$$S_1 = \sum_{n=1}^N (T_n - T_{n-1}).$$

By Wald's identity, we have

$$E(S_1) = E(N)E(T_1).$$

Since for $n \in \mathbb{N}$,

$$\begin{aligned} P(N = n) &= P(T_n - T_{n-1} < T) \prod_{k=1}^{n-1} P(T_k - T_{k-1} \geq T) \\ &= P(T_1 < T) \prod_{k=1}^{n-1} P(T_1 \geq T) \\ &= F(T) (1 - F(T))^{n-1}, \end{aligned}$$

$$E(N) = \sum_{n=1}^{\infty} nP(N = n) = F(T) \sum_{n=1}^{\infty} n(1 - F(T))^{n-1}.$$

We know the Maclaurin series of $\frac{1}{(1-x)^2}$ is

$$\sum_{n=1}^{\infty} nx^{n-1}$$

for $-1 < x < 1$.

The density f is continuous on $(0, \infty)$, so $0 < F(T) < 1$ for $T \in \mathbb{R}^+$.

Then we have $0 < 1 - F(T) < 1$.

So, replace x with $1 - F(T)$, we have

$$\sum_{n=1}^{\infty} n(1 - F(T))^{n-1} = \frac{1}{1 - (1 - F(T))^2} = \frac{1}{F^2(T)}.$$

Then

$$E(N) = F(T) \frac{1}{F^2(T)} = \frac{1}{F(T)}.$$

Thus, the average time between two successive in-use failures is

$$E(S_1) = E(N)E(T_1) = \frac{\int_0^T xf(x)dx + T(1 - F(T))}{F(T)}.$$

□

3.

1. Assume $p \neq q$.

Let $\{X_n\}$ be the winning of the n^{th} bet for $n \geq 1$.

Then for $n \geq 1$,

$$X_n = \begin{cases} 1, & \text{with probability } p; \\ -1, & \text{with probability } 1 - p. \end{cases}$$

So $\{X_n\}_{n=1}^{\infty}$ is an *iid* sequence of nonnegative random variables.

Then $E(X_n) = p - q$.

Let $T = \inf\{n \geq 1 : \sum_{k=1}^n X_k = N \text{ or } \sum_{k=1}^n X_k = 0\}$.

Since

$$\begin{aligned} \mathbb{1}(T \leq n) &= 1 - \mathbb{1}(T > n) \\ &= 1 - \mathbb{1}(0 < \sum_{k=1}^1 X_k < N, 0 < \sum_{k=1}^2 X_k < N, \dots, 0 < \sum_{k=1}^n X_k < N), \end{aligned}$$

T is a stopping time w.r.t $\{\sum_{k=1}^n X_k\}_{n=1}^{\infty}$.

Let Y be the final winning of the gambler when he stops.

So given the initial fortune i \$,

$$Y = \begin{cases} N - i, & \text{with probability } P_i = \frac{1-(q/p)^i}{1-(q/p)^N}; \\ -i, & \text{with probability } 1 - P_i = \frac{(q/p)^i - (q/p)^N}{1-(q/p)^N}. \end{cases}$$

So

$$E\left(\sum_{n=1}^T X_n\right) = E(Y) = (N - i)P_i - i(1 - P_i).$$

By Wald's identity, we also have

$$E\left(\sum_{n=1}^T X_n\right) = E(T)E(X_1) = (p - q)E(T).$$

Thus,

$$E(T) = \frac{(N - i)P_i - i(1 - P_i)}{p - q}.$$

2. Assume $p = q$.

Let $E_i(T)$ be $E(T)$ when given initial fortune is i \$.

Then

$$\begin{aligned} E_i(T) &= E(E_i(T|X_1)) \\ &= ET|X_1 = 1 \frac{1}{2} + ET|X_1 = -1 \frac{1}{2} \\ &= \frac{1}{2}(1 + E_{i+1}(T)) + \frac{1}{2}(1 + E_{i-1}(T)) \\ &= 1 + \frac{1}{2}E_{i+1}(T) + \frac{1}{2}E_{i-1}(T) \end{aligned}$$

Let $d_i = E_i(T) - E_{i-1}(T)$.

Then $d_{i+1} = d_i - 2$.

We can find that

$$\begin{aligned} \begin{pmatrix} d_{i+1} \\ -2 \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} d_i \\ -2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^2 \begin{pmatrix} d_{i-1} \\ -2 \end{pmatrix} \\ &= \dots \\ &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^i \begin{pmatrix} d_1 \\ -2 \end{pmatrix} \end{aligned}$$

We claim

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^i = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}.$$

Basic case:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$$

Assume

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^i = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}$$

Then

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{i+1} &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^i \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & i+1 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

So the claim holds. Then we get

$$\begin{pmatrix} d_{i+1} \\ -2 \end{pmatrix} = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix} \begin{pmatrix} d_1 \\ -2 \end{pmatrix}$$

So

$$d_{i+1} = d_1 - 2i.$$

So

$$d_N = d_1 - 2(N - 1).$$

We know $d_1 = E_1(T) - E_0(T) = E_1(T)$, and

$d_N = E_N(T) - E_{N-1}(T) = -E_{N-1}(T)$. Since $p = q$, by symmetry, we have

$$E_1(T) = E_{N-1}(T).$$

So

$$d_N = -d_1.$$

Combine it with $d_N = d_1 - 2(N - 1)$, we have

$$d_1 = N - 1.$$

Then

$$d_i = d_1 - 2(i - 1) = N - 2i + 1.$$

As a result, given initial fortune i \$, we have

$$\begin{aligned} E_T = E_i(T) &= d_i + E_{i-1}^T \\ &= d_i + d_{i-1} + E_{i-2}^T \\ &= \dots \\ &= d_i + d_{i-1} + \dots + d_1 + E_0^T \\ &= (N - 2i + 1) + (N - 2(i - 1) + 1) + \dots + N - 1 + 0 \\ &= Ni - i^2. \end{aligned}$$

4.