

Assignment - 1

Shubham Ramesh Dangat
SM21MTECH14003

PROBLEM

1. The co-ordinates of points A,B are (r_1, θ_1) , (r_2, θ_2) referred to O as pole. The internal bisector of angle AOB meets the line AB in D. Find the co-ordinates of D.

SOLUTION

Let the co-ordinates of point D be $\begin{pmatrix} r \\ \theta \end{pmatrix}$.

Now , $\theta = \frac{\theta_1 + \theta_2}{2}$

\therefore Converting into rectangular coordinates,

$$\mathbf{A} = \begin{pmatrix} r_1 \sin(\theta_1) \\ r_1 \cos(\theta_1) \end{pmatrix}, \mathbf{B} = \begin{pmatrix} r_2 \sin(\theta_2) \\ r_2 \cos(\theta_2) \end{pmatrix}, \quad (1)$$

$$\mathbf{O} = \begin{pmatrix} 0 \sin(0) \\ 0 \cos(0) \end{pmatrix}, \mathbf{D} = \begin{pmatrix} r \sin(\theta) \\ r \cos(\theta) \end{pmatrix} \quad (2)$$

For two vectors,

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} \quad (3)$$

$$\|\mathbf{a} \times \mathbf{b}\| = |(a_1 b_2 - a_2 b_1)| \quad (4)$$

$$\|\mathbf{A} - \mathbf{O}\| = \begin{pmatrix} r_1 \sin(\theta_1) \\ r_1 \cos(\theta_1) \end{pmatrix},$$

$$\|\mathbf{A} - \mathbf{D}\| = \begin{pmatrix} r_1 \sin(\theta_1) - r \sin(\theta) \\ r_1 \cos(\theta_1) - r \cos(\theta) \end{pmatrix},$$

$$\|\mathbf{B} - \mathbf{D}\| = \begin{pmatrix} r_2 \sin(\theta_2) - r \sin(\theta) \\ r_2 \cos(\theta_2) - r \cos(\theta) \end{pmatrix},$$

$$\|\mathbf{B} - \mathbf{O}\| = \begin{pmatrix} r_2 \sin(\theta_2) \\ r_2 \cos(\theta_2) \end{pmatrix},$$

$$\|\mathbf{A} - \mathbf{B}\| = \begin{pmatrix} r_1 \sin(\theta_1) - r_2 \sin(\theta_2) \\ r_1 \cos(\theta_1) - r_2 \cos(\theta_2) \end{pmatrix}$$

$$\text{Area } (\triangle AOD) + \text{Area}(\triangle DOB) = \text{Area}(\triangle AOB)$$

$$\frac{1}{2} \|(\mathbf{A} - \mathbf{O}) \times (\mathbf{A} - \mathbf{D})\| + \frac{1}{2} \|(\mathbf{B} - \mathbf{D}) \times (\mathbf{B} - \mathbf{O})\| = \frac{1}{2} \|(\mathbf{A} - \mathbf{O}) \times (\mathbf{A} - \mathbf{B})\| \quad (5)$$

\therefore Substituting A-O , A-D , B-D , B-O , A-B and simplifying further we get,

$$\begin{aligned} \frac{1}{2} r_1 r \sin\left(\frac{\theta}{2}\right) + \frac{1}{2} r_2 r \sin\left(\frac{\theta}{2}\right) \\ = \frac{1}{2} r_1 r_2 \sin\left(\frac{\theta}{2}\right) \end{aligned} \quad (6)$$

$$\begin{aligned} \text{Now , R.H.S} &= \frac{1}{2} r_1 (r_2 \sin(\theta)) \\ &= \frac{1}{2} 2 r_1 r_2 \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{2}\right) \end{aligned} \quad (7)$$

On Comparing (6) and (7),

$$\therefore \frac{1}{2} r (r_1 + r_2) = r_1 r_2 \cos\left(\frac{\theta}{2}\right) \quad (8)$$

$$\therefore r = \frac{2 r_1 r_2}{r_1 + r_2} \cos\left(\frac{\theta}{2}\right) \quad (9)$$

Now , D has co-ordinates as $\begin{pmatrix} r \\ \theta \end{pmatrix}$

$$\therefore \text{Co-ordinates of D are } \begin{pmatrix} \frac{2 r_1 r_2}{r_1 + r_2} \cos\left(\frac{\theta}{2}\right) \\ \frac{\theta_1 + \theta_2}{2} \end{pmatrix}$$

As required , co-ordinates of point D in terms of r, θ are found

Since ,

To find coordinates of point 'D' which intersects line joining 'A' and 'B' and angle bisector of angle AOB
 $A(r_1, \theta_1) = (12, 90)$
 $O(r', \theta') = (0, 0)$
 $B(r_2, \theta_2) = (10, 45)$
 $D(r, \theta) = (10.07865889214038, 67.5)$




Fig. 1. Triangles AOB, AOD, DOB