

Cardevia (Cardevastatin) 10 mg / 20 mg / 40 mg film-coated tablet

FICTIONAL DEMONSTRATION DOCUMENT — Qorvexa Healthcare is an invented company. This product, its molecule and every clinical statement here are fabricated for software testing. Not medical information. Do not use for any clinical purpose.

Qorvexa Healthcare · Summary of Product Characteristics · Cardiology

1. Name of the medicinal product

Cardevia 10 mg / 20 mg / 40 mg film-coated tablet.

2. Qualitative and quantitative composition

Each film-coated tablet contains Cardevastatin as the active substance. Cardevia is a HMG-CoA reductase inhibitor.

4.1 Therapeutic indications

- Primary hypercholesterolaemia in adults inadequately controlled by diet alone.
- Reduction of cardiovascular risk in adults with established atherosclerotic disease.

4.2 Posology and method of administration

Population	Dose	Notes
Adults, initial	10 mg once daily	Evening dose, with or without food
Adults, maintenance	20–40 mg once daily	Titrate at 4-week intervals
Elderly (>65 years)	10 mg once daily	Monitor renal function at 8 weeks
With ciclosporin	10 mg once daily	Do not exceed 10 mg

4.2.1 Renal impairment

eGFR 30–59 mL/min: maximum 20 mg once daily. eGFR below 30 mL/min: maximum 10 mg once daily and monitor creatine kinase every 12 weeks.

4.2.2 Hepatic impairment

Contraindicated in active hepatic disease or persistent transaminase elevation above three times the upper limit of normal.

4.3 Contraindications

- Hypersensitivity to Cardevastatin or any excipient.
- Active hepatic disease.
- Concomitant use of gemfibrozil.

4.5 Interaction with other medicinal products

Interacting substance	Effect and action required
Ciclosporin	Marked increase in Cardevastatin exposure. Restrict to 10 mg daily.
Gemfibrozil	Increased myopathy risk. Concomitant use not recommended.
Clarithromycin	Suspend Cardevia for the duration of the antibiotic course.
Metformin	No clinically significant interaction has been observed. Monitor renal function in patients over 65 years.

Interacting substance	Effect and action required
Warfarin	Monitor INR when initiating or changing dose.

4.8 Undesirable effects

- **Common (1–10%):** Headache, myalgia, nausea, elevated transaminases
- **Uncommon (0.1–1%):** Rash, dizziness, insomnia
- **Rare (<0.1%):** Rhabdomyolysis, immune-mediated necrotising myopathy

6.4 Special precautions for storage

Store below 25 °C in the original blister. Do not refrigerate.