

Hepatoval (Hepatovaline) 10 mg / 20 mg enteric-coated tablet

FICTIONAL DEMONSTRATION DOCUMENT — Qorvexa Healthcare is an invented company. This product, its molecule and every clinical statement here are fabricated for software testing. Not medical information. Do not use for any clinical purpose.

Qorvexa Healthcare · Summary of Product Characteristics · Hepatology

1. Name of the medicinal product

Hepatoval 10 mg / 20 mg enteric-coated tablet.

2. Qualitative and quantitative composition

Each enteric-coated tablet contains Hepatovaline as the active substance. Hepatoval is a hepatoprotective aminothiol.

4.1 Therapeutic indications

- Adjunctive treatment of non-alcoholic fatty liver disease in adults.
- Supportive therapy during hepatotoxic chemotherapy regimens.

4.2 Posology and method of administration

Population	Dose	Notes
Adults	20 mg twice daily	Before meals; swallow whole
Elderly	20 mg once daily	Increase only if tolerated at 6 weeks
Child-Pugh A	20 mg once daily	No further adjustment
Child-Pugh B	10 mg once daily	Monitor ammonia weekly

4.2.1 Renal impairment

No dose adjustment required down to eGFR 30 mL/min. Below 30 mL/min, safety has not been established.

4.2.2 Hepatic impairment

Child-Pugh C: not recommended. Child-Pugh B: halve the dose and monitor serum ammonia weekly for the first month.

4.3 Contraindications

- Hypersensitivity to Hepatovaline.
- Decompensated cirrhosis (Child-Pugh C).

4.5 Interaction with other medicinal products

Interacting substance	Effect and action required
Paracetamol	Additive glutathione depletion at doses above 2 g/day. Advise patients to limit paracetamol.
Rifampicin	Reduces Hepatovaline exposure by approximately 40%.
Oral contraceptives	No interaction observed.

4.6 Fertility, pregnancy and lactation

Not recommended during pregnancy. Animal studies showed no teratogenicity, but human data are insufficient. If treatment is essential, discuss with a hepatologist before continuing.

4.8 Undesirable effects

- **Common (1–10%):** Nausea, abdominal discomfort, transient pruritus
- **Uncommon (0.1–1%):** Taste disturbance, mild neutropenia

6.4 Special precautions for storage

Store below 30 °C. Protect from moisture.