

Nephrocine (Nephrocitine) 10 mg / 20 mg prolonged-release tablet

FICTIONAL DEMONSTRATION DOCUMENT — Qorvexa Healthcare is an invented company. This product, its molecule and every clinical statement here are fabricated for software testing. Not medical information. Do not use for any clinical purpose.

Qorvexa Healthcare · Summary of Product Characteristics · Nephrology

1. Name of the medicinal product

Nephrocine 10 mg / 20 mg prolonged-release tablet.

2. Qualitative and quantitative composition

Each prolonged-release tablet contains Nephrocitine as the active substance. Nephrocine is a renal tubular protectant.

4.1 Therapeutic indications

- Slowing of progression in adults with chronic kidney disease stages 2–3.

4.2 Posology and method of administration

Population	Dose	Notes
eGFR $\geq$ 60 mL/min	20 mg once daily	Morning, with food
eGFR 45–59 mL/min	10 mg once daily	Reassess at 12 weeks
eGFR 30–44 mL/min	10 mg alternate days	Monitor potassium fortnightly
eGFR < 30 mL/min	Not recommended	Insufficient data

4.2.1 Renal impairment

Dosing is defined by eGFR band; see section 4.2. Potassium must be checked before initiation and at 2, 4 and 12 weeks.

4.2.2 Hepatic impairment

No adjustment required.

4.3 Contraindications

- Serum potassium above 5.5 mmol/L at baseline.
- Acute kidney injury.

4.5 Interaction with other medicinal products

Interacting substance	Effect and action required
ACE inhibitors	Additive hyperkalaemia risk. Check potassium at 2 weeks.
NSAIDs	May blunt the renoprotective effect. Avoid chronic use.
Metformin	No pharmacokinetic interaction. Follow standard metformin renal thresholds independently.

4.8 Undesirable effects

- **Common (1–10%):** Hyperkalaemia, fatigue, mild hypotension
- **Uncommon (0.1–1%):** Metallic taste, peripheral oedema

6.4 Special precautions for storage

Store below 30 °C in the original container.